1 **Polymerase chain reaction-based assays facilitate the breeding and study of mouse**
2 **models of Klinefelter syndrome**

3 Haixia Zhang ^{1§}, Wenyan Xu ^{1§}, Yulin Zhou ^{1§}, Xiaolu Chen ¹, Jiayang Jiang ², Xiaoman
4 Zhou ¹, Zengge Wang ³, Rongqin Ke ², and Qiwei Guo ^{1*}

5 ¹United Diagnostic and Research Center for Clinical Genetics, Women and Children's
6 Hospital, School of Medicine & School of Public Health, Xiamen University. Xiamen,
7 Fujian 361102, China

8 ²School of Biomedical Sciences & School of Medicine, Huaqiao University, Quanzhou,
9 Fujian 362021, China

10 ³Beijing Children's Hospital, Capital Medical University, Beijing 100045, China

11 [§]These authors contributed equally to this work.

12 ***Correspondence to:** Qiwei Guo, United Diagnostic and Research Center for Clinical
13 Genetics, Women and Children's Hospital, School of Medicine & School of Public
14 Health, Xiamen University. Xiamen, Fujian 361102, China

15 Email: guoqiwei@xmu.edu.cn

16 Running title: PCR assays for breeding and study of KS mouse

17

18 **Abstract**

19 Klinefelter syndrome (KS) is one of the most frequent genetic abnormalities and the
20 leading genetic cause of non-obstructive azoospermia. The breeding of mouse models of
21 KS and their study are essential to advance our knowledge of the pathologic mechanism.
22 Karyotyping and fluorescence *in situ* hybridization are reliable methods for identifying
23 chromosomal contents. However, technical issues associated with these methods can
24 decrease the efficiency of breeding KS mouse models and limit studies that require
25 rapid identification of target mice. To overcome these limitations, we developed three
26 polymerase chain reaction-based assays to measure the specific genetic information,
27 including the presence or absence of *Sry*, copy number of *Amelx*, and Xist RNA
28 transcript levels. Through a combined analysis of the assay results, we can infer the
29 karyotype of target mice. We confirmed the utility of our assays with the successful
30 generation of KS mouse models. Our assays are rapid, inexpensive, high capacity, easy
31 to perform, and require small amounts of sample. Therefore, they facilitate the breeding
32 and study of KS mouse models and help advance our knowledge of the pathologic
33 mechanism underlying KS.

34 **Keywords:**

35 Klinefelter syndrome; mouse model; 40,XX^{Y*} mouse; 41,XXY mouse

36 **Introduction**

37 Klinefelter syndrome (KS), a set of symptoms that results from an extra X chromosome
38 in males, is one of the most frequent genetic abnormalities and the leading genetic cause
39 of non-obstructive azoospermia.¹ The prevalence of KS in newborn, infertile, and
40 azoospermic males is approximately 0.15%, 3–4%, and 10–12%, respectively.^{2,3} The
41 phenotypic spectrum of KS is wide, ranging from presenting only small testes and
42 infertility to the classical traits and comorbidities such as hypergonadotropic
43 hypogonadism, infertility, neurocognitive deficits, psychiatric disorders, obesity,
44 diabetes, osteoporosis, and autoimmune disorders.⁴ Early diagnosis of KS and
45 subsequent treatment and intervention, such as testosterone replacement therapy,
46 testicular sperm extraction, early speech and occupation therapy, and educational
47 assistance, have been shown to improve the long-term quality of life for patients with
48 KS and alleviate later complications.⁵⁻¹⁰ However, the data was limited and more
49 rigorous scientific investigations are needed.¹¹ In fact, the pathologic mechanism
50 between abovementioned phenotypes and the extra X chromosome is much more
51 complicated than previously envisioned and remain largely unknown, in large part
52 because in-depth investigations, such as developmental studies and experimental
53 manipulations, are almost impossible to perform in patients with KS for ethical reasons.
54^{11,12} Therefore, studies of animal models are essential to advance our knowledge of the
55 pathologic mechanism of this prevalent syndrome.¹²

56 The discovery of a mutant mouse line, the Y* mouse, has enabled us to generate KS
57 mouse models.¹³ As shown in Fig. 1, the Y* chromosome contains the X-centromere,

partial pseudoautosomal region (PAR) of the X chromosome, partial PAR of the Y chromosome, and the entire non-recombining region of the Y chromosome (NRY). In addition, a small fragment of the non-pseudoautosomal region of the X chromosome (NPX), containing eight genes including *Amelx*, is located between the X-centromere and the duplicated PAR. Through cross-breeding, 40,XX^{Y*} and 41,XXY mice could be generated in the second and fourth generations, respectively.¹⁴ These two mouse breeds have many of the characteristics of KS, including small firm testes, germ cell loss, hypergonadotropic hypogonadal endocrine changes, altered body proportions, behavioral and cognitive issues, and so on.^{12, 14} Therefore, these mouse breeds have been widely used as animal models to overcome the limitations of human studies to explore the pathological mechanisms underlying KS.^{15, 16}

One of the most important steps in the successful breeding of 40,XX^{Y*} and 41,XXY mice is the accurate identification of the karyotypes and chromosomal fragments of the targeted mouse breeds (Fig. 1). This is typically performed through cytogenetic methods, such as karyotyping or fluorescence *in situ* hybridization (FISH). To ensure that the mice are alive for subsequent breeding after testing, culture cells derived from blood samples or tissue biopsies are used.¹⁷ However, identification of target mice by karyotyping or FISH is labor-intensive, time-consuming (5–10 days), and requires a relatively high level of expertise. These drawbacks reduce their usefulness of large-scale detection, which is generally required to generate enough model mice for study use. Moreover, the pathological changes in the KS model mice, such as germ cell loss, occur prenatally and progress rapidly after birth.^{15, 18} Therefore, studies of these early

pathological changes require immediate identification of target mice prenatally or neonatally; this poses a challenge for karyotyping and FISH, which require a relatively long turnaround time.

To overcome these limitations, we developed three polymerase chain reaction (PCR)-based assays to identify the chromosomal contents of target mice and confirmed the utility of our assays for reliable breeding of KS mouse models.

Materials and Methods

Study design

As shown in Fig. 1, each karyotype has specific sex chromosome contents that could provide specific genetic information such as gene copy number and transcriptional activity. Theoretically, by analyzing this genetic information using molecular methods, instead of cytogenetic analyses such as karyotyping and FISH, we would be able to infer the karyotype of a mouse breed. Based on this concept, three genes, *Sry*, *Amelx*, and *Xist*, could be informative. *Sry* is located in the NRY, and its presence or absence determines the gender of the mouse. Since *Amelx* is located in the NPX and is also present on the Y* chromosome, the copy number of *Amelx* implies the copy number of the NPX fragment.¹⁹ *Xist* is also located on the NPX and is transcribed into a long noncoding RNA that can transcriptionally silence one of the two X chromosomes to achieve dosage equivalence between males and females. Thus, a high level of *Xist* RNA is expected when there are two X chromosomes in a single cell, whereas an absence or low level of *Xist* RNA is expected when there is only one X chromosome in a single cell.²⁰ Corresponding to the breeding protocol shown in Fig. 1, the genetic information

of the mouse breeds in each generation is shown in Fig. 2. This genetic information was determined using the PCR-based assays designed to measure the presence or absence of *Sry*, copy number of *Amelx*, and the transcript levels of Xist RNA. Notably, the combined information for two genes was sufficient to identify a target karyotype (Fig. 2).

Mice and samples

To establish the PCR assays, tail tissue samples from 16 pairs of 40,XX and 40,XY mice (C57BL/6J) were obtained from the Experimental Teaching Department, School of Medicine, Xiamen University. To generate 40,XX^{Y*} and 41,XXY mice, breeding pairs of 40,XY* (C57BL/6JeiJ) and 40,XX (C57BL/6J) mice were purchased from The Jackson Laboratory (Bar Harbor, ME, USA). The 40,XY mice (C57BL/6J) were purchased from Xiamen University Laboratory Animal Center. The breeding was performed in the standard animal facility of the Xiamen University Laboratory Animal Center.

Nucleic acid purification

Approximately 20 mg of tail tissue was used to purify DNA or RNA with the QIAamp Fast DNA Tissue Kit (Qiagen, Valencia, CA, USA) or the RNeasy[®] mini Kit (Qiagen), respectively, according to the manufacturer's protocols. The purity and concentration of the purified nucleic acid were determined by measuring the absorbance at 260 nm and 280 nm using a NanoDrop 2000 spectrophotometer (Thermo Fisher, Waltham, MA, USA).

Reaction conditions

The PCR conditions for all assays were identical: each 25 μ L reaction contained 10 mmol/L Tris-HCl (pH 8.3), 50 mmol/L KCl, 1 U of TaqHS (Takara, Dalian, China), 2 mmol/L Mg^{2+} , 0.2 mmol/L each dNTP, 0.8 $\times$ LightCycler 480 ResoLight Dye (Roche Applied Science GmbH, Mannheim, Germany), 0.2 mmol/L each forward and reverse primer, and a specific amount of DNA or cDNA template. The amplification cycling conditions were as follows: 95 $^{\circ}$ C for 3 min, followed by 40 cycles of 95 $^{\circ}$ C for 20 s, 60 $^{\circ}$ C for 20 s, and 72 $^{\circ}$ C for 20 s. Fluorescence was recorded at the end of each annealing step. The sequences of the primers used in each assay are listed in Table 1. For the measurement of *Sry*, a melting analysis was performed after the amplification, which began with denaturation at 95 $^{\circ}$ C for 1 min and renaturation at 35 $^{\circ}$ C for 3 min, followed by melting from 40 $^{\circ}$ C to 95 $^{\circ}$ C, with a ramp rate of 0.04 $^{\circ}$ C/s. Fluorescence was recorded every 0.3 $^{\circ}$ C. PCR and melting analysis were performed on a SLAN-96S thermocycler (Zeesan, Xiamen, China). Before PCR, RNA samples were reverse transcribed with the GoScript Reverse Transcription System (Promega, Beijing, China) according to the manufacturer's protocol, and 1 μ L of the resulting cDNA was used as template for subsequent PCR.

Data analysis

For the analysis of *Sry*, a melting profile was used to determine its presence or absence. To quantify the copy number of *Amelx*, relative quantification was performed by measuring the difference between the quantification cycle (Cq) values obtained for *Amelx* and *Actb* ($\Delta Cq_{Amelx - Actb}$). Similarly, to evaluate Xist RNA transcript levels, relative quantification was performed by measuring the difference between the Cq

values for the *Xist* and *Actb* transcripts ($\Delta Cq_{Xist - Actb}$). *Actb* was used as a quantitative reference because it has consistently has two copies and a stable transcriptional pattern in mouse cells. The *Cq* value is defined as the amplification cycle when the fluorescence intensity of an amplicon reaches a specific threshold.²¹ Based on a previous study, the threshold was set to 30% of the plateau fluorescence intensity as the optimal *Cq* value for differentiation of different karyotypes.²²

152 **Karyotyping and histologic analysis**

To confirm the accuracy of our assays, after breeding, the karyotypes of the target mice were examined by karyotyping using hemopoietic lineage cells derived from the bone marrow according to a previous protocol.¹⁷ For histologic analysis of the testes of KS model mice, periodic acid-Schiff staining was performed on cross-sections of the seminiferous tubules according to a previous protocol.²³

158 **Ethics statement**

Mouse breeding and experimentation were performed in accordance with international, national, and institutional guidelines for the ethical use of animals.²⁴ The study was approved by the Ethics Committee of Xiamen University Laboratory Animal Center.

162 **Results**

163 **Establishing an assay for the detection of *Sry***

To determine the optimal amount of DNA template for the *Sry* detection assay, serial dilutions of DNA isolated from 40,XY (*Sry* positive) and 40,XX (*Sry* negative) mice were evaluated. A no template control was also examined to determine the melting profiles of primer dimers. As shown in Fig. 3A, the melting profiles changed with the

amount of DNA template. To obtain only the *Sry* amplicon, at least 2 ng of DNA template was required for samples from the 40,XY mouse. However, for the 40,XX mouse samples, the use of a larger amount of DNA template (≥ 20 ng) generated non-specific amplicons that presented similar melting profiles to those of the *Sry* amplicons. Therefore, 2 ng was considered as the optimal amount of template to specifically detect the presence or absence of *Sry* and was used for subsequent experiments.

Establishing an assay for quantification of *Amelx*

We first evaluated the influence of the amount of DNA template on the effectiveness of the *Amelx* quantification assay by examining serial dilutions of DNA isolated from a 40,XX mouse as a template. As shown in Fig. 3B, amplification in both the *Amelx* and *Actb* detection reactions was robust across a wide range of template concentrations (20 pg–200 ng). However, a repressive effect was observed when a relatively large amount of DNA template (≥ 20 ng) was used. Therefore, 2 ng was chosen as the amount of template to evaluate the ranges of the $\Delta Cq_{Amelx - Actb}$ values for the 40,XX (two copies of *Amelx*) and 40,XY (one copy of *Amelx*) mouse samples. As shown in Fig. 3C, the ranges for 16 samples from 40,XX and 40,XY mice were distinct from each other. The mean $\Delta Cq_{Amelx - Actb}$ value (1.65) of the maximum of the 40,XX samples and the minimum of the 40,XY samples was used as the cut-off value for classifying mice as having one or two copies of *Amelx* in subsequent experiments.

Establishing an assay for evaluating Xist RNA transcript levels

We first evaluated the influence of different amounts (2000, 200, and 20 ng) of RNA

190 template from a 40,XX mouse (high level of Xist RNA transcripts) in a reverse
191 transcription reaction on PCR effectiveness. As shown in Fig. 3D, both 2 000 ng and
192 200 ng of RNA had a repressive effect. Therefore, 20 ng of RNA template was used for
193 reverse transcription to evaluate the range of the $\Delta Cq_{Xist - Actb}$ values for 40,XX and
194 40,XY mouse samples (no or low level of Xist RNA transcript). As shown in Fig. 3E,
195 the range of the $\Delta Cq_{Xist - Actb}$ values for the 16 40,XX and 40,XY mouse samples were
196 distinct from each other. The mean $\Delta Cq_{Xist - Actb}$ value (6.05) of the maximum for the
197 40,XX samples and the minimum for the 40,XY samples was used as the cut-off value
198 for differentiating Xist RNA transcript levels in subsequent experiments.

199 **Breeding KS mouse models using these PCR assays**

200 Using the established PCR assays, we bred 40,XX^{Y*} and 41,XXY mice from a pair of
201 40,XY* and 40,XX mice following the protocol in Fig. 1, which confirmed the
202 effectiveness of our method. Ultimately, we obtained one 40,XX^{Y*} mouse and four
203 41,XXY mice. Subsequent karyotyping and histological analysis confirmed the
204 reliability of our assays (Figs. 4 and 5).

205 **Discussion**

206 Karyotyping and FISH are reliable methods that have been used for decades to identify
207 chromosomal abnormalities, which are essential for the breeding and study of KS
208 mouse models. However, these methods have some technical issues, e.g., a relatively
209 low capacity and long turnaround time, that decrease the breeding efficiency and limit
210 studies which require rapid identification of chromosomal contents.

211 PCR is a basic tool of the molecular laboratory that has demonstrated capability for

detecting various types of genetic abnormalities through different data analysis strategies. For example, we can detect the presence or absence of a specific gene region (e.g., *AZF* on the Y chromosome) by analyzing amplification profiles,²⁵ and we can detect numerical changes (e.g., trisomy 21, 47,XXY, or a *SMA* carrier) by analyzing melting profiles or ΔCq values.^{5, 22, 26-28} Based on previous studies, we developed three PCR-based assays to detect genetic information related to KS, including the presence or absence of *Sry*, changes in the copy number of *Amelx*, and Xist RNA transcript levels in mouse tissue samples. Using these assays, we were able to identify target mice for the breeding of KS models without performing karyotyping or FISH (Figs. 1 and 2). Compared to karyotyping or FISH, our method is rapid, as it takes less than 1 h for nucleic acid purification, less than 2 h for reverse transcription, and less than 2 h for PCR and melting analysis. A short turnaround time for mouse identification is essential for studies focused on rapid physical and pathologic changes in fetal or newborn mice, e.g., testicular degeneration.^{15, 18} Our method is also inexpensive (<0.5 US dollar per PCR assay), high capacity (up to 96 samples per assay), and easy to perform. Moreover, only 2 ng of DNA and 20 ng of RNA was sufficient to obtain reliable results (Fig. 3), suggesting that less than 0.2 mg of tissue or 1 μ l of peripheral blood would be a sufficient sample, further revealing its utility, especially for the detection of fetal or newborn mice.

An ideal assay for *Sry* identification should only yield a product from male mouse samples with *Sry* but not from female mouse samples without *Sry*, so that we can identify male mouse samples directly by their amplification profiles. In practice,

however, non-specific amplicons derived from paralogous sequences and/or primer dimers were difficult to eliminate from reactions with female mouse samples. Because of this, melting profiles are more specific than the amplification profiles since the melting profiles of the *Sry* amplicons could be readily differentiated from those of non-specific amplicons (Fig. 3A). The amount of DNA template used plays an important role in the melting profiles, and 2 ng was confirmed to be an optimal amount of template in our study (Fig. 3A). However, if different PCR reagents are used, the optimal amount of DNA template should be re-evaluated.

Theoretically, after equalizing the DNA template input for the *Amelx* assay, samples with different *Amelx* copy numbers should generate different ranges of Cq values. Therefore, we could infer the *Amelx* copy number in a mouse sample by its Cq value. In practice, we confirmed this assumption (data not shown). However, to further diminish the potential impacts of slight differences (e.g., differences in the quantity or quality of the DNA template) between samples, *Actb* was used as a reference gene to calibrate the *Amelx* quantification. Accordingly, instead of ranges of Cq values, ranges of $\Delta Cq_{Amelx - Actb}$ values were established to quantify *Amelx* in unknown samples (Fig. 3C). Moreover, rather than using confidence intervals for the different ranges of $\Delta Cq_{Amelx - Actb}$ values, we used a single cut-off value to determine the *Amelx* copy number (Fig. 3C), which simplified the data analysis.

The underlying principle of the assay used to determine Xist RNA transcript levels was identical to that of *Amelx* quantification. Similarly, a large amount of nucleic acid template had a repressive effect on PCR amplification (Fig. 3D). We further evaluated

whether the repressive effect was derived from the large amount of RNA or the resulting cDNA. The results suggested that the repressive effect was greater for large amounts of RNA (data not shown). Therefore, a smaller amount of RNA was preferable for reverse transcription, which was sufficient to reliably determine Xist RNA transcript levels (Fig. 3E).

In conclusion, we developed three PCR-based assays for facilitating the breeding and study of KS mouse models. Our method is rapid, inexpensive, high capacity, easy to perform, and requires small amounts of sample. We confirmed the utility of our assays for the successful generation of KS mouse models. We believe our method will advance the knowledge of the pathologic mechanism of KS.

Authors' Contributions

HXZ, WYX and YLZ participated in study design and the establishment of PCR assays. XLC, JYJ, XMZ, ZGW, and RQK participated in breeding of KS mouse models. QWG conceived of the study, and participated in its design and coordination and draft the manuscript. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Acknowledgements

We thank the Experimental Teaching Department, School of Medicine, Xiamen University and the Xiamen University Laboratory Animal Center for their kind supports to this study.

Conflict of Interest

None of the authors declare competing financial interests.

278 **References**

- 279 1. Klinefelter HF, Reifenstein, E.C., Albright, F. Syndrome characterized by gynecomastia
280 aspermatogenesis without A-Leydigism and increased excretion of follicle stimulating hormone. *J Clin*
281 *Endocrinol Metab* 1942; **2**: 615-27.
- 282 2. Berglund A, Stochholm K, Gravholt CH. The epidemiology of sex chromosome abnormalities. *Am J*
283 *Med Genet C Semin Med Genet* 2020; **184**: 202-15.
- 284 3. Forti G, Corona G, Vignozzi L, Krausz C, Maggi M. Klinefelter's syndrome: a clinical and
285 therapeutical update. *Sex Dev* 2010; **4**: 249-58.
- 286 4. Gravholt CH, Chang S, Wallentin M, Fedder J, Moore P, et al. Klinefelter Syndrome: Integrating
287 Genetics, Neuropsychology, and Endocrinology. *Endocr Rev* 2018; **39**: 389-423.
- 288 5. Fu DM, Zhou YL, Zhao J, Hu P, Xu ZF, et al. Rapid screening for Klinefelter syndrome with a
289 simple high-resolution melting assay: a multicenter study. *Asian J Androl* 2018; **20**: 349-54.
- 290 6. Pizzocaro A, Vena W, Condorelli R, Radicioni A, Rastrelli G, et al. Testosterone treatment in male
291 patients with Klinefelter syndrome: a systematic review and meta-analysis. *J Endocrinol Invest* 2020; **43**:
292 1675-87.
- 293 7. Graham JM, Jr., Bashir AS, Stark RE, Silbert A, Walzer S. Oral and written language abilities of
294 XXY boys: implications for anticipatory guidance. *Pediatrics* 1988; **81**: 795-806.
- 295 8. Gies I, Unuane D, Velkeniers B, De Schepper J. Management of Klinefelter syndrome during
296 transition. *Eur J Endocrinol* 2014; **171**: R67-77.
- 297 9. Skakkebaek A, Wallentin M, Gravholt CH. Neuropsychology and socioeconomic aspects of
298 Klinefelter syndrome: new developments. *Curr Opin Endocrinol Diabetes Obes* 2015; **22**: 209-16.
- 299 10. Tournaye H, Krausz C, Oates RD. Concepts in diagnosis and therapy for male reproductive

-
- 300 impairment. *Lancet Diabetes Endocrinol* 2017; **5**: 554-64.
- 301 11. Gravholt CH, Tartaglia N, Distech C. Sex chromosome aneuploidies in 2020-The state of care and
302 research in the world. *Am J Med Genet C Semin Med Genet* 2020; **184**: 197-201.
- 303 12. Wistuba J, Beumer C, Brehm R, Gromoll J. 41,XX(Y) * male mice: An animal model for Klinefelter
304 syndrome. *Am J Med Genet C Semin Med Genet* 2020; **184**: 267-78.
- 305 13. Eicher EM, Hale DW, Hunt PA, Lee BK, Tucker PK, et al. The mouse Y* chromosome involves a
306 complex rearrangement, including interstitial positioning of the pseudoautosomal region. *Cytogenet Cell*
307 *Genet* 1991; **57**: 221-30.
- 308 14. Cox KH, Bonthuis PJ, Rissman EF. Mouse model systems to study sex chromosome genes and
309 behavior: relevance to humans. *Front Neuroendocrinol* 2014; **35**: 405-19.
- 310 15. Lue Y, Jentsch JD, Wang C, Rao PN, Hikim AP, et al. XXY mice exhibit gonadal and behavioral
311 phenotypes similar to Klinefelter syndrome. *Endocrinology* 2005; **146**: 4148-54.
- 312 16. Wistuba J, Luetjens CM, Stukenborg JB, Poplinski A, Werler S, et al. Male 41, XXY* mice as a
313 model for klinefelter syndrome: hyperactivation of leydig cells. *Endocrinology* 2010; **151**: 2898-910.
- 314 17. Lee JJ, Warburton D, Robertson EJ. Cytogenetic methods for the mouse: preparation of
315 chromosomes, karyotyping, and in situ hybridization. *Anal Biochem* 1990; **189**: 1-17.
- 316 18. Hunt PA, Worthman C, Levinson H, Stallings J, LeMaire R, et al. Germ cell loss in the XXY male
317 mouse: altered X-chromosome dosage affects prenatal development. *Mol Reprod Dev* 1998; **49**: 101-11.
- 318 19. Vernet N, Mahadevaiah SK, Yamauchi Y, Decarpentrie F, Mitchell MJ, et al. Mouse Y-linked Zfy1
319 and Zfy2 are expressed during the male-specific interphase between meiosis I and meiosis II and promote
320 the 2nd meiotic division. *PLoS Genet* 2014; **10**: e1004444.
- 321 20. Wolstenholme JT, Rissman EF, Bekiranov S. Sexual differentiation in the developing mouse brain:

-
- 322 contributions of sex chromosome genes. *Genes Brain Behav* 2013; **12**: 166-80.
- 323 21. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, et al. The MIQE guidelines: minimum
- 324 information for publication of quantitative real-time PCR experiments. *Clin Chem* 2009; **55**: 611-22.
- 325 22. Guo Q, Zhou Y, Wang X, Li Q. Simultaneous detection of trisomies 13, 18, and 21 with multiplex
- 326 ligation-dependent probe amplification-based real-time PCR. *Clin Chem* 2010; **56**: 1451-9.
- 327 23. Keeney S. Meiosis. Volume 2, cytological methods. Preface. *Methods Mol Biol* 2009; **558**: v-vi.
- 328 24. Committee for the Update of the Guide for the Care and Use of Laboratory Animals. Guide for the
- 329 Care and Use of Laboratory Animals, 8th ed. *Washington (DC): National Academies Press (US)* 2011.
- 330 25. Guo Q, Lan F, Xu L, Jiang Y, Xiao L, et al. Quadruplex real-time polymerase chain reaction assay
- 331 for molecular diagnosis of Y-chromosomal microdeletions. *Fertil Steril* 2012; **97**: 864-9.
- 332 26. Guo Q, Xiao L, Zhou Y. Rapid diagnosis of aneuploidy by high-resolution melting analysis of
- 333 segmental duplications. *Clin Chem* 2012; **58**: 1019-25.
- 334 27. Xia Z, Zhou Y, Fu D, Wang Z, Ge Y, et al. Carrier screening for spinal muscular atrophy with a
- 335 simple test based on melting analysis. *J Hum Genet* 2019; **64**: 387-96.
- 336 28. Zhou Y, Xu W, Jiang Y, Xia Z, Zhang H, et al. Clinical Utility of a High-Resolution Melting Test for
- 337 Screening Numerical Chromosomal Abnormalities in Recurrent Pregnancy Loss. *J Mol Diagn* 2020; **22**:
- 338 523-31.

339

340

Table 1. Primer information of PCR-based assays.

Target gene	Primer sequence	Amplification region (assembly: GRCm39/mm39)
<i>Sry</i>	F: GCAGCTTACCTACTTACTAACA R: CTGAGGTGCTCCTGGTA	chrY:2662499-2662564
<i>Amelx</i>	F: CCTTCAGCCTCATCACCA R: TTGGGTTGGAGTCATGGA	chrX:167965017-167965122
<i>Xist</i>	F: CATCCTACCATCATCTGCTTT R: GGGAAGATGACTCCAGTCT	chrX:102504207-102504284
<i>Actb</i>	F: CCATGAAACTACATTCAATTCCAT R: TGTGTTGGCATAGAGGTCTT	chr5:142889915-142889984

344 **Figure 1. Breeding of Klinefelter syndrome (KS) mouse models.** The mouse in the
 345 blue circle is a 40,XX^{Y*} mouse, i.e., one of the KS mouse models. The mice in the red
 346 circles are the target mice in each generation for breeding 41,XXY mice, i.e., another
 347 KS mouse model. This figure was modified from figures in a previous study (Front
 348 Neuroendocrinol 2014; 35: 405-19). NPX: non-pseudoautosomal region of the X
 349 chromosome; NRY: non-recombining region of the Y chromosome; PAR:
 350 pseudoautosomal region.

351

352 **Figure 2. Genetic information of the mouse breeds obtained during the breeding of**
353 **Klinefelter syndrome mouse models.** +: presence of *Sry* or a high Xist RNA transcript
354 level; -: absence of *Sry* or low/no Xist RNA transcript. Pink labels indicate the target
355 mice in each generation for the breeding of 41,XXY mice. Green labels indicate the
356 assays required to identify the target mice in each generation. For example, in the
357 second generation, each mouse was first tested for *Sry*, and those negative for *Sry* were
358 further tested for Xist RNA transcript levels. Mice with low/no Xist RNA transcript, i.e.,
359 negative results in both assays, were identified as 40,XY*^X mice.

360

361 **Figure 3. PCR-based assays for the breeding of Klinefelter syndrome mouse**

362 **models.** (a) Melting profiles for detecting *Sry* using different amounts of DNA template.

363 (b) Assays to quantify *Amelx* using different amounts of DNA template. (c) Establishing

364 a cut-off value for *Amelx* quantification. (d) Assays for evaluating Xist RNA transcript

365 levels using different amounts of RNA template. (e) Establishing a cut-off value for Xist

366 RNA quantification. Each sample was tested in triplicate, and the Cq value of each

367 sample is the mean Cq value of three replicates. NTC: no template control.

368

369 **Figure 4. Karyotypes of the target mice for breeding Klinefelter syndrome mouse**

370 **models.** (a) A 40,XX^{Y*} mouse from the second generation. The arrow indicates the X^{Y*}
 371 chromosome. (b) A 40,XY^{X*} mouse from the second generation. The arrow indicates the
 372 Y^{X*} chromosome. (c) A 41,XXY^{X*} mouse from the third generation. The arrow
 373 indicates the Yx* chromosome. (d) A 41,XXY mouse from the fourth generation.

374

375 **Figure 5. Testicular phenotypes of Klinefelter syndrome mouse models. (a-c)**

376 Respective appearance, anatomic analysis, and histologic analysis of the testes of a
 377 40,XX^{Y*} mouse and 40,X^{Y*} littermate. (d-f) Respective appearance, anatomic analysis,
 378 and histologic analysis of the testes of a 41,XXY mouse and 40,XY littermate. The
 379 arrows indicate the position of the testes. As expected, the 40,XX^{Y*} and 41,XXY mice
 380 presented small, firm testes. Moreover, the tubule diameters of 40,X^{Y*} and 41,XXY
 381 mice were smaller than those of their reference littermates, germ cells were absent, and
 382 only Sertoli cells were observed within the tubules.

First crossbreeding

Second crossbreeding

Third crossbreeding

Breed	F2-1	F2-2	F2-3	F2-4
Karyotype	40 , XX	40 , XY*	40 , XX ^{Y*}	40 , XY ^{*X}
<i>Sry</i>	-	+	+	-
<i>Amelx</i>	2	2	2	2
Xist RNA	+	-	+	-

Breed	F3-1	F3-2	F3-3	F3-4	F3-5
Karyotype	40 , XX	40 , XY	40 , XY ^{*X}	41 , XXY ^{*X}	41 , XYY ^{*X}
<i>Sry</i>	-	+	-	-	+
<i>Amelx</i>	2	1	2	3	2
Xist RNA	+	-	-	+	-

Breed	F4-1	F4-2	F4-3	F4-4	F4-5	F4-6
Karyotype	40 , XX	40 , XY	40 , XY ^{*X}	41 , XXY ^{*X}	41 , XYY ^{*X}	41 , XXY
<i>Sry</i>	-	+	-	-	+	+
<i>Amelx</i>	2	1	2	3	2	2
Xist RNA	+	-	-	+	-	+

a

b

c

d

a

b

c

d

e

f