1 When to depart from a stopover site? Time since arrival matters more than current weather
2 conditions

3 Sébastien Roques^{1,2}, Pierre-Yves Henry^{3,4}, Gaétan Guyot⁵, Bruno Bargain⁵, Emmanuelle
4 Cam^{2,6}, Roger Pradel¹

5 ¹CEFE, CNRS, Université de Montpellier, Université Paul-Valéry Montpellier - EPHE,
6 Montpellier Cedex 5 France

7 ² Laboratoire Evolution et Diversité Biologique, UMR CNRS 5171, Bât 4R1, 118 route de
8 Narbonne, Université Toulouse 3, Paul Sabatier, 31062 Toulouse Cedex 09, France

9 ³ Mécanismes adaptatifs et Evolution (MECADEV UMR 7179), Muséum National d'Histoire
10 Naturelle, Centre National de la Recherche Scientifique, 1 avenue du Petit Château, F-91800
11 Brunoy, France

12 ⁴ Centre de Recherches sur la Biologie des Populations d'Oiseaux (CRBPO), Centre
13 d'Ecologie et des Sciences de la Conservation (CESCO UMR 7204), Muséum National
14 d'Histoire Naturelle, Centre National de la Recherche Scientifique, Sorbonne Université,
15 Bâtiment 135, CP135, 43 rue Buffon, F-75005 Paris, France

16 ⁵ Bretagne Vivante, 19 rue de Gouesnou - BP 62132, 29221 BREST CEDEX 2, France

17 ⁶ Univ Brest, CNRS, IRD, Ifremer, LEMAR, F-29280 Plouzané, France

18 Correspondence author: Sébastien Roques

19 E-mail: se.roques@gmail.com

20

21

22   **Abstract**

23   On the journey to wintering sites, migratory birds usually alternate between flights and
 24   stopovers where they rest and refuel. Migration strategies are assumed to differ according to
 25   season: a time-minimization pre-breeding migration strategy towards breeding locations, and
 26   an energy-minimization post-breeding migration strategy to wintering ones. The duration of
 27   flights and stopovers determines the energy requirements and the total duration of the journey.
 28   Since migrating birds actually spend most of the time at stopovers, selection to minimize the
 29   amount of energy or time spent on migration is very likely to operate on the effectiveness of
 30   stopover rest and refueling. Here we address the relative contribution of factors to departure
 31   decisions from stopover sites during the post-breeding migration in a long-distance migratory
 32   songbird. When capture probability is low it is impossible to assess the variation in body
 33   condition over the entire duration of the stopover. To get around this, we use Time Since
 34   Arrival (TSA) as a proxy for the changes in the state of individuals during the stopover. We
 35   propose that TSA is an integrative proxy for resting, feeding and fattening efficiency. We
 36   develop a capture-recapture model to address the relationship between departure probability,
 37   estimated TSA, and weather conditions. Using a 20-year dataset from sedge warblers, we
 38   show that TSA has a larger effect on departure probability than weather conditions. Low
 39   humidity and an increase in atmospheric pressure in the days preceding departure are
 40   associated with higher departure probability, but these effects are smaller than that of TSA.

41

42   **Keywords:** Stopover, Time Since Arrival, Capture-Recapture, Sedge warbler, Bayesian
 43   inference

44

45 **Introduction**

46 Each year migratory birds species commute between breeding and wintering areas (Alerstam
47 1990, Somveille et al. 2015). Most species cannot go from breeding to wintering grounds in a
48 single flight of thousands of kilometers. They must stop to rest, feed and refill energy stores
49 regularly (Alerstam 1990, Åkesson and Hedenström 2007, Schmaljohann and Eikenaar 2017,
50 Schmaljohann 2018). The duration of migratory flights and stopovers determines the energy
51 requirements and also the total duration of the journey. To maximize survival, individuals
52 must optimize their journey to match with seasonal variation in food availability (and their
53 energy requirements) at stopover sites and to arrive at wintering areas when resource
54 availability is sufficient (Alerstam 1990, Somveille et al. 2015, 2018, 2019; Zúñiga et al.
55 2017, Schmaljohann 2018). Birds spend much more time at stopover sites than in migratory
56 flight, with almost 85% of the journey spent on stopover (Hedenström and Alerstam 1997,
57 Green et al. 2002, Schmaljohann et al. 2012, Schmaljohann 2018). Consequently, selection to
58 minimize the amount of energy spent (energy minimization) or the total time spent on
59 migration (time minimization) likely operates mainly on the effectiveness of stopover rest and
60 refueling (Hedenström and Alerstam 1997, Schmaljohann 2018).

61 Deciding to leave the stopover site normally means a new flight of hundreds of kilometers.
62 This movement itself, and its termination, are highly constrained by the maximal flight
63 capacity (given the size of energy reserves and body size) and atmospheric conditions
64 (Alerstam, 1990; Jenni & Schaub, 2003; Schmaljohann & Eikenaar, 2017; Wikelski et al.,
65 2003). Once settled at a stopover site, the probability of departing (and the duration of the
66 stay) depends on factors associated with the initiation of movement (i.e. to depart from the
67 site) on fuel store, resting state, food availability, weather conditions and migratory
68 experience (Jenni and Schaub 2003, Schmaljohann and Eikenaar 2017). Recent studies have

69 tried to disentangle the environmental factors driving departure decision (Schaub et al. 2008,
70 Ktitorov et al. 2010, Arizaga et al. 2011, Deppe et al. 2015, Dossman et al. 2016,
71 Schmaljohann and Eikenaar 2017). Weather conditions are usually considered through the
72 constraint they impose on flight. Departure decision depends on wind speed and direction
73 (tailwind assistance reduces the cost of flight; Tsvey et al. 2007, Arizaga et al. 2011, Ma et al.
74 2011, Dossman et al. 2016). This decision also depends on rainfall or humidity which usually
75 force birds to stay at the stopover site (Tsvey et al. 2007, Arizaga et al. 2011, Deppe et al.
76 2015, Dossman et al. 2016). Last, cloud cover influences departure decision by decreasing
77 visibility and the ability of birds to navigate (Zehnder et al. 2001, Åkesson and Hedenström
78 2007).

79 But the internal state of the individual (resting state, fuel store, migratory experience) also
80 largely influences departure decision: birds need to rest and reach a sufficient level of fuel
81 store to perform the next migratory flight (Alerstam 1990, Hedenström and Alerstam 1997,
82 Schaub et al. 2008, Goymann et al. 2010, Schmaljohann et al. 2012, Dossman et al. 2016,
83 Schmaljohann and Eikenaar 2017, Moore et al. 2017, Anderson et al. 2019). Apart from
84 resting to recover from extreme physical exercise and sleeping to recover from sleep
85 deprivation during migratory flight (Schwilch et al. 2002), birds at stopover allocate most of
86 their time and energy to foraging (Hedenström and Alerstam 1997, Cohen et al. 2014, Smith
87 and McWilliams 2014). They are assumed to refill their energy stores as fast as possible, and
88 to continue their migratory journey if their energy stores and weather conditions are favorable
89 (Schmaljohann 2018). In addition, the ability to refuel and rest may depend on age or
90 experience, which can lead individuals migrating for the first time (juveniles) and those that
91 already traveled to wintering areas in the past (adults) to depart from stopover sites after stays
92 of different durations or under different weather conditions (Deppe et al. 2015, Dossman et al.
93 2016, Schmaljohann et al. 2018).

94 To our knowledge, the relative importance of individual internal state (resting state, fuel
95 store) and weather conditions at determining departure probability has never been assessed
96 within a single modelling framework. The different results of previous studies concerning the
97 relationship between fuel store and departure probability could be due to different local
98 constraints or to the migration strategy (time or energy-minimizing; Hedenström and
99 Ålerstam 1997; Schmaljohann 2018; Anderson et al. 2019). But more importantly, it could be
100 due to the fact that fuel store can only be measured when individuals are captured (i.e., not
101 necessarily at the very beginning and end of their stay). The lack of an effect of fuel store on
102 departure probability could also arise because the method to jointly incorporate internal
103 individual state and environmental covariates in the same capture-recapture (CR) framework
104 was not available (Jenni and Schaub 2003, Schmaljohann and Eikenaar 2017). Here we define
105 and estimate a variable that cannot be measured without using electronic devices: the time the
106 individual has spent at the stopover site since its arrival: Time Since Arrival (Pledger et al.
107 2008). Our goal is to assess the influence of TSA on departure probability using a long-term
108 dataset created when electronic devices were not available, and to include a large number of
109 individuals and years in the analysis. Despite limitations of retrospective analyses of long-
110 term datasets (modern electronic techniques were not available), such datasets offer
111 interesting opportunities: (i) large sample sizes to draw inference about departure probability
112 using statistical approaches, and (ii) long time series of observations to characterize long-term
113 patterns and yearly deviations from deep-rooted trends. Birds need to rest after a long distant
114 migratory flight and fattening increases with the duration of the stay at stopover (Ålerstam
115 1990, Schwilch and Jenni 2001, Jenni and Schaub 2003, Schaub et al. 2008, Schmaljohann
116 and Eikenaar 2017). We propose that Time Since Arrival (TSA) is an integrative proxy for the
117 overall change in the internal state of stopovering birds. If this assumption is true, then
118 departure probability should decrease as the TSA increases, and TSA should be a major

determinant of departure probability. Importantly, TSA must be estimated using an analytical technique accounting for the daily capture probability of marked individuals (i.e., the probability of capturing an individual that is alive and present in the site). This is a methodological challenge because investigators know neither the day when a bird arrives nor the day when it departs from the site. Many capture-recapture stopover studies have relied on the assumption that birds are captured for the first and last time on the exact days of arrival and departure from the site (Schmaljohann and Eikenaar 2017). This assumption is unrealistic: the probability that a bird is captured in a given day usually lies between 0.1 and 0.2 (Schaub and Jenni 1999, Schaub et al. 2001, Moore et al. 2017), and the resulting “minimal stopover duration” (duration between first and last captures) strongly underestimates the actual stopover duration (by a factor that may be as large as 3; Schaub, Pradel, Jenni, & Lebreton, 2001). Estimating TSA requires the development of a statistical model accounting of imperfect detection probability.

Here we evaluate the contribution of estimated TSA and weather conditions to departure probability. We develop a capture-recapture model and analyze a long-term CR dataset from a long-distance migrant songbird at a stopover site. Since refueling seems to be the primary biological function of stopover (Schmaljohann and Eikenaar 2017, Schmaljohann 2018, Klinner et al. 2020), we hypothesize that TSA is be the main driver of departure probability in this long-distant migrant and that wind, humidity, cloud cover and atmospheric pressure have a secondary, but significant effect on the departure probability. Estimated TSA should be closer to the genuine duration of the individual’s stay than the time elapsed between the first and the last physical captures of the individual. If TSA integrates the overall changes in the individual internal state, this should help us detect the effect of stay duration on departure probability, if any. Moreover, we hypothesize that stopover duration differs according to

migratory experience, as documented in other studies (Deppe et al. 2015) and test for differences between juveniles and adults.

Material and methods

Study area, sampling and dataset

The study site is the Trunvel ringing station (Tréogat, Brittany, France; 47.8964859°N, 4.3698274°W). Data from marked individuals have been collected using a standardized mist-netting protocol (including tape-luring; B. Bargain, C. Vansteenwegen, & J. Henry, 2002). Each captured bird was identified, ringed and aged. We used data collected from 1990 to 2014, from the 1st to the 30th of August because 90% of the captures occur in this month. The years when the number of recaptures was too low were not used in analyses (less than 10 birds recaptured only once). The Sedge Warbler (*Acrocephalus schoenobaenus*) is the most abundant species that stopovers at this site during its journey to winter quarters in sub-Saharan Africa. This 12-g songbird strictly depends on reedbeds where it essentially forages on one aphid species (*Hyalopterus prunii*) to refill energy stores (Bibby and Green 1981).

Between 1990 and 2014, 79,700 individuals have been marked (Dehorter and CRBPO 2015). Among all migrant songbirds that use this site, only a small fraction stays several days to rest and refuel (i.e. actual stopover; Warnock, 2010). The majority either continues migration by the following night, or moves to another stopover place (i.e., transients; Bächler & Schaub, 2007; Schaub et al., 2008). As we aim to study the departure probability of birds that stayed over at the site, we analyzed only capture-recapture data from birds that were caught at least twice during a season (including recaptures during the same day). Hence, the estimated stopover duration applies only to the part of the population passing by the site and that stays for at least some hours or days. The sample we used included data from 683 adults and 4927

juveniles; their latest recapture occurred at the site on average 3.4 ± 3.6 (SD) days after their first capture. The mean mass gain between first and last capture on individuals was $0.48\text{g} \pm 1.48$ (SD).

Weather conditions

Weather variables expected to influence daily departure probability (between day $t-1$ and t) were: (i) wind (on day $t-1$), (ii) relative humidity (on day $t-1$), (iii) cloud cover during the night [i.e. between day $t-1$ and t ; scale from 0 (no cloud) to 8 (complete sky cover)] (iv) atmospheric pressure; as birds likely perceive changes in pressure rather than pressure itself, we used the change in atmospheric pressure between day t and day $t-1$ as a covariate (denoted as $\Delta Pressure$ in hPa). Depending on its direction, wind can either facilitate flight (tailwind) or increase the cost of flight (headwind). To integrate both wind effects, we computed the wind covariate as in (Arizaga et al. 2011):

$$wind = V \cos[\alpha_T - (180^\circ + \alpha_W)],$$

where V is wind speed (in $m \cdot s^{-1}$), α_T is the assumed departure direction (120° according to recovery data; note that birds don't cross the Bay of Biscay, Dehorter and CRBPO 2015, so 120° is almost the coast direction), and α_W is the direction the wind (Åkesson et al. 2002). Since birds depart on migration at the end of the day (Müller et al. 2018), we used wind speed and direction observed during a period of time starting 2 hours before sunset on day $t-1$ and ending in the middle of the night on day t . All weather covariates were scaled prior to the analyses. The weather data were provided by the Penmarch meteorological station ($47.797537, -4.374768$).

Modelling & statistical analyses

We used a formulation of the Jolly-Seber (JS) model (Jolly 1965, Seber 1965) parameterized with entry probability in the sampling area (Crosbie and Manly 1985, Schwarz and Arnason 1996). This allows modeling the arrival of birds at the stopover site and estimating stopover duration (Lyons et al. 2016, Lok et al. 2019, Roques et al. 2020). The parameters of the model are:

ϕ_t Probability of staying in the sampling area from day t to $t+1$,

η_t Probability of arriving at the stopover area on day t given that the individual was not present in the site before,

p_t Probability of capturing the individual on day t given that the individual has arrived and has not yet left the site.

We used the Bayesian, state-space formulation of the JS model (Gimenez et al. 2007, Royle 2008). This model contains a submodel for the state process: true, partially unobservable states are “not yet arrived”, “present in the study area”, and “departed”. The model also includes a submodel for the observations (conditional on true state) directly encoded in the individual capture histories. For each individual capture history h_i , the true state history is accounted for by the vector z_i . This vector of binary state variables describes if an individual i is present in the stopover area on day t , $z_{i,t} = 1$, or not, $z_{i,t} = 0$.

The state process is defined as:

$$z_{i,t} | z_{i,t-1} \sim \text{Bernoulli} \left(\phi_{t-1} z_{i,t-1} + \eta_t \prod_{j=1}^{t-1} (1 - z_{i,j}) \right)$$

The term $\prod_{j=1}^{t-1}(1 - z_{i,j})$ accounts for the availability of the individual to enter the stopover area and is equal to 1 when the individual has not yet entered the stopover area, and 0 when it has already entered.

As the binary observations are conditionally independent Bernoulli random variables, the link between the state and observation processes is given by the following equation:

$$h_{i,t} | z_{i,t} \sim \text{Bernoulli}(z_{i,t}, p_t)$$

This means that if individual i has not yet entered the stopover area or has left ($z_{i,t} = 0$), then $h_{i,t} = 0$ with probability equal to 1. If $z_{i,t} = 1$, then the capture history $h_{i,t}$ is a Bernoulli trial with probability p_t , which is the probability of capturing the individual on day t . This formulation allows us to estimate TSA for each individual. The TSA covariate is a partially- or non-observable variable computed using the sum of true states $z_{i,t}$. TSA accounts for the time individual i has already spent in the stopover area on day t :

$$TSA_{i,t} = \sum_{s=1}^{t-1} z_{i,s}$$

The state vector z_i also allows us to use a new formulation of the stopover duration described in Lyons et al. (2016). We computed the mean stopover duration (in days) as follows:

$$SOD = \frac{\sum_i \sum_t z_{i,t}}{n}$$

where n is the number of individuals and z the true state variable (whether individual i was present or not at the stopover site on day t).

To account for heterogeneity in detection probability among capture occasions and limit the number of parameters to estimate, we modeled detection probability as a random effect. Hence, we modeled p as:

$$\text{logit}(p) \sim \text{Norm}(\text{logit}(\mu_p), \sigma_p),$$

where μ_p is the mean recapture probability and σ_p the standard deviation of the random effect.

We expressed the probability of remaining at the site as a function of the previously defined weather covariates and TSA. We considered effects as ‘statistically significant’ when the estimated slope corresponding to these covariates had a 95% credible interval excluding 0 (Kéry and Schaub 2011). We analyzed the 20 years of data simultaneously, but accounted for potential differences among years (Péron et al. 2007) by means of a random year effect, where y is the number of the year:

$$\text{Year}_y \sim \text{Norm}(\mu_{\text{year}}, \sigma_{\text{year}})$$

To account for the effect of the experience on an individual’s departure probability from the stopover site, we used age-dependent random effects with 2 age classes (Adult and Juvenile), where a is the age class:

$$\text{Age}_a \sim \text{Norm}(\mu_{\text{Age}}, \sigma_{\text{Age}})$$

Using a logit link, the probability of staying at the stopover area between $t - 1$ and t was formulated as:

$$\begin{aligned} \text{Logit}(\phi_{i,t-1}) &= \alpha + \beta_1 * TSA_{i,t-1} + \beta_2 * wind_{t-1} + \beta_3 * temperature_{t-1} + \beta_4 \\ &* cloudcover_{t-1} + \beta_5 * humidity_{t-1} + \beta_6 * \Delta Pressure + Year_y \\ &+ Age_a \end{aligned}$$

Here we included all the covariates in the model despite a non-null collinearity between most of the environmental covariates (see online Supplementary material for details). However, this collinearity appeared sufficiently small to allow the inclusion of all effects in a same model (Hair et al. 2006). Also, because TSA is computed at each occasion for each individual, TSA cannot be standardized prior to analyses. To compare the effect of TSA on departure probability to that of weather covariates, we calculated the effect of a standardized TSA by multiplying the estimated values of the TSA slopes (β_1) by the standard deviation of all estimated TSA values.

We performed analyses with JAGS (with the package R2jags, Hornik et al. 2003; Su et al. 2015) using R version 3.6.1 (R Development Core Team 3.0.1. 2013). We used 60 000 iterations with a burnin of 30 000, and we checked chain mixing and convergence (Kéry and Schaub 2011). The JAGS code is available in the online Supplementary material.

Results

The mean estimated stopover duration for the whole study period is 12.5 ± 2.2 days CI [12.2; 12.8], with unstructured variation among years (Figure 1). Adults (experienced birds) stay on average 1.6 days more than juveniles (naive birds) (13.8 ± 2.2 for adults, and 12.2 ± 2.1 days for juveniles; this age difference is robust through years; Figure 1). However, the 95% credible intervals and standard deviation of estimated stopover duration for juveniles and adults overlap (Figure 1) and therefore age accounts for a very limited part of the variation in

stopover duration between individuals. This suggests that factors other than migratory experience and covariates accounted for in this study also determine departure probability.

Departure probability from the stopover site between two days ($1-\phi_t$, i.e. the complement of the probability of staying at the stopover site) is positively related to TSA (Figure legends and Figure 2). In other words, the longer a bird has already stayed at the site, the higher its probability of resuming migration flight by the following night. TSA is the most important predictor of departure probability compared to other variables, based on effect sizes (Table 1). TSA effect is also the effect that is estimated with the largest precision, which suggests a small variability of the TSA effect among individuals (see CI in Table 1).

Both humidity and Δ Pressure have an effect on the departure probability from the stopover site (Table 1). Departure probability increases with drier conditions and large changes in pressure (Δ pressure). We did not find evidence of an effect of wind on departure probability (Table 1). We found slight evidence of a negative effect of cloud cover on the departure probability from the stopover site (Table 1), but the robustness of this result is weak because one boundary of the CI overlaps 0. Estimates of TSA and humidity effects on departure probability (Figure 2A) or of Δ pressure (Figure 2B) show that departure probability primarily depends on TSA, and that weather covariates have a smaller influence on this probability.

Discussion

Effect of the Time Since Arrival on departure probability

Our statistical framework to address covariates influencing the probability of leaving a stopover site allowed us to provide evidence that TSA is the major determinant of this probability for the Sedge Warbler at our studied area. TSA is a reliable indicator of the

propensity of an individual to leave the stopover site. We acknowledge that TSA is only a proxy of all the changes in the individual ‘state’ (*sensu* Clark and Mangel 2000) during the stopover, but it is reasonable to think that TSA reflects the progressive change in the individual internal state. On the first day after arrival, birds are supposed to be exhausted, starving and to lack fuel stores. The longer they stay, the more opportunities they have to rest, feed and fatten (Schmaljohann and Eikenaar 2017).

TSA encompasses different functions of the stopover behavior: (i) resting after a migratory flight (McWilliams et al. 2004, Skrip et al. 2015), and (ii) reaching a sufficient level of fuel load to perform the following migratory flight (Schmaljohann and Eikenaar 2017). TSA also reflects (iii) the refueling rate, which depends on environmental conditions and physiological processes involved in refueling (Jenni and Schaub 2003, Schmaljohann and Eikenaar 2017). Here, TSA has a positive effect on departure probability: birds need to stay a sufficient number of days before leaving. For birds that do fatten, the fattening (or fuel deposition) increases through time: the longer individuals stopover, the larger their last measured body mass or fuel load (mean mass gain of 0.48g in this study; T. Alerstam, 1990; Schmaljohann & Eikenaar, 2017) and the larger the daily mass gain (Péron et al. 2007). Consequently, it is reasonable to think that birds need to rebuild a sufficient level of fuel store to perform another migratory flight (Alerstam 1990). Moreover, the resting time after a long migratory flight is apparently brief and confined to the first hours or days of the stopover (Fuchs et al. 2006, 2009; Németh 2009). This suggests that resting is not the physiological process that requires a 12-day stopover. Rather, most of the time spent in stopover is allocated to foraging in order to refill energy reserves; the latter is a long and progressive process. Hence, for the fraction of birds that stopover at this place, we believe that TSA reflects the time required by the physiological processes involved in refueling. Determining the precise relationship between

TSA and fuel store will be an important area of future research in stopover ecology and more specifically for the application of the present modelling framework.

Traditional measures of fuel store such as size-scaled body mass or fat score have some limitations (extensively discussed in Schmaljohann & Eikenaar, 2017; Schilch & Jenni, 2001), of which we want to highlight two. (i) In many long-term migration monitoring programs of marked birds, especially in old datasets, body mass was not systematically recorded at each recapture event. This drastically reduces the sample size available for long term analyses where we need a body mass measurement at each recapture. In the French dataset for the Sedge Warbler, in the twentieth century, individuals were weighed on only 60% of the capture events (Dehorter and CRBPO 2015). Nowadays (2000-2016 period), individuals are weighed on nearly all the capture events (90%). (ii) Since the probability of being captured in a given day can be low in routine trapping protocols (0.161 [0.058, 0.376] in this study; Schaub et al. 2001, Schmaljohann and Eikenaar 2017), the body mass measured at the latest capture is unlikely to be representative of the body mass that actually triggers departure. Imperfect detection probability is a common situation where modeling the individual history before the first capture is required to estimate arrival date. This imprecision in the assessment of the body mass change through time can mask the effect of body mass on departure probability in datasets where daily capture probability is low. Overall, the proposed analytical method, relying on TSA, allows analyzing individual variations in stopover duration over long time series, even in absence of biometric data, and even when body mass or fat score information are too sparse to reliably document the progress through time of the energetic state of each monitored individual.

Using TSA as a proxy for the internal state of the bird just before departure make some critical assumptions. When birds stay only few days, they may not improve condition with time spent at the stopover site because they may first continue to degrade upon arrival waiting for their digestive system to redevelop to refuel after a long-distant migratory flight (McWilliams et al. 2004). To overcome this situation, further research on the refueling process during stopover may be helpful to better model the TSA effect. The relevance of TSA may also be limited when birds relocate in the vicinity of the study site (Bächler and Schaub 2007): in this case the departure from the study site does not mean that individuals are resuming a long-distance migratory flight. Birds relocating a few kilometers away from the study site do not need to wait until their internal state improves before leaving the site. To get around these problems, birds captured only once are usually removed from capture-recapture datasets. This removes most of the transient individuals (Mills et al. 2011, Taylor et al. 2011, Sjöberg et al. 2015). Including both estimated TSA and fuel store data (body mass, fat score) into a model would be an interesting avenue to revisit previous analyses. This will help understand the contrasted results of previous studies about fuel store effects on departure probability (Tsvey et al. 2007, Schaub et al. 2008, Arizaga et al. 2011, Smith and McWilliams 2014, Schmaljohann and Eikenaar 2017). Also, other limitations, more related to modelling and data, appear when we use this type of model. It requires a large amount of recapture to test for more precise effects of the various determinants of departure probability. Interactions between TSA and date or TSA and age could have been tested. However, to correctly estimate the interaction, it is possible that it would have been necessary to drastically reduce the dataset to keep individuals captured at least three times, which does not exist for some years of our dataset. This would be interesting avenues to test on a dataset with a higher detection probability, for example with a capture-mark-resight shorebirds dataset (Lok et al. 2019).

Effect of weather conditions on departure probability

In relatively humid days, birds tend to postpone departure. Birds wait for dryer conditions to resume their migratory flight. Humidity can be high even in absence of precipitation in Western Brittany. The negative effect of high humidity can reflect not only the inhibitory effect of rain, but also the increased flight cost when the air is very humid (Åkesson et al. 2001, Deppe et al. 2015). The probability of departing from the stopover site increases when atmospheric pressure increases between the day before the night of departure and two days before. When birds perceive an increase in pressure (indicative of improving, anticyclonic conditions), this could encourage them to resume migration flight by the following night. Unexpectedly, departure probability does not depend on wind, probably because wind was too rare and weak in the study area in August to be influential (mean wind force during the study period was 5 m.s⁻¹, see Supplementary material for a summary of the weather covariates).

Effect of migratory experience on departure probability

Juveniles (birds migrating for the first time) stay on average 1.6 days (11.6%) less than the older, more experienced birds at the stopover area. Even though this account for a limited variation in stopover duration in our study and also that there is no consensus about the age effect on stopover strategy (Hake et al. 2003, Moore et al. 2017), this result is consistent with former studies that have shown that juvenile and experienced birds behave differently regarding departure probability from stopover sites: juveniles make shorter and more frequent stopovers than adults (Reilly and Reilly 2009, McKinnon et al. 2014). However, other studies reached opposite conclusions: telemetry studies of departure decisions in songbirds in the Gulf of Mexico during autumn did not find any effect of age on the decision to cross the Gulf of Mexico, or of the selection of weather conditions (McKinnon et al. 2014, Deppe et al. 2015). Again, it is legitimate to ask whether imperfect detection probability is involved in

inconsistencies among studies concerning the relationship between age and departure probability from stopover sites.

Respective effects of TSA and weather conditions on departure probability

In this study, we show that the contribution of TSA to departure probability in a long-distant migrant is larger than that of weather conditions. This suggests that even when weather conditions are favorable to departure, birds need to stay before departing. To our knowledge, this is the first time that the contributions of these factors were estimated with a capture-recapture model. Nevertheless, our result is consistent with numerous studies that have highlighted the key role of the improvement of the individual internal state (fuel store, fuel deposition rate, body condition, body mass, resting) on departure decisions from a stopover site (see Schmaljohann and Eikenaar 2017 for a review and Anderson et al. (2019) for a recent study with telemetry tools).

The improvement of the internal state during a stopover could be indirectly related to environmental conditions. Indeed, harsh weather can decrease the ability of individuals to feed and food abundance in the stopover area. This likely leads to a decrease in the rate at which birds accumulate energy (Jenni and Schaub 2003). If this holds, weather conditions can affect TSA and stopover duration. Interestingly, here we found slight evidence of an effect of weather conditions on departure probability while TSA was taken into account in the analysis. The relationship between TSA and weather conditions during the stopover should be addressed in future work to better understand the processes involved in departure decisions from a stopover site.

Recent studies also highlighted that migration distance affects the stopover strategy of birds (Anderson et al. 2019) and that birds also behave differently between stopover sites along

their journey (Schmaljohann et al. 2017). Concerning the Sedge Warbler at our studied site, we have no clues from a specific origin (controls indicate birds from Great Britain, Scandinavia, Eastern Europe; Dehorter and CRBPO 2015) or a final destination. However, as we only study one species which is strictly trans-saharian, the variability of stopover strategy induced by migration distance may be limited.

Conclusions

We incorporated TSA, a partially hidden individual state, and weather conditions in the same capture-recapture modeling framework to disentangle the factors playing a part in the decision to depart from a stopover site. Using a long-term dataset, we showed that TSA is the main driver of departure probability (and of stopover duration) in a long-distant migrant songbird. This approach will allow investigating the determinants of stopover duration and departure probability (not only weather variables but also some hidden physiological processes accounted for by TSA) in hundreds of existing long-term datasets, where there is no, or scattered information about mass or fat score. We demonstrated the feasibility and relevance of this analytical approach using data from one site, one species and over a large period of time. Our modeling approach will have to be used with data from several species, at several sites, to assess the robustness and generality of our conclusion about the major influence of TSA on the time when individuals decide to leave stopover sites.

TSA also has broader implications outside migration ecology in situations where the time individuals spend on sites is a partially observable variable. For example, in behavioral ecology and foraging ecology, the probability of an individual changing foraging site could also depend on the time spent in a site, the number of competitors, or food availability. TSA opens large perspectives when behaviour depends on the time spent in a site, in a specific

state, and when detection probability is imperfect at the time scale relevant to the research topic addressed.

Acknowledgements

This study was made possible thanks to tens of bird ringers and hundreds of assistants who volunteered in this long-term monitoring over three decades, and the continued support of the Société d'Etude de la Protection et de la Nature de Bretagne / Bretagne Vivante. This study was supervised by the Centre de Recherches sur la Biologie des Populations d'Oiseaux (CRBPO), the Museum National d'Histoire Naturelle, the CNRS and the Ministère de l'Environnement. This work was carried out in accordance with standard animal care protocols approved by the CRBPO. Emmanuelle Cam was supported by the French Laboratory of Excellence project "TULIP" (ANR-10-LABX-41; ANR-11-IDEX-0002-02).

Data availability statement

Data used to conduct the analyses is available on the github account of the corresponding author at the following link https://github.com/sebroques/Article_Sedge_warbler

References

- Åkesson, S., and A. Hedenström (2007). How Migrants Get There: Migratory Performance and Orientation. *BioScience* 57:123–133.
- Åkesson, S., G. Walinder, L. Karlsson, and S. Ehnborn (2001). Reed warbler orientation: Initiation of nocturnal migratory flights in relation to visibility of celestial cues at dusk. *Animal Behaviour* 61:181–189.
- Åkesson, S., G. Walinder, L. Karlsson, and S. Ehnborn (2002). Nocturnal migratory flight initiation in reed warblers *Acrocephalus scirpaceus*: Effect of wind on orientation and

443 timing of migration. *Journal of Avian Biology* 33:349–357.

444 Alerstam, T. (1990). *Bird migration*. Cambridge University Press.

445 Anderson, A. M., S. Duijns, P. A. Smith, C. Friis, and E. Nol (2019). Migration Distance and
446 Body Condition Influence Shorebird Migration Strategies and Stopover Decisions
447 During Southbound Migration. *Frontiers in Ecology and Evolution* 7:1–14.

448 Arizaga, J., E. J. Belda, and E. Barba (2011). Effect of fuel load, date, rain and wind on
449 departure decisions of a migratory passerine. *Journal of Ornithology* 152:991–999.

450 B. Bargain, C. Vansteenwegen, and J. Henry (2002). Importance des marais de la baie
451 d’Audierne (Bretagne) pour la migration du phragmite des joncs *Acrocephalus*
452 *schoenobaenus*. *Alauda* 70:37–55.

453 Bächler, E., and M. Schaub (2007). the Effects of Permanent Local Emigration and Encounter
454 Technique on Stopover Duration Estimates As Revealed By Telemetry and Mark-
455 Recapture. *The Condor* 109:142–154.

456 Bibby, C. J., and R. E. Green (1981). Autumn migration strategies of Reed and Sedge
457 Warblers. *Ornis Scandinavica* 12:1–12.

458 Clark, C. W., and M. Mangel (2000). *Dynamic state variable models in ecology: methods and*
459 *applications*. Oxford University Press on Demand.

460 Cohen, E. B., F. R. Moore, and R. A. Fischer (2014). Fuel stores, time of spring, and
461 movement behavior influence stopover duration of Red-eyed Vireo *Vireo olivaceus*.
462 *Journal of Ornithology* 155:785–792.

463 Crosbie, S. F., and B. F. J. Manly (1985). *Parsimonious Modelling of Capture-Mark-*

464 Recapture Studies. *Biometrics* 41:385.

465 Dehorter, O., and CRBPO (2015). Bird ringing and movement database for France. Centre de
466 Recherches sur la Biologie des Populations d'Oiseaux, Muséum National d'Histoire
467 Naturelle, Paris, France. <http://crbpo.mnhn.fr/>. Consulté le / Accessed on 21 August
468 2015.

469 Deppe, J. L., M. P. Ward, R. T. Bolus, R. H. Diehl, A. Celis-Murillo, T. J. Zenzal, F. R.
470 Moore, T. J. Benson, J. A. Smolinsky, L. N. Schofield, D. A. Enstrom, et al. (2015). Fat,
471 weather, and date affect migratory songbirds' departure decisions, routes, and time it
472 takes to cross the Gulf of Mexico. *Proceedings of the National Academy of Sciences*
473 112:E6331–E6338.

474 Dossman, B. C., G. W. Mitchell, D. R. Norris, P. D. Taylor, C. G. Guglielmo, S. N.
475 Matthews, and P. G. Rodewald (2016). The effects of wind and fuel stores on stopover
476 departure behavior across a migratory barrier. *Behavioral Ecology* 27:567–574.

477 Fuchs, T., A. Haney, T. J. Jechura, F. R. Moore, and V. P. Bingman (2006). Daytime naps in
478 night-migrating birds: behavioural adaptation to seasonal sleep deprivation in the
479 Swainson's thrush, *Catharus ustulatus*. *Animal Behaviour* 72:951–958.

480 Fuchs, T., D. Maury, F. R. Moore, and V. P. Bingman (2009). Daytime micro-naps in a
481 nocturnal migrant: An EEG analysis. *Biology Letters* 5:77–80.

482 Gimenez, O., V. Rossi, R. Choquet, C. Dehais, B. Doris, H. Varella, J. P. Vila, and R. Pradel
483 (2007). State-space modelling of data on marked individuals. *Ecological Modelling*
484 206:431–438.

485 Goymann, W., F. Spina, A. Ferri, and L. Fusani (2010). Body fat influences departure from

486 stopover sites in migratory birds: evidence from whole-island telemetry. *Biology Letters*
487 6:478–481.

488 Green, M., T. Alerstam, P. Clausen, R. Drent, and B. S. Ebbinge (2002). Dark-bellied Brent
489 Geese *Branta bernicla bernicla*, as recorded by satellite telemetry, do not minimize flight
490 distance during spring migration. *Ibis* 144:106–121.

491 Hair, J. F., W. C. Black, B. J. Babin, R. E. Anderson, and R. L. Tatham (2006). Multivariate
492 data analysis (Vol. 6).

493 Hake, M., N. Kjellén, and T. Alerstam (2003). Age-dependent migration strategy in honey
494 buzzards *Pernis apivorus* tracked by satellite. *Oikos*. pp. 385–396.

495 Hedenström, A., and T. Alerstam (1997). Optimum fuel loads in migratory birds:
496 Distinguishing between time and energy minimization. *Journal of Theoretical Biology*
497 189:227–234.

498 Hornik, K., F. Leisch, A. Zeileis, and M. Plummer (2003). JAGS: A Program for Analysis of
499 Bayesian Graphical Models Using Gibbs Sampling.

500 Jenni, L., and M. Schaub (2003). Behavioural and Physiological Reactions to Environmental
501 Variation in Bird Migration: a Review. *Avian Migration*. [Online.] Available at
502 http://link.springer.com/10.1007/978-3-662-05957-9_10.

503 Jolly, G. M. (1965). Explicit Estimates from Capture-Recapture Data with Both Death and
504 Immigration-Stochastic Model. *Biometrika* 52:225.

505 Kéry, M., and M. Schaub (2011). Bayesian Population Analysis using WinBUGS: A
506 hierarchical perspective. Academic Press.

- 507 Klinner, T., J. Buddemeier, F. Bairlein, and H. Schmaljohann (2020). Decision-making in
508 migratory birds at stopover: an interplay of energy stores and feeding conditions.
509 Behavioral Ecology and Sociobiology 74:1–14.
- 510 Ktitorov, P., A. Tsvey, and A. Mukhin (2010). The good and the bad stopover: Behaviours of
511 migrant reed warblers at two contrasting sites. Behavioral Ecology and Sociobiology
512 64:1135–1143.
- 513 Lok, T., C. J. Hassell, T. Piersma, R. Pradel, and O. Gimenez (2019). Accounting for
514 heterogeneity when estimating stopover duration, timing and population size of red knots
515 along the Luannan Coast of Bohai Bay, China. Ecology and Evolution 9:6176–6188.
- 516 Lyons, J. E., W. L. Kendall, J. A. Royle, S. J. Converse, B. A. Andres, and J. B. Buchanan
517 (2016). Population size and stopover duration estimation using mark-resight data and
518 Bayesian analysis of a superpopulation model. Biometrics 72:262–271.
- 519 Ma, Z., N. Hua, X. Zhang, H. Guo, B. Zhao, Q. Ma, W. Xue, and C. Tang (2011). Wind
520 conditions affect stopover decisions and fuel stores of shorebirds migrating through the
521 south Yellow Sea. Ibis 153:755–767.
- 522 McKinnon, E. A., K. C. Fraser, C. Q. Stanley, and B. J. M. Stutchbury (2014). Tracking from
523 the tropics reveals behaviour of juvenile songbirds on their first spring migration. PLoS
524 ONE 9:e105605.
- 525 McWilliams, S. R., C. Guglielmo, B. Pierce, and M. Klaassen (2004). Flying, fasting, and
526 feeding in birds during migration: A nutritional and physiological ecology perspective.
527 *Journal of Avian Biology*.
- 528 Mills, A. M., B. G. Thurber, S. A. Mackenzie, and P. D. Taylor (2011). Passerines Use

Nocturnal Flights for Landscape-Scale Movements during Migration Stopover. The
Condor 113:597–607.

Moore, F. R., K. M. Covino, W. B. Lewis, T. J. Zenzal, and T. J. Benson (2017). Effect of
fuel deposition rate on departure fuel load of migratory songbirds during spring stopover
along the northern coast of the Gulf of Mexico. *Journal of Avian Biology* 48:123–132.

Müller, F., C. Eikenaar, Z. J. Crysler, P. D. Taylor, and H. Schmaljohann (2018). Nocturnal
departure timing in songbirds facing distinct migratory challenges. *Journal of Animal
Ecology* 87:1102–1115.

Németh, Z. (2009). Observation of Daytime Sleep-like Behavior in a Migratory Songbird
during Stopover. *The Wilson Journal of Ornithology* 121:644–646.

Péron, G., P. Y. Henry, P. Provost, O. Dehorter, and R. Julliard (2007). Climate changes and
post-nuptial migration strategy by two reedbed passerines. *Climate Research* 35:147–
157.

Pledger, S., M. Efford, K. Pollock, J. Collazo, and J. Lyons (2008). Stopover Duration
Analysis with Departure Probability Dependent on Unknown Time Since Arrival. In
Modeling Demographic Processes in Marked Populations. pp. 349–363.

R Development Core Team 3.0.1. (2013). A Language and Environment for Statistical
Computing. R Foundation for Statistical Computing 2:<https://www.R-project.org>.

Reilly, J. R., and R. J. Reilly (2009). Bet-hedging and the orientation of juvenile passerines in
fall migration. *Journal of Animal Ecology* 78:990–1001.

Roques, S., D. B. Lank, E. Cam, and R. Pradel (2020). More Than Just Refuelling: Lengthy
Stopover and Selection of Departure Weather by Sandpipers Prior to Transoceanic and

551 Transcontinental Flights. *Ibis*:ibi.12891.

552 Royle, J. A. (2008). Modeling Individual Effects in the Cormack-Jolly-Seber Model: A State-
553 Space Formulation. *Biometrics* 64:364–370.

554 Schaub, M., and L. Jenni (1999). Does Tape-Luring of Migrating Eurasian Reed-Warblers
555 Increase Number of Recruits or Capture Probability? *The Auk* 116:1047–1053.

556 Schaub, M., L. Jenni, and F. Bairlein (2008). Fuel stores, fuel accumulation, and the decision
557 to depart from a migration stopover site. *Behavioral Ecology* 19:657–666.

558 Schaub, M., R. Pradel, L. Jenni, and J. D. Lebreton (2001). Migrating birds stop over longer
559 than usually thought: An improved capture-recapture analysis. *Ecology* 82:852–859.

560 Schmaljohann, H. (2018). Proximate mechanisms affecting seasonal differences in migration
561 speed of avian species. *Scientific Reports* 8:1–10.

562 Schmaljohann, H., and C. Eikenaar (2017). How do energy stores and changes in these affect
563 departure decisions by migratory birds? A critical view on stopover ecology studies and
564 some future perspectives. *Journal of Comparative Physiology A: Neuroethology,*
565 *Sensory, Neural, and Behavioral Physiology* 203:411–429.

566 Schmaljohann, H., J. W. Fox, and F. Bairlein (2012). Phenotypic response to environmental
567 cues, orientation and migration costs in songbirds flying halfway around the world.
568 *Animal Behaviour* 84:623–640.

569 Schmaljohann, H., S. Lisovski, and F. Bairlein (2017). Flexible reaction norms to
570 environmental variables along the migration route and the significance of stopover
571 duration for total speed of migration in a songbird migrant. *Frontiers in Zoology* 14:17.

572 Schmaljohann, H., F. Müller, T. Klinner, and C. Eikenaar (2018). Potential age differences in
573 the migratory behaviour of a nocturnal songbird migrant during autumn and spring.
574 *Journal of Avian Biology* 49:1–16.

575 Schwarz, C. J., and A. N. Arnason (1996). A General Methodology for the Analysis of
576 Capture-Recapture Experiments in Open Populations. *Biometrics* 52:860.

577 Schwilch, R., and L. Jenni (2001). Low initial refueling rate at stopover sites: a
578 methodological effect? *The Auk* 118:698–708.

579 Schwilch, R., L. Jenni, T. Piersma, and N. M. A. Holmgren (2002). Do migratory birds need a
580 nap after a long non-stop flight? *Ardea* 90:149–154.

581 Seber, G. A. F. (1965). A Note on the Multiple-Recapture Census. *Biometrika* 52:249.

582 Sjöberg, S., T. Alerstam, S. Åkesson, A. Schulz, A. Weidauer, T. Coppack, and R. Muheim
583 (2015). Weather and fuel reserves determine departure and flight decisions in passerines
584 migrating across the Baltic Sea. *Animal Behaviour* 104:59–68.

585 Skrip, M. M., U. Bauchinger, W. Goymann, L. Fusani, M. Cardinale, R. R. Alan, and S. R.
586 McWilliams (2015). Migrating songbirds on stopover prepare for, and recover from,
587 oxidative challenges posed by long-distance flight. *Ecology and Evolution* 5:3198–3209.

588 Smith, A. D., and S. R. McWilliams (2014). What to do when stopping over: Behavioral
589 decisions of a migrating songbird during stopover are dictated by initial change in their
590 body condition and mediated by key environmental conditions. *Behavioral Ecology*
591 25:1423–1435.

592 Somveille, M., A. Manica, and A. S. L. Rodrigues (2019). Where the wild birds go:
593 explaining the differences in migratory destinations across terrestrial bird species.

594   Ecography 42:225–236.

595   Somveille, M., A. S. L. Rodrigues, and A. Manica (2015). Why do birds migrate? A
596   macroecological perspective. *Global Ecology and Biogeography* 24:664–674.

597   Somveille, M., A. S. L. Rodrigues, and A. Manica (2018). Energy efficiency drives the global
598   seasonal distribution of birds. *Nature Ecology and Evolution* 2:962–969.

599   Su, Y.-S., M. Yajima, M. Y.-S. Su, and J. SystemRequirements (2015). Package ‘R2jags.’ R
600   package version 0.03-08, URL <http://CRAN.R-project.org/package=R2jags>.

601   Taylor, P. D., S. A. Mackenzie, B. G. Thurber, A. M. Calvert, A. M. Mills, L. P. McGuire,
602   and C. G. Guglielmo (2011). Landscape movements of migratory birds and bats reveal
603   an expanded scale of stopover. *PLoS ONE* 6:e27054.

604   Tsvey, A., V. N. Bulyuk, and V. Kosarev (2007). Influence of body condition and weather on
605   departures of first-year European robins, *Erithacus rubecula*, from an autumn migratory
606   stopover site. *Behavioral Ecology and Sociobiology* 61:1665–1674.

607   Warnock, N. (2010). Stopping vs. staging: The difference between a hop and a jump. *Journal*
608   *of Avian Biology* 41:621–626.

609   Wikelski, M., A. Raim, E. M. Tarlow, R. P. Larkin, G. H. Visser, and R. H. Diehl (2003).
610   Costs of migration in free-flying songbirds. *Nature* 423:704–704.

611   Zehnder, S., S. Åkesson, F. Liechti, and B. Bruderer (2001). Nocturnal autumn bird migration
612   at Falsterbo, South Sweden. *Journal of Avian Biology* 32:239–248.

613   Zúñiga, D., Y. Gager, H. Kokko, A. M. Fudickar, A. Schmidt, B. Naef-Daenzer, M. Wikelski,
614   and J. Partecke (2017). Migration confers winter survival benefits in a partially

615   migratory songbird. eLife 6:1–12.

616

Table 1: Mean value, standard deviation and credible intervals (CI) for each covariate effect on the probability of staying at the stopover area. Bold: significant effects.

	<i>Mean</i>	<i>Sd</i>	<i>CI</i>
(α) <i>Intercept</i>	6.159	0.175	5.843, 6.578
(β_1) <i>Time Since Arrival</i>	-1.538	0.039	-1.652, -1.432
(β_2) <i>Wind</i>	0.116	0.1	-0.121, 0.196
(β_3) <i>Temperature</i>	-0.207	0.89	-0.387, 0.03
(β_4) <i>Cloud cover</i>	0.142	0.101	-0.052, 0.207
(β_5) <i>Humidity</i>	0.497	0.121	0.261, 0.735
(β_6) Δ <i>Pressure</i>	-0.702	0.23	-1.176, -0.196

Figure 1: Between-year variation in individual stopover duration estimates (in days) per age category for the Sedge Warbler (*Acrocephalus schoenobaenus*).

Figure 2: Departure probability as a function of the number of days since the bird arrived. **(A)** For different humidity conditions, dashed line: low humidity conditions (~70%); plain: high humidity conditions (~90%). **(B)** For different Δ Pressure conditions, dashed line: a substantial decrease of atmospheric pressure (-5Hpa); plain: a substantial increase of atmospheric pressure (+5Hpa). The grey area represents departure probability values for humidity between 70 and 90% (a) and Δ Pressure between -5Hpa and +5Hpa.