Increased cortical volume without increased neuron number in heterozygous *Chd8* mutant mouse cortex

Cesar P. Canales^{1,2,4}, Samuel Frank^{2,4}, Jeffrey Bennett², Paris Beauregard¹, Pierre Lavenex³,
David G. Amaral² and Alex. S. Nord^{1,2}

¹ Center for Neuroscience, UC Davis, Davis, CA, USA. ² Department of Psychiatry and Behavioral Sciences, The M.I.N.D. Institute, UC Davis, Davis, CA, USA. ³ Laboratory of Brain and Cognitive Development, Institute of Psychology, University of Lausanne, Switzerland.

⁴ These authors contributed equally to this work.

Corresponding author:

Alex S. Nord
Center for Neuroscience
UC Davis
1544 Newton Court
Davis CA 95618
Phone: 530-754-5022
Email: asnord@ucdavis.edu

Running title: **Cortical stereology in a *Chd8* mouse model**

Key Words: ASD, Macrocephaly, Stereology, Chd8, mouse model

Acknowledgements

This work and ASN and CPC were supported by NIH NIMH (MH120513-02) and NIGMS (R35 GM119831-05). CPC is recipient of NIMH Institutional National Research Service Award Autism Research Training Program fellowship (T32 MH073124-16). This research was also supported by NIH Grants 1R01MH121447-01 and 1R01MH103371 to DGA and Swiss National Science Foundation grant 310030_143956 to PL. Authors declare no conflict of interest.

Lay Summary

We measured the size and neuron number in the neocortex in mice with heterozygous *Chd8* mutation, a model relevant to Autism Spectrum Disorder. We found an increased cortical volume in male mutants, which was not accompanied by increased neuron number or soma size. Our results indicate that the enlarged brain in *Chd8* mutant mice is complex, more evident here in males, and is due to factors other than increased neuron number.

Abstract

De novo mutations in the chromatin-remodeling factor *CHD8* (Chromodomain-Helicase DNA-binding protein 8) have emerged as a key genetic risk factor for Autism Spectrum Disorder (ASD) and, more generally, neurodevelopmental disorders. Individuals with heterozygous mutations in *CHD8* typically present hallmarks of ASD with comorbid cognitive disability and macrocephaly. Knockdown or haploinsufficiency of *Chd8* in animal models has recapitulated phenotypes observed in patients, including increased head circumference and brain size. Here, we aimed to determine whether increased neuron numbers or soma size drives increased cortical volume. We performed design-based stereological analyses of cortical structure in adult male and female heterozygous *Chd8* mice and wild-type littermate controls. *Chd8* haploinsufficient male mice displayed a ~8-12% increase in cortical volume, no differences in cortical neuron number and comparable neuronal soma size. Our study reproduced previous reports of increased brain size associated with *CHD8* mutation in humans and mice and are consistent with reported sex-specific impacts of *Chd8* mutations in mice and increased burden of *CHD8* mutations in human males with ASD. These findings suggest that the nature of the cortical enlargement due to *Chd8* haploinsufficiency is complex and appears to be due to a factor other than an increased neuron number or soma size.

Introduction

Autism spectrum disorder (ASD) is a genetically heterogeneous disorder that involves the interaction of both common and rare variants and produces a broad phenotypic spectrum (Geschwind, 2009). Despite the clinical and genetic heterogeneity that typifies ASD, some anatomical features appear to be either present in many cases or so dramatically altered in some that their presence is now reasonably well replicated as characteristic of ASD among a subset of cases across a number of studies. One such finding is the tendency towards increased size of the brain during the early postnatal period causing macrocephaly (Albores-Gallo et al., 2017; Amaral et al., 2017; Bernier et al., 2014, 2016; Donovan & Basson, 2017; Li et al., 2017; Libero et al., 2016; Nordahl et al., 2011; Woodbury-Smith et al., 2017; Wu et al., 2020).

Mutations in chromatin remodeler factors (CRFs) are strongly associated with causal roles across Neurodevelopmental Disorders (NDDs) (De Rubeis et al., 2014; Gilissen et al., 2014; Iossifov et al., 2014; Nishiyama et al., 2004; O’Roak et al., 2012; O’Roak et al., 2012; Sanders et al., 2015; Vissers et al., 2016; Vogel-Ciernia & Wood, 2014). Of the top ASD risk genes ranked by enrichment of *de novo* case mutations, approximately 40% are CRFs (Sanders et al., 2015; Satterstrom et al., 2020). Heterozygous mutations to different CRFs cause similar phenotypes (Bögershausen & Wollnik, 2018), suggesting shared mechanisms of brain pathology. The gene *CHD8*, which encodes the CRF chromodomain helicase DNA-binding protein 8, has been implicated in genetic analyses profiling rare and *de novo* coding variation in ASD and intellectual disability (ID) (An et al., 2020; Barnard et al., 2015; De Rubeis et al., 2014; O’Roak et al., 2012; Tatton-Brown et al., 2017). Along with ASD, humans with *CHD8* mutations

frequently exhibit ID, macrocephaly, craniofacial dysmorphology, and gastrointestinal dysfunction (Barnard et al., 2015; Beighley et al., 2020; Bernier et al., 2014).

The proportion of individuals with *CHD8* mutations who present macrocephaly is much higher than that seen in the general ASD population, and NDD-relevant behaviors and macrocephaly have been reported across multiple *Chd8* mutant mouse models (Barnard et al., 2015; Bernier et al., 2014; Gompers et al., 2017; Katayama et al., 2016; Platt et al., 2017a; Suetterlin et al., 2018; Zhao et al., 2018). We previously reported the impact of germline heterozygous frameshift *Chd8* mutation (*Chd8*^{+/del5}) on neurodevelopment in mice (Gompers et al., 2017). *Chd8*^{+/del5} mice exhibited cognitive impairment correlated with increased regional brain volume. To date, the differences underlying increased brain and cortical volume in mature mutant *Chd8* mice remain undefined. Towards advancing understanding of megalencephaly and increased brain volume associated with *Chd8* haploinsufficiency, we applied design-based stereology to characterize the neocortex of *Chd8*^{+/del5} mice. Design-based stereology is now the gold-standard method to obtain reliable quantitative measurements of brain structure (Hilgetag et al., 2018; Kipp et al., 2017). We estimated cortical volume, neuron number and neuronal soma size in male and female *Chd8*^{+/del5} mice and their wild-type littermates.

Methods

Animals

Generation of Cas9-mediated 5bp frameshift deletion in exon 5 of *Chd8* in mice was previously described (Gompers et al., 2017). Thirty-two 42-day-old mice were included in this analysis (16

brains per genotype, 16 per sex). All protocols utilized in the generation of brain histological samples were approved by the Institutional Animal Care and Use Committees (IACUC) at the University of California Davis. Mice were housed in a temperature-controlled vivarium maintained on a 12-h light–dark cycle. We made efforts to minimize pain, distress and the number of animals used in the study.

Tissue preparation and Nissl Stain

Mice were anesthetized using Isoflurane and perfused transcardially using room temperature saline solution (0.9% NaCl) followed by 4% paraformaldehyde (PFA) in 0.1 M phosphate buffered (PB) saline using a peristaltic pump (Minipuls 3, Gilson) at 20 rpm flow rate. Brains were removed and post-fixed in 4% PFA at 4°C for 24 h. Fixed brains were cryoprotected in 50 ml of 10% and 20% glycerol solutions with 2% DMSO in 0.1 M phosphate buffer (PB; pH 7.4; for 24 and 72 hours, respectively). Brains were blocked at -2.00mm posterior to the interaural line prior to flash freezing. Sections were cut using a freezing, sliding microtome in 4 series at 60 µm (Microm HM 440E, Microm International, Germany). Every 4th section was collected in 10% formaldehyde in 0.1 M PB (pH 7.4) and post-fixed at 4°C for 1 week (Chareyron et al., 2011). Sections were rinsed twice for 1 hour each in 1ml of 0.1M PB, mounted on gelatin-coated slides, and air-dried overnight at 37°C. Defatting was performed in a mixture of chloroform/ethanol (1:1, vol) for 2 x 1 hour. The brains were rehydrated in graded alcohols; 100% ethanol, 100% ethanol, 95% ethanol, 70%, 50% and dipped in deionized water (Chareyron et al., 2011; Lavenex et al., 2009). Sections were stained 40 seconds in a 0.25% thionin solution (Fisher Scientific, Waltham, MA, cat. no. T-409), dehydrated through graded alcohols,

differentiated in 95% ethanol and 5 drops of glacial acetic acid, cleared in xylene and cover-slipped with DPX (BDH Laboratories, Poole, UK) (Chareyron et al., 2011).

Stereology

The inferior boundary of the neocortex was defined as the bottom of the agranular insular area (AIA), and the superior boundary was defined as the anterior cingulate area (ACA) or the retrosplenial area (RSA) and the white matter was then followed to the ACA/RSA border (Franklin, 2013). Neuron number, neuronal soma size, and brain volume were estimated using Stereoinvestigator V10.50 (MBF Biosciences, Williston, VT) attached to a Zeiss Imager.Z2 Vario with a Zeiss AxioCam MRc. An EC PlanNeoFluar 2.5 x objective was used to trace the outline of the neocortex and a PlanApoChromat 100x oil objective was used for cell counts and soma size determinations. The volume of the neocortex was determined using the Cavalieri method (Gundersen & Jensen, 1987; Lavenex et al., 2000; Mark J. West & Gundersen, 1990). For each mouse, 15-18 sections (480 μ m apart) were analyzed with the first section chosen randomly amongst the first through the neocortex. Estimates of neuron numbers were obtained using the Optical Fractionator method (Gundersen, 1986; M. J. West et al., 1991). Since no lateralization was found in the volumetric estimates ($p = 0.65$), we sampled the left hemisphere of half of the mice and the right hemisphere of the other half for estimates of neuron numbers. We used the same sections that were used for volumetric estimates. We used a counting frame size of 18x18x10 μ m with 2 μ m guard zones, and a scan grid of 480x480 μ m placed at a random angle. Neurons were defined following established protocols (Gundersen, 1986; Gundersen & Jensen, 1987; Lavenex et al., 2000; Long et al., 1998; M. J. West et al., 1991; Mark J. West & Gundersen, 1990), based on: 1) dark staining, 2) a well-defined nucleolus and 3) a round, non-

jagged cell body (Chareyron et al., 2011; García-Cabezas et al., 2016; Haida et al., 2019; Lauber et al., 2018; Long et al., 1998). Section thickness was measured at every counting site. The volume of neuronal somas was determined using the Nucleator method (Gundersen, 1988) during the Optical Fractionator analysis. Comparison of stereological measures (volume, neuron number, and soma size) were performed via ANOVA to examine global effects of sex and genotype and test for sex by genotype interaction, and then via t-test to compare the effect of genotype in males and females. In addition to test results, effect sizes are reported for ANOVA (partial eta squared, η_p^2) and t-test (Cohen's d). Statistical analyses were performed using SPSS (version 26), plots were generated using *ggplots2* package in R programming language (Wickham, 2009). All stereological analyses were carried out blinded to the experimental condition or sex of the animal.

Results

Nissl-stained brain sections were used to perform design-based stereological analyses of cortical structure in *Chd8^{+/-del5}* and wild-type mice at postnatal day 42, an age that may represent the end of the adolescence period in mice (Spear, 2000) (**Fig 1**). Considering cortical volume (**Fig 2a**), there was an effect of sex ($F_{(1,28)} = 4.881$, $p = 0.036$, $\eta_p^2 = 0.148$), an effect of genotype ($F_{(1,28)} = 6.190$, $p = 0.019$, $\eta_p^2 = 0.181$), and an interaction between sex and genotype that neared statistical significance ($F_{(1,28)} = 2.956$, $p = 0.097$, $\eta_p^2 = 0.095$). In males, cortical volume was larger in *Chd8^{+/-del5}* mice than in wild-type littermates ($t_{(14)} = 3.103$, $p = 0.008$, $d = 1.539$). In contrast in females, cortical volume did not differ between *Chd8^{+/-del5}* and wild-type mice ($t_{(14)} = 0.523$, $p = 0.609$, $d = 0.249$). Given the larger cortical volume in male *Chd8^{+/-del5}* mice, we sought

to determine whether this was due to differences in neuronal number (**Fig 2b**). There was no effect of sex ($F_{(1,28)} = 2.619$, $p = 0.117$, $\eta_p^2 = 0.086$), no effect of genotype ($F_{(1,28)} = 0.107$, $p = 0.747$, $\eta_p^2 = 0.004$), and no interaction between sex and genotype ($F_{(1,28)} = 1.968$, $p = 0.172$, $\eta_p^2 = 0.066$). The number of cortical neurons did not differ between *Chd8*^{+/del5} and wild-type littermates in either males ($t_{(14)} = 0.921$, $p = 0.373$, $d = 0.460$) or females ($t_{(14)} = 1.065$, $p = 0.305$, $d = 0.532$). We also sought to determine whether a larger cortical volume in male *Chd8*^{+/del5} mice was associated with differences in neuronal soma volume (**Fig 2c**). There was an effect of sex ($F_{(1,28)} = 4.611$, $p = 0.041$, $\eta_p^2 = 0.141$; males > females), but no effect of genotype ($F_{(1,28)} = 1.705$, $p = 0.202$, $\eta_p^2 = 0.057$) or interaction between sex and genotype ($F_{(1,28)} = 1.121$, $p = 0.299$, $\eta_p^2 = 0.038$). Neuronal soma size did not differ between *Chd8*^{+/del5} and wild-type littermates in either males ($t_{(14)} = 0.242$, $p = 0.812$, $d = 0.120$) or females ($t_{(14)} = 1.375$, $p = 0.191$, $d = 0.687$).

Discussion

Germline heterozygous *CHD8* mutations in humans, mice, and zebrafish all have been reported to produce macrocephaly/megalencephaly (Barnard et al., 2015; Beighley et al., 2020; Bernier et al., 2014; Gompers et al., 2017; Katayama et al., 2016; Platt et al., 2017b; Suetterlin et al., 2018; Sugathan et al., 2014; Zhao et al., 2018). As *Chd8* haploinsufficiency has also been tied to altered neurogenesis in vitro and in vivo (Cotney et al., 2015; Durak et al., 2016; Gompers et al., 2017; Suetterlin et al., 2018), one possible explanation for a larger brain size is an increased proliferation and greater numbers of neurons. Using design-based stereology, we demonstrated a larger cortical volume in male *Chd8*^{+/del5} mice. Our findings further indicate that the underlying

cause of this difference, as observed in male *Chd8*^{+/*del5*} mice here, is not an increase in neuron number or neuronal soma size in the neocortex.

Recent comparisons of *CHD8* mutations in ASD case populations indicate a higher number of male carriers, suggesting that there are sex-specific differences in susceptibility and phenotype (An et al., 2020; Bernier et al., 2016; T. Wang et al., 2016). One study of *Chd8* mutant mice similarly reported sex differences in transcriptomic and behavioral phenotypes (Jung et al., 2018). Our findings add to the evidence that males may be differentially impacted by *CHD8* mutations, as we only observed increased cortical volume in male mice in the current study. However, the literature indicates that both male and female ASD cases with *CHD8* mutations demonstrate consistent clinical phenotypes, including macrocephaly (An et al., 2020; Barnard et al., 2015; Beighley et al., 2020; Bernier et al., 2014; Douzgou et al., 2019; Li et al., 2017; T. Wang et al., 2016; Wu et al., 2020). Additionally, publications of *Chd8* mouse models, including earlier work on this line, reported relevant phenotypes in both males and females (Gompers et al., 2017; Suetterlin et al., 2018; Zhao et al., 2018). The specific *Chd8* loss-of-function mutation characterized here occurs on the C57Cl/6N inbred line, and differences across mutant alleles and inbred mouse genetic backgrounds could influence phenotype penetrance and expressivity. Future studies that are well-powered to test for sex-specific differences and incorporate genetic background are needed to determine differences in penetrance and expressivity of NDD-relevant molecular, neuroanatomical, and behavioral phenotypes in *Chd8* mouse models.

We previously reported increased embryonic neurogenesis via Edu labeling and analysis of neural progenitor and postmitotic populations at E14.5 in this same *Chd8* mutant mouse line

(Gompers et al., 2017). Other studies have similarly found evidence for macrocephaly, altered neural proliferation, and altered regulation of cell cycle genes (Durak et al., 2016; Marie et al., 2018; Suetterlin et al., 2018; Sugathan et al., 2014; P. Wang et al., 2017; Xu et al., 2018). However, the findings here suggest that any altered proliferative output during embryonic corticogenesis does not directly explain lasting postnatal increase in cortical volume in *Chd8*^{+/del5} mice. A study of *Chd8* knockdown via in utero electroporation of siRNA found decreased neurogenesis at E16 (Durak et al., 2016), and increased or decreased proliferation could be transient or compensated via increased neuronal apoptosis, resulting in no overall change in total neuron numbers (Chen et al., 2015; Naruse & Keino, 1995). Factors independent of neurogenesis could contribute to increased cortical volume in mature mice, including changes in neuronal dendritic morphology, and altered number or morphology of glial cells (Matta et al., 2019; Schumann & Nordahl, 2011; Winden et al., 2015). Heterozygous mutant *Pten* mice, another ASD risk gene associated with macrocephaly, exhibit increased neurogenesis and a greater number of neurons in early postnatal brain, but change to excess glial cells in the mature brain (Chen et al., 2015). Further, changes in brain volume are dynamic across *Pten* mutant mice and with differential impacts across brain regions (Chen et al., 2015; Clipperton-Allen et al., 2019). While mutation to *Chd8* and *Pten* represent distinct models, mutations in both lead to comorbid ASD and macrocephaly and *Chd8* has been implicated in *Pten* signaling pathways (Sugathan et al., 2014; P. Wang et al., 2017). Future efforts will be needed to test for the dynamic changes in corticogenesis and relevant neuronal populations across development and for differences in populations of non-neuronal cells as a factor in *Chd8*-associated macrocephaly.

References

- Albores-Gallo, L., Fritsche-García, L., Miranda-Aguirre, A. P., & Avila-Acosta, M. (2017). Brief Report: Macrocephaly Phenotype and Psychiatric Comorbidity in a Clinical Sample of Mexican Children and Adolescents with Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, 47(9), 2911–2917. <https://doi.org/10.1007/s10803-017-3175-4>
- Amaral, D. G., Li, D., Libero, L., Solomon, M., Van de Water, J., Mastergeorge, A., Naigles, L., Rogers, S., & Wu Nordahl, C. (2017). In pursuit of neurophenotypes: The consequences of having autism and a big brain. In *Autism Research* (Vol. 10, Issue 5, pp. 711–722). John Wiley and Sons Inc. <https://doi.org/10.1002/aur.1755>
- An, Y., Zhang, L., Liu, W., Jiang, Y., Chen, · Xue, Lan, X., Li, G., Hang, Q., Wang, J., James, ·, Gusella, F., Du, Y., & Shen, Y. (2020). De novo variants in the Helicase-C domain of CHD8 are associated with severe phenotypes including autism, language disability and overgrowth. *Human Genetics*, 139, 499–512. <https://doi.org/10.1007/s00439-020-02115-9>
- Barnard, R. A., Pomaville, M. B., & O’Roak, B. J. (2015). Mutations and modeling of the chromatin remodeler CHD8 define an emerging autism etiology. In *Frontiers in Neuroscience* (Vol. 9, Issue DEC). Frontiers Media S.A. <https://doi.org/10.3389/fnins.2015.00477>
- Beighley, J. S., Hudac, C. M., Arnett, A. B., Peterson, J. L., Gerdts, J., Wallace, A. S., Mefford, H. C., Hoekzema, K., Turner, T. N., O’Roak, B. J., Eichler, E. E., & Bernier, R. A. (2020). Clinical Phenotypes of Carriers of Mutations in CHD8 or Its Conserved Target Genes. *Biological Psychiatry*, 87(2), 123–131. <https://doi.org/10.1016/j.biopsych.2019.07.020>
- Bernier, R., Golzio, C., Xiong, B., Stessman, H. A., Coe, B. P., Penn, O., Witherspoon, K., Gerdts, J., Baker, C., Vulto-Van Silfhout, A. T., Schuurs-Hoeijmakers, J. H., Fichera, M., Bosco, P., Buono, S., Alberti, A., Failla, P., Peeters, H., Steyaert, J., Vissers, L. E. L. M., ... Eichler, E. E. (2014). Disruptive CHD8 mutations define a subtype of autism early in development. *Cell*, 158(2), 263–276. <https://doi.org/10.1016/j.cell.2014.06.017>
- Bernier, R., Steinman, K. J., Reilly, B., Wallace, A. S., Sherr, E. H., Pojman, N., Mefford, H. C., Gerdts, J., Earl, R., Hanson, E., Goin-Kochel, R. P., Berry, L., Kanne, S., Snyder, L. G., Spence, S., Ramocki, M. B., Evans, D. W., Spiro, J. E., Martin, C. L., ... Wenegrat, J. (2016). Clinical phenotype of the recurrent 1q21.1 copy-number variant. *Genetics in Medicine*, 18(4), 341–349. <https://doi.org/10.1038/gim.2015.78>
- Bögershausen, N., & Wollnik, B. (2018). Mutational Landscapes and Phenotypic Spectrum of SWI/SNF-Related Intellectual Disability Disorders. *Frontiers in Molecular Neuroscience*, 11, 252. <https://doi.org/10.3389/fnmol.2018.00252>
- Chareyron, L. J., Banta Lavenex, P., Amaral, D. G., & Lavenex, P. (2011). Stereological analysis of the rat and monkey amygdala. *Journal of Comparative Neurology*, 519(16), 3218–3239. <https://doi.org/10.1002/cne.22677>
- Chen, Y., Huang, W. C., Séjourné, J., Clipperton-Allen, A. E., & Page, D. T. (2015). Pten mutations alter brain growth trajectory and allocation of cell types through elevated β -

- Catenin signaling. *Journal of Neuroscience*, 35(28), 10252–10267.
<https://doi.org/10.1523/JNEUROSCI.5272-14.2015>
- Clipperton-Allen, A. E., Cohen, O. S., Aceti, M., Zucca, A., Levy, J., Ellegood, J., Lerch, J. P., & Page, D. T. (2019). Pten haploinsufficiency disrupts scaling across brain areas during development in mice. *Translational Psychiatry*, 9(1), 1–14. <https://doi.org/10.1038/s41398-019-0656-6>
- Cotney, J., Muhle, R. A., Sanders, S. J., Liu, L., Willsey, A. J., Niu, W., Liu, W., Klei, L., Lei, J., Yin, J., Reilly, S. K., Tebbenkamp, A. T., Bichsel, C., Pletikos, M., Sestan, N., Roeder, K., State, M. W., Devlin, B., & Noonan, J. P. (2015). The autism-associated chromatin modifier CHD8 regulates other autism risk genes during human neurodevelopment. *Nature Communications*, 6. <https://doi.org/10.1038/ncomms7404>
- De Rubeis, S., He, X., Goldberg, A. P., Poultney, C. S., Samocha, K., Cicek, A. E., Kou, Y., Liu, L., Fromer, M., Walker, S., Singh, T., Klei, L., Kosmicki, J., Fu, S. C., Aleksic, B., Biscaldi, M., Bolton, P. F., Brownfeld, J. M., Cai, J., ... Buxbaum, J. D. (2014). Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature*, 515(7526), 209–215.
<https://doi.org/10.1038/nature13772>
- Donovan, A. P. A., & Basson, M. A. (2017). The neuroanatomy of autism – a developmental perspective. In *Journal of Anatomy* (Vol. 230, Issue 1, pp. 4–15). Blackwell Publishing Ltd.
<https://doi.org/10.1111/joa.12542>
- Douzgou, S., Liang, H. W., Metcalfe, K., Somarathi, S., Tischkowitz, M., Mohamed, W., Kini, U., McKee, S., Yates, L., Bertoli, M., Lynch, S. A., Holder, S., & Banka, S. (2019). The clinical presentation caused by truncating CHD8 variants. *Clinical Genetics*, 96(1), 72–84.
<https://doi.org/10.1111/cge.13554>
- Durak, O., Gao, F., Kaeser-Woo, Y. J., Rueda, R., Martorell, A. J., Nott, A., Liu, C. Y., Watson, L. A., & Tsai, L. H. (2016). Chd8 mediates cortical neurogenesis via transcriptional regulation of cell cycle and Wnt signaling. *Nature Neuroscience*, 19(11), 1477–1488.
<https://doi.org/10.1038/nn.4400>
- Franklin, K. (2013). *Paxinos and Franklin's The mouse brain in stereotaxic coordinates* (Fourth edition.). Academic Press an imprint of Elsevier.
- García-Cabezas, M., John, Y. J., Barbas, H., & Zikopoulos, B. (2016). Distinction of neurons, glia and endothelial cells in the cerebral cortex: An algorithm based on cytological features. *Frontiers in Neuroanatomy*, 10(NOV). <https://doi.org/10.3389/fnana.2016.00107>
- Geschwind, D. H. (2009). Advances in autism. In *Annual Review of Medicine* (Vol. 60, pp. 367–380). Annu Rev Med. <https://doi.org/10.1146/annurev.med.60.053107.121225>
- Gilissen, C., Hehir-Kwa, J. Y., Thung, D. T., Van De Vorst, M., Van Bon, B. W. M., Willemsen, M. H., Kwint, M., Janssen, I. M., Hoischen, A., Schenck, A., Leach, R., Klein, R., Tearle, R., Bo, T., Pfundt, R., Yntema, H. G., De Vries, B. B. A., Kleefstra, T., Brunner, H. G., ... Veltman, J. A. (2014). Genome sequencing identifies major causes of severe intellectual disability. *Nature*, 511(7509), 344–347. <https://doi.org/10.1038/nature13394>
- Gompers, A. L., Su-Feher, L., Ellegood, J., Copping, N. A., Riyadh, M. A., Stradleigh, T. W.,

- Pride, M. C., Schaffler, M. D., Wade, A. A., Catta-Preta, R., Zdilar, I., Louis, S., Kaushik, G., Mannion, B. J., Plajzer-Frick, I., Afzal, V., Visel, A., Pennacchio, L. A., Dickel, D. E., ... Nord, A. S. (2017). Germline Chd8 haploinsufficiency alters brain development in mouse. *Nature Neuroscience*, 20(8), 1062–1073. <https://doi.org/10.1038/nn.4592>
- Gundersen, H. J. G. (1986). Stereology of arbitrary particles. In *Journal of Microscopy* (Vol. 143, Issue 1, pp. 3–45). J Microsc. <https://doi.org/10.1111/j.1365-2818.1986.tb02764.x>
- Gundersen, H. J. G. (1988). The nucleator. *Journal of Microscopy*, 151(1), 3–21. <https://doi.org/10.1111/j.1365-2818.1988.tb04609.x>
- Gundersen, H. J. G., & Jensen, E. B. (1987). The efficiency of systematic sampling in stereology and its prediction. *Journal of Microscopy*, 147(3), 229–263. <https://doi.org/10.1111/j.1365-2818.1987.tb02837.x>
- Haida, O., Al Sagheer, T., Balbous, A., Francheteau, M., Matas, E., Soria, F., Fernagut, P. O., & Jaber, M. (2019). Sex-dependent behavioral deficits and neuropathology in a maternal immune activation model of autism. *Translational Psychiatry*, 9(1). <https://doi.org/10.1038/s41398-019-0457-y>
- Hilgetag, C. C., Dell, L.-A., Jedlicka, P., Keller, D., Erö, C., & Markram, H. (2018). Cell Densities in the Mouse Brain: A Systematic Review. *Frontiers in Neuroanatomy* / *Www.Frontiersin.Org*, 12, 83. <https://doi.org/10.3389/fnana.2018.00083>
- Iossifov, I., O’Roak, B. J., Sanders, S. J., Ronemus, M., Krumm, N., Levy, D., Stessman, H. A., Witherspoon, K. T., Vives, L., Patterson, K. E., Smith, J. D., Paepier, B., Nickerson, D. A., Dea, J., Dong, S., Gonzalez, L. E., Mandell, J. D., Mane, S. M., Murtha, M. T., ... Wigler, M. (2014). The contribution of de novo coding mutations to autism spectrum disorder. *Nature*, 515(7526), 216–221. <https://doi.org/10.1038/nature13908>
- Jung, H., Park, H., Choi, Y., Kang, H., Lee, E., Kweon, H., Roh, J. D., Ellegood, J., Choi, W., Kang, J., Rhim, I., Choi, S. Y., Bae, M., Kim, S. G., Lee, J., Chung, C., Yoo, T., Park, H., Kim, Y., ... Kim, E. (2018). Sexually dimorphic behavior, neuronal activity, and gene expression in Chd8-mutant mice. *Nature Neuroscience*, 21(9), 1218–1228. <https://doi.org/10.1038/s41593-018-0208-z>
- Katayama, Y., Nishiyama, M., Shoji, H., Ohkawa, Y., Kawamura, A., Sato, T., Suyama, M., Takumi, T., Miyakawa, T., & Nakayama, K. I. (2016). CHD8 haploinsufficiency results in autistic-like phenotypes in mice. *Nature*, 537(7622), 675–679. <https://doi.org/10.1038/nature19357>
- Kipp, M., Kiessling, M. C., Hochstrasser, T., Roggenkamp, C., & Schmitz, C. (2017). Design-Based Stereology for Evaluation of Histological Parameters. *Journal of Molecular Neuroscience*, 61(3), 325–342. <https://doi.org/10.1007/s12031-016-0858-9>
- Lauber, E., Filice, F., & Schwaller, B. (2018). Dysregulation of Parvalbumin Expression in the Cntnap2^{-/-} Mouse Model of Autism Spectrum Disorder. *Frontiers in Molecular Neuroscience*, 11, 262. <https://doi.org/10.3389/fnmol.2018.00262>
- Lavenex, P., Lavenex, P. B., Bennett, J. L., & Amaral, D. G. (2009). Postmortem changes in the neuroanatomical characteristics of the primate brain: Hippocampal formation. *Journal of*

- Comparative Neurology*, 512(1), 27–51. <https://doi.org/10.1002/cne.21906>
- Lavenex, P., Steele, M. A., & Jacobs, L. F. (2000). The seasonal pattern of cell proliferation and neuron number in the dentate gyrus of wild adult eastern grey squirrels. *European Journal of Neuroscience*, 12(2), 643–648. <https://doi.org/10.1046/j.1460-9568.2000.00949.x>
- Li, J., Wang, L., Guo, H., Shi, L., Zhang, K., Tang, M., Hu, S., Dong, S., Liu, Y., Wang, T., Yu, P., He, X., Hu, Z., Zhao, J., Liu, C., Sun, Z. S., & Xia, K. (2017). Targeted sequencing and functional analysis reveal brain-size-related genes and their networks in autism spectrum disorders. *Molecular Psychiatry*, 22(9), 1282–1290. <https://doi.org/10.1038/mp.2017.140>
- Libero, L. E., Nordahl, C. W., Li, D. D., Ferrer, E., Rogers, S. J., & Amaral, D. G. (2016). Persistence of megalencephaly in a subgroup of young boys with autism spectrum disorder. *Autism Research*, 9(11), 1169–1182. <https://doi.org/10.1002/aur.1643>
- Long, J. M., Kalehua, A. N., Muth, N. J., Hengemihle, J. M., Jucker, M., Calhoun, M. E., Ingram, D. K., & Mouton, P. R. (1998). Stereological estimation of total microglia number in mouse hippocampus. *Journal of Neuroscience Methods*, 84(1–2), 101–108. [https://doi.org/10.1016/S0165-0270\(98\)00100-9](https://doi.org/10.1016/S0165-0270(98)00100-9)
- Marie, C., Clavairol, A., Frah, M., Hmidan, H., Yan, J., Zhao, C., Van Steenwinckel, J., Daveau, R., Zalc, B., Hassan, B., Thomas, J. L., Gressens, P., Ravassard, P., Moszer, I., Martin, D. M., Richard Lu, Q., & Parras, C. (2018). Oligodendrocyte precursor survival and differentiation requires chromatin remodeling by Chd7 and Chd8. *Proceedings of the National Academy of Sciences of the United States of America*, 115(35), E8246–E8255. <https://doi.org/10.1073/pnas.1802620115>
- Matta, S. M., Hill-Yardin, E. L., & Crack, P. J. (2019). The influence of neuroinflammation in Autism Spectrum Disorder. In *Brain, Behavior, and Immunity* (Vol. 79, pp. 75–90). Academic Press Inc. <https://doi.org/10.1016/j.bbi.2019.04.037>
- Naruse, I., & Keino, H. (1995). Apoptosis in the developing CNS. In *Progress in Neurobiology* (Vol. 47, Issue 2). Prog Neurobiol. [https://doi.org/10.1016/0301-0082\(95\)00024-P](https://doi.org/10.1016/0301-0082(95)00024-P)
- Nishiyama, M., Nakayama, K., Tsunematsu, R., Tsukiyama, T., Kikuchi, A., & Nakayama, K. I. (2004). Early Embryonic Death in Mice Lacking the β -Catenin-Binding Protein Duplin. *Molecular and Cellular Biology*, 24(19), 8386–8394. <https://doi.org/10.1128/mcb.24.19.8386-8394.2004>
- Nordahl, C. W., Lange, N., Li, D. D., Barnett, L. A., Lee, A., Buonocore, M. H., Simon, T. J., Rogers, S., Ozonoff, S., & Amaral, D. G. (2011). Brain enlargement is associated with regression in preschool-age boys with autism spectrum disorders. *Proceedings of the National Academy of Sciences of the United States of America*, 108(50), 20195–20200. <https://doi.org/10.1073/pnas.1107560108>
- O’Roak, B. J., Vives, L., Fu, W., Egerton, J. D., Stanaway, I. B., Phelps, I. G., Carvill, G., Kumar, A., Lee, C., Ankenman, K., Munson, J., Hiatt, J. B., Turner, E. H., Levy, R., O’Day, D. R., Krumm, N., Coe, B. P., Martin, B. K., Borenstein, E., ... Shendure, J. (2012). Multiplex targeted sequencing identifies recurrently mutated genes in autism spectrum disorders. *Science*, 338(6114), 1619–1622. <https://doi.org/10.1126/science.1227764>

- O'Roak, B. J., Vives, L., Girirajan, S., Karakoc, E., Krumm, N., Coe, B. P., Levy, R., Ko, A., Lee, C., Smith, J. D., Turner, E. H., Stanaway, I. B., Vernot, B., Malig, M., Baker, C., Akey, J. M., Borenstein, E., Rieder, M. J., Nickerson, D. A., ... Eichler, E. E. (2012). Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations. *Nature*, 485(7397), 246–250. <https://doi.org/10.1038/nature10989>
- Platt, R. J., Zhou, Y., Slaymaker, I. M., Shetty, A. S., Weisbach, N. R., Kim, J. A., Sharma, J., Desai, M., Sood, S., Kempton, H. R., Crabtree, G. R., Feng, G., & Zhang, F. (2017a). Chd8 Mutation Leads to Autistic-like Behaviors and Impaired Striatal Circuits. *Cell Reports*, 19(2), 335–350. <https://doi.org/10.1016/j.celrep.2017.03.052>
- Platt, R. J., Zhou, Y., Slaymaker, I. M., Shetty, A. S., Weisbach, N. R., Kim, J. A., Sharma, J., Desai, M., Sood, S., Kempton, H. R., Crabtree, G. R., Feng, G., & Zhang, F. (2017b). Chd8 Mutation Leads to Autistic-like Behaviors and Impaired Striatal Circuits. *Cell Reports*, 19(2), 335–350. <https://doi.org/10.1016/j.celrep.2017.03.052>
- Sanders, S. J., He, X., Willsey, A. J., Ercan-Sencicek, A. G., Samocha, K. E., Cicek, A. E., Murtha, M. T., Bal, V. H., Bishop, S. L., Dong, S., Goldberg, A. P., Jinlu, C., Keaney, J. F., Klei, L., Mandell, J. D., Moreno-De-Luca, D., Poultney, C. S., Robinson, E. B., Smith, L., ... State, M. W. (2015). Insights into Autism Spectrum Disorder Genomic Architecture and Biology from 71 Risk Loci. *Neuron*, 87(6), 1215–1233. <https://doi.org/10.1016/j.neuron.2015.09.016>
- Satterstrom, F. K., Kosmicki, J. A., Wang, J., Breen, M. S., De Rubeis, S., An, J. Y., Peng, M., Collins, R., Grove, J., Klei, L., Stevens, C., Reichert, J., Mulhern, M. S., Artomov, M., Gerges, S., Sheppard, B., Xu, X., Bhaduri, A., Norman, U., ... Buxbaum, J. D. (2020). Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiology of Autism. *Cell*, 180(3), 568-584.e23. <https://doi.org/10.1016/j.cell.2019.12.036>
- Schumann, C. M., & Nordahl, C. W. (2011). Bridging the gap between MRI and postmortem research in autism. In *Brain Research* (Vol. 1380, pp. 175–186). NIH Public Access. <https://doi.org/10.1016/j.brainres.2010.09.061>
- Spear, L. P. (2000). The adolescent brain and age-related behavioral manifestations. *Neuroscience and Biobehavioral Reviews*, 24(4), 417–463. [https://doi.org/10.1016/S0149-7634\(00\)00014-2](https://doi.org/10.1016/S0149-7634(00)00014-2)
- Suetterlin, P., Hurley, S., Mohan, C., Riegman, K. L. H., Pagani, M., Caruso, A., Ellegood, J., Galbusera, A., Crespo-Enriquez, I., Michetti, C., Yee, Y., Ellingford, R., Brock, O., Delogu, A., Francis-West, P., Lerch, J. P., Scattoni, M. L., Gozzi, A., Fernandes, C., & Basson, M. A. (2018). Altered neocortical gene expression, brain overgrowth and functional over-connectivity in chd8 haploinsufficient mice. *Cerebral Cortex*, 28(6), 2192–2206. <https://doi.org/10.1093/cercor/bhy058>
- Sugathan, A., Biagioli, M., Golzio, C., Erdin, S., Blumenthal, I., Manavalan, P., Ragavendran, A., Brand, H., Lucente, D., Miles, J., Sheridan, S. D., Stortchevoi, A., Kellis, M., Haggarty, S. J., Katsanis, N., Gusella, J. F., & Talkowski, M. E. (2014). CHD8 regulates neurodevelopmental pathways associated with autism spectrum disorder in neural progenitors. *Proceedings of the National Academy of Sciences of the United States of*

- America*, 111(42), E4468–E4477. <https://doi.org/10.1073/pnas.1405266111>
- Tatton-Brown, K., Loveday, C., Yost, S., Clarke, M., Ramsay, E., Zachariou, A., Elliott, A., Wylie, H., Ardisson, A., Rittinger, O., Stewart, F., Temple, I. K., Cole, T., Mahamdallie, S., Seal, S., Ruark, E., & Rahman, N. (2017). Mutations in Epigenetic Regulation Genes Are a Major Cause of Overgrowth with Intellectual Disability. *American Journal of Human Genetics*, 100(5), 725–736. <https://doi.org/10.1016/j.ajhg.2017.03.010>
- Vissers, L. E. L. M., Gilissen, C., & Veltman, J. A. (2016). Genetic studies in intellectual disability and related disorders. In *Nature Reviews Genetics* (Vol. 17, Issue 1, pp. 9–18). Nature Publishing Group. <https://doi.org/10.1038/nrg3999>
- Vogel-Ciernia, A., & Wood, M. A. (2014). Neuron-specific chromatin remodeling: A missing link in epigenetic mechanisms underlying synaptic plasticity, memory, and intellectual disability disorders. In *Neuropharmacology* (Vol. 80, pp. 18–27). Elsevier Ltd. <https://doi.org/10.1016/j.neuropharm.2013.10.002>
- Wang, P., Mokhtari, R., Pedrosa, E., Kirschenbaum, M., Bayrak, C., Zheng, D., & Lachman, H. M. (2017). CRISPR/Cas9-mediated heterozygous knockout of the autism gene CHD8 and characterization of its transcriptional networks in cerebral organoids derived from iPS cells. *Molecular Autism*, 8(1). <https://doi.org/10.1186/s13229-017-0124-1>
- Wang, T., Guo, H., Xiong, B., Stessman, H. A. F., Wu, H., Coe, B. P., Turner, T. N., Liu, Y., Zhao, W., Hoekzema, K., Vives, L., Xia, L., Tang, M., Ou, J., Chen, B., Shen, Y., Xun, G., Long, M., Lin, J., ... Eichler, E. E. (2016). De novo genic mutations among a Chinese autism spectrum disorder cohort. *Nature Communications*, 7. <https://doi.org/10.1038/ncomms13316>
- West, M. J., Slomianka, L., & Gundersen, H. J. G. (1991). Unbiased stereological estimation of the total number of neurons in the subdivisions of the rat hippocampus using the optical fractionator. *The Anatomical Record*, 231(4), 482–497. <https://doi.org/10.1002/ar.1092310411>
- West, Mark J., & Gundersen, H. J. G. (1990). Unbiased stereological estimation of the number of neurons in the human hippocampus. *Journal of Comparative Neurology*, 296(1), 1–22. <https://doi.org/10.1002/cne.902960102>
- Wickham, H. (2009). ggplot2, Elegant Graphics for Data Analysis. In *ggplot2*. Springer New York. <https://doi.org/10.1007/978-0-387-98141-3>
- Winden, K. D., Yuskaitis, C. J., & Poduri, A. (2015). Megalencephaly and macrocephaly. *Seminars in Neurology*, 35(3), 277–287. <https://doi.org/10.1055/s-0035-1552622>
- Woodbury-Smith, M., Deneault, E., Yuen, R. K. C., Walker, S., Zarrei, M., Pellicchia, G., Howe, J. L., Hoang, N., Uddin, M., Marshall, C. R., Chrysler, C., Thompson, A., Szatmari, P., & Scherer, S. W. (2017). Mutations in RAB39B in individuals with intellectual disability, autism spectrum disorder, and macrocephaly. *Molecular Autism*, 8(1). <https://doi.org/10.1186/s13229-017-0175-3>
- Wu, H., Li, H., Bai, T., Han, L., Ou, J., Xun, G., Zhang, Y., Wang, Y., Duan, G., Zhao, N., Chen, B., Du, X., Yao, M., Zou, X., Zhao, J., Hu, Z., Eichler, E. E., Guo, H., & Xia, K.

(2020). Phenotype-to-genotype approach reveals head-circumference-associated genes in an autism spectrum disorder cohort. *Clinical Genetics*, 97(2), 338–346.
<https://doi.org/10.1111/cge.13665>

Xu, Q., Liu, Y. Y., Wang, X., Tan, G. H., Li, H. P., Hulbert, S. W., Li, C. Y., Hu, C. C., Xiong, Z. Q., Xu, X., & Jiang, Y. H. (2018). Autism-associated CHD8 deficiency impairs axon development and migration of cortical neurons. *Molecular Autism*, 9(1).
<https://doi.org/10.1186/s13229-018-0244-2>

Zhao, C., Dong, C., Frah, M., Deng, Y., Marie, C., Zhang, F., Xu, L., Ma, Z., Dong, X., Lin, Y., Koenig, S., Nait-Oumesmar, B., Martin, D. M., Wu, L. N., Xin, M., Zhou, W., Parras, C., & Lu, Q. R. (2018). Dual Requirement of CHD8 for Chromatin Landscape Establishment and Histone Methyltransferase Recruitment to Promote CNS Myelination and Repair. *Developmental Cell*, 45(6), 753-768.e8. <https://doi.org/10.1016/j.devcel.2018.05.022>

Figure legends:

Figure 1: Nissl-stained coronal sections through the mouse brain (arranged from rostral A, to caudal F) illustrating the portion of the neocortex that was evaluated in this study. Not all levels are shown and the analyzed neocortex extended to the rostral and caudal poles of the brain. Before the study was started, a template of cortical areas was designed that was based on identifiable neuroanatomical landmarks. The template was used to demarcate the extent of analyzed neocortex in all experimental animals. The analysis was carried out blinded to the experimental condition or sex of the animal. Calibration bar equals 1 mm.

Figure 2: Increased cortical volume without increased neuron number in heterozygous *Chd8* mutant mice. Stereological estimations of Cortical volume (a), Neuron count (b), and Neuron Soma Size (c) measured on Nissl-stained sections from *Chd8*^{+/del5} mice and wild type littermates. P values resulting from t-test comparisons of the effect of genotype and sex stratified dispersion plots are shown. All data expressed as mean ± SEM.

Figure 4

Figure 2

A

B

C