Original Research Article

Title: Moderate increases in channel discharge are positively related to ecosystem respiration in forested Ozark streams

Short title: Stream discharge is positively related to ecosystem respiration

Authors: Allyn K. Dodd¹, Daniel D. Magoulick², and Michelle A. Evans-White³

¹Division of Natural Sciences, Lyon College, Batesville, AR 72501

²U.S. Geological Survey, Arkansas Cooperative Fish and Wildlife Research Unit, Department of Biological Sciences, University of Arkansas, Fayetteville, AR 72701

³Department of Biological Sciences, University of Arkansas, Fayetteville, AR 72701

This draft manuscript is distributed solely for the purposes of scientific peer review. Its content is deliberative and predecisional, so it must not be disclosed or released by reviewers. Because the manuscript has not yet been approved for publication by the U.S. Geological Survey (USGS), it does not represent any official USGS findings or policy.

Corresponding author: Email: allyn.dodd@lyon.edu, Phone: 501-307-7216

ABSTRACT

The natural flow regime is considered the “master variable” in lotic systems, controlling structure and function at organismal, population, community, and ecosystem levels. We sought to estimate forested headwater stream metabolism across two dominant flow regimes (*Runoff* and *Groundwater*) in northern Arkansas and evaluate potential differences in, and drivers of, gross primary production, ecosystem respiration, and net ecosystem metabolism. Flow regimes differed in intermittency, substrate heterogeneity, hyporheic connectivity, and dominant water source (subsurface runoff vs. groundwater), which we expected to result in differences in primary production and respiration. Average daily gross primary production (GPP) and ecosystem respiration (ER) estimated from field data collected from May 2015-June 2016 tended to be greater in *Groundwater* streams. Respiration was positively related to discharge ($R^2 = 0.98$ $p < 0.0001$) and net metabolism became more heterotrophic with increasing average annual discharge across sites ($R^2 = 0.94$, $p = 0.0008$). Characterizing ecosystem-level responses to differences in flow can reveal mechanisms governing stream metabolism and, in turn, provide information regarding trophic state and energy inputs as efforts continue to determine global trends in aquatic carbon sources and fates.

INTRODUCTION

The natural flow regime exerts primacy over water quality and quantity, habitat structure, disturbance regime, and, in turn, ecological processes and functions in lotic systems. Flow regime is characterized by the timing, duration, magnitude, frequency, and rate of change of water flowing through a channel over various temporal scales (Poff et al. 1997), arranging habitat space and thereby creating a unique template for life history strategies and community interactions (Southwood 1977, Poff and Ward 1990). Natural disturbances, such as flooding and drought, serve as life cycle prompts for many fishes and macroinvertebrates, whose reproductive cues are intimately linked with predictable, seasonal changes in flow (Poff and Ward 1989, Huryn and Wallace 2000, Humphries and Baldwin 2003, Lytle and Poff 2004). Flow regime is ultimately a byproduct of landscape-level processes and variation, as climate, topography, geology, vegetation, and soils interact to determine primary water sources (e.g. groundwater vs. subsurface runoff), quantity of within-channel flow, and geomorphology. Indeed, flow regime is typically a region- and land cover-specific phenomenon; streams reflect the diverse biomes that generate and sustain their flows as well as the relative contributions of groundwater, surface water, soil water, and precipitation (Hynes 1975, Poff et al. 1997, Carlisle et al. 2010).

Stream metabolism is an estimator of carbon (C) dynamics and an indicator of of nutrient cycling and trophic status that is sensitive to natural and anthropogenic disturbances, revealing ecosystem-level responses to changes in hydrology and geomorphology. Metabolism is comprised of gross primary production (GPP) and ecosystem respiration (ER), which yields the net amount of carbon fixed into biomass, or net ecosystem metabolism (NEM) (Hall and Hotchkiss 2017). Metabolism can reveal whole-stream responses to landscape changes as well as predict potential bottom-up effects on higher trophic levels. Ecosystem metabolism is driven by

proximal factors such as light and nutrients, which are influenced by the surrounding watershed (Bernot et al. 2010, Yates et al. 2013). The direct and indirect susceptibility of primary production and respiration to landscape-level variation makes it a useful metric for assessing impacts at the ecosystem level. Additionally, daily metabolism can vary temporally due to changes in light levels, organic matter inputs, algal biomass, and hydrology (Acuña et al. 2004, Roberts et al. 2007). Previous work assessing annual metabolism across multiple streams has focused primarily on the effects of biome and land use (Bott et al. 1985, Mulholland 2001, Bernot et al. 2010). The large dependence of other metabolism estimates on flow timing and magnitude (e.g. Uehlinger et al. 2003, Roberts et al. 2007, Qasem et al. 2018) suggests they will vary significantly across differing flow regimes within the same biome. Further, comparing function within and among hydrologic classifications provides insight into processes and variables controlling ecosystem function. A growing body of work has begun to address how stream metabolic regimes vary in time and space at multiple hierarchical scales (Appling et al. 2018, Koenig et al. 2019, Savoy et al. 2019), and a subset of these efforts have focused specifically on hydrologic influences impacting production and respiration (Jones et al. 1995, Dodds et al. 1996, Battin 1999, Uehlinger 2000, Uehlinger et al. 2003, Vilches and Giorgi 2010, Leggieri et al. 2013, Cook et al. 2015, Rovelli et al. 2017, Reisinger et al. 2017, Demars et al. 2019, O'Donnell and Hotchkiss 2019).

Existing conceptual models of headwater stream metabolism have posited that factors controlling metabolism differ by biome (Mulholland et al. 2001), land use category (Bernot et al. 2010), and season (Roberts et al. 2007). In reference systems, biome and season are considered the primary drivers of differences across streams (Bott et al. 1985, Mulholland et al. 2001, Hornbach et al. 2015). However, others have shown distinct hydroecological regions at

hierarchical spatial scales characterized by significant variation in flow dynamics within a biome (Poff et al. 2006, Leasure et al. 2016). This variation arises from changes in geology and water sources across basins and sub-basins, which can result in differences in ecosystem function and carbon availability (Thoms and Parsons 2002).

Flow variability within a stream can be a determinant of annual metabolism, as flow extremes can exert a strong influence on organic matter movement through the system (Acuña et al. 2004, Roberts et al. 2007, Demars 2019). High flows can depress primary production (Uehlinger 2000, Uehlinger 2006) while increasing production rates in autumn by removing abscised leaves covering the benthos (Argerich et al. 2011). High flows can also influence respiration rates by reducing respiration initially due to loss of autotrophic biomass, then increasing rates as the autotrophic community recovers from scouring (Roberts et al. 2007, Izagirre et al. 2008). Consistently higher discharge, or higher discharge in one year compared to another, can depress primary production rates by preventing regrowth of algal biomass, while others have found clear relationships between hydrologic regime and benthic organic matter that supports respiration (Acuña et al. 2007, Demars 2019). Hot, dry summers that increase water temperature but reduce depth can support extensive algal production and may lead to an overall reduction in metabolic rates over summer in the absence of scouring floods (Izagirre et al. 2008).

Drying and flooding can both temporarily depress primary production and respiration, while the weeks following these disturbances are typically marked by high rates of production and respiration as algae recolonize the benthos (Uehlinger 2000, Uehlinger 2006). Specifically, the number of dry days, number of days experiencing high flows (defined as >75% average daily flow), and number of flood events affect production and respiration; the strength of this effect

would be dependent upon the magnitude, frequency, and duration of the disturbance (Biggs et al. 2005, Palmer and Ruhi 2019).

Several natural flow categories have been identified for streams within the Ozark and Ouachita Interior Highlands in northern/western Arkansas, eastern Oklahoma, and southern Missouri (Leasure et al. 2016), but efforts to characterize these systems based on their unique hydrology in the field have only recently begun. Leasure et al. (2016) revealed distinct geographic areas demarcated by dominant flow types that are likely functionally unique. These flow classifications consist of subsurface runoff- and groundwater-fed systems, known as *Runoff Flashy* and *Groundwater Flashy* streams (hereafter *Runoff* and *Groundwater*), that are dominant in the Ozark and Boston Mountains ecoregions. Key differences between flow regimes are frequency and duration of low flow days, frequency of floods, channel substrate heterogeneity and size, and dominant water sources (e.g. groundwater versus subsurface runoff).

Both flow regimes experience flash floods marked by high magnitudes but short duration (Leasure et al. 2016). *Runoff* stream flow originates primarily from subsurface runoff originating from precipitation and tend to dry for several weeks during autumn. Mixing analysis has shown that, on average, 89 (± 6)% of *Runoff* base flows derive from precipitation in the form of interflow. *Groundwater* streams tend to flow perennially, with 79 (± 3)% of base flow coming from groundwater contributions (Dodd et al. 2020) coming through the hyporheic zone. *Groundwater* streams tend to exhibit less variable annual flows (Leasure et al. 2016).

Additionally, channel geomorphology and dominant substrate are markedly different between flow regimes. *Groundwater* streams are dominated by a mixture of substrate types, dominated by sand, gravel, and pebble with some cobble and boulder mixed in; substantial bed movement is common during high flow events. Conversely, *Runoff* streambeds are often

comprised of bedrock overlain in some areas by cobble and boulders. Substrate heterogeneity can alter near-bed flow and turbulence intensity, and in turn influence production rates (Cardinale et al. 2002), and a greater diversity of benthic substrate sizes may foster greater production in *Groundwater* streams. Our primary objective was to determine whether differences exist in stream gross primary production, ecosystem respiration, and net ecosystem production between *Runoff* and *Groundwater* streams and characterize natural variation in ecological-flow responses across seasons. We expected average GPP to be greater in *Groundwater* streams, as these streams tend to exhibit perennial flow, have relatively stable hydrology over the year, and exhibit low turbidity. We expected ER to be greater in *Groundwater* systems as well given that the heterogeneous benthic substrate may result in higher rates of respiration by providing spaces for organic matter to settle and pockets in the benthic and hyporheic zones for heterotrophic microbes to colonize.

We also sought to confirm that differences exist in stream discharge and flow metrics. We hypothesized that *Groundwater* streams would have greater annual discharge since these streams do not dry and have more sustained, less variable flows from groundwater.

We also examined potential drivers of metabolism across sites. We predicted positive relationships between GPP and light, nutrients (total N and total P), algal biomass, and a negative relationship between GPP and discharge, number of high flow days, number of floods, number of low flow days, and number of no flow days. We hypothesized that positive relationships would exist between ER and biofilm ash-free dry mass, nutrients, floods, high flow days, and discharge (Figure 1).

Figure 1. Conceptual diagram illustrating factors controlling stream metabolism, which incorporates the natural flow regime of an area as influenced by biome, season, and land use. This, in turn, can influence hydrologic responses to land use change. GPP and ER denote gross primary production and ecosystem respiration, respectively.

METHODS

STUDY SITES

This study was conducted in six temperate, minimally impacted headwater streams in the deciduous forests of Arkansas (Figure 2). We chose three *Groundwater* streams and three *Runoff* streams categorized based on flow classifications modeled by Leasure et al. (2016). Study sites were chosen based on membership in *Runoff Flashy* or *Groundwater Flashy* flow classes as well as stream surface area, surrounding land cover, and presence of downstream USGS stream gauges. Streams selected for the study were all headwater systems with forested land cover ranging from 84 to 95% of total watershed area (Stroud Water Research Center 2017) (Table 1).

Table 1. Site abbreviations and flow metrics for May 2015 through June 2016. Discharge estimates reflect year-round measurements calculated from daily gauge data (Gauge) or from discharge measured monthly in the field, which were taken at or near base flow (Field).

Flow Classification	Site	Abbreviation	%Forest	Watershed Area (km ²)	Mean Discharge	Mean Discharge	No Flow Days	High Flow Days	Floods	Annual Rain (cm)
					(Gauge) (m ³ s ⁻¹)	(Field) (m ³ s ⁻¹)				
Runoff	Big Piney	BPC	92.23	87	3.65	0.41	33	146	11	154
Runoff	Little Piney	LPC	94.73	86	1.49	0.90	47	106	11	141
Runoff	Murray	MRY	94.04	65	1.16	0.42	0	121	11	141
Groundwater	Roasting Ear	REC	83.97	91	1.87	1.21	9	107	12	151
Groundwater	Spring	SPR	85.28	101	1.54	0.48	0	137	13	137
Groundwater	Sylamore	SYL	92.94	81	5.54	1.44	0	107	15	151

Figure 2. Map of flow regimes in the Ozark and Ouachita Interior Highlands based on Leasure et al. (2016). Highlighted area shows individual study sites sampled from 2015-2016 across northern Arkansas. Teal lines represent *Groundwater* streams. Light green lines represent *Runoff* streams.

All streams are located within the Ozark National Forest in northern Arkansas and were chosen based on the amount of surrounding forest in the watershed and a lack of tributaries or

ephemeral drains feeding into accessible reaches to prevent extraneous variables influencing GPP and ER. Vegetation surrounding both stream types was primarily oak and hickory trees forest (Woods et al. 2004). While all streams had similar surrounding landcover and vegetation, the geology underlying *Runoff* and *Groundwater* streams differ. *Groundwater* streams are found within a highly dissected limestone plateau. Conversely, the *Runoff* streams are found in an area dominated by green/gray shale and sandstone. Vegetation surrounding both stream types was primarily oak and hickory trees forest (Woods et al. 2004, Chapman et al. 2006, Stephenson et al. 2007).

METABOLISM RATES

We calculated reach-scale metabolism using the open-channel single-station method (Odum 1956, Riley and Dodds 2012). Dissolved oxygen (DO) and temperature were measured every 15 minutes by Hydrolab DS5X multiparameter sondes (Hach Company, Loveland, CO) from May 2015 to June 2016 in a well-mixed area at the bottom of each study reach. Data were corrected as necessary by comparing with DO concentrations determined via Winkler titrations (Dodds et al. 2018). Reaeration coefficients as estimates of air-water gas exchange were determined via propane release in five out of six streams, while nighttime regression was utilized in one *Runoff* stream, Murray Creek (Hall and Hotchkiss 2017). Propane release was necessary in five streams because nighttime regressions yielded significant relationships between ER and air-water gas exchange, or reaeration, coefficients (K_{600}), whereas no such relationship was present at Murray Creek. Corrections for groundwater contributions to reaches receiving appreciable inputs were made according to Hall and Tank (2005). Photosynthetically-active radiation (PAR) measurements were logged concurrently with metabolism parameters using an Odyssey light meter positioned in an area near the stream with open canopy. Stream metabolism was estimated based

on diel changes in DO, temperature, depth, and light measurements; we used R package StreamMetabolizer (Appling et al. 2018) to solve for GPP and ER utilizing a single-station metabolism maximum likelihood model with the core equation:

$$O_2(t) = O_2(t - \Delta t) + \left(\frac{GPP_{Total}}{\underline{z}} \cdot \frac{PAR(t)}{\int_{u=t_0}^{t_1} PAR} \right) + \left(\frac{ER_{Total}}{\underline{z}} \times \Delta t \right) + K(t)(O_{2sat}(t) - O_2(t))\Delta t$$

where t is time and Δt is the time step between measurements (15 minutes), $\underline{z}$ is mean reach depth, $\int_{u=t_0}^{t_1} PAR$ is daily photosynthetically-active radiation, and $K_{(t)}$ is air-water gas exchange corrected for temperature. Estimates of GPP_{Total} and ER_{Total} fitted by StreamMetabolizer yielded daily rates for every day that a sonde was deployed at each stream (from 158 to 215 days). Data were not collected every day of the 422-day study due to flash floods and/or unwadeable conditions, drying, and/or equipment failure. Seasonal GPP and ER were calculated by averaging daily rates from the beginning of the respective season (e.g. equinox or solstice). Overall average daily rates were calculated by averaging daily rates over the study period. Summing daily rates by season would hinder comparisons since we did not have equal numbers of sampling dates across streams, so averages of daily rates over each season are reported. Average daily rates for each flow class were computed by computing the mean of the three streams in that flow class' overall average daily rates.

PHYSICOCHEMICAL VARIABLES

To evaluate relationships between metabolism and flow metrics, we quantified high flow days, number of floods, and number of days with no flow. High flow days were defined as exceeding the 75th percentile of mean annual discharge at a site. Floods were quantified as distinct hydropeaks greater than 100% of mean annual flow calculated from gauge data and upstream-downstream discharge relationships between USGS gauges and discharge measured in each study reach. We measured discharge in each reach monthly from May 2015 to June 2016

using velocity-area gauging (Gore 2006). We measured median diameter (d₅₀) of randomly selected stones across each discharge transect. One stone was randomly selected at each meter across every discharge transect during four spring and summer 2016 monthly sampling events.

Persulfate digests of unfiltered water samples taken at base or near-base flow during monthly sampling events were followed by colorimetric analyses to determine nutrient concentrations. Total nitrogen (TN) was measured monthly by automated cadmium reduction on a Lachat Quikchem 8500 (Hach Company, Loveland, CO). Total phosphorus (TP) was measured monthly using the ascorbic acid method (APHA 2005).

Canopy cover was determined for each stream channel once in summer and once following abscission using a densiometer to calculate percent coverage.

PERIPHYTON

For algal biomass, we collected six cobbles per reach at six equidistant transects down the stream reach. Monthly algal biomass was determined beginning in summer 2015 by scrubbing rocks of algae by extracting chlorophyll *a* each from algal slurry in ethanol, then wrapping with aluminum foil to determine rock area from mass-area relationships (Steinman et al. 2006).

STATISTICS

We used 2-way repeated-measures ANOVA to determine whether differences existed in metabolism, flow metrics, and periphyton biomass and biofilm ash-free dry mass between flow regimes by season. Linear regressions were employed to examine relationships between daily metabolism variables (GPP, ER, and NEP), physicochemical parameters (e.g. total nitrogen and phosphorus, temperature), biological metrics (e.g. chlorophyll *a*, biofilm ash-free dry mass), and flow metrics (e.g. gauge-calculated discharge, discharge measured by hand in each reach, number of low flow days, number of high flow days, coefficient of variation of daily flow,

number of floods). We evaluated relationships between variables with both gauge-calculated discharge, which yielded estimates of flow for all days of the study including extreme floods and times during which sondes were not deployed, and discharge measured in the reach monthly, which represented conditions under which sondes were deployed and did not include extreme high flows (Table 1). All statistical analyses were performed in R version 3.4.3. Statistical significance threshold was $p \leq 0.05$.

RESULTS

METABOLISM

There was no significant effect of flow regime on average daily GPP across seasons (Table 2), though *Groundwater* streams tended to show greater GPP all year except during winter (Figure 3). Average daily GPP was highest in summer 2016 in *Groundwater* streams, while *Runoff* streams exhibited the greatest average daily GPP in summer 2015. Total GPP ranged from 99.7 to 435.1 g O₂ m⁻² y⁻¹ in *Runoff* streams (days of record= 158-188 days), and 289.5 to 405.4 g O₂ m⁻² y⁻¹ in *Groundwater* streams (days of record= 159-202 days). Big Piney and Murray Creek, two *Runoff* streams, had fairly constant rates of GPP over the study year, with very small increases in GPP during the spring and summer. Conversely, the third *Runoff* site, Little Piney, showed marked increases in production during the spring and summer months, with daily production remaining high until the stream dried in October. Similarly high daily production rates were found at Sylamore, a *Groundwater* stream, throughout the spring and summer, though production decreased from July to September. The other two *Groundwater* sites exhibited spring and summer production rates like Big Piney and Murray, both *Runoff* streams. However, *Groundwater* stream production appeared to be stimulated by floods that occurred at the end of 2015, while *Runoff* streams showed little to no such response. Additionally, all

285 *Groundwater* streams and one *Runoff* (Little Piney) revealed greater primary production in
286 summer 2016 compared to the previous year.

287 Table 2. Results of repeated measures Analysis of Variance (ANOVA) for variables of interest. *

288 denote statistical significance.

Dependent Variable (2-way RM-ANOVA)	F-statistic	p- value
GPP		
Flow	$F_{(1,12)} = 1.06$	0.32
Season	$F_{(3,12)} = 0.96$	0.31
Flow x Season	$F_{(5,12)} = 1.26$	0.48
ER		
Flow	$F_{(1,12)} = 0.08$	0.79
Season	$F_{(3,12)} = 0.18$	0.91
Flow x Season	$F_{(5,12)} = 1.05$	0.39
Discharge		
Flow	$F_{(1,12)} = 2.15$	0.17
Season	$F_{(3,12)} = 2.86$	0.04*
Flow x Season	$F_{(5,12)} = 1.41$	0.29
Total Nitrogen		
Flow	$F_{(1,12)} = 0.12$	0.75
Season	$F_{(3,12)} = 2.35$	0.17
Flow x Season	$F_{(5,12)} = 2.54$	0.21
Total Phosphorus		
Flow	$F_{(1,12)} = 0.04$	0.85
Season	$F_{(3,12)} = 2.35$	0.17
Flow x Season	$F_{(5,12)} = 0.30$	0.61
Algal Biomass		
Flow	$F_{(1,12)} = 1.39$	0.27
Season	$F_{(3,12)} = 8.34$	0.0004*
Flow x Season	$F_{(5,12)} = 2.78$	0.09
Ash-Free Dry Mass		
Flow	$F_{(1,12)} = 18.56$	0.0005*
Season	$F_{(3,12)} = 6.60$	0.004*
Flow x Season	$F_{(5,12)} = 2.21$	0.13

289

290

291

Figure 3. Seasonal average daily gross primary production (A) and respiration (B) in *Runoff* (white) and *Groundwater* (gray) streams. Error bars denote ± 1 standard error. $n=3$ per flow regime.

We found no difference in average daily ER between flow regimes across seasons, though ER also tended to be greater in *Groundwater* streams. Average daily ER was greatest in spring 2016 across streams. Average daily GPP in *Runoff* streams was $1.2 (\pm 0.54)$ g O₂ m⁻² d⁻¹, while *Groundwater* streams averaged $1.9 (\pm 0.27)$ g O₂ m⁻² d⁻¹. Daily ER in *Runoff* streams averaged $-2.1 (\pm 0.99)$ g O₂ m⁻² d⁻¹ and $-5.2 (\pm 1.0)$ g O₂ m⁻² d⁻¹ in *Groundwater* streams. Daily rates of GPP and ER for each stream are shown in Figure 4. Total ER in *Runoff* streams ranged from -151.4 to -771.2 g O₂ m⁻² y⁻¹ and from -280.2 to -1340.3 g O₂ m⁻² y⁻¹ in *Groundwater* streams. Peaks in respiration rates over the year varied across all six sites. While Big Piney and Murray (*Runoff* streams) had low respiration rates (similar to GPP), trends in respiration in Little Piney were similar to those observed in *Groundwater* streams. Respiration was especially high in Little Piney and *Groundwater* streams during the summer of 2015 and spring 2016, though *Groundwater* streams overall showed greater rates over the year and elevated respiration extended well into the autumn of 2015. Respiration was also stimulated by the heavy flooding that occurred in December 2015. In *Groundwater* streams, respiration was greater in summer 2016 than the previous year, while respiration- though low overall in comparison- was greater in summer 2015 than 2016 in *Runoff* streams.

Figure 4. Daily rates of gross primary production (GPP) (black) and ecosystem respiration (ER) (gray) in Big Piney, Little Piney, Murray, Roasting Ear, Spring, and Sylamore from May 2015 to June 2016. *Runoff* streams are shown in panels on the left, *Groundwater* streams are represented in panels on the right. n= number of days of record at each site. “Flood” denotes large floods that prevented data collection.

PHYSICOCHEMICAL VARIABLES

Stream discharge did not differ significantly across flow regimes but differed by season ($F_{(5, 12)} = 2.86$, $p = 0.04$) (Table 2), with discharge peaking in the spring following substantial heavy winter rains that resulted in extreme flash floods in four out of six sites (Figure 5). Discharge tended to be greater in *Groundwater* streams over the study period; however, these systems showed substantially elevated flows compared to *Runoff* streams during the spring across both 2015 and 2016. Between March and June 2016, *Groundwater* streams experienced one to two more high-flow events than *Runoff* streams. *Runoff* streams experienced two moderate storm events during summer 2015, while *Groundwater* streams experienced three smaller events during the same period. Little to no rain fell across Arkansas from early August to November 7, 2015, causing two *Runoff* streams and one *Groundwater* stream to dry. However, *Runoff* streams that dried experienced longer drought periods, as Little Piney and Big Piney dried for 47 and 33 days, respectively, whereas Roasting Ear (a *Groundwater* stream) dried for only 9 days. Precipitation inputs did not differ significantly across flow regimes ($p = 0.85$). Over the year, *Runoff* sites received 145 ± 4 cm of rainfall, while *Groundwater* sites received 146 ± 4 cm. Flow metrics at each site are listed in Table 1.

Figure 5. Hydrographs for study sites from May 2015 to June 2016. Note differences in y-axis scales; scales were not standardized to preserve details of individual site hydrology.

Greater discharge stimulated respiration ($R^2 = 0.98$, $p < 0.0001$) and drove streams to be more heterotrophic over the year ($R^2 = 0.94$, $p = 0.0008$) (Figure 6). Discharge is often related to watershed area and stream size, though we did not find a relationship between watershed area and discharge ($R^2 = 0.004$, $p = 0.89$). Cooler *Groundwater* streams had greater rates of primary production; GPP was negatively related to water temperature across sites ($R^2 = 0.63$, $p = 0.04$). We found no further relationships between GPP or ER and physicochemical or biological variables.

Figure 6. Ecosystem respiration (A) and net ecosystem metabolism (B) compared with mean annual discharge measured in the field across flow regimes. These discharge measurements were taken only when streams were wadeable and thus represent variation in discharge across excluding high flows.

Runoff stream substrate types were primarily bedrock and cobble, while *Groundwater* streams were dominated by pebbles with a variety of other substrate sizes present (i.e. sand, gravel, cobble, and some boulders). Median particle size (d_{50}) was 19 mm (range = 10- 82 mm) in *Runoff* streams and 13 mm (range = <1-110 mm) in *Groundwater* streams.

Nutrient concentrations were not significantly different across sites and seasons, though both TN and TP tended to be greater in *Groundwater* streams (Table 2). Total N averaged 0.10 ± 0.03 mg/L in *Runoff* streams and 0.56 ± 0.26 mg/L in *Groundwater* streams. Mean total P was 6.21 ± 0.63 μ g/L in *Runoff* streams and 8.70 ± 1.38 μ g/L in *Groundwater* streams.

We explored potential relationships between watershed land cover and nutrient concentrations to determine whether even forested systems could be experiencing impacts of surrounding anthropogenic activity. Instream phosphorus concentrations decreased as surrounding forested land cover increased within a watershed ($R^2 = 0.77$, $p = 0.01$). We observed

a similar negative trend with respect to TN concentrations and forested land cover, though this relationship was not statistically significant ($R^2 = 0.56$, $p = 0.09$).

PERIPHYTON AND ORGANIC MATTER

Algal biomass was not significantly different between flow regimes, though there was a seasonal effect ($F_{(5, 12)} = 8.33$, $p = 0.0004$) (Table 2) and algal biomass tended to be greater in *Groundwater* streams throughout the year. Algal biomass peaked in the spring of 2016 in all streams. Groundwater stream algal biomass over the year was $3.4 \pm 1.4 \mu\text{g}/\text{cm}^2$ and Runoff stream algal biomass was less than half of that, averaging $1.5 \pm 0.1 \mu\text{g}/\text{cm}^2$. Organic matter measured as ash-free dry mass differed across seasons ($F_{(3, 12)} = 6.60$, $p = 0.004$) and was greater in *Groundwater* streams across all seasons except winter ($F_{(1, 12)} = 18.56$, $p = 0.0005$) (Figure 7). Similar to algal biomass, mean *Groundwater* stream biofilm ash-free dry mass was 11.7 ± 1.5 mg, while Runoff streams yielded 5.9 ± 0.5 mg.

Figure 7. Seasonal algal biomass (A) and biofilm slurry ash-free dry mass (B) in *Runoff* and *Groundwater* streams. Error bars denote ± 1 standard error. $n = 3$ per flow regime.

We evaluated whether instream nutrient concentrations could be driving differences between algal biomass and ash-free dry mass. Neither TN nor TP were related to algal biomass (TP: $R^2 = 0.47$, $p = 0.13$, TN: $R^2 = 0.38$, $p = 0.19$) or ash-free dry mass (TP: $R^2 = 0.10$, $p = 0.56$, TN: $R^2 = 0.39$, $p = 0.18$).

DISCUSSION

METABOLISM

Primary production tended to be greater in *Groundwater* streams during spring and summer 2016, possibly resulting from *Groundwater* streams' greater spring flows, slightly smaller, more mobile benthic substrate, and limestone karst rather than shale and sandstone that

characterize the geology of *Runoff* sites. Discharge was greater in *Groundwater* streams during the spring, resulting from heavy rains that fell at the end of 2015. *Groundwater* streams experienced floods of much greater magnitude, and experienced elevated flows for the duration of the spring that were not the product of differences in rainfall between *Runoff* and *Groundwater* sites. While floods can depress primary production through scouring of the benthos (Grimm and Fisher 1984, Uehlinger et al. 2003, Roberts et al. 2007), moderately elevated discharge can stimulate production by reducing competition for light and nutrients through continuous thinning of the biofilm while more nutrients are transported downstream (Stevenson 1990, Humphrey and Stevenson 1992). Further, channel substrate can influence GPP; in particular, a greater diversity of substrate sizes, as found in *Groundwater* streams, has been shown to increase primary production (Cardinale et al. 2002). The highly variable rock sizes included more mobile substrate that may have further stimulated GPP by providing more surface area for algal colonization and allowing for slight bed movement during spring rain events, which contrasts with the relatively homogeneous boulder and bedrock substrate of *Runoff* streams. Greater GPP in *Groundwater* streams may be a byproduct of channel substrate coupled with greater water clarity during the spring's higher flows; the benthos of each *Runoff* stream was difficult to observe if the water was more than approximately 0.5 meters deep (such as during the spring) due to the presence of minerals from glauconitic, easily-weathered green shale common to the Boston Mountains ecoregion (Caplan 1957, Caplan 1960), where *Runoff* streams are the dominant flow type. Conversely, *Groundwater* streams were clear year-round. While *Runoff* streams were warmer than *Groundwater* streams and contained more stable substrate for algal colonization, mineral effects on water clarity may have restricted primary production potential.

Similar to patterns reported in other forested systems and larger rivers, daily *Groundwater* stream GPP was greatest in spring and summer 2016, while summer 2015 had the greatest average GPP for *Runoff* streams (Mulholland et al. 2001, Acuña et al. 2004, Genzoli and Hall 2016). However, these trends differ from Roberts et al. (2007) in that GPP at Walker Branch was greatest during the spring and comparatively low during two consecutive summers. Uehlinger (2006) also reported GPP to be greatest in May over a 15-year period and lower in summers. *Runoff* stream GPP was lowest in spring 2015, coinciding with maximum algal biomass that may have reduced production rates due to competition. Daily *Groundwater* stream GPP was lowest in autumn 2015, which may have arisen from competition between benthic producers and microbes for resources, as reduced flows are common during early-to-mid autumn in Arkansas. It is worth noting that maximum algal biomass and production rates will not necessarily coincide. Algal biomass represents the state/structure of the algal community at a specific timepoint, and structure does not always directly reflect function (such as rates of production). At high levels of algal biomass, competition for light and nutrients may slow production rates (Sumner and Fisher 1979, Morin et al. 1999). While algae are allocating energy to maximize resource exploitation or resist the deleterious impacts of nutrient or light limitation due to lower competitive ability, there is less energy available for production, thus reducing primary production despite high biomass (McCormick 1996).

Respiration tended to be greater in *Groundwater* streams all year except during winter. This trend may have resulted from increased microbial activity post-abscission during the fall and a lack of large flood events through the summers and autumn. *Runoff* streams dried around the time of abscission, but even when rainfall replenished channel flow, respiration rates remained lower; a lack of substantial benthic storage in *Runoff* streams may have led to a net

export of organic matter once rains returned in November. More variable substrate size and a hyporheic corridor in *Groundwater* streams could have facilitated greater community respiration rates by providing spaces for organic matter storage and colonization of heterotrophic microbes (Demars 2019).

While our analyses did not reveal statistically significant differences in metabolism across flow regimes, *Groundwater* streams tended to have greater rates of GPP and ER that are likely biologically relevant to instream food webs and carbon dynamics. Production rates were two times greater in *Groundwater* streams in nearly every season, while respiration was, in some seasons, three and four times greater. *Groundwater* streams held significantly more biofilm ash-free dry mass, revealing that even marginally greater production and respiration rates can have noticeable effects on basal resources and the biological community. The tendency for *Groundwater* streams to have higher rates of GPP and ER through much of the year almost certainly influences algal and consumer community structure. Others have found relationships between metabolism and fish assemblages (Munn et al. 2020), and further work measuring invertebrate and fish assemblages across systems simultaneously with metabolism would provide insight into the potential link between biologically significant differences in stream metabolism and food web effects. Thus, while our statistical results did not detect an effect of flow regime on ecosystem function, that does not discount flow regime as an important variable shaping stream metabolic regime.

Our results show a stimulatory effect of discharge on ER, just as others have found (Roley et al. 2014). Greater discharge over the year may drive streams toward heterotrophy by transporting greater amounts of organic matter from riparian soils for microbes to consume (Demars 2019). Additionally, discharge, watershed area, and stream size are often strongly

correlated, and others have found GPP and ER to respond positively to drainage area (Mejia et al. 2019). While we did not find a direct relationship between watershed area and discharge in this study, this positive relationship between discharge and ER could potentially be an artifact of stream size. However, there was no relationship between discharge and GPP; thus, if stream size were affecting metabolism, it would be through its influence on heterotrophic and autotrophic carbon processing rather than production. Another possibility may be that grazers are mediating this relationship; herbivores may be more active in streams with lower discharge, thinning algal mats and increasing P:R ratios by ameliorating competition among periphyton, removing senescent cells, and ingesting microbes in the mat (Peckarsky et al. 2015).

The observed relationships between discharge and metabolism appear to be driven more by *Groundwater* than *Runoff* streams. Additionally, *Runoff* streams showed increased respiration then decreased at higher discharges. This could reflect potential flow class-specific differences in algal community composition, in which *Runoff* stream communities respond more variably to increases in discharge, though algal community data and flow-metabolism measurements from a greater number of streams would be needed to support this with certainty.

Greater discharge drove streams to be more heterotrophic due to streamflow stimulating ER with no detectable effect on GPP. This relationship between net ecosystem metabolism (NEM) and discharge may also have resulted from differences in hyporheic connectivity and channel substrate type and heterogeneity, reflecting an important point with respect to flow classifications: even during times of seemingly similar hydrologic conditions, ecosystem structure is an important mediator of functional responses from autotrophs and heterotrophs.

Mean annual discharge was the only flow variable we found to be related to metabolism in this study. This was likely through effects on organic matter transport coupled with the

presence or absence of a hyporheic corridor that provided additional space for biota to colonize as well as a source of continuous flow by groundwater intrusion. The importance of hyporheic connectivity and channel substrate to stream metabolism is well-documented, as carbon is utilized from surface water organic matter to support hyporheic respiration (Jones et al. 1995), and while greater discharge could have reduced microbial abundance in *Runoff* streams with no hyporheic refuge, it could have had a stimulatory effect in *Groundwater* streams with subsurface microbes benefitting from organic matter subsidies coming from upstream and the immediate riparian in elevated flows. The hyporheos is a crucial refuge for biota in streams that are susceptible to flash floods as well as drying (Dole-Olivier 2011, Stubbington 2012), and these two flow classifications exhibited distinct disturbance regimes over the course of the study; namely, *Runoff* streams are susceptible to drying, while *Groundwater* streams tend to experience more flood events of greater magnitude. *Groundwater* streams' connectivity to groundwater may have facilitated greater production and respiration in *Groundwater* streams during certain times of the year by providing asylum for biota to resist to the effects of drying (in the one *Groundwater* stream that dried for nine days) and flooding. In short, discharge and disturbance regimes are different across flow regimes, but hyporheic connectivity may mitigate impacts of drying and flooding on *Groundwater* stream metabolism.

Temperature and light have been shown to synergistically enhance the photosynthetic capacity of primary producers. Temperature can also be predictive of respiration, as it exerts control on the speed of organismal metabolism (Hill et al. 2000, Mulholland et al. 2001, Acuña et al. 2008, Beaulieu et al. 2013). Interestingly, we found the opposite trend- cooler streams had greater rates of respiration. However, this effect of temperature may be related to the volume of

groundwater flowing into the system and actually be an indirect measure of groundwater influence (Constantz 1998).

Similar to Bernot et al. (2010), we identified no relationships between phosphorus and GPP. However, other inter-regional studies have determined P to be a driver of GPP, and P concentrations were similar to those reported by others in forested systems (Lamberti and Steinman 1997, Mulholland et al. 2001). Our study sites exhibited low P concentrations; there was not a large gradient in TP concentrations, as these streams were minimally-impacted forested systems. The same was true of nitrogen; total N was also not related to GPP or ER. Others have also found N to be a significant predictor of GPP (Bernot et al. 2010). However, nutrient concentrations were similarly low across sites throughout the study, and our results indicate that the primary influence on metabolism across *Groundwater* and *Runoff* sites was channel discharge.

CONCLUSION

This comparison of forested systems across flow types provides a foundation for refining comparisons of stream metabolism across systems that may be similar in surrounding land cover, but differ in intermittency, discharge, dominant water source, and hyporheic connectivity. This provides insight into natural variation based on differences in flow versus anthropogenic hydrologic alteration but can also inform predictions regarding how depletion of groundwater resources can influence flow-function relationships. Loss of groundwater inputs will render stream biota more susceptible to droughts and will reduce the capacity to transport organic matter downstream for heterotrophic processing and sustenance of microbial carbon stocks that are important for lotic communities (Battin 1999, Demars 2019).

Our efforts reveal that forested stream metabolism rates are dependent upon discharge and exhibit seasonal trends based on flow regime classification. While biome and land use play key roles in determining stream metabolism, other factors such as streambed stability, hyporheic connectivity, and dominant water source may interact with landscape-level variables to produce variation in carbon dynamics across streams with similar land cover. These controls and flow-function relationships must be integrated across ecosystems as we work to understand sources, fates, and drivers of carbon transformation and transport in aquatic systems (Bernhardt et al. 2017, Hotchkiss et al. 2018).

ACKNOWLEDGEMENTS

We are grateful to Blake Lefler, Tyler Fletcher, Delaney Hall, Kayla Sayre, Hal Halvorson, Brad Austin, and Richard Walker for field assistance. We warmly thank Brian Haggard, Kristofor Brye, Kusum Naithani for suggestions on earlier versions on the manuscript. This work was funded by the USGS 104b program through the Arkansas Water Resources Center in FY2015. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

REFERENCES

- Acuña V, Giorgi A, Muñoz I, Uehlinger UR, Sabater S. Flow extremes and benthic organic matter shape the metabolism of a headwater Mediterranean stream. *Freshwater Biology*. 2004 Jul;49(7):960-71.
- Acuña V, Giorgi A, Muñoz I, Sabater F, Sabater S. Meteorological and riparian influences on organic matter dynamics in a forested Mediterranean stream. *Journal of the North American Benthological Society*. 2007 Mar;26(1):54-69.
- Acuña V, Wolf A, Uehlinger U, Tockner K. Temperature dependence of stream benthic respiration in an Alpine river network under global warming. *Freshwater Biology*. 2008 Oct;53(10):2076-88.
- Allan JD. Landscapes and riverscapes: the influence of land use on stream ecosystems. *Annu. Rev. Ecol. Evol. Syst.*. 2004 Dec 15;35:257-84.

545 Appling AP, Hall RO, Arroita M, Yackulic CB. streamMetabolizer: Models for estimating
546 aquatic photosynthesis and respiration. R package version 0.10. 2017;9.
547 <https://github.com/USGS-R/streamMetabolizer>
548

549 Appling AP, Read JS, Winslow LA, Arroita M, Bernhardt ES, Griffiths NA, Hall Jr RO, Harvey
550 JW, Heffernan JB, Stanley EH, Stets EG. The metabolic regimes of 356 rivers in the United
551 States. *Scientific data*. 2018 Dec 11;5:180292.

552 Argerich A, Martí E, Sabater F, Ribot M. Temporal variation of hydrological exchange and
553 hyporheic biogeochemistry in a headwater stream during autumn. *Journal of the North*
554 *American Benthological Society*. 2011 Sep;30(3):635-52.

555 Arthington AH, Bunn SE, Poff NL, Naiman RJ. The challenge of providing environmental flow
556 rules to sustain river ecosystems. *Ecological applications*. 2006 Aug;16(4):1311-8.

557 Arthington AH. *Environmental flows: saving rivers in the third millennium*. Univ of California
558 Press; 2012 Oct 15.

559 Austin BJ. Effects of landscape disturbances on autotrophic processes within Arkansas Ozark
560 streams. Doctoral Dissertation, University of Arkansas. 2015.

561 Beaulieu JJ, Arango CP, Balz DA, Shuster WD. Continuous monitoring reveals multiple controls
562 on ecosystem metabolism in a suburban stream. *Freshwater Biology*. 2013 May;58(5):918-
563 37.

564 Bernhardt ES, Heffernan JB, Grimm NB, Stanley EH, Harvey JW, Arroita M, Appling AP,
565 Cohen MJ, McDowell WH, Hall Jr RO, Read JS. The metabolic regimes of flowing waters.
566 *Limnology and Oceanography*. 2018 Mar;63(S1):S99-118.

567 Bernot MJ, Sobota DJ, Hall Jr RO, Mulholland PJ, Dodds WK, Webster JR, Tank JL, Ashkenas
568 LR, Cooper LW, Dahm CN, Gregory SV. Inter-regional comparison of land-use effects on
569 stream metabolism. *Freshwater Biology*. 2010 Sep;55(9):1874-90.

570 Biggs BJ, Nikora VI, Snelder TH. Linking scales of flow variability to lotic ecosystem structure
571 and function. *River research and applications*. 2005 Feb;21(2-3):283-98.

572 Blum AG, Ferraro PJ, Archfield SA, Ryberg KR. Causal effect of impervious cover on annual
573 flood magnitude for the United States. *Geophysical Research Letters*. 2020 Mar
574 16;47(5):no-.

575 Bott TL, Brock JT, Dunn CS, Naiman RJ, Ovink RW, Petersen RC. Benthic community
576 metabolism in four temperate stream systems: an inter-biome comparison and evaluation of
577 the river continuum concept. *Hydrobiologia*. 1985 Apr 1;123(1):3-45.

578 Bruckerhoff LA, Magoulick DD. Hydrologic regimes as potential drivers of morphologic
579 divergence in fish. *Evolutionary Ecology*. 2017 Aug 1;31(4):517-31.

580 Caplan, WM. Subsurface geology of northwestern Arkansas. 1957 Little Rock, Arkansas
581 Geological and Conservation Commission Information Circular 19:14.

582 Caplan, WM. Subsurface geology of Pre-Everton rocks in northern Arkansas. 1960. Little Rock,
583 Arkansas Geological and Conservation Commission Information Circular 21:17.

584 Cardinale BJ, Palmer MA, Swan CM, Brooks S, Poff NL. The influence of substrate
585 heterogeneity on biofilm metabolism in a stream ecosystem. *Ecology*. 2002 Feb;83(2):412-
586 22.

587 Carlisle DM, Falcone J, Wolock DM, Meador MR, Norris RH. Predicting the natural flow
588 regime: models for assessing hydrological alteration in streams. *River Research and*
589 *Applications*. 2010 Feb;26(2):118-36.

590 Chapman RA, Heitzman E, Shelton MG. Long-term changes in forest structure and species
591 composition of an upland oak forest in Arkansas. *Forest Ecology and Management*. 2006
592 Nov 15;236(1):85-92.

593 Constantz J. Interaction between stream temperature, streamflow, and groundwater exchanges in
594 alpine streams. *Water resources research*. 1998 Jul;34(7):1609-15.

595 Cook RA, Gawne B, Petrie R, Baldwin DS, Rees GN, Nielsen DL, Ning NS. River metabolism
596 and carbon dynamics in response to flooding in a lowland river. *Marine and Freshwater*
597 *Research*. 2015 Oct 19;66(10):919-27.

598 Dasmann RF. A system for defining and classifying natural regions for purposes of conservation.
599 A progress report. UICN, Morges (Suiza); 1973.

600 Demars BO. Hydrological pulses and burning of dissolved organic carbon by stream respiration.
601 *Limnology and Oceanography*. 2019 Jan;64(1):406-21.

602 Detenbeck NE, DeVore PW, Niemi GJ, Lima A. Recovery of temperate-stream fish communities
603 from disturbance: a review of case studies and synthesis of theory. *Environmental*
604 *management*. 1992 Jan 1;16(1):33.

605 Dodd, AK, Pollock, ED, and Evans-White MA. Quantifying flow sources and their impact on
606 water quality in Ozark streams. 2020; *Arkansas Bulletin of Water Research* 2020: 18-23.

607 Dodds WK, Hutson RE, Eiche AC, Evans MA, Gudder DA, Fritz KM, Gray L. The
608 relationship of floods, drying, flow and light to primary production and producer biomass in
609 a prairie stream. *Hydrobiologia*. 1996 Oct 1;333(3):151-9.

610 Dodds WK, Higgs SA, Spangler MJ, Guinnip J, Scott JD, Hedden SC, Frenette BD, Taylor R,
611 Schechner AE, Hoeninghaus DJ, Evans-White MA. Spatial heterogeneity and controls of
612 ecosystem metabolism in a Great Plains river network. *Hydrobiologia*. 2018 May
613 1;813(1):85-102.

614 Dole-Olivier MJ. The hyporheic refuge hypothesis reconsidered: a review of hydrological
615 aspects. *Marine and Freshwater Research*. 2011 Nov 23;62(11):1281-302.

616 Dugan PJ, Barlow C, Agostinho AA, Baran E, Cada GF, Chen D, Cowx IG, Ferguson JW,
617 Jutagate T, Mallen-Cooper M, Marmulla G. Fish migration, dams, and loss of ecosystem
618 services in the Mekong basin. *Ambio*. 2010 Jun 1;39(4):344-8.

619 Eaton AD, Franson MA. American Public Health Association (APHA) (2005) Standard Methods
620 for the Examination of Water & Wastewater. American Public Health Association,
621 American Water Works Association, Water Environment Federation, USA, Washington.
622 2005.

623 Elser JJ, Bracken ME, Cleland EE, Gruner DS, Harpole WS, Hillebrand H, Ngai JT, Seabloom
624 EW, Shurin JB, Smith JE. Global analysis of nitrogen and phosphorus limitation of primary
625 producers in freshwater, marine and terrestrial ecosystems. *Ecology letters*. 2007
626 Dec;10(12):1135-42.

627 Fisher SG, Gray LJ, Grimm NB, Busch DE. Temporal succession in a desert stream ecosystem
628 following flash flooding. *Ecological monographs*. 1982 Feb;52(1):93-110.

629 Genzoli L, Hall Jr RO. Shifts in Klamath River metabolism following a reservoir cyanobacterial
630 bloom. *Freshwater Science*. 2016 Sep 1;35(3):795-809.

631 Gore JA, Banning J. Discharge measurements and streamflow analysis. In *Methods in Stream
632 Ecology*, Volume 1 2017 Jan 1 (pp. 49-70). Academic Press.

633 Grimm, NB and Fisher SG. Stability of periphyton and macroinvertebrates to disturbance by
634 flash floods in a desert stream. 1989. *J N Am Benthol Soc* 8: 293–307.

635 Hall Jr RO, Tank JL. Correcting whole-stream estimates of metabolism for groundwater input.
636 *Limnology and Oceanography: Methods*. 2005 Apr;3(4):222-9.

637 Hall Jr RO, Hotchkiss ER. Stream metabolism. In *Methods in stream ecology* 2017 Jan 1 (pp.
638 219-233). Academic Press.

639 Hill BH, Hall RK, Husby P, Herlihy AT, Dunne M. Interregional comparisons of sediment
640 microbial respiration in streams. *Freshwater Biology*. 2000 Jun 1;44(2):213-22.

641 Hornbach DJ, Beckel R, Hustad EN, McAdam DP, Roen IM, Wareham AJ. The influence of
642 riparian vegetation and season on stream metabolism of Valley Creek, Minnesota. *Journal
643 of Freshwater Ecology*. 2015 Oct 2;30(4):569-88.

644 Hotchkiss ER, Sadro S, Hanson PC. Toward a more integrative perspective on carbon
645 metabolism across lentic and lotic inland waters. *Limnology and Oceanography Letters*.
646 2018 Jun 1;3(3):57-63.

647 Humphrey KP, Stevenson RJ. Responses of benthic algae to pulses in current and nutrients
648 during simulations of subscouring spates. *Journal of the North American Benthological*
649 *Society*. 1992 Mar 1;11(1):37-48.

650 Humphries P, Baldwin DS. Drought and aquatic ecosystems: an introduction. *Freshwater*
651 *Biology*. 2003 Jul;48(7):1141-6.

652 Hurn AD, Wallace JB. Life history and production of stream insects. *Annual review of*
653 *entomology*. 2000 Jan;45(1):83-110.

654 Hynes HB. The stream and its valley: With 4 figures and 2 tables in the text. *Internationale*
655 *Vereinigung für theoretische und angewandte Limnologie: Verhandlungen*. 1975 Oct
656 1;19(1):1-5.

657 Izagirre O, Agirre U, Bermejo M, Pozo J, Elozegi A. Environmental controls of whole-stream
658 metabolism identified from continuous monitoring of Basque streams. *Journal of the North*
659 *American Benthological Society*. 2008 Jun;27(2):252-68.

660 Jones JA, Swanson FJ, Wemple BC, Snyder KU. Effects of roads on hydrology, geomorphology,
661 and disturbance patches in stream networks. *Conservation Biology*. 2000 Feb;14(1):76-
662 85.

663
664 Jones Jr JB, Fisher SG, Grimm NB. Vertical hydrologic exchange and ecosystem metabolism in
665 a Sonoran Desert stream. *Ecology*. 1995 Apr;76(3):942-52.

666
667 Koenig LE, Helton AM, Savoy P, Bertuzzo E, Heffernan JB, Hall Jr RO, Bernhardt ES.
668 Emergent productivity regimes of river networks. *Limnology and Oceanography Letters*.
669 2019 Oct;4(5):173-81. Lajoie F, Assani AA, Roy AG, Mesfioui M. Impacts of dams on
670 monthly flow characteristics. The influence of watershed size and seasons. *Journal of*
671 *hydrology*. 2007 Feb 28;334(3-4):423-39.

672 Lamberti GA, Steinman AD. A comparison of primary production in stream ecosystems. *Journal*
673 *of the North American Benthological Society*. 1997 Mar 1;16(1):95-104.

674 Leasure DR, Magoulick DD, Longing SD. Natural flow regimes of the Ozark–Ouachita interior
675 highlands region. *River Research and Applications*. 2016 Jan;32(1):18-35.

676 Ligon FK, Dietrich WE, Trush WJ. Downstream ecological effects of dams. *BioScience*. 1995
677 Mar 1;45(3):183-92.

678 Lytle DA, Poff NL. Adaptation to natural flow regimes. *Trends in ecology & evolution*. 2004
679 Feb 1;19(2):94-100.

680 McCormick PV. Resource competition and species coexistence in freshwater benthic algal
681 assemblages. In *Algal ecology* 1996 Jan 1 (pp. 229-252). Academic Press.

682 Mejia FH, Fremier AK, Benjamin JR, Bellmore JR, Grimm AZ, Watson GA, Newsom M.
683 Stream metabolism increases with drainage area and peaks asynchronously across a stream
684 network. *Aquatic Sciences*. 2019 Jan 1;81(1):9.

685 Morin A, Lamoureux W, Busnarda J. Empirical models predicting primary productivity from
686 chlorophyll a and water temperature for stream periphyton and lake and ocean
687 phytoplankton. *Journal of the North American Benthological Society*. 1999 Sep
688 1;18(3):299-307.

689 Mulholland PJ, Marzolf ER, Webster JR, Hart DR, Hendricks SP. Evidence that hyporheic zones
690 increase heterotrophic metabolism and phosphorus uptake in forest streams. *Limnology and*
691 *oceanography*. 1997 May;42(3):443-51.

692 Mulholland PJ, Fellows CS, Tank JL, Grimm NB, Webster JR, Hamilton SK, Martí E, Ashkenas
693 L, Bowden WB, Dodds WK, McDowell WH. Inter-biome comparison of factors controlling
694 stream metabolism. *Freshwater biology*. 2001 Nov 1;46(11):1503-17.

695 Mulholland PJ, Helton AM, Poole GC, Hall RO, Hamilton SK, Peterson BJ, Tank JL, Ashkenas
696 LR, Cooper LW, Dahm CN, Dodds WK. Stream denitrification across biomes and its
697 response to anthropogenic nitrate loading. *Nature*. 2008 Mar;452(7184):202-5.

698 Munn MD, Sheibley RW, Waite IR, Meador MR. Understanding the relationship between stream
699 metabolism and biological assemblages. *Freshwater Science*. 2020 Dec 1;39(4):680-92.

700 Naiman RJ. The annual pattern and spatial distribution of aquatic oxygen metabolism in boreal
701 forest watersheds. *Ecological monographs*. 1983 Feb;53(1):73-94.

702 Odum HT. Primary production in flowing waters 1. *Limnology and oceanography*. 1956
703 Apr;1(2):102-17.

704 O'Donnell B, Hotchkiss ER. Coupling Concentration-and Process-Discharge Relationships
705 Integrates Water Chemistry and Metabolism in Streams. *Water Resources Research*. 2019
706 Dec;55(12):10179-90.

707 Palmer M, Ruhi A. Linkages between flow regime, biota, and ecosystem processes: Implications
708 for river restoration. *Science*. 2019 Sep 20;365(6459):eaaw2087.

709 Paul MJ, Meyer JL. Streams in the urban landscape. *Annual review of Ecology and Systematics*.
710 2001 Nov;32(1):333-65.

711 Peckarsky BL, McIntosh AR, Álvarez M, Moslemi JM. Disturbance legacies and nutrient
712 limitation influence interactions between grazers and algae in high elevation streams.
713 *Ecosphere*. 2015 Nov;6(11):1-5.

714 Poff NL, Ward JV. Implications of streamflow variability and predictability for lotic community
715 structure: a regional analysis of streamflow patterns. *Canadian journal of fisheries and*
716 *aquatic sciences*. 1989 Oct 1;46(10):1805-18.

717 Poff NL, Ward JV. Physical habitat template of lotic systems: recovery in the context of
718 historical pattern of spatiotemporal heterogeneity. *Environmental management*. 1990 Sep
719 1;14(5):629.

720 Poff NL, Allan JD, Bain MB, Karr JR, Prestegard KL, Richter BD, Sparks RE, Stromberg JC.
721 The natural flow regime. *BioScience*. 1997 Dec 1;47(11):769-84.

722 Poff NL, Bledsoe BP, Cuhaciyan CO. Hydrologic variation with land use across the contiguous
723 United States: geomorphic and ecological consequences for stream ecosystems.
724 *Geomorphology*. 2006 Sep 30;79(3-4):264-85.

725 Poff NL, Richter BD, Arthington AH, Bunn SE, Naiman RJ, Kendy E, Acreman M, Apse C,
726 Bledsoe BP, Freeman MC, Henriksen J. The ecological limits of hydrologic alteration
727 (ELOHA): a new framework for developing regional environmental flow standards.
728 *Freshwater Biology*. 2010 Jan;55(1):147-70.

729 Poff NL, Zimmerman JK. Ecological responses to altered flow regimes: a literature review to
730 inform the science and management of environmental flows. *Freshwater Biology*. 2010
731 Jan;55(1):194-205.

732 Qasem K, Vitousek S, O'Connor B, Hoellein T. The effect of floods on ecosystem metabolism in
733 suburban streams. *Freshwater Science*. 2019 Jun 1;38(2):412-24.

734 Reisinger AJ, Rosi EJ, Bechtold HA, Doody TR, Kaushal SS, Groffman PM. Recovery and
735 resilience of urban stream metabolism following Superstorm Sandy and other floods.
736 *Ecosphere*. 2017 Apr;8(4):e01776.

737 Richter BD. Re-thinking environmental flows: from allocations and reserves to sustainability
738 boundaries. *River Research and Applications*. 2010 Oct;26(8):1052-63.

739 Riley AJ, Dodds WK. Whole-stream metabolism: strategies for measuring and modeling diel
740 trends of dissolved oxygen. *Freshwater Science*. 2013 Mar;32(1):56-69.

741 Roberts BJ, Mulholland PJ, Hill WR. Multiple scales of temporal variability in ecosystem
742 metabolism rates: results from 2 years of continuous monitoring in a forested headwater
743 stream. *Ecosystems*. 2007 Jun 1;10(4):588-606.

744 Roley SS, Tank JL, Griffiths NA, Hall Jr RO, Davis RT. The influence of floodplain restoration
745 on whole-stream metabolism in an agricultural stream: insights from a 5-year continuous
746 data set. *Freshwater Science*. 2014 Dec 1;33(4):1043-59.

747 Rovelli L, Attard KM, Binley A, Heppell CM, Stahl H, Trimmer M, Glud RN. Reach-scale river
748 metabolism across contrasting sub-catchment geologies: Effect of light and hydrology.
749 *Limnology and oceanography*. 2017 Nov;62(S1):S381-99.

750 Sauer VB, Thomas Jr WO, Stricker VA, Wilson KV. Flood characteristics of urban watersheds
751 in the United States. USGPO,; 1983.

752 Savoy P, Appling AP, Heffernan JB, Stets EG, Read JS, Harvey JW, Bernhardt ES. Metabolic
753 rhythms in flowing waters: An approach for classifying river productivity regimes.
754 *Limnology and Oceanography*. 2019 Sep;64(5):1835-51.

755 Sinsabaugh RL. Large-scale trends for stream benthic respiration. *Journal of the North American*
756 *Benthological Society*. 1997 Mar 1;16(1):119-22.

757 Southwood TR. Habitat, the templet for ecological strategies? *Journal of animal ecology*. 1977
758 Jun 1;46(2):337-65.

759 Steinman AD, Lamberti GA, Leavitt PR, Uzarski DG. Biomass and pigments of benthic algae. In
760 *Methods in Stream Ecology, Volume 1* 2017 Jan 1 (pp. 223-241). Academic Press.

761 Stephenson SL, Adams HS, Huebner CD. Upland forest vegetation of the Ozark Mountains in
762 northwestern Arkansas. *Rhodora*. 2007 Jun;109(938):197-221.

763 Stevenson RJ. Benthic algal community dynamics in a stream during and after a spate. *Journal of*
764 *the North American Benthological Society*. 1990 Sep 1;9(3):277-88.

765 Stroud Water Research Center. Model My Watershed [Software]. 2017. Available from
766 <https://wikiwatershed.org/>

767 Stubbington R. The hyporheic zone as an invertebrate refuge: a review of variability in space,
768 time, taxa and behaviour. *Marine and Freshwater Research*. 2012 Apr 24;63(4):293-311.

769 Sumner WT, Fisher SG. Periphyton production in Fort River, Massachusetts. *Freshwater*
770 *Biology*. 1979 Jun;9(3):205-12.

771 Thoms MC, Parsons M. Eco-geomorphology: an interdisciplinary approach to river science.
772 *International Association of Hydrological Sciences, Publication*. 2002 Sep;276:113-9p.

773 Uehlinger U. Resistance and resilience of ecosystem metabolism in a flood-prone river system.
774 *Freshwater Biology*. 2000 Nov;45(3):319-32.

775 Uehlinger U, Kawecka B, Robinson CT. Effects of experimental floods on periphyton and
776 stream metabolism below a high dam in the Swiss Alps (River Spöl). *Aquatic Sciences*.
777 2003 Sep 1;65(3):199-209.

778 Uehlinger U. Annual cycle and inter-annual variability of gross primary production and
779 ecosystem respiration in a floodprone river during a 15-year period. *Freshwater Biology*.
780 2006 May;51(5):938-50.

781 Vilches C, Giorgi A. Metabolism in a macrophyte-rich stream exposed to flooding.
782 *Hydrobiologia*. 2010 Oct 1;654(1):57-65.

783 Walsh CJ, Roy AH, Feminella JW, Cottingham PD, Groffman PM, Morgan RP. The urban
784 stream syndrome: current knowledge and the search for a cure. *Journal of the North*
785 *American Benthological Society*. 2005 Sep;24(3):706-23.

786 Webster JR, Wallace JB, Benfield EF. Organic processes in streams of the eastern United States.
787 *River stream Ecosyst. world*. 1995:117-87.

788 Williamson CE, Dodds W, Kratz TK, Palmer MA. Lakes and streams as sentinels of
789 environmental change in terrestrial and atmospheric processes. *Frontiers in Ecology and the*
790 *Environment*. 2008 Jun;6(5):247-54.

791 Woods AJ. Ecoregions of Arkansas. US Geological Survey; 2004.

792 Yates AG, Brua RB, Culp JM, Chambers PA. Multi-scaled drivers of rural prairie stream
793 metabolism along human activity gradients. *Freshwater Biology*. 2013 Apr;58(4):675-89.

794 Young RG, Huryn AD. Effects of land use on stream metabolism and organic matter turnover.
795 *Ecological Applications*. 1999 Nov;9(4):1359-76.

796

797

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 1