Rapid screening of pest resistance genes in maize using a sugarcane mosaic virus vector

Seung Ho Chung¹, Mahdiyeh Bigham¹, Ryan R. Lappe³, Barry Chan², Ugrappa Nagalakshmi², Steven A. Whitham³, Savithramma P. Dinesh-Kumar², and Georg Jander¹

¹Boyce Thompson Institute for Plant Research, Ithaca, New York 14853

²Department of Plant Biology and The Genome Center, College of Biological Sciences, University of California, Davis, CA 95616

³Department of Plant Pathology and Microbiology, Iowa State University, Ames, Iowa 50011

Abstract:

Spodoptera frugiperda (fall armyworm) is a notorious pest that threatens maize production world-wide. Current control measures involve the use of chemical insecticides and transgenic maize expressing *Bacillus thuringiensis* (*Bt*) toxins. Although several additional transgenes have confirmed insecticidal activity in other plants, limited research has been conducted in maize, at least partially due to the technical difficulty of maize transformation. Here, we describe implementation of a sugarcane mosaic virus (SCMV) vector for rapidly testing the efficacy of transgenes for the control of *S. frugiperda* in maize. Four categories of proteins were tested using the SCMV vector: (i) maize defense signaling proteins: peptide elicitors (Pep1 and Pep3) and jasmonate acid conjugating enzymes (JAR1a and JAR1b); (ii) maize defensive proteins: the previously identified ribosome-inactivating protein (RIP2) and maize proteinase inhibitor (MPI), and two proteins with predicted but unconfirmed anti-insect activities, an antimicrobial peptide (AMP) and a lectin (JAC1); (iii) lectins from other plant species: *Allium cepa* agglutinin (ACA) and *Galanthus nivalis* agglutinin (GNA); and (iv) spider and scorpion toxins: peptides from *Urodacus yaschenkoi* (UyCT3 and UyCT5) and *Hadronyche versuta* (Hvt). In most cases, *S. frugiperda* larval growth on maize was reduced by transient SCMV-mediated overexpression of genes encoding these proteins. Additionally, experiments with some of the SCMV-expressed genes showed effectiveness against two aphid species, *Rhopalosiphum maidis* (corn leaf aphid) and *Myzus persicae* (green peach aphid). Together, these results demonstrate that SCMV vectors can be exploited as a rapid screening method for testing the efficacy and insecticidal activity of candidate genes in maize.

Keywords:

Maize, *Zea mays*, *Spodoptera frugiperda*, fall armyworm, *Rhopalosiphum maidis*, *Myzus persicae*, sugarcane mosaic virus, peptide elicitors, venom toxins, lectins

Introduction:

Maize (*Zea mays*) is one of the world's most important cereal crops, serving not only as a food source for humans and livestock, but also as a raw material for the production of ethanol and other industrial products (Ai and Jane, 2016; Chaudhary *et al.*, 2014). The needs of an ever-expanding population will lead to increasing demands on maize production in the coming years. Therefore, maintaining adequate maize yields will require reducing not only reducing the cost of agricultural inputs, but also the negative impacts of biotic and abiotic stresses that limit maize productivity.

More than 100 species of insect pests limit maize productivity in agricultural fields (McMullen *et al.*, 2009; Meihls *et al.*, 2012). Among these pests, one of the most damaging is *Spodoptera frugiperda* (fall armyworm; Figure 1a), a lepidopteran species that is indigenous to the Americas but recently has become invasive in Africa and Asia (Food and Agriculture Organization of the United Nations, 2018; Goergen *et al.*, 2016). By consuming all above-ground plant parts, *S. frugiperda* larvae reduce photosynthetic area, cause developmental delays, and decrease yield.

Currently available *S. frugiperda* control methods, both application of chemical insecticides (Togola *et al.*, 2018) and transgenic maize producing *Bacillus thuringiensis* (Bt) toxins (Huang *et al.*, 2014; Tabashnik and Carrière, 2017), are becoming less effective as the insects develop resistance. Therefore, there is a need to screen for additional transgenes that can be used to enhance maize resistance to *S. frugiperda* feeding. Broadly, such approaches can include upregulation of maize defense signaling, overexpression of individual maize defensive proteins, and expression of exogenous insecticidal proteins.

Plant elicitor peptides (Peps) trigger anti-herbivore defense responses (Huffaker, 2015; Huffaker *et al.*, 2013; Poretsky *et al.*, 2020). In maize, *ZmPep1* and *ZmPep3* upregulate defenses, at least in part by induction of the jasmonic acid (JA) signaling pathway (Huffaker *et al.*, 2013, 2011). A key step in the JA pathway is the conjugation of JA with isoleucine by JAR1 (JASMONATE RESISTANT 1) enzymes (Koo and Howe, 2009; Staswick *et al.*, 2002) to form JA-isoleucine. Expression of *JAR1a* and *JAR1b*, two of the five predicted *JAR* genes in maize (Borrego and Kolomiets, 2016), is highly induced by *Spodoptera exigua* (beet armyworm) herbivory (Tzin *et al.*, 2017). Thus, these maize genes are good targets for overexpression to enhance resistance against *S. frugiperda*.

Maize and other plants produce ribosome-inactivating proteins (RIPs) that block ribosome function by depurinating a specific adenine residue of the large ribosomal RNA (Bass *et al.*, 2004; Zhu *et al.*, 2018). These proteins, which are toxic for a variety of insects, including Lepidoptera (Shahidi-Noghabi *et al.*, 2009) and Hemiptera (Hamshou *et al.*, 2016), have been used previously in transgenic approaches. For instance, the expression of a maize kernel RIP1 in *Nicotiana tabacum* (tobacco) increased resistance to *Helicoverpa zea* (corn earworm) feeding (Dowd *et al.*, 2003). The *RIP2* gene is expressed in all maize tissues except the kernels (Bass *et al.*, 2004). *RIP2* expression was induced by *S. frugiperda* herbivory and recombinant *RIP2* protein decreased caterpillar growth on artificial diet (Chuang, Herde, *et al.*, 2014).

Two additional classes of maize proteins with anti-herbivore activity are proteinase inhibitors and antimicrobial peptides (Campos *et al.*, 2018; Koiwa *et al.*, 1997). Proteinase inhibitors, which are produced by many plant families, impair the growth and survival of insects by disrupting the function of digestive enzymes. Maize proteinase inhibitor (*MPI*) expression was induced by both caterpillar herbivory and JA signaling (Cordero *et al.*, 1994; Shivaji *et al.*, 2010; Tamayo *et al.*, 2000). Heterologous expression of *MPI* in rice increased resistance to *Chilo*

suppressalis (striped stem borer) (Vila *et al.*, 2005). Cyclotides are macrocyclic insecticidal peptides with the length of about 30 amino acids and a conserved cystine knot motif containing three disulfide bonds (Campos *et al.*, 2018; Craik *et al.*, 1999; Weidmann and Craik, 2016). Cyclotide Kalata B1 from *Oldenlandia affinis* decreased the growth of *Helicoverpa armigera* (corn earworm) larvae by rupturing epithelial cells in the midgut (Barbeto *et al.*, 2008). Cycloviolacins, cyclotides from *Viola odorata*, negatively affected the probing and feeding behavior of *Mysus persicae* (green peach aphid), suggesting that cycloviolacins limit aphid population growth (Dancewicz *et al.*, 2020). Among predicted antimicrobial peptides in maize, a few belong to the cyclotide family (Mulvenna *et al.*, 2006; Noonan *et al.*, 2017), but their efficacy against insects has not been confirmed.

Plant lectins, carbohydrate-binding proteins that interact with glycoproteins and glycan structures in insect guts, have antinutritional or insecticidal effects (Macedo *et al.*, 2015). For instance, snowdrop lectin (*Galanthus nivalis* agglutinin; GNA), onion lectin (*Allium cepa* agglutinin; ACA) and garlic (*Allium sativum*) leaf lectin reduce nutrient uptake and growth in wide range of insects (Vandenborre *et al.*, 2011). Expression of a maize lectin gene, Jacalin 1 (*JAC1*), is induced by JA, an indication that it may provide protection against herbivory (Van Damme *et al.*, 2004). There can be additive or even synergistic effects if lectins are co-expressed or fused to scorpion or spider venom peptides. For instance, the insecticidal efficacy of GNA was increased by fusions to Hvt (Fitches *et al.*, 2012), ButaIT from *Mesobuthus tamulus* (Fitches *et al.*, 2010), AaIT from *Androctonus australis* (Liu *et al.*, 2016), and δ -amaurobitoxin-P1la from *Pireneitega luctuosus* (Yang *et al.*, 2014).

Scorpion and spider venoms, which contain numerous insecticidal toxins (King and Hardy, 2013; Ortiz *et al.*, 2015), have been explored as sources of insecticidal peptides. UyCT3 and UyCT5, two antimicrobial peptides that are produced in the venom glands of *Urodacus yaschenkoi* (inland robust scorpion) decrease the fitness of *Acyrtosiphon pisum* (pea aphid) and the density of the primary symbiont, *Buchnera aphidicola*, suggesting that those are promising candidates for the production of insect-resistant transgenic plants (Luna-Ramirez *et al.*, 2017). Similarly, ω -hexatoxin Hv1a (Hvt, also called ω -ACTX Hv1a) from *Hadronyche versuta* (Blue Mountains funnel web spider) is broadly effective against both lepidopteran and hemipteran pests when expressed in transgenic plants (Javaid *et al.*, 2016; Ullah *et al.*, 2015; Rauf *et al.*, 2019). Given the broad insecticidal activity of these spider and scorpion venom proteins, we hypothesized that they would also be effective in reducing *S. frugiperda* growth.

Testing the effectiveness of transgenes for controlling pest insects on maize is limited by the high cost of maize transformation and the often greater than one-year timeline that is required to obtain transgenic maize plants for experiments. Therefore, we are proposing an alternate approach, whereby the efficacy of transgenes that enhance maize pest tolerance is tested by transient expression in maize using a virus vector. Sugarcane mosaic virus (SCMV), a positive-sense single-stranded RNA virus, which has been adapted for efficient transgene expression in maize (Mei *et al.*, 2019), is attractive vector for such experiments. Genes of interest are inserted between the SCMV *PI* and *HC-Pro* cistrons in SCMV-CS3, a newly created plasmid vector, maize seedlings are infected by particle bombardment or *Agrobacterium* inoculation, and the pest resistance of the infected plants can be assessed after three weeks. Here we show that virus-mediated expression of maize defense-regulating proteins, maize insecticidal proteins, and exogenous toxins can reduce the growth of insect pests. We demonstrate the insect-controlling properties of not only known proteins but also two maize proteins that were not previously confirmed to have anti-herbivore properties. Furthermore, we show that expression of transgenes

using SCMV also is effective in reducing production by two hemipteran pests, *Rhopalosiphum maidis* (corn leaf aphid; Figure 1b) and *M. persicae* (Figure 1c).

Materials and methods

Plants and insects

Maize (*Zea mays*) plants, sweet corn variety Golden Bantam (West Coast Seeds, British Columbia, Canada) and inbred lines P39 and B73, were grown in a maize mix [0.16 m³ Metro-Mix 360 (Scotts, Marysville, OH, USA), 0.45 kg finely ground lime, 0.45 kg Peters Unimix (Griffin Greenhouse Supplies, Auburn, NY, USA), 68 kg Turface MVP (Banfield-Baker Corp., Horseheads, NY, USA), 23 kg coarse quartz sand, and 0.018 m³ pasteurized field soil]. All plants, including those used for SCMV propagation and insect bioassays, were maintained in a growth chamber at 23°C with a 16:8 light:dark cycle. Unless specified otherwise, Golden Bantam maize was used for the described experiments.

Eggs of *S. frugiperda* (fall armyworm) were purchased from Benzon Research (Carlisle, PA, USA) and maintained on an artificial diet (Fall Armyworm Diet, Southland Products Inc, Lake Village, AR, USA) in an incubator at 28°C. A colony of a genome-sequenced *R. maidis* lineage (W. Chen *et al.*, 2019) was maintained on maize (Golden Bantam or P39) and a colony of a previously described tobacco-adapted strain of *M. persicae* (Ramsey *et al.*, 2007) was maintained on *Nicotiana tabacum* (tobacco) at 23°C under 16:8 light:dark cycle. Both *R. maidis* and *M. persicae* were originally collected in USDA-ARS greenhouses by Steward Gray (Robert W. Holley Center for Agriculture & Health, Ithaca, NY, USA).

Cloning of candidate genes into *Sugarcane mosaic virus* for protein expression

The pSCMV-CS3 expression vector used in this work was derived from pSCMV-CS2 (Mei *et al.*, 2019), which was modified to contain the CS3 restriction sites in the MCS between the P1 and HC-Pro cistrons (Figure 2a). The modified pSCMV-CS2 genome plus the flanking 35S promoter and NOS terminator was amplified with SuperFi polymerase (ThermoFisher Scientific, Waltham, MA) using primers DCPacI 1380 F and DCPacI 1380 R (Supplementary Table S1). The pCambia1380 backbone (www.cambia.org) was amplified with SuperFi polymerase using primers 1380F and 1380R (Supplementary Table S1). The two PCR fragments were subsequently assembled into pCambia1380-SCMV-CS3, hereafter referred to as pSCMV-CS3, by Gibson Assembly (New England Biolabs, Ipswich, MA). pSCMV-GFP was created by amplifying the mEGFP coding sequence (Zacharias *et al.*, 2002) using SuperFi polymerase with the GFP-Psp and GFP-SbfI primers (Supplementary Table S1). The resulting amplicon was digested with *Psp*OMI and *Sbf*I, gel purified and ligated into similarly digested pSCMV-CS3.

Genes encoding maize defense regulators and protein toxins were amplified from the B73 cDNA template with gene-specific primers containing restriction sites at the 5' end for cloning into SCMV vector. PCR products were gel purified, digested with the corresponding restriction enzymes and cloned into the SCMV vector (Figure 2a). The *UyCT3* and *UyCT5* genes (codon optimized for maize), as well as the maize *Pep1* and *Pep3* genes, were synthesized by Genscript Biotech (Piscataway, NJ, USA) and cloned into the SCMV vector. To generate SCMV with *Hvt*, *ACA*, *GNA*, *RIP2*, *Hvt-ACA*, *Hvt*-ACA*, *RIP2-GNA*, and *Hvt-GNA*, primers listed in Supplementary Table S1 and DNA synthesized by Genewiz (South Plainfield, NJ, USA) were used as template and the resulting PCR products were gel purified and cloned into *Psp*OMI-SbfI cut SCMV by following the NEBuilder HiFi DNA Assembly method (NEB, Ipswich, MA,

USA). Primers, restriction sites, and cloning methods used in this study are listed in Supplementary Table S1.

Inoculation of maize with SCMV constructs

SCMV constructs were delivered into maize plants by particle bombardment using a Biolistic PDS-1000/He system (Bio-Rad, Hercules, CA, USA) as described previously (Mei and Whitham, 2018). One μg of the plasmid DNA was coated on 3 mg 1.0 μm diameter gold particles, and the coated gold particles were distributed onto five microcarriers and allowed to air dry. Plants were placed in the dark 12 h before the particle bombardment. Two leaves of one-week-old plants were bombarded with using 1,100 psi rupture disks at a distance of 6 cm (between stopping screen and leaves).

Agroinoculation was used to initiate maize infections with the SCMV constructs containing *Hvt*, *ACA*, and *UyCT3*. The constructs were transformed into the *Agrobacterium tumefaciens* strain GV3101 and an *A. tumefaciens* suspension with optical density at 600 nm (OD_{600}) = 1.0 in infiltration buffer (200 μM acetosyringone, 10 mM MES, pH 5.6, and 10 mM MgCl_2) was injected above the coleoptile node of one-week-old plants.

After the initial infection by particle bombardment or Agroinoculation, SCMV constructs were further propagated by rub-inoculation. Leaf sap of SCMV-infected plants was prepared by grinding 0.5 g leaf tissue in 5 ml of 50 mM pH 7.0 potassium phosphate buffer. One-week-old maize plants were dusted with 600-mesh carborundum and mechanically inoculated by rubbing leaf sap from virus-infected maize plants on two leaves.

Confocal microscopy

Three weeks post-inoculation, leaf samples were collected from the seventh or eighth leaves of maize plants infected with SCMV-*GFP*. The samples were observed at an excitation of 488 nm. The emitted fluorescence signal was monitored from 505 to 545 nm using a SP5 Leica Confocal Microscope in the Plant Cell Imaging Center of Boyce Thompson Institute. A scan of fluorescence across a range of wavelengths (lambda scan) was used to confirm that the observed signal was derived from GFP rather than endogenous maize fluorescence.

Insect bioassays

To determine the effect of defensive proteins on the growth of *S. frugiperda*, SCMV-infected plants, three weeks post-inoculation, were used for caterpillar bioassays. Five two-day-old caterpillars were placed on each plant and enclosed using perforated plastic bags (13 cm x 61 cm, <https://www.clearbags.com>). The caterpillar fresh mass was measured one week later. For aphid bioassays, eight 10-day-old apterous adult *R. maidis* or ten 10-day-old apterous adult *M. persicae* were placed on each virus-infected plant 15-18 days after SCMV inoculation and enclosed using perforated plastic bags. The total numbers of aphids were counted one week later.

RNA extraction, cDNA synthesis, RT-PCR, and quantitative real-time PCR (qRT-PCR)

Three weeks post-inoculation and prior to the insect bioassays, leaf tissue was collected from the seventh or eighth leaves of infected plants, flash-frozen in liquid nitrogen, and stored at -80°C . RNA was extracted using TRIzol Reagent (Invitrogen, Carlsbad, CA, USA) and treated with RQ1 RNase-free DNase (Promega, Madison, WI, USA). One microgram of RNA was used to synthesize first-strand cDNA using High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA, USA) with random primers. To verify the expression of insect

resistance genes cloned into the SCMV vector, qRT-PCR was conducted with gene-specific primers (Supplementary Table S1). For the inserts of less than 75 bp (*Pep1*, *Pep3*, *UyCT3*, and *UyCT5*), one primer was designed to bind to the insert and another primer was designed to bind to the region flanking the cloning site, with the amplification products ranging in size from 100-150 bp. The reactions consisted of 5.0 µl of the PowerUp SYBR Green PCR master mix (Applied Biosystems), 0.6 µl primer mix (300 nM for the final concentration of each primer) and 2 µl of cDNA (1:10 dilution with nuclease-free H₂O) in 10 µl total volume. Template-free reactions were included as negative controls. The PCR amplification was performed on QuantStudio 6 Flex Real-Time PCR Systems (Applied Biosystems, Foster City, CA, USA) with the following conditions: 2 min at 50°C, 2 min at 95°C, 40 cycles of 95°C for 15 sec and 60°C for 1 min. Primer specificity was confirmed by melting curve analysis. Mean cycle threshold values of duplicates of each sample were normalized using two reference genes, *Actin* and *EF1-α*. Relative gene expression values were calculated using 2^{-ΔΔCt} method (Livak and Schmittgen, 2001).

Statistical analysis

All statistical analyses were conducted using R (R Core Team, 2017). Data for gene expression and larval mass of *S. frugiperda* larvae, and aphid fecundity were compared using analysis of variance (ANOVA) followed by Tukey's test or Dunnett's test relative to the GFP control or *t*-test. Gene expression data were log2 transformed before the statistical analysis to meet the assumptions of ANOVA but untransformed data are presented in the figures. Survival of *S. frugiperda* larvae was analyzed using non-parametric Kruskal-Wallis tests. Raw data underlying the bar graphs are presented in Supplemental Tables S2-S8.

Results

SCMV-GFP does not affect *S. frugiperda* growth on maize

A previously described SCMV cloning vector (Mei *et al.*, 2019) was modified to produce SCMV-CS3 (Figure 2a) and the mEGFP coding sequence (Zacharias *et al.*, 2002) was placed in the multiple cloning site to produce SCMV-*GFP*. An empty vector control virus (SCMV-EV) and SCMV-*GFP* were used to inoculate one-week-old maize seedlings. Both SCMV-*GFP* and SCMV-EV caused mosaic symptoms within two weeks after inoculation and continued to spread in newly emerging leaves of the infected plants (Figure 2b). Three weeks post-inoculation, the *GFP* expression level was significantly higher in plants infected by SCMV-*GFP* than SCMV-EV (Figure 2c). Infected leaves were examined using confocal microscopy and the green fluorescence signal was only detected in leaves infected with SCMV-*GFP* (Figure 2d). Neither *S. frugiperda* larval survival (Figure 2e) nor larval growth (Figure 2f) differed significantly between plants infected by SCMV-*GFP* and SCMV-EV. SCMV-*GFP* was used as a transgene-expressing virus control treatment in subsequent experiments to test the efficacy of insect growth-inhibiting proteins (Table 1) against *S. frugiperda*, *R. maidis*, and *M. persicae* (Figure 1a-c).

Expression of endogenous maize genes using SCMV enhances *S. frugiperda* resistance

We cloned the 69 bp sequences of the maize *Pep1* and *Pep3* defense elicitors (Huffaker *et al.*, 2011, 2013) into pSCMV-CS3 and inoculated seedlings of sweet corn variety Golden Bantam. *Pep1* and *Pep3* expression was confirmed by qRT-PCR three weeks post-inoculation (Figure 3a,b). Relative to SCMV-*GFP*, infection with SCMV-*Pep1* and SCMV-*Pep3* caused increased

transcript accumulation of maize proteinase inhibitor (*MPI*; Figure 3c), a JA pathway marker gene with antiherbivore activity (Chuang, Ray, *et al.*, 2014). *Pep1* and *Pep3* expression decreased the growth of *S. frugiperda* larvae on Golden Bantam maize by 25% and 51%, respectively, compared to GFP control plants (Figure 3d), but did not affect survival of the larvae (Table S4). Similar results were obtained for *Pep3* using popcorn (P39) and field corn (B73) inbred lines (Figure 3e,f), showing that the effect on *S. frugiperda* growth is not specific to Golden Bantam.

We targeted previously investigated maize insect resistance genes (*JAR1a*, *JAR1b*, *RIP2*, and *MPI*) and two predicted resistance genes (*AMP* and *JAC1*) for overexpression using SCMV. A previously uncharacterized gene (GRMZM2G032198) encoding a maize cyclotide antimicrobial peptides was designated as *AMP*. Expression of both *AMP* and *JAC1* was induced by *S. frugiperda* herbivory (Figure S1), suggesting that these genes are involved in maize insect resistance. The expression levels of *JAR1a*, *JAR1b*, *RIP2*, *JAC1*, *AMP*, and *MPI* were significantly higher in plants infected by the corresponding SCMV constructs than in SCMV-*GFP* control plants (Figure 3g-j,m,n). SCMV-*JAR1a* and SCMV-*JAR1b* also significantly increased *MPI* expression (Figure 3k), confirming upregulation of JA-related defense pathways. More importantly, the growth of *S. frugiperda* larvae was up to 60% lower on plants expressing *JAR1a*, *JAR1b*, *RIP2*, *JAC1*, *AMP*, and *MPI* than on Golden Bantam control plants infected with SCMV-*GFP* (Figure 3l,o).

Expression of scorpion insecticidal proteins reduces *S. frugiperda* growth

To determine whether SCMV can be used to express heterologous insect resistance genes in maize, we cloned *UyCT3* and *UyCT5*, which encode *U. yaschenkoi* venom toxins, into the SCMV vector. Transgene expression in maize was confirmed by qRT-PCR (Figure 4a,b). Both SCMV-*UyCT3* and SCMV-*UyCT5* reduced *S. frugiperda* weight on P39 and B73 plants compared to SCMV-*GFP* control plants (Figure 4c,d).

Expression of fusion proteins using SCMV has additive effects on *S. frugiperda*

Fusion of spider and scorpion neurotoxins with plant lectins can improve their toxicity (Fitches *et al.*, 2012; Liu *et al.*, 2016; Rauf *et al.*, 2019). We investigated this effect in maize using SCMV constructs. Because SCMV vectors produce a polyprotein precursor from which functional proteins are cleaved by the NIa protease (Mei *et al.*, 2019), we generated SCMV constructs with and without an NIa cleave site between the venom toxin and lectin to determine whether the proteins are more efficacious separately or as fusion proteins. The expression levels of *H. versuta* toxin (*Hvt*) and *A. cepa* agglutinin (*ACA*) genes were significantly higher in plants infected by each corresponding SCMV constructs than SCMV-*GFP* control (Figure 5a and 5b), and the presence of protease cleavage site between *Hvt* and *ACA* did not affect *Hvt* and *ACA* transcript accumulation. Expression of single proteins did not significantly decrease *S. frugiperda* growth relative to the GFP control (Figure 5c). However, the expression of fusion protein by SCMV-*Hvt-ACA* or two individual proteins of *Hvt* and *ACA* by SCMV-*Hvt-cleavage site-ACA* reduced the larval growth by 39% and 46%, relative to the GFP control (Figure 5c), respectively, suggesting an additive effect from the expression the two genes.

As the presence of a cleavage site between the two protein components did not increase efficacy (Figure 5c), we tested insecticidal activity of fusion proteins without viral protease cleavage sites in subsequent experiments. SCMV constructs were made with a maize defense gene, *RIP2*, and a spider insecticidal protein, *Hvt*, fused to *G. nivalis* agglutinin (*GNA*). We

confirmed that each gene was expressed in plants infected by the corresponding construct (Figure 5d,e,g,h). Although the expression of *GNA* alone did not affect *S. frugiperda* larval growth, infection with SCMV-*RIP2* and SCMV-*RIP2-GNA* reduced *S. frugiperda* growth by 28% and 27%, respectively (Figure 5f). In the case of *Hvt* and *GNA*, neither gene by itself significantly reduced caterpillar growth. However, the SCMV-*Hvt-GNA* fusion construct significantly decreased *S. frugiperda* larval growth compared to GFP control protein (Figure 5i). Together, these results indicate that fusion proteins combining lectins and maize defense proteins or venom toxins can improve resistance against *S. frugiperda*.

Expression of maize and scorpion genes enhances resistance to phloem-feeding herbivores

To investigate whether our SCMV constructs also provide protection against phloem-feeding insects, we conducted aphid bioassays using P39 plants infected with a subset of the previously tested constructs: SCMV-*GFP*, SCMV-*Pep3*, SCMV-*RIP2*, SCMV-*UyCT3*, SCMV-*UyCT5*, SCMV-*AMP*, and SCMV-*JAC1*. GFP expression did not affect aphid numbers compared to the empty vector control (Figure 6a,b). By contrast, expression of maize defense proteins and scorpion toxins significantly decreased progeny production by both *R. maidis* (Figure 6a,c) and *M. persicae* (Figure 6b,d) compared to SCMV-*GFP* control.

Discussion

Artificial diet assays are commonly employed for testing the oral efficacy of novel insecticidal proteins against insect herbivores (Panwar *et al.*, 2018; Yao *et al.*, 2003; Fitches *et al.*, 2012). However, growth inhibition on artificial diet does not always correlate well with the effects that are observed when the same insecticidal proteins are subsequently expressed in transgenic plants (Khan *et al.*, 2020). Both the context of the surrounding plant tissue and the localization of the insecticidal proteins in the plants could affect their toxicity against insect herbivores. Therefore, rather than pre-screening insecticidal proteins by cloning in microbial systems, purification, and artificial diet assays, we propose that transient expression using a viral vector such as SCMV will be a more effective approach for rapidly testing the *in planta* efficacy of novel insecticidal proteins.

Reduced weight gain of *S. frugiperda* in response to SCMV-mediated expression of *Pep1*, *Pep3*, *JAR1a*, *JAR1b*, *RIP2*, *MPI*, *UyCT3*, and *UyCT5* as single-gene constructs is consistent with previous reports of these genes providing protection against insect herbivory. Additionally, *AMP* and *JAC1*, two maize genes that are upregulated in response to *S. frugiperda* feeding (Figure S1), reduced caterpillar weight gain when overexpressed in maize. Although transcriptomic studies have identified numerous maize genes that are upregulated in response to arthropod feeding (Bui *et al.*, 2018; Pan *et al.*, 2020; Tzin *et al.*, 2015, 2017; Yang *et al.*, 2020; Guo *et al.*, 2019; Song *et al.*, 2017; Wang *et al.*, 2017; Zhang *et al.*, 2016), the majority of these genes have not been investigated for their role in plant defense against herbivory. This is at least in part due to the time and cost of creating maize lines that have individual genes overexpressed. Transient expression using SCMV, as we have done for *AMP* and *JAC1*, will accelerate the process of testing the defending functions of maize genes that are induced in response to herbivory.

Although these genes increased resistance in other plant-insect studies (Liu *et al.*, 2016; Ullah *et al.*, 2015; Vandenborre *et al.*, 2011), overexpression of *Hvt*, *ACA*, and *GNA* as single-gene SCMV constructs in maize did not significantly reduce *S. frugiperda* weight gain. Nevertheless, expression of gene fusions, *Hvt-ACA* and *Hvt-GNA*, reduced *S. frugiperda* weight

gain, indicating that there are additive effects of the spider venom and the lectin (Figure 5c,i). This is consistent with other studies that the toxicity of spider and scorpion toxins was improved by combining with a lectin, which may facilitate the transfer of the venom proteins across the gut lumen (Fitches *et al.*, 2002; Javaid *et al.*, 2016; Nakasu *et al.*, 2014; Rauf *et al.*, 2019). In addition to these synergistic effects on larval growth, it is likely that the stacking of multiple toxic proteins with different modes of action in one viral construct will delay the development of resistance in insects (Head *et al.*, 2017; Ni *et al.*, 2017). In contrast to Hvt-lectin fusions, we did not observe an additive effect when RIP2 was linked to GNA (Figure 5f). This difference may be attributed to the differing origins of the toxin proteins expressed in the SCMV constructs. Whereas spider venoms like Hvt are injected directly into the hemolymph, the maize RIP2 protein would be consumed orally by lepidopteran larvae. As an endogenous maize insecticidal protein, RIP2 may bind to an as yet unknown receptor and thereby enter the midgut cells and/or the insect hemolymph. Thus, these results suggest that the fusion proteins of non-maize toxins and lectins enhance the insecticidal activity of the fusion protein.

Our observation of increased defense gene expression (Figure 3c) and reduced *S. frugiperda* weight gain (Figure 3d-f) on plants infected with SCMV-*Pep1* and SCMV-*Pep3*, are consistent with experiments showing that pre-treatment of maize plants with *Pep1* and *Pep3* increases JA levels, defense gene expression, and defensive metabolites, leading to reduced growth of *S. exigua* larvae (Huffaker *et al.*, 2011, 2013). Under normal circumstances, peptide signaling is initiated when plasma membranes are disrupted and elicitor peptides are released from the cytoplasm into the apoplastic space (Bartels and Boller, 2015). Receptors on neighboring intact cells recognize these peptides and elicit downstream defense pathways (Lori *et al.*, 2015). However, in our experiments, the expression of *MPI* was induced in plants infected by SCMV-*Pep1* and SCMV-*Pep3*, even before initiation of caterpillar feeding (Figure 3c). Pathogen attack can cause plant proteins without secretory signals to be released into apoplast (Agrawal *et al.*, 2010), and is possible that SCMV infection initiated *Pep1* and *Pep3* signaling in this manner.

Induced defenses, in particular those regulated by the jasmonic acid pathway, typically are turned on after perception of insect herbivory by maize and other plants (Erb and Reymond, 2019; Howe and Jander, 2008). However, initiation of jasmonate-regulated defenses takes time and some lepidopteran herbivores may have the ability to suppress jasmonate signaling (C.-Y. Chen *et al.*, 2019). Therefore, targeted initiation of maize defense responses by expression of regulatory proteins in SCMV may be an approach for increasing pest resistance. Such virus-mediated induction could be deployed in maize fields when there is the specific threat of insect pests such as *S. frugiperda*.

Currently available insect-resistant transgenic maize varieties, in particular those expressing *Cry*, *Cyt*, or *Vip* genes from *B. thuringiensis* (Bravo *et al.*, 2011; Chakraborty *et al.*, 2016), are not effective against phloem-feeding insects such as *Rhopalosiphum maidis* (corn leaf aphid; Figure 1b) or *Myzus persicae* (green peach aphid; Figure 1c). Although the toxicity of *Bt* toxins can be increased by incorporating peptide sequences that bind to aphid guts (Chougule *et al.*, 2013), this has not yet resulted in a commercially viable *Bt* toxin directed at hemipteran herbivores. Thus, there is an interest in identifying additional insecticidal proteins that can be expressed in the plant phloem to enhance aphid resistance. Our demonstration that transient expression of both endogenous maize proteins and insecticidal proteins from other species can reduce reproduction of *R. maidis* and *M. persicae* (Figure 6), suggests that SCMV-mediated

overexpression can be used to rapidly screen proteins for their effectiveness against hemipteran pests of maize.

Our results demonstrate the utility of SCMV-mediated overexpression for screening the efficacy of proteins that reduce insect growth on maize plants. Virus-mediated transient expression assays included genes encoding maize regulatory proteins, endogenous maize defensive proteins, and non-maize insecticidal proteins. The main advantage of the SCMV overexpression system is a timeline that makes it possible to test the effectiveness of single- and multi-gene constructs in actual maize plants in only two months. Although the main focus of our efforts was a lepidopteran herbivore, *S. frugiperda*, we also showed efficacy of SCMV constructs against two aphid species, *R. maidis* and *M. persicae*, suggesting that the SCMV-mediated transient expression approach will be broadly useful for experiments with both chewing and piercing/sucking herbivores of maize.

Acknowledgements

We thank Mamta Srivastava for help with confocal microscopy. This work was supported by agreement HR0011-17-2-0053 from the Defense Advanced Research Projects Agency (DARPA) Insect Allies Program with the Boyce Thompson Institute. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of DARPA or the US Government. The US Government is authorized to reproduce and distribute reprints for Government purposes notwithstanding any copyright notation hereon.

Short legends for Supporting Material

Supplementary Figure S1. Gene expression of *AMP* and *JAC1* in P39 maize plants damaged by *Spodoptera frugiperda* caterpillars.

Supplementary Table S1. Primers used in this study.

Supplementary Table S2. Raw data for the graphs in Figure 2.

Supplementary Table S3. Raw data for the graphs in Figure 3.

Supplementary Table S4. Raw data for the graphs in Figure 4.

Supplementary Table S5. Raw data for the graphs in Figure 5.

Supplementary Table S6. Raw data for the graphs in Figure 6.

Supplementary Table S7. Raw data for the graphs in Figure S1.

References

- Agrawal, G.K., Jwa, N.-S., Lebrun, M.-H., Job, D., and Rakwal, R. (2010) *Plant secretome: unlocking secrets of the secreted proteins*. *Proteomics*, **10**, 799–827.
- Ai, Y. and Jane, J. (2016) *Macronutrients in Corn and Human Nutrition*. *Comprehensive Reviews in Food Science and Food Safety*, **15**, 581–598.
- Barbeta, B.L., Marshall, A.T., Gillon, A.D., Craik, D.J., and Anderson, M.A. (2008) *Plant cyclotides disrupt epithelial cells in the midgut of lepidopteran larvae*. *Proc. Natl. Acad. Sci. U.S.A.*, **105**, 1221–1225.
- Bartels, S. and Boller, T. (2015) *Quo vadis, Pep? Plant elicitor peptides at the crossroads of immunity, stress, and development*. *J. Exp. Bot.*, **66**, 5183–5193.
- Bass, H.W., Krawetz, J.E., OBrian, G.R., Zinselmeier, C., Habben, J.E., and Boston, R.S. (2004) *Maize ribosome-inactivating proteins (RIPs) with distinct expression patterns have similar requirements for proenzyme activation*. *J. Exp. Bot.*, **55**, 2219–2233.
- Borrego, E.J. and Kolomiets, M.V. (2016) *Synthesis and Functions of Jasmonates in Maize*. *Plants (Basel)*, **5**.
- Bravo, A., Likitvivanavong, S., Gill, S.S., and Soberón, M. (2011) *Bacillus thuringiensis: A story of a successful bioinsecticide*. *Insect Biochem Mol Biol*, **41**, 423–431.
- Bui, H., Greenhalgh, R., Ruckert, A., Gill, G.S., Lee, S., Ramirez, R.A., and Clark, R.M. (2018) *Generalist and specialist mite herbivores induce similar defense responses in maize and barley but differ in susceptibility to benzoxazinoids*. *Front Plant Sci*, **9**, 1222.
- Campos, M.L., de Souza, C.M., de Oliveira, K.B.S., Dias, S.C., and Franco, O.L. (2018) *The role of antimicrobial peptides in plant immunity*. *J. Exp. Bot.*, **69**, 4997–5011.
- Chakroun, M., Banyuls, N., Bel, Y., Escriche, B., and Ferré, J. (2016) *Bacterial Vegetative Insecticidal Proteins (Vip) from Entomopathogenic Bacteria*. *Microbiol Mol Biol Rev*, **80**, 329–350.
- Chaudhary, D. P., Kumar, S., and Yadav, O.P. (2014) *Nutritive Value of Maize: Improvements, Applications and Constraints*. In: *Maize: Nutrition Dynamics and Novel Uses* (Chaudhary, Dharam Paul, Kumar, S., and Langyan, S., eds), pp. 3–17. New Delhi: Springer India.
- Chen, C.-Y., Liu, Y.-Q., Song, W.-M., Chen, D.-Y., Chen, F.-Y., Chen, Xue-Ying, et al. (2019) *An effector from cotton bollworm oral secretion impairs host plant defense signaling*. *Proc Natl Acad Sci U S A*, **116**, 14331–14338.
- Chen, W., Shakir, S., Biggam, M., Richter, A., Fei, Z., and Jander, G. (2019) *Genome sequence of the corn leaf aphid (Rhopalosiphum maidis Fitch)*. *Gigascience*, **8**, doi: 10.1093/gigascience/giz033.
- Chougule, N.P., Li, H., Liu, S., Linz, L.B., Narva, K.E., Meade, T., and Bonning, B.C. (2013) *Retargeting of the Bacillus thuringiensis toxin Cyt2Aa against hemipteran insect pests*. *Proc Natl Acad Sci U S A*, **110**, 8465–8470.
- Chuang, W.-P., Herde, M., Ray, S., Castano-Duque, L., Howe, G.A., and Luthe, D.S. (2014) *Caterpillar attack triggers accumulation of the toxic maize protein RIP2*. *New Phytol.*, **201**, 928–939.
- Chuang, W.-P., Ray, S., Acevedo, F.E., Peiffer, M., Felton, G.W., and Luthe, D.S. (2014) *Herbivore cues from the fall armyworm (Spodoptera frugiperda) larvae trigger direct defenses in maize*. *Mol. Plant Microbe Interact.*, **27**, 461–470.

- Cordero, M.J., Raventos, D., and San Segundo, B. (1994) *Expression of a maize proteinase inhibitor gene is induced in response to wounding and fungal infection: systemic wound-response of a monocot gene*. *Plant J.*, **6**, 141–150.
- Craik, D.J., Daly, N.L., Bond, T., and Waine, C. (1999) *Plant cyclotides: A unique family of cyclic and knotted proteins that defines the cyclic cystine knot structural motif*. *J. Mol. Biol.*, **294**, 1327–1336.
- Dancewicz, K., Slazak, B., Kielkiewicz, M., Kapusta, M., Bohdanowicz, J., and Gabryś, B. (2020) *Behavioral and physiological effects of Viola spp. cyclotides on Myzus persicae (Sulz.)*. *J. Insect Physiol.*, **122**, 104025.
- Dowd, P.F., Zuo, W.N., Gillikin, J.W., Johnson, E.T., and Boston, R.S. (2003) *Enhanced resistance to Helicoverpa zea in tobacco expressing an activated form of maize ribosome-inactivating protein*. *Journal of Agricultural and Food Chemistry*, **51**, 3568–3574.
- Erb, M. and Reymond, P. (2019) *Molecular Interactions Between Plants and Insect Herbivores*. *Annu Rev Plant Biol*, **70**, 527–557.
- Fitches, E., Audsley, N., Gatehouse, J.A., and Edwards, J.P. (2002) *Fusion proteins containing neuropeptides as novel insect control agents: snowdrop lectin delivers fused allatostatin to insect haemolymph following oral ingestion*. *Insect Biochem. Mol. Biol.*, **32**, 1653–1661.
- Fitches, E.C., Bell, H.A., Powell, M.E., Back, E., Sargiotti, C., Weaver, R.J., and Gatehouse, J.A. (2010) *Insecticidal activity of scorpion toxin (ButaIT) and snowdrop lectin (GNA) containing fusion proteins towards pest species of different orders*. *Pest Management Science*, **66**, 74–83.
- Fitches, E.C., Pyati, P., King, G.F., and Gatehouse, J.A. (2012) *Fusion to Snowdrop Lectin Magnifies the Oral Activity of Insecticidal ω -Hexatoxin-Hv1a Peptide by Enabling Its Delivery to the Central Nervous System*. *PLoS ONE*, **7**, e39389.
- Food and Agriculture Organization of the United Nations (2018) *Integrated management of the fall armyworm on maize: a guide for farmer field schools in Africa*.
- Goergen, G., Kumar, P.L., Sankung, S.B., Togola, A., and Tamò, M. (2016) *First Report of Outbreaks of the Fall Armyworm Spodoptera frugiperda (J E Smith) (Lepidoptera, Noctuidae), a New Alien Invasive Pest in West and Central Africa*. *PLoS ONE*, **11**, e0165632.
- Guo, J., Qi, J., He, K., Wu, J., Bai, S., Zhang, T., et al. (2019) *The Asian corn borer Ostrinia furnacalis feeding increases the direct and indirect defence of mid-whorl stage commercial maize in the field*. *Plant Biotechnol J*, **17**, 88–102.
- Hamshou, M., Shang, C., Smagghe, G., and Van Damme, E.J.M. (2016) *Ribosome-inactivating proteins from apple have strong aphicidal activity in artificial diet and in planta*. *Crop Protection*, **87**, 19–24.
- Head, G.P., Carroll, M.W., Evans, S.P., Rule, D.M., Willse, A.R., Clark, T.L., et al. (2017) *Evaluation of SmartStax and SmartStax PRO maize against western corn rootworm and northern corn rootworm: efficacy and resistance management*. *Pest Manag. Sci.*, **73**, 1883–1899.
- Howe, G.A. and Jander, G. (2008) *Plant immunity to insect herbivores*. *Ann Rev Plant Biol*, **59**, 41–66.

- Huang, F., Qureshi, J.A., Meagher, R.L., Reisig, D.D., Head, G.P., Andow, D.A., et al. (2014) *CryIF Resistance in Fall Armyworm Spodoptera frugiperda: Single Gene versus Pyramided Bt Maize*. *PLoS One*, **9**.
- Huffaker, A. (2015) *Plant elicitor peptides in induced defense against insects*. *Current Opinion in Insect Science*, **9**, 44–50.
- Huffaker, A., Dafoe, N.J., and Schmelz, E.A. (2011) *ZmPep1, an ortholog of Arabidopsis elicitor peptide 1, regulates maize innate immunity and enhances disease resistance*. *Plant Physiol.*, **155**, 1325–1338.
- Huffaker, A., Pearce, G., Veyrat, N., Erb, M., Turlings, T.C.J., Sartor, R., et al. (2013) *Plant elicitor peptides are conserved signals regulating direct and indirect antiherbivore defense*. *PNAS*, **110**, 5707–5712.
- Javaid, S., Amin, I., Jander, G., Mukhtar, Z., Saeed, N.A., and Mansoor, S. (2016) *A transgenic approach to control hemipteran insects by expressing insecticidal genes under phloem-specific promoters*. *Sci Rep*, **6**, 34706.
- Khan, M.H., Jander, G., Mukhtar, Z., Arshad, M., Sarwar, M., and Asad, S. (2020) *Comparison of in Vitro and in Planta Toxicity of Vip3A for Lepidopteran Herbivores*. *J Econ Entomol*, **113**, 2959–2971.
- King, G.F. and Hardy, M.C. (2013) *Spider-venom peptides: structure, pharmacology, and potential for control of insect pests*. *Annu. Rev. Entomol.*, **58**, 475–496.
- Koiwa, H., Bressan, R.A., and Hasegawa, P.M. (1997) *Regulation of protease inhibitors and plant defense*. *Trends in Plant Science*, **2**, 379–384.
- Koo, A.J.K. and Howe, G.A. (2009) *The wound hormone jasmonate*. *Phytochemistry*, **70**, 1571–1580.
- Liu, S.-M., Li, J., Zhu, J.-Q., Wang, X.-W., Wang, C.-S., Liu, S.-S., et al. (2016) *Transgenic plants expressing the AaIT/GNA fusion protein show increased resistance and toxicity to both chewing and sucking pests*. *Insect Sci.*, **23**, 265–276.
- Livak, K.J. and Schmittgen, T.D. (2001) *Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2- $\Delta\Delta CT$ Method*. *Methods*, **25**, 402–408.
- Lori, M., van Verk, M.C., Hander, T., Schatowitz, H., Klauser, D., Flury, P., et al. (2015) *Evolutionary divergence of the plant elicitor peptides (Peps) and their receptors: interfamily incompatibility of perception but compatibility of downstream signalling*. *J Exp Bot*, **66**, 5315–5325.
- Luna-Ramirez, K., Skaljic, M., Grotmann, J., Kirfel, P., and Vilcinskas, A. (2017) *Orally Delivered Scorpion Antimicrobial Peptides Exhibit Activity against Pea Aphid (Acyrtosiphon pisum) and Its Bacterial Symbionts*. *Toxins*, **9**, 261.
- Macedo, M.L.R., Oliveira, C.F.R., and Oliveira, C.T. (2015) *Insecticidal Activity of Plant Lectins and Potential Application in Crop Protection*. *Molecules*, **20**, 2014–2033.
- McMullen, M., Frey, M., and Degenhardt, J. (2009) *Genetics and biochemistry of insect resistance in maize*. In: *Handbook of Maize: its Biology* (Bennetzen, J.L. and Hake, S., eds), p. 587. New York: Springer.
- Mei, Y., Liu, G., Zhang, C., Hill, J.H., and Whitham, S.A. (2019) *A Sugarcane mosaic virus vector for gene expression in maize*. *Plant Direct*, **3**, e00158.
- Mei, Y. and Whitham, S.A. (2018) *Virus-Induced Gene Silencing in Maize with a Foxtail mosaic virus Vector*. *Methods Mol. Biol.*, **1676**, 129–139.
- Meihls, L.N., Kaur, H., and Jander, G. (2012) *Natural variation in maize defense against insect herbivores*. *Cold Spring Harbor Symp Quant Biol*, **77**, 269–283.

- Mulvenna, J.P., Mylne, J.S., Bharathi, R., Burton, R.A., Shirley, N.J., Fincher, G.B., et al. (2006) *Discovery of cyclotide-like protein sequences in graminaceous crop plants: ancestral precursors of circular proteins?* *Plant Cell*, **18**, 2134–2144.
- Nakasu, E.Y.T., Edwards, M.G., Fitches, E., Gatehouse, J.A., and Gatehouse, A.M.R. (2014) *Transgenic plants expressing ω -ACTX-Hv1a and snowdrop lectin (GNA) fusion protein show enhanced resistance to aphids.* *Front Plant Sci*, **5**, 673.
- Ni, M., Ma, W., Wang, Xiaofang, Gao, M., Dai, Y., Wei, X., et al. (2017) *Next-generation transgenic cotton: pyramiding RNAi and Bt counters insect resistance.* *Plant Biotechnol. J.*, **15**, 1204–1213.
- Noonan, J., Williams, W.P., and Shan, X. (2017) *Investigation of Antimicrobial Peptide Genes Associated with Fungus and Insect Resistance in Maize.* *Int J Mol Sci*, **18**.
- Ortiz, E., Gurrola, G.B., Schwartz, E.F., and Possani, L.D. (2015) *Scorpion venom components as potential candidates for drug development.* *Toxicon*, **93**, 125–135.
- Pan, Y., Zhao, S.-W., Tang, X.-L., Wang, S., Wang, X., Zhang, X.-X., et al. (2020) *Transcriptome analysis of maize reveals potential key genes involved in the response to belowground herbivore Holotrichia parallela larvae feeding.* *Genome*, **63**, 1–12.
- Panwar, B.S., Kaur, J., Kumar, P., and Kaur, S. (2018) *A novel cry52Ca1 gene from an Indian Bacillus thuringiensis isolate is toxic to Helicoverpa armigera (cotton boll worm).* *J Invertebr Pathol*, **159**, 137–140.
- Poretsky, E., Dressano, K., Weckwerth, P., Ruiz, M., Char, S.N., Shi, D., et al. (2020) *Differential activities of maize plant elicitor peptides as mediators of immune signaling and herbivore resistance.* *Plant J*, **104**, 1582–1602.
- R Core Team (2017) *R: A Language and Environment for Statistical Computing.* Vienna, Austria: R Foundation for Statistical Computing.
- Ramsey, J.S., Wilson, A.C., De Vos, M., Sun, Q., Tamborindoguy, C., Winfield, A., et al. (2007) *Genomic resources for Myzus persicae: EST sequencing, SNP identification, and microarray design.* *BMC Genomics*, **8**, 423.
- Rauf, I., Javaid, S., Naqvi, R.Z., Mustafa, T., Amin, I., Mukhtar, Z., et al. (2019) *In-planta expression of insecticidal proteins provides protection against lepidopteran insects.* *Sci Rep*, **9**, 6745.
- Shahidi-Noghabi, S., Van Damme, E.J.M., and Smagghe, G. (2009) *Expression of Sambucus nigra agglutinin (SNA-I') from elderberry bark in transgenic tobacco plants results in enhanced resistance to different insect species.* *Transgenic Res.*, **18**, 249–259.
- Shivaji, R., Camas, A., Ankala, A., Engelberth, J., Tumlinson, J.H., Williams, W.P., et al. (2010) *Plants on constant alert: elevated levels of jasmonic acid and jasmonate-induced transcripts in caterpillar-resistant maize.* *J Chem Ecol*, **36**, 179–191.
- Song, J., Liu, H., Zhuang, H., Zhao, C., Xu, Y., Wu, S., et al. (2017) *Transcriptomics and alternative splicing analyses reveal large differences between maize lines B73 and Mo17 in response to aphid Rhopalosiphum padi infestation.* *Front Plant Sci*, **8**, 1738.
- Staswick, P.E., Tiriyaki, I., and Rowe, M.L. (2002) *Jasmonate response locus JAR1 and several related Arabidopsis genes encode enzymes of the firefly luciferase superfamily that show activity on jasmonic, salicylic, and indole-3-acetic acids in an assay for adenylation.* *Plant Cell*, **14**, 1405–1415.
- Tabashnik, B.E. and Carrière, Y. (2017) *Surge in insect resistance to transgenic crops and prospects for sustainability.* *Nature Biotechnology*, **35**, 926–935.

- Tamayo, M.C., Rufat, M., Bravo, J.M., and San Segundo, B. (2000) *Accumulation of a maize proteinase inhibitor in response to wounding and insect feeding, and characterization of its activity toward digestive proteinases of Spodoptera littoralis larvae*. *Planta*, **211**, 62–71.
- Togola, A., Meseka, S., Menkir, A., Badu-Apraku, B., Boukar, O., Tamò, M., and Djouaka, R. (2018) *Measurement of Pesticide Residues from Chemical Control of the Invasive Spodoptera frugiperda (Lepidoptera: Noctuidae) in a Maize Experimental Field in Mokwa, Nigeria*. *Int J Environ Res Public Health*, **15**.
- Tzin, V., Fernandez-Pozo, N., Richter, A., Schmelz, E.A., Schoettner, M., Schaefer, M., et al. (2015) *Dynamic maize responses to aphid feeding are revealed by a time series of transcriptomic and metabolomic assays*. *Plant Physiol*, **169**, 1727–1743.
- Tzin, V., Hojo, Y., Strickler, S.R., Bartsch, L.J., Archer, C.M., Ahern, K.R., et al. (2017) *Rapid defense responses in maize leaves induced by Spodoptera exigua caterpillar feeding*. *J Exp Bot*, **68**, 4709–4723.
- Ullah, I., Hagenbucher, S., Álvarez-Alfageme, F., Ashfaq, M., and Romeis, J. (2015) *Target and non-target effects of a spider venom toxin produced in transgenic cotton and tobacco plants*. *Journal of Applied Entomology*, **139**, 321–332.
- Van Damme, E.J.M., Zhang, W., and Peumans, W.J. (2004) *Induction of cytoplasmic mannose-binding jacalin-related lectins is a common phenomenon in cereals treated with jasmonate methyl ester*. *Commun Agric Appl Biol Sci*, **69**, 23–31.
- Vandenborre, G., Smagghe, G., and Van Damme, E.J.M. (2011) *Plant lectins as defense proteins against phytophagous insects*. *Phytochemistry*, **72**, 1538–1550.
- Vila, L., Quilis, J., Meynard, D., Breitler, J.C., Marfà, V., Murillo, I., et al. (2005) *Expression of the maize proteinase inhibitor (mpi) gene in rice plants enhances resistance against the striped stem borer (Chilo suppressalis): effects on larval growth and insect gut proteinases*. *Plant Biotechnol. J.*, **3**, 187–202.
- Wang, H., Li, S., Teng, S., Liang, H., Xin, H., Gao, H., et al. (2017) *Transcriptome profiling revealed novel transcriptional regulators in maize responses to Ostrinia furnacalis and jasmonic acid*. *PLoS One*, **12**, e0177739.
- Weidmann, J. and Craik, D.J. (2016) *Discovery, structure, function, and applications of cyclotides: circular proteins from plants*. *J. Exp. Bot.*, **67**, 4801–4812.
- Yang, L., Gao, J., Zhang, Y., Tian, J., Sun, Y., and Wang, C. (2020) *RNA-Seq identification of candidate defense genes by analyzing Mythimna separata feeding-damage induced systemic resistance in balsas teosinte*. *Pest Manag Sci*, **76**, 333–342.
- Yang, S., Pyati, P., Fitches, E., and Gatehouse, J.A. (2014) *A recombinant fusion protein containing a spider toxin specific for the insect voltage-gated sodium ion channel shows oral toxicity towards insects of different orders*. *Insect Biochemistry and Molecular Biology*, **47**, 1–11.
- Yao, J.H., Zhao, X.Y., Liao, Z.H., Lin, J., Chen, Z.H., Chen, F., et al. (2003) *Cloning and molecular characterization of a novel lectin gene from Pinellia ternata*. *Cell Research*, **13**, 301–308.
- Zacharias, D.A., Violin, J.D., Newton, A.C., and Tsien, R.Y. (2002) *Partitioning of lipid-modified monomeric GFPs into membrane microdomains of live cells*. *Science*, **296**, 913–916.

- Zhang, Y., Huang, Q., Pennerman, K.K., Yu, J., Liu, Z., Guo, A., and Yin, G. (2016) *Datasets for transcriptomic analyses of maize leaves in response to Asian corn borer feeding and/or jasmonic acid*. *Data Brief*, **7**, 1010–1014.
- Zhu, F., Zhou, Y.-K., Ji, Z.-L., and Chen, X.-R. (2018) *The Plant Ribosome-Inactivating Proteins Play Important Roles in Defense against Pathogens and Insect Pest Attacks*. *Front. Plant Sci.*, **9**.

Table 1. Insect resistance genes that were tested in maize by transience expression using *Sugarcane mosaic virus*

Source species	Gene name	GenBank ID	MaizeGDB ID	Description
<i>Zea mays</i>	<i>Pep1</i>	XM_008670579	GRMZM2G055447	Peptide elicitor 1
<i>Zea mays</i>	<i>Pep3</i>	XM_008670581	GRMZM2G339117	Peptide elicitor 3
<i>Zea mays</i>	<i>JAR1a</i>	NM_001174342	GRMZM2G091276	Jasmonate-isoleucine conjugating enzyme
<i>Zea mays</i>	<i>JAR1b</i>	NM_001361064	GRMZM2G162413	Jasmonate-isoleucine conjugating enzyme
<i>Zea mays</i>	<i>RIP2</i>	NM_001137489	GRMZM2G119705	Ribosome-inactivating protein 2
<i>Zea mays</i>	<i>JAC1</i>	NM_001148875	GRMZM2G050412	Jacalin 1, maize lectin
<i>Zea mays</i>	<i>AMP</i>	NM_001371023	GRMZM2G032198	Cyclotide antimicrobial peptide
<i>Zea mays</i>	<i>MPI</i>	X78988	GRMZM2G028393	Maize proteinase inhibitor
<i>Allium cepa</i>	<i>ACA</i>	DQ255944		Onion lectin
<i>Galanthus nivalis</i>	<i>GNA</i>	M55556		Snowdrop lectin
<i>Hadronyche versuta</i>	<i>Hvt</i>	AJ938032		ω -hexatoxin-Hv1a, spider venom toxin
<i>Urodacus yaschenkoi</i>	<i>UyCT3</i>	JX274241		Scorpion venom toxin, antimicrobial peptide
<i>Urodacus yaschenkoi</i>	<i>UyCT5</i>	JX274242		Scorpion venom toxin, antimicrobial peptide

(a) *Spodoptera frugiperda*   (b) *Rhopalosiphum maidis*   (c) *Myzus persicae*

Figure 1. Insect species used in this study. (a) *Spodoptera frugiperda*, fall armyworm, (b) *Rhopalosiphum maidis*, corn leaf aphid, (c) *Myzus persicae*, green peach aphid.

Figure 2. Use of a sugarcane mosaic virus (SCMV) to transiently express *GFP* in maize. (a) Schematic diagram of the SCMV-CS3 cloning vector. A multiple cloning site (MCS) was inserted between P1 and HC-Pro genes of SCMV. 35S: cauliflower mosaic virus 35S promoter; T: NOS terminator. (b) Mosaic infection symptom of SCMV vector in maize three weeks post inoculation. (c) Transient overexpression of *GFP* in plants infected by SCMV-empty vector (EV) or -*GFP*. Gene expression was determined by qRT-PCR three weeks post-inoculation. Means $\pm$ s.e. of N = 5, ***P < 0.001, *t*-test. (d) Confocal image of foliar GFP expression. (e, f) *S. frugiperda* survival and larval growth on plants infected by EV or SCMV-*GFP*. Two-day-old caterpillars were fed on infected plants for one week and caterpillar mass was determined. Means $\pm$ s.e. of N = 10, NS= non-significant, ND = not detected, EV = empty vector, MCS = multiple cloning site.

Figure 3. Gene expression and *Spodoptera frugiperda* larval growth on plants expressing maize defense genes. (a,b) Transient overexpression of *Pep1* and *Pep3* in Golden Bantam plants infected by SCMV-GFP, -*Pep1* or -*Pep3*. (c) The expression level of *MPI* in plants expressing SCMV-encoded GFP, *Pep1*, or *Pep3*. Gene expression was determined by qRT-PCR three weeks post inoculation. (d) Performance of *S. frugiperda* on Golden Bantam plants expressing GFP, *Pep1*, or *Pep3*. (e,f) Performance of *S. frugiperda* on P39 and B73 plants expressing GFP or *Pep3*. (g-j, m,n) Transgene expression in Golden Bantam plants infected with SCMV-GFP, -*JAR1a*, -*JAR1b*, -*RIP2*, -*JAC1*, *AMP*, or *MPI*. (k) The expression level of *MPI* in Golden Bantam plants expressing GFP, *JAR1a*, or *JAR1b*. (l, o) The performance of *S. frugiperda* on plants expressing GFP, *JAR1a*, *JAR1b*, *RIP2*, *JAC1*, *AMP*, or *MPI*. Caterpillars were confined on infected plants three weeks post inoculation and mass was measured one week later. Means $\pm$ s.e. ²⁰ N = 5 for gene expression; N = 10-12 for insect bioassays, *P < 0.05; **P < 0.01; ***P < 0.001 relative to GFP control, Dunnett's test.

Figure 4. Gene expression and larval growth on P39 plants expressing heterologous insecticidal genes. (a, b) Transient overexpression of *UyCT3* and *UyCT5* in P39 plants infected by SCMV-*GFP*, -*UyCT3* or -*UyCT5*. Gene expression was determined by qRT-PCR three weeks post inoculation. (c) The performance of *S. frugiperda* on P39 plants expressing *GFP*, *UyCT3* or *UyCT5*. (d) The performance of *S. frugiperda* on B73 plants expressing *GFP* or *UyCT3*. Five caterpillars were confined on infected plants three weeks post inoculation. Caterpillar weight was measured one week later. Means $\pm$ s.e. of $N = 5$ for gene expression; $N = 7-14$ for insect bioassays, * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ relative to *GFP* control, Dunnett's test.

Figure 5. Gene expression and larval growth on plants expressing heterologous insecticidal genes. (a, b) Transient overexpression of *Hvt* and *ACA* in plants infected by SCMV-GFP, -*Hvt*, -*ACA*, -*Hvt-ACA* or -*Hvt*-ACA*. * represents a protease cleavage site. (c) The performance of *Spodoptera frugiperda* on plants expressing GFP, *Hvt*, *ACA*, *Hvt-ACA* or *Hvt*-ACA*. (d, e) Transient overexpression of *RIP2* and *GNA* in plants infected by SCMV-GFP, -*RIP2*, -*GNA* or -*RIP2-GNA*. (f) Performance of *S. frugiperda* on plants expressing GFP, *RIP2*, *GNA*, or *RIP2-GNA*. (g, h) Transient overexpression of *GNA* and *Hvt* in plants infected by SCMV-GFP, -*Hvt*, -*GNA*, or -*Hvt-GNA*. (i) Performance of *S. frugiperda* on plants expressing GFP, *Hvt*, *GNA*, or *Hvt-GNA*. Gene expression was determined by qRT-PCR three weeks post-inoculation. Caterpillars were confined on infected plants three weeks post-inoculation and caterpillar weight was measured one week later. Means $\pm$ s.e. of N = 5 for gene expression and N = 10-12 for insect bioassays. Different letters indicate significant differences, $P < 0.05$, ANOVA followed by Tukey's HSD test.

Figure 6. Total number of aphids on P39 plants expressing maize defense genes and scorpion toxin genes. (a,c) *Rhopalosiphum maidis* (b,d) *Myzus persicae*. Eight *R. maidis* adults or ten *M. persicae* adults were confined on plants infected by SCMV-EV, -GFP, -Pep3, -RIP2, -UyCT3, UyCT5, AMP or JAC1 three weeks post inoculation. Surviving aphids and their progeny were counted one week later. Means $\pm$ s.e. of N = 8-10, *P < 0.05; **P < 0.01; ***P < 0.001 relative to GFP control, Dunnett's test.

Figure S1. Gene expression of (a) *AMP* and (b) *JAC1* in P39 maize plants damaged by *Spodoptera frugiperda* caterpillars. Two caterpillars were confined on plants. Gene expression was determined by qRT-PCR one day after placing the caterpillars. Means $\pm$ s.e. of $N = 6$, ** $P < 0.01$; *** $P < 0.001$ relative to GFP control, t -test.