

**Polymorphic SNPs, short tandem repeats and structural variants are responsible
for differential gene expression across C57BL/6 and C57BL/10 substrains**

Milad Mortazavi¹, Yangsu Ren¹, Shubham Saini², Danny Antaki^{1,3}, Celine St. Pierre⁴,
April Williams⁵, Abhishek Sohni⁶, Miles Wilkinson^{6,7}, Melissa Gymrek^{2,7,8}, Jonathan
Sebat^{*1,3,7}, and Abraham A. Palmer^{*1,7}

¹Department of Psychiatry, University of California San Diego, La Jolla, CA

²Department of Computer Science and Engineering, University of California San Diego,
La Jolla, CA

³Department of Cellular and Molecular Medicine and Pediatrics, University of California
San Diego, La Jolla, CA

⁴Department of Genetics, Washington University School of Medicine, St. Louis MO

⁵Salk Institute for Biological Studies, La Jolla, CA

⁶Department of Obstetrics, Gynecology and Reproductive Sciences, University of
California San Diego, La Jolla, CA

⁷Institute for Genomic Medicine, University of California San Diego, La Jolla, CA

⁸Department of Medicine, University of California San Diego, La Jolla, CA

*** Corresponding Authors:**

Abraham A. Palmer (aap@ucsd.edu)

Jonathan Sebat (jsebat@ucsd.edu)

Summary

Mouse substrains are an invaluable model for understanding disease. We compared C57BL/6J, which is the most commonly used inbred mouse strain, with several closely related substrains. We performed whole genome sequencing and RNA-sequencing analysis on 9 C57BL/6 and 5 C57BL/10 substrains. We identified 352,631 SNPs, 109,096 INDELS, 150,344 short tandem repeats (STRs), 3,425 structural variants (SVs) and 2,826 differentially expressed genes (DEGenes) among these 14 strains. 312,981 SNPs (89%) distinguished the B6 and B10 lineages. These SNPS were clustered into 28 short segments that are likely due to introgressed haplotypes rather than new mutations. Outside of these introgressed regions, we identified 53 SVs, protein-truncating SNPs and frameshifting INDELS that were associated with DEGenes. Our results can be used for both forward and reverse genetic approaches, and illustrate how introgression and mutational processes give rise to differences among substrains.

Keywords: C57BL/6, C57BL/10, Reduced Complexity Cross, substrain, mutational processes, gene expression, introgression, *Wdfy1*, *Lpp*, *Srp54*

1. Introduction

Since Clarence C. Little generated the C57BL/6 inbred strain a century ago, the C57BL/6J has become the most commonly used inbred mouse strain. Closely-related C57BL/10 substrains^{1,2}, which were separated from C57BL/6 in about 1937, are also commonly used in specific fields such as immunology³ and muscular dystrophy⁴. The popularity of C57BL strains has led to the establishment of many substrains (defined as >20 generations of separation from the parent colony). Among the C57BL/6 branches, the two predominant lineages are based on C57BL/6J (from The Jackson Laboratory; **JAX**) and C57BL/6N (from the National Institutes of Health; **NIH**^{5,6}). Subsequently, several additional substrains have been derived from the JAX and the NIH branches.

Genetic differences between closely-related laboratory strains have been assumed to be the result of accumulated spontaneous mutations⁷. For those that are selectively neutral, genetic drift dictates that some new mutations will be lost, others will

maintain an intermediate frequency, and others will become fixed, replacing the ancestral allele⁸. Because of historical bottlenecks and small breeding populations over many generations, fixation of new mutations can be relatively rapid.

Numerous studies have reported phenotypic differences among various C57BL/6- and C57BL/10-derived substrains, which are likely attributable to genetic variation. For C57BL/6 substrains, these differences include learning behavior⁹, prepulse inhibition¹⁰, anxiety and depression¹¹, fear conditioning¹²⁻¹⁴, glucose tolerance¹⁵, alcohol-related behaviors^{16,17}, and responses to other various drugs¹⁸⁻²¹. For C57BL/10 substrains, these differences include seizure traits²² and responses to drugs²³. Crosses between two phenotypically divergent strains can be used for quantitative trait mapping. Because crosses among closely related substrains segregate fewer variants than crosses of more divergent strains, identification of causal alleles is greatly simplified²¹. Such crosses have been referred to as a reduced complexity cross (**RCC**)²⁴ and have been further simplified by the recent development of an inexpensive microarray explicitly designed for mapping studies that use RCCs²⁵.

Whole Genome Sequencing (**WGS**) technology provides a deep characterization of Single Nucleotide Polymorphisms (**SNPs**), small insertions and deletions (**INDELS**), Short Tandem Repeats (**STRs**), and Structural Variations (**SVs**). SNPs that differentiate a few of the C57BL/6 substrains have been previously reported^{21,26}. While most SNPs are expected to have no functional consequences, a subset will; for example, SNPs in regulatory and coding regions, which can profoundly alter gene expression and function. STRs have never been systematically studied in C57BL substrains. STRs are highly variable elements that play a pivotal role in multiple genetic diseases, population genetics applications, and forensic casework. STRs exhibit rapid mutation rates of $\sim 10^{-5}$ mutations per locus per generation²⁷, orders of magnitude higher than that of point mutations ($\sim 10^{-8}$)²⁸, and are known to play a key role in more than 30 Mendelian disorders²⁹; recent evidence has underscored their profound regulatory role suggesting widespread involvement in complex traits³⁰. SVs include deletions, duplications, insertions, inversions, and translocations. SVs are individually less abundant than SNPs and STRs, but collectively account for a similar proportion of overall sequence difference between genomes³¹. In addition, SVs can have greater functional consequences because they can

result in large changes to protein coding exons or regulatory elements³². Large SVs among C57BL/6 (but not C57BL10) substrains were identified using array comparative genomic hybridization⁷, and have also been identified in more diverse panels of inbred strains using WGS³³. Although some genetic variants that differ between closely related C57BL substrains have been previously reported^{7,34-36}, a comprehensive, genomewide map of SNPs, INDELs, STRs, SVs and gene expression differences among C57BL6 and C57BL10 substrains does not exist.

In an effort to create such a resource, we performed whole genome sequencing in a single male individual from 9 C57BL/6 and 5 C57BL/10 substrains (~30x per substrain) and called SNPs, INDELs, STRs and SVs. In addition, to identify functional consequences of these polymorphisms, we performed RNA-sequencing of the hippocampal transcriptome in 6-11 male mice from each substrain, which allowed us to identify genes that were differentially expressed (**Figure 1A**). This approach has two advantages: first it provides a large number of molecular phenotypes that may be caused by substrain specific polymorphisms. Second, we assumed that the gene expression differences would often reflect the action of cis regulatory variants, making it possible to narrow the number of potentially causal mutations without requiring the creation of intercrosses.

2. Results

Processing WGS data, we identified 352,631 SNPs, 109,096 INDELs, 150,344 STRs and 3,425 SVs in nine C57BL/6 and five C57BL/10 substrains. 5.6% of SNPs and 17.2% of INDELs are singletons (only occur in one substrain). 89% of SNPs and 58% of INDELs separated the C57BL/6 and C57BL/10 branches. The fraction of variants in each category observed in different number of substrains is plotted in **Figure S1**. RNA-sequencing analysis on 106 hippocampal samples identified 16,400 expressed genes and 2,826 DEGenes (17.2%) in C57BL/6 and C57BL/10 substrains (FDR<0.05). These data are available in the **Supplementary material**.

2.1 Genetic evidence for origin of C57BL/6 and C57BL/10 substrain differences

Figure 1B shows the relationships among C57BL/6 and C57BL/10 substrains based on historical records³⁷⁻³⁹. **Figure 1C** shows a dendrogram that was produced using SNPs, STRs and bi-allelic SVs. Comparison of these two figures shows that the records about the relationships among C57BL/6 and C57BL/10 substrains are consistent with our sequencing results.

2.2 Distribution of genomic variants across the genome

The distribution of variants across the genome is shown in **Figure 1D**. Several dense clusters of variants common in all categories (SNPs, INDELS, STRs and SVs) are evident (e.g. on chromosomes 4, 8, 11 and 13 for example). The non-uniformity of these polymorphisms was inconsistent with our expectation that polymorphisms were due to new mutations and genetic drift. To further explore this observation, we examined the distribution of SNPs for each of the 14 different substrains (**Figure 1E**). This figure demonstrates that these clusters consist of a series of highly divergent haplotypes that differentiated the C57BL/6 and C57BL/10 lineages. In total 312,981 SNPs (89% of SNP variants detected in this study and 99.6% of the C57BL/6 vs C57BL/10 SNPs) reside in several C57BL/10-specific clusters that represent just 5% of the genome (28 segments on 11 chromosomes: 1, 2, 4, 6, 8, 9, 11, 13, 14, 15, 18) with a SNP density of $\sim 1/425$ bp. Across the remaining 95% of the genome SNPs do not appear to be clustered and have a density of $\sim 1/67,000$ bp (more than 100-fold less dense). We found that many of the SNPs in these intervals were present in the Mouse Genome Informatics (MGI) database ([URL](#)), which suggested they were not due to new mutations in the C57BL/10 lineage. We used the MGI database to identify strains that were similar to these 28 segments. No single strain matched all 28 segments. However, **Figure S2** shows that for 24 of the 28 segments, at least one strain in the database had greater than 90% concordance (we only considered strains for which a minimum of 300 SNPs was available in that segment). Based on these data, we hypothesize that one or more inbred or outbred mice were accidentally introduced into either the C57BL/6 or C57BL/10 lineage. Another possibility is that their last common ancestor was not fully inbred, and that these regions were differentially fixed after their separation. Most of the concordant strains have *domesticus* origin; however, two large segments on chromosomes 4 and 11 showed apparent

musculus origin. Additionally, 9,218 SNPs (2.6% of all SNPs) did not match the C57BL/6J reference genome (mm10) for all substrains, even for the sample we obtained from C57BL/6J. This may be because the cryopreserved embryo stock used for the modern C57BL/6J mice at JAX are more than 20 generations separated from the samples used to generate the mouse reference genome, GRCm38⁴⁰. The remaining SNP appeared to be uniformly distributed and are likely due to new mutations. The distributions of other variants (INDELs, STRs and SVs) mirrored SNPs and are plotted in **Figure S3**.




Figure 1. Study design, genetic distance analysis and distribution of genomic variants across the genome. **A:** This figure shows the design of our study. Mice from nine C57BL/6 and five C57BL/10 substrains were purchased from four vendors. Between six and eleven male offspring from the first generation born in our colony from each substrain were chosen for hippocampal RNA-sequencing. One male offspring per substrain was chosen for whole genome sequencing from spleen tissue. **B:** Historical development of C57BL/6 and C57BL/10 substrains is illustrated as a tree³⁷⁻³⁹. The year in which each substrain is separated from its branch is shown at its junction. **C:** Dendrogram showing the similarity of the substrains based on genomic variants including: SNPs (LD-pruned, INDELs not included), STRs and SVs.

D: Circus plot showing the SNPs, INDELS, STRs, SVs, and DEGenes across the genome for 14 C57BL/6 and C57BL/10 substrains. Regions with a high density of polymorphisms (hot spots) on chromosomes 4, 8, 9, 11 and 13 are obvious. **E:** Circus plot showing SNPs with non-reference genotypes for each substrain. This plot shows that most hot spots in panel D are due to regions where all C57BL/6 differ from all C57BL/10 substrains. A few regions where all substrains (including C57BL/6J) do not match the reference are also evident.

2.3 Identification of candidate genomic variants causing differential gene expression

We found that 2,826 of 16,400 genes (17.2%) were differentially expressed among the 14 substrains (FDR 0.05), we call these DEGenes. We assumed that many of the DEGenes were due to local (*cis*) polymorphisms.

In order to identify genomic variants that might be causally related differential gene expression, we tested all identified variants (SNPs, INDELS, STRs and SVs) in the *cis*-window (1Mb upstream of gene start and 1Mb downstream of gene end) for association with the corresponding DEGene. Specifically, we tested the association between the *cis*-variants and the median of DEGene expression by a linear regression test, using Limix⁴¹. The resulting p-values are reported in the **Supplementary material**.

Our cohort consists of 14 substrains. As expected, all variants with the same strain distribution pattern have identical p-values in the association tests. For example, the gene *Kcnc2*, which had significantly reduced expression in C57BL/6JEiJ (**Figure 2A**), there was an equally strong correlation with four SNPs, one INDEL and one STR in the *cis*-region (**Figure 2B**), and many more variants outside the *cis*-region (**Figure S4**). The INDEL was annotated as a frameshift loss-of-function variant by Variant Effect Predictor (VEP)⁴², therefore, it had a strong prior to be the causal variant. Thus, even in this small cohort, we found a number of examples in which a variant within the *cis*-window had a strong prior and therefore appeared likely to explain a DEGene. We describe several such examples in the next section. However, for the majority of DEGenes, there were no polymorphisms that had strong priors, meaning that any of the variants with the smallest p-values in the *cis*-window, or a combination of them, or *trans*-acting variants elsewhere in the genome, could be causal.

2.4 Differential expression of genes is associated with multiple categories of functional variants

Genomic variants that disrupt protein coding exons or nearby *cis*-regulatory elements have strong potential to cause differential gene expression. We investigated the causal role of variants in the *cis*-window by quantifying the strength of effects for multiple functional categories of variants. SNPs and INDELS were annotated using VEP⁴², which identified 555 loss-of-function variants (frameshift, stopgain or splice variant). SVs were annotated by intersecting with the gene features including exons, Transcription Start Site (TSS), Untranslated Regions (UTRs), promoters, enhancers, and introns. When an SV intersected with multiple types of functional elements, it was categorized according to the order mentioned above. The same gene annotations were applied to STRs. Intergenic SVs and STRs, which are defined as those that did not intersect with any gene features, were paired with the gene that had the nearest TSS. In addition, we assessed multiallelic copy number variation of genes by quantifying sequence coverage of all Segmental Duplications mapped to the mm10 reference genome ([URL](#)) that intersected with genes.

Genomic variants of the above categories that intersected with DEGenes were tested for association with gene expression by a Linear Mixed Model (LMM) using Limix⁴¹. We controlled for the complex relationships among inbred strains (a form of population structure) by using a Genomic Relatedness Matrix (GRM), derived from SNP genotypes, as a random effect, and parent strain (C57BL/6 or C57BL/10) as a fixed effect. **Figure 2C** shows the QQ plot for the p-values obtained from the data versus the uniform distribution. The black dots show the deciles of the data in each category. SegDups that intersected genes were strongly correlated with the expression of those genes, as would be expected for gene copy number variation. Loss-of-Function SNPs and INDELS also showed a significant inflation of correlated DEGenes, followed by the genic SVs. The genic STRs showed a slight inflation, which was not as significant as other variant types. The missense SNPs, intergenic SV and intergenic STR p-values followed the uniform distributions.

For each category, the p-values obtained by the LMM model are corrected by the Benjamini-Hochberg procedure to obtain an FDR. We identified 53 significant (FDR < 0.05) associations between DEGenes and features, which are reported in **Tables S2 and S3**. The majority of associations (41 of 53 genes) reflected segmental duplications. In **Table S3** we report the genotype pattern in the substrains for each variant; notably, there

are several clusters of significant associations with the same genotype pattern. For example, one extensive region on Chromosome 2 that clearly distinguishes C57BL/6NJ from all other substrains accounts for 18 of the 53 genes identified. Another cluster with a more complex genotype pattern on Chromosome 4 accounts for 11 of the 53 identified genes.




Figure 2. Association of gene expressions with genomic variants. **A:** Expression of *Kcnc2* is lower for C57BL/6JEIj compared to the other substrains. **B:** *Cis*-variants of *Kcnc2* are tested for association with the median expression by the linear regression model. One INDEL is a frameshift loss-of-function variant and therefore has a strong prior to be the causal variant. In addition to that, four SNPs and one STR also have the same strain distribution pattern, and therefore the same $-\log_{10}(p)$ value. For most of the DEGenes, no variant belonged to a class that had a strong prior for causality. In those cases, any of the variants with the smallest p-values (or a combination thereof) are equally likely to be causal. **C:** The distribution of the p-values of variants in different categories is compared against the uniform distribution in a QQ plot. A Linear Mixed Model is used with the Genomic Relatedness Matrix (GRM) as a random effect to control for population structure and the parental strain (C57BL/6 or C57BL/10) is used as a fixed effect to identify associations within C57BL/6 and C57BL/10 substrains. The SegDup category includes associations between the copy number variation of the DEGenes intersecting with SegDup regions (obtained by read depth across the segmental duplication regions of the reference genome) and the gene expression. Loss-of-function and missense mutations are two categories of SNP/INDELS. Genic SVs include those intersecting with gene features such as exons, TSS, UTRs, promoters, enhancers and introns, and genic STRs include those intersecting with exons, TSS, 5'UTRs and promoters. Intergenic SVs and STRs are those not intersecting with any gene features, and are paired with a gene with the closest TSS.

2.5 Distinct mechanisms of differential gene expression caused by SVs

SVs can affect gene function by (1) varying the dosage of a full-length gene (2) deletion or insertion of exons producing alternative isoforms of a gene, or (3) rearrangement of the cis-regulatory elements of genes. For example, there are three copies of the gene *Srp54* in the mouse reference genome, but we found significant variability in the number of copies across the substrains; the number of copies was strongly associated with expression of *Srp54* (**Figure 3A**). Thus, in this example, copy number variation in highly mutable segmental duplication regions is the likely cause of differential gene expression. An example of a SV that likely impacts expression is the *Lpp* gene. The *Lpp* gene has a tandem duplication of the first two exons in two substrains (C57BL/10ScCr and C57BL/10ScNHsd) that creates two copies of the TSS, which probably accounts for its ~2-fold increased expression (**Figure 3B**).

An intriguing example of altered expression caused by a SV is the *Wdfy1* gene. This gene has a tandem duplication of exons 4-6 in the C57BL/6J substrain, which is also present in the mm10 reference genome. We found this duplication is associated with a paradoxical decrease in *Wdfy1* gene expression (**Figure 3C**). Given that a frameshift is incurred by this tandem duplication, we hypothesized that reduced *Wdfy1* expression is caused by the nonsense-mediated RNA decay (NMD) pathway. This highly conserved RNA-turnover pathway promotes the turnover of mRNAs harboring premature termination codons, including those generated by frameshifts⁴³. We reasoned that if a major spliced isoform of *Wdfy1* contains this tandem duplication and associated frameshift, its decay by NMD would be detectable by examining expression in NMD-deficient cells. In support, RNA-seq analysis showed that NMD-deficient (*Upf2*^{-/-}) C57BL/6J ES cells expressed ~2-fold higher levels of *Wdfy1* than control (sibling) C57BL/6J ES cells (**Figure 3F**). Analysis of splice junctions from RNA-seq analysis confirmed the existence of an aberrant isoform in all C57BL/6J lines that includes splicing of exon 6 to the downstream (duplicated) exon 4 (**Figure 3D**), which we refer to as the “6a->4b” junction. This splice junction was unique to C57BL/6J strains, and the ratio of 6a->4b to all splice junctions was increased by ~2-fold in the *Upf2*^{-/-} ES cells relative to control ES cells (**Figure 3E**). These results demonstrate that major isoforms transcribed from *Wdfy1* in C57BL/6J mice are degraded by NMD, and the transcripts that are retained are alternative splice forms that exclude the 6a->4b junction.




Figure 3. Structural variations affecting gene expression. Structural variations in *Srp54*, *Lpp* and *Wdfy1* associate with gene expression of these three genes. **A:** Various copy numbers in a SegDup region in different substrains is associated with expression of *Srp54*. The read coverage in these SegDup regions is used to infer the number of copies of the intersecting genes. **B:** A duplication involving the first two exons of *Lpp* in two substrains of C57BL/10 is associated with increasing the gene expression. Duplicating the TSS and the promoter site seems to be the cause of enhancing the expression level of *Lpp*. **C:** A segmental

duplication region which intersects with three exons in *Wdfy1*, and is present in C57BL/6J and the mouse reference genome, is associated with reduction of expression of *Wdfy1* in C57BL/6J. All the other substrains lack this duplication. **D:** Sashimi plots for C57BL/6BomTac (a closely related substrain to C57BL/6J), C57BL/6J, *Upf2*^{+/+}, and *Upf2*^{-/-} cell lines from C57BL/6J highlighting the junction between 6a and 4b exons across the segmental duplication region. Since C57BL/6BomTac lacks the duplication, it does not have any junctions between those exons, while the relative number of junctions in the *Upf2*^{-/-} cell line is significantly larger than the other wild type C57BL/6J samples. **E:** The bar plot shows the ratio of the normalized number of junctions between 6a and 4b exons (normalized by the total number of junctions in each sample) in *Upf2*^{-/-} over *Upf2*^{+/+} cell lines. It shows a significant increase in the relative number of junctions between the two segmental duplications in the *Upf2*^{-/-} cell line. The numbers on top of the bars show p-values obtained by the Chi-squared test. **F:** The expression level of *Wdfy1* in the *Upf2*^{-/-} cell line is significantly higher than the *Upf2*^{+/+} cell line. This supports our hypothesis that the reduction of gene expression in C57BL/6J is due to the nonsense mediated decay (NMD) mechanism.

3. Discussion

We have performed a large-scale multi-omics analysis of 14 C57BL substrains. We identified 352,631 SNPs, 109,096 INDELs, 150,344 STRs and 3,425 SVs; furthermore, of the 16,400 genes that were expressed in the hippocampus, 2,826 were significantly differentially expressed (FDR<0.05). Unexpectedly, many of the polymorphisms that differentiated the C57BL/6 and C57BL/10 substrains were concentrated in a few haplotypes, comprising just 5% of the genome. These polymorphisms appear to be due to either introgression of an unrelated individual, or incomplete inbreeding at the time that the C57BL/6 and C57BL/10 lineages diverged. Setting these introgressed regions aside, we tried to identify variants that were causally related to differential gene expression by focusing on the *cis* regions around DEGenes. This allowed us to identify 53 genes in which a variant with high prior probability to be causal was significantly associated with gene expression. While the majority of these 53 instances were caused by segmental duplications, several of which spanned many adjacent genes, a smaller proportion were due to SVs and INDELs (see **Supplemental Tables S2** and **S3**). Inflation of test statistics for these categories of variants further underscores their likely causal roles and highlights the fact that a relaxed FDR threshold would have identified more than 53 variant/DEGene associations.

An unexpectedly large subset of variants (89% of all SNPs) were concentrated in 28 highly diverged haplotypes that were present in all C57BL/10 strains and represented just 5% of the genome. These dense clusters of genetic variation (1 SNP/425 bp) perfectly differentiated C57BL/10 from C57BL/6, and likely reflect introgression from another strain.

Intriguingly, the smaller haplotypes appeared to be of *domesticus* origin, and were similar to haplotypes found in multiple non-C57BL inbred strains. The two largest haplotypes appeared to be of *musculus* origin and were also similar to multiple non-C57BL inbred strains. The exact sequence of events that led to this situation are impossible to deduce, but these patterns are clearly due to breeding errors rather than spontaneous mutations; this conclusion is based on several observations: 1) the density of the polymorphisms, 2) the abrupt boundaries of the regions/haplotypes and 3) the fact that the SNPs in these introgressed regions are found in other inbred strains, which would not be the case if they were due to spontaneous mutations. A previous microarray study performed on 198 inbred mouse strains also identified SNP differences between C57BL/6J and three C57BL/10 substrain (C57BL/10J, C57BL/10ScNJ and C57BL/10ScSnJ) for all the 28 introgressed segments that we identified^{44,45}; however, that study did not highlight the significance of that finding, and did not have sufficiently dense coverage to define the boundaries of the introgressed regions. While a majority of C57BL/10-specific genetic variants lie within these introgressed regions, they contained only a small fraction (~13%) of DEGenes; however, given that the introgressed regions represent only 5% of the genome, this is still about 2.6-fold greater density of DEGenes that would be expected if they were randomly distributed across the genome.

Outside of these apparently introgressed regions we identified 37,745 SNPs that were distributed throughout the genome in a Poisson fashion with more than 100-fold lower density (~1 SNP/67,000 bp). These SNPs are apparently due to the accumulation of new mutations and their identification was the original goal of our study. Dendrograms based on these SNPs recapitulated the historically recorded relationships among the substrains (**Figure 1B**). For the relatively large number of DEGenes (>2,000) that were located outside of the introgressed regions, we considered the association between different categories of nearby (*cis*) variants, and expression of DEGenes. Variable copy number segmental duplicated regions were shown to be highly enriched for significant associations as were genic SVs and loss of function SNP/INDELs (**Figure 2C**).

We presented several examples to highlight how different classes of variants underlie DEGenes. For example, variable copy number segmental duplications led to both increased and decreased expression of *Srp54* (**Fig 3A**). In another example,

duplication of transcription start sites led to increased expression of *Lpp* (**Fig 3B**). In the case of *Wdfy1*, duplication of several exons led to down-regulation of expression (**Fig 3C**), which we showed was due to NMD-mediated mRNA decay (**Fig 3D**). *Wdfy1* was previously reported to be differentially expressed between C57BL/6J and C57BL/6NCrl, and was identified as one of the candidate genes for reduced alcohol preference in C57BL/6NCrl⁴⁶. This gene is also within the QTL named *Emo4* (location: Chr1:68,032,186-86,307,305 bp, [URL](#)); mice which are homozygous for C57BL/6J allele are more active in the open field test. Whether *Wdfy1* is actually the cause of either association cannot be resolved by our study.

Despite the numerous examples in which likely causal variants were identified, a majority of the causal variants underlying DEGenes remain unknown. Many are likely to be due to variants in regulatory regions that have not been distinguished from other nearby variants with the same strain distribution pattern (and thus identical p-values). Although we focused on the possibility that DEGenes were due to nearby variants (*cis*-eQTLs), the large fraction of differentially expressed genes (17.2% of all expressed genes) could indicate that many DEGenes are due to *trans*-eQTLs. Producing crosses between pairs of strains will be necessary to address the relative importance of *cis*- versus *trans*-eQTLs in the observed DEGenes; it is possible that such crosses could identify one or more major *trans*-regulatory hot spots.

Our results create a resource for future efforts to identify genes and causal polymorphisms that give rise to phenotypic differences among C57BL strains using the increasingly popular reduced complexity cross (**RCC**) approach in which two phenotypically divergent nearly isogenic inbred substrains are crossed to produce an F₂ population²⁴. Because of the low density of polymorphisms, identifying the causal allele is much more tractable. For example, the gene *Cyfp2* was identified as the cause of differential sensitivity to cocaine and methamphetamine in a cross between C57BL/6J and C57BL/6N²¹. In the **Supplementary material** we have provided genomic variants (SNPs, INDELs, STRs and SVs), differentially expressed genes in the hippocampus, as well as association tests between DEGenes and nearby variants. In addition, we have provided the VEP annotated SNP/INDELs, which distinguishes loss of function, missense and synonymous mutations. Our data also identify some regions that have a high density

of polymorphisms that may complicate the RCC approach. For example, phenotypic differences between C57BL/6 and C57BL/10 strains might frequently map to the introgressed regions, which have a high density of polymorphisms that would significantly hinder gene identification and negate the advantage of RCCs. Furthermore, crosses between two C57BL/6 or between two C57BL/10 strains may map to large segmental duplication regions such as those on Chromosomes 2 and 4 (see **Figure 1E** and **Supplemental Table S3**), which would again hinder gene identification. Thus, one key observation from this study is that genetic differences among C57BL/6 and C57BL/10 strains are not uniformly distributed. Furthermore, our study used a single individual to represent each strain for whole genome sequencing. Therefore, we did not explore the extent to which the polymorphic regions we identified may be segregating versus fixed within each inbred strain. If some of these polymorphic regions are not fixed, it would further complicate the analysis of RCCs.

Whereas the RCCs represent a forward genetic approach (starting with a phenotypic difference, searching for the genetic cause), another novel application of our dataset would be to select two strains that are divergent for a coding or expression difference and to use that cross to study gene function. This reverse genetic approach (starting with a genetic difference, searching for the phenotypic consequences) has not been attempted using closely related substrains, but is conceptually similar to characterization of a knockout mouse. This approach is limited by the available polymorphisms. Although it would be necessary to account for the impact of linked polymorphisms, most of the polymorphisms would be unlinked and would not confound the interpretation of results.

In summary, we have created a dataset that elucidates the differences among C57BL strains and can be used for both forward genetic (RCC) and reverse genetic approaches. We identify previously unknown introgressed segments that differentiate the C57BL/6 and C57BL/10 lineages. Our results can also be used to explore mutational processes and highlight the tendency of inbred strains to change over time due to mutational processes.

Acknowledgements: M.M., Y.R., C.L.S.P., A.W. and A.A.P. were supported by P50DA037844. Additionally, Y.R. was supported by T32MH018399 and A.W. was supported by T32MH020065.

Author Contributions

A.P. designed the study. C.L.S.P. performed the animal breeding, dissection and the preparation of WGS and RNASeq libraries. Y.R. and A.W. performed initial analyses of the WGS and RNASeq data. M.M. carried out all statistical genetic and functional genomic analyses. M.G. and S.S. performed STR calling. M.M. and J.S. performed SV calling and analysis of SV eQTLs. M.W. developed the *Upf2* mouse model, A.S. derived the ES cells and performed the corresponding RNA-Seq. M.M., J.S., A.P. wrote the paper.

Declaration of interest

The authors declare no competing interests.

4. Methods

4.1 Mice

We obtained a panel of 14 C57BL substrains from four vendors. The panel included 9 C57BL/6 substrains: C57BL/6J, C57BL/6NJ, C57BL/6ByJ, C57BL/6NTac, C57BL/6JBomTac, B6N-TyrC/BrdCrCrI, C57BL/6NCrI, C57BL/6NHsd, C57BL/6JEiJ, and 5 C57BL/10 substrains: C57BL/10J, C57BL/10ScCr, C57BL/10ScSnJ, C57BL/10SnJ, C57BL/10ScNHsd (**Table 1**). All of the substrains were bred for one generation at the University of Chicago before tissue was collected for whole genome sequencing and RNA-sequencing; this avoided gene expression differences that were secondary to environmental differences among the four vendors. All procedures were approved by the University of Chicago IACUC. One hundred and ten male mice in total, with six to eleven mice per substrain, were chosen for RNA-sequencing from hippocampus, and one male mouse per substrain was chosen for whole genome sequencing from spleen (**Figure 1A**).

4.2 Whole-genome sequencing (WGS)

DNA from one male animal per substrain (n=14) was extracted from spleens using a standard “salting-out” protocol. Sequencing libraries were prepared using a TruSeq DNA LT kit, as per the manufacturer’s instructions. Subsequently, sequencing data was generated by Novogene at an average depth of ~30X coverage on an Illumina HiSeq X Ten (paired-end 150bp) (**Table 1**).

Table 1. Strain IDs, vendors and results from sequencing and RNA-sequencing.

Strain	Vendor	Strain ID	Whole Genome Sequencing		RNA-sequencing	
			Reads	Coverage	Average reads	# samples
C57BL/6NTac	Taconic	B6	336,470,599	33.65	13,459,345	8
C57BL/6NJ	JAX	#005304	333,094,625	33.31	15,767,477	9
C57BL/6NHsd	Harlan	#044	349,305,241	34.93	14,656,480	7
C57BL/6NCrl	Charles River	#027	301,990,053	30.20	19,408,390	8
C57BL/6JEiJ	JAX	#000924	442,604,628	44.26	16,334,723	8
C57BL/6JBomTac	Taconic	B6JBom	309,053,911	30.91	17,340,684	7
C57BL/6J	JAX	#000664	326,208,826	32.62	18,555,108	8
C57BL/6ByJ	JAX	#001139	294,061,881	29.41	15,552,591	9
B6N-TyrC/BrdCrCrl	Charles River	#493	316,655,519	31.67	15,139,469	7
C57BL/10SnJ	JAX	#000666	284,191,454	28.42	14,012,862	6
C57BL/10ScSnJ	JAX	#000476	285,201,231	28.52	18,947,235	7
C57BL/10ScNHsd	Taconic	#046	326,586,425	32.66	12,453,407	11
C57BL/10ScCr	JAX	#003752	289,068,061	28.91	16,384,792	8
C57BL/10J	JAX	#000665	311,292,991	31.13	17,550,456	7
Average			321,841,818	32.18	16,013,858	7.86

4.3 RNA-sequencing and data processing

Total RNA was extracted from 110 hippocampal samples using Trizol reagent (Invitrogen, Carlsbad, CA). RNA was treated with DNase (Invitrogen) and purified using RNeasy columns (Qiagen, Hilden, Germany). RNA-sequencing library prep and sequencing was performed by the University of California San Diego Sequencing Core using Illumina TruSeq prep and Illumina HiSeq 4000 machine (single-end 50bp; **Table 1**). Reads were mapped to mouse reference transcriptome (mm10) using the splice-aware alignment software HiSat2⁴⁷, and counts were normalized using HTSeq⁴⁸. Only genes that had at least one Count Per Million (CPM), for at least two samples were included in our analysis. We further removed four outlier samples identified by PCA analysis. This left us with gene expression data for 16,400 genes across 106 samples in 14 substrains.

To identify Differentially Expressed Genes (DEGenes) we performed analysis of variance using the *anova* function in R, and adjusted the p-values by computing the false-

discovery rate (FDR) using the *p.adjust* function in R, with the Benjamini-Hochberg procedure. We obtained 2,826 DEGenes among C57BL/6 and C57BL/10 substrains combined, 1,210 DEGenes within C57BL/6, and 104 DEGenes within C57BL/10 substrains with FDR<0.05.

4.3.1 Nonsense mediated decay assay

To determine whether SVs of the *Wdfy1* gene in C57BL/6J create novel mRNA isoforms that are degraded by the Nonsense-Mediated Decay (NMD) pathway, we performed RNA-seq on mouse embryonic stem cells (mESCs) from a *Upf2*^{-/-} strain of C57BL/6J that has impaired NMD and control mouse mESCs from C57BL/6J. Samples with an RNA integrity index of >8 (as determined by a BioAnalyzer) were used for RNA-seq analysis. The University of California San Diego Sequencing Core performed library preparation using ribosomal RNA depletion protocol followed by paired-end sequencing (100 cycles) using a HiSeq4000. Reads from three replicates of *Upf2*^{-/-} samples and three controls were mapped to the mouse reference genome (mm10) by HiSat2⁴⁷, and counts were normalized using HTSeq⁴⁸. We kept all genes with CPM>1 and normalized the counts with *edgeR* function in R, however, we only analyzed *Wdfy1* expression in an effort to detect differences in NMD between *Upf2*^{-/-} and control samples.

4.4 SNPs and INDELS

We used SpeedSeq⁴⁹ to process the WGS paired-end reads. SpeedSeq uses BWA-mem (v.0.7.8) to map the reads to the mm10 reference genome, SAMBLAST⁵⁰ to mark duplicates, Sambamba⁵¹ to sort the BAM files, and FreeBayes⁵² to jointly call SNPs and INDELS. INDELS are defined as insertions or deletions which are relatively short in length. The length range for the detected INDELS in our study is between one and 64 base pairs, which is approximately the lower bound for SV length scales. We restricted our analysis to variants that were fixed within individual substrains by including homozygous SNPs and INDELS only, resulting in a callset consisting of 352,631 SNPs and 109,096 INDELS. These variants are provided in the **Supplementary material**.

When computing the identity-by-state (IBS) matrix for dendrograms, we LD-pruned the SNP panel with Plink⁵³ (--indep-pairwise 50 5 0.5) yielding 16,739 SNPs. This pruned

SNP set was augmented by STRs and all bi-allelic SVs, followed by computing the distance matrix with *dist* and plotting the dendrograms with *hclust* in *R* v3.6.1.

4.5 Short Tandem Repeat (**STR**)

We used HipSTR v0.6 with default parameters⁵⁴ to call STRs from mapped reads using the mm10 reference STR set available from the HipSTR website ([URL](#)). The reference STR set was generated using Tandem Repeats Finder⁵⁵ allowing a maximum repeat unit length of 6bp. STRs for the substrains were jointly genotyped on a single node of a local server in batches of 500 STRs. Resulting VCF files from each batch were merged to create a genome-wide callset in VCF format. We filtered out calls with missing genotypes, as well as calls with reference alleles for all substrains, resulting in a total of 150,344 polymorphic STRs. The STR calls are available in the **Supplementary material**.

4.6 Structural Variations (**SV**)

SVs were detected using a combination of approaches. First, we called SVs with LUMPY⁵⁶ and CNVnator⁵⁷, two complementary methods that rely on discordant and split read signals or coverage respectively. Second, because SV calling accuracy by the above methods is low in regions that are dense in segmental duplications, copy number variation within annotated segmental duplications was quantified directly from coverage, and these coverage values were used for the correlation of gene copy numbers with gene expression.

We filtered out SV calls that overlapped 50% or more with the gap regions of the mouse reference genome, as well as the calls with length smaller than 50 bp and larger than 1 Mbp. A more stringent >1000 bp length filter was applied to CNVnator calls. We then filtered out non-homozygous calls and calls that were homozygous for the alternative allele in all substrains.

Concordant calls from LUMPY and CNVnator with 50% or greater reciprocal overlap and the same genotypes were merged and the breakpoints reported by LUMPY were used. Consensus calls that overlapped with annotated segmental duplications (SegDup) in the reference genome were excluded, and instead SegDup copy number was assessed directly from read depth signal using *mosdepth* v0.2.6⁵⁸ with window size

100 bp. SegDup annotations from the mm10 genome with at least 98% similarity were intersected with gene annotations, and the median read coverage across SegDups which intersect with genes was normalized by the median coverage of the corresponding chromosome. These normalized coverage values were used to correlate gene copy numbers with gene expression. The final set of SVs included 3,425 deletions, duplications and inversions in nine C57BL/6 and five C57BL/10 substrains. The distribution of SVs in each category and substrain is summarized in **Table S1**. The VCF file of the SV calls, and the read coverage data for the SegDup regions are provided in the **Supplementary material**.

4.7 Resource availability

4.7.1 Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Milad Mortazavi (miladm@alumni.stanford.edu).

4.7.2 Material availability

This study does not generate new unique reagents.

4.7.3 Data and code availability

The datasets (supplementary material) generated during this study are available at Mendeley Data, DOI: 10.17632/k6tkmm6m5h.1

5. Supplemental information

Strain	Substrain	CNVnator			Lumpy	
		DEL	DUP	DEL	DUP	INV
C57BL/6	C57BL/6J	27	461	11	97	2
	C57BL/6NJ	5	456	65	105	2
	C57BL/6ByJ	11	426	82	90	5
	C57BL/6JEiJ	14	469	42	91	4
	C57BL/6NTac	27	465	64	77	4
	C57BL/6NCrI	9	420	66	92	2
	C57BL/6NHsd	127	488	73	86	3
	B6N-TyrC/BrdCrCrI	7	445	63	96	3
	C57BL/6JBomTac	13	447	50	78	3
C57BL/10	C57BL/10ScSnJ	95	430	1204	129	15
	C57BL/10SnJ	89	420	1195	125	13
	C57BL/10ScCr	269	481	1196	103	15
	C57BL/10J	74	437	1193	115	16
	C57BL/10ScNHsd	72	426	1192	123	14
	TOTAL	448	1308	1369	279	21

Table S1. Number of SVs found in substrains of C57BL/6 and C57BL/10.




Figure S1. Fraction of variants observed in substrains (five C57BL/10 and nine C57BL/6 substrains) in each variant category. The spike at 5 reflects polymorphisms that separated C57BL/10 (n=5) from C57BL/6 (n=9) substrains. The smaller spike at 14 represents instances where none of the substrains (including C57BL/6J, which is the basis for mm10) matched the mm10 reference genome.




Figure S2. Concordance of 24 C57BL/10-specific haplotypes (SNP hotspots) with SNPs of other strains from *domesticus* and *musculus* origin. Y-axis shows segments with C57BL/10-specific SNP hotspots. X-axis shows strains which have at least 300 common loci and at least 90% concordance with C57BL/10-

specific SNPs in each segment. The SNP data for the strains is obtained from MGI ([URL](#)). The segments are color coded with the concordance value. The strains on the x-axis are color coded with blue: *domesticus* origin, and red: *musculus* origin⁴⁴.




Figure S3. A: SNP distribution, **B:** INDEL distribution, **C:** STR distribution, and **D:** SV distribution for nine C57BL/6 and five C57BL/10 substrains show clusters of variants which are specific to C57BL/10 substrains on chromosomes two, four, eight, nine, eleven, thirteen, fourteen and fifteen.




Figure S4. Association tests of median expressions of C57BL/6 and C57BL/10 substrains with all genomic variants (SNPs, INDELS, STRs and SVs) are performed by linear regression model with Limix⁴¹ **A:** Association of DEGene median expressions with all variants (SNPs, INDELS, STRs and SVs) in the *cis*-region defined as 1Mb upstream and 1Mb downstream of the DEGene. The p-values are plotted at the genomic locations of the corresponding DEGenes. **B:** Association of *Kcnc2* expression with all genomic variants across the genome shows that variants with the same strain distribution pattern have identical p-values. The flat horizontal line at about $-\log_{10}(p)=8.4$ reflects features that have the same strain distribution pattern and therefore all yield identical p-values when tested for association with the gene expression data.

Gene	Variant type	feature	Chrom	Start	End	FDR
<i>Tlr4</i>	SV_DEL	exon	chr4	66815109	66889841	2.73E-02
<i>Sh2b3</i>	SV_DUP	exon	chr5	121767693	121843545	2.73E-02
<i>Arf5</i>	SV_DUP	exon	chr6	28420693	28493886	3.05E-02
<i>Gcc1</i>	SV_DUP	exon	chr6	28420693	28493886	3.21E-02
<i>Lpp</i>	SV_DUP	exon	chr16	24378934	24444153	4.53E-02
<i>Btaf1</i>	SV_DUP	exon	chr19	36911370	37379559	4.53E-02
<i>Fgfbp3</i>	SV_DUP	exon	chr19	36911370	37379559	2.73E-02

<i>Ide</i>	SV_DUP	exon	chr19	36911370	37379559	2.73E-02
<i>Crb1</i>	INDEL	frameshift	chr1	139237085	139237087	2.02E-02
<i>Tmem108</i>	INDEL	frameshift	chr9	103499764	103499767	4.25E-02
<i>Kcnc2</i>	INDEL	frameshift	chr10	112455818	112455821	2.02E-02
<i>Clec16a</i>	INDEL	frameshift	chr16	10741835	10741842	4.25E-02
<i>Wdfy1</i>	SD		chr1	79715430	79722709	1.63E-03
<i>Gm14444</i>	SD		chr2	174958961	175012684	1.09E-02
<i>Gm14391</i>	SD		chr2	175176236	175476161	3.35E-03
<i>Gm4631</i>	SD		chr2	175176236	175476161	1.63E-03
<i>Gm6710</i>	SD		chr2	175600000	175900699	2.60E-03
<i>Gm14288</i>	SD		chr2	175675829	175804756	1.95E-03
<i>Zfp966</i>	SD		chr2	175926930	175959022	4.45E-03
<i>Zfp973</i>	SD		chr2	176598872	176698447	2.38E-03
<i>Gm14305</i>	SD		chr2	176702042	176745817	1.67E-03
<i>Gm14295</i>	SD		chr2	176742479	176876976	1.63E-03
<i>Gm14296</i>	SD		chr2	176834443	177060914	1.63E-03
<i>Gm14401</i>	SD		chr2	176936335	177079562	2.50E-03
<i>Gm14410</i>	SD		chr2	177131439	177214443	1.63E-03
<i>Gm14418</i>	SD		chr2	177332425	177443112	1.63E-03
<i>Zfp970</i>	SD		chr2	177382170	177490852	1.63E-03
<i>Gm14322</i>	SD		chr2	177714087	177825086	1.91E-03
<i>Gm14325</i>	SD		chr2	177837513	177900857	1.63E-03
<i>Gm14326</i>	SD		chr2	177900735	177965213	1.63E-03
<i>Zfp971</i>	SD		chr2	177954479	178027444	1.63E-03
<i>Gm13302</i>	SD		chr4	41882892	42033952	1.63E-03
<i>Gm20878</i>	SD		chr4	41882892	42033952	1.63E-03
<i>Gm21093</i>	SD		chr4	41882892	42033952	1.63E-03
<i>Gm37530</i>	SD		chr4	41888084	42115916	1.63E-03
<i>Gm13306</i>	SD		chr4	41967677	42175681	3.35E-03
<i>Gm17167</i>	SD		chr4	42035272	42214541	1.63E-03
<i>Gm10600</i>	SD		chr4	42232133	42238731	1.63E-03
<i>Gm13301</i>	SD		chr4	42232133	42238731	1.63E-03
<i>Gm5859</i>	SD		chr4	42232133	42238731	1.63E-03
<i>Gm12394</i>	SD		chr4	42652191	42844101	1.63E-03
<i>Gm21955</i>	SD		chr4	42715617	42730325	1.34E-02

SD_chr17_6253548_6260484	<i>Tmem181a</i>	
SD_chr17_6325771_6473130	<i>Tmem181b-ps</i>	
SD_chr17_6497992_6645878	<i>Dynlt1c</i>	
SD_chr17_15014588_15045963	<i>Ermard</i>	
SD_chrX_169969770_170000000	<i>G530011O06Rik</i>	
SD_chrX_169969770_170000000	<i>Gm15726</i>	

Table S3. Genotype patterns for variants with significant association with DEGene expression. For bi-allelic variants (SVs and SNP/INDELs), red and blue colors represent two genotypes, while for multiallelic copy number variants the normalized read depth varies between 0 and 1 where 0: blue, 0.5: white, and 1: red represent three genotypes. The same genotype patterns are clustered together for chromosomes such as 2 and 4, which shows that nearby genes in these regions have been affected by the same copy number variation patterns. Bold horizontal lines segregate nearby variants with similar genotype patterns.

References

1. Festing, M.F.W. (1979). Inbred strains in biomedical research (United Kingdom: Macmillan Education, Limited).
2. Lyon, M.F., Searle, A.G., and International Committee on Standardized Genetic Nomenclature for Mice. (1989). Genetic variants and strains of the laboratory mouse, 2nd edn (Oxford England ; New York; Stuttgart: Oxford University Press ; G. Fischer Verlag).
3. von Kockritz-Blickwede, M., Rohde, M., Oehmcke, S., Miller, L.S., Cheung, A.L., Herwald, H., Foster, S., and Medina, E. (2008). Immunological mechanisms underlying the genetic predisposition to severe *Staphylococcus aureus* infection in the mouse model. *The American journal of pathology* 173, 1657-1668.
4. Kincaid, A. (2007). Muscular Dystrophy. In *xPharm: The Comprehensive Pharmacology Reference*, S.J.E.a.D.B. Bylund, ed. (Elsevier Inc.).
5. Bailey, D.W. (1978). Sources of Subline Divergence and Their Relative Importance for Sublines of Six Major Inbred Strains of Mice. In *Origins of inbred mice*, H.C. Morse, ed. (New York: Academic Press).
6. Altman, P.L., and Katz, D.D. (1979). Inbred and genetically defined strains of laboratory animals (Bethesda, Md.: Federation of American Societies for Experimental Biology).
7. Egan, C.M., Sridhar, S., Wigler, M., and Hall, I.M. (2007). Recurrent DNA copy number variation in the laboratory mouse. *Nature genetics* 39, 1384-1389.
8. Reed, C., Baba, H., Zhu, Z., Erk, J., Mootz, J.R., Varra, N.M., Williams, R.W., and Phillips, T.J. (2017). A Spontaneous Mutation in *Taar1* Impacts Methamphetamine-Related Traits Exclusively in DBA/2 Mice from a Single Vendor. *Frontiers in pharmacology* 8, 993.
9. Clapcote, S.J., and Roder, J.C. (2004). Survey of embryonic stem cell line source strains in the water maze reveals superior reversal learning of 129S6/SvEvTac mice. *Behavioural brain research* 152, 35-48.
10. Grottick, A.J., Bagnol, D., Phillips, S., McDonald, J., Behan, D.P., Chalmers, D.T., and Hakak, Y. (2005). Neurotransmission- and cellular stress-related gene expression associated with prepulse inhibition in mice. *Brain research Molecular brain research* 139, 153-162.
11. Mayorga, A.J., and Lucki, I. (2001). Limitations on the use of the C57BL/6 mouse in the tail suspension test. *Psychopharmacology* 155, 110-112.
12. Radulovic, J., Kammermeier, J., and Spiess, J. (1998). Generalization of fear responses in C57BL/6N mice subjected to one-trial foreground contextual fear conditioning. *Behavioural brain research* 95, 179-189.
13. Stiedl, O., Radulovic, J., Lohmann, R., Birkenfeld, K., Palve, M., Kammermeier, J., Sananbenesi, F., and Spiess, J. (1999). Strain and substrain differences in context- and tone-dependent fear conditioning of inbred mice. *Behavioural brain research* 104, 1-12.
14. Siegmund, A., Langnaese, K., and Wotjak, C.T. (2005). Differences in extinction of conditioned fear in C57BL/6 substrains are unrelated to expression of alpha-synuclein. *Behavioural brain research* 157, 291-298.
15. Toye, A.A., Lippiat, J.D., Proks, P., Shimomura, K., Bentley, L., Hugill, A., Mijat, V., Goldsworthy, M., Moir, L., Haynes, A., *et al.* (2005). A genetic and physiological study of impaired glucose homeostasis control in C57BL/6J mice. *Diabetologia* 48, 675-686.
16. Khisti, R.T., Wolstenholme, J., Shelton, K.L., and Miles, M.F. (2006). Characterization of the ethanol-deprivation effect in substrains of C57BL/6 mice. *Alcohol* 40, 119-126.
17. Green, M.L., Singh, A.V., Zhang, Y., Nemeth, K.A., Sulik, K.K., and Knudsen, T.B. (2007). Reprogramming of genetic networks during initiation of the Fetal Alcohol Syndrome. *Developmental dynamics : an official publication of the American Association of Anatomists* 236, 613-631.
18. Diwan, B.A., and Blackman, K.E. (1980). Differential susceptibility of 3 sublines of C57BL/6 mice to the induction of colorectal tumors by 1,2-dimethylhydrazine. *Cancer letters* 9, 111-115.
19. Roth, D.M., Swaney, J.S., Dalton, N.D., Gilpin, E.A., and Ross, J., Jr. (2002). Impact of anesthesia on cardiac function during echocardiography in mice. *American journal of physiology Heart and circulatory physiology* 282, H2134-2140.
20. Kumar, V., Kim, K., Joseph, C., Kourrich, S., Yoo, S.H., Huang, H.C., Vitaterna, M.H., de Villena, F.P., Churchill, G., Bonci, A., *et al.* (2013). C57BL/6N mutation in cytoplasmic FMRP interacting protein 2 regulates cocaine response. *Science* 342, 1508-1512.

21. Akinola, L.S., McKiver, B., Toma, W., Zhu, A.Z.X., Tyndale, R.F., Kumar, V., and Damaj, M.I. (2019). C57BL/6 Substrain Differences in Pharmacological Effects after Acute and Repeated Nicotine Administration. *Brain sciences* 9.
22. Kadiyala, S.B., Papandrea, D., Herron, B.J., and Ferland, R.J. (2014). Segregation of seizure traits in C57 black mouse substrains using the repeated-flurothyl model. *PloS one* 9, e90506.
23. Markham, B.E., Kernodle, S., Nemzek, J., Wilkinson, J.E., and Sigler, R. (2015). Chronic Dosing with Membrane Sealant Poloxamer 188 NF Improves Respiratory Dysfunction in Dystrophic Mdx and Mdx/Utrophin-/- Mice. *PloS one* 10, e0134832.
24. Bryant, C.D., Smith, D.J., Kantak, K.M., Nowak, T.S., Jr., Williams, R.W., Damaj, M.I., Redei, E.E., Chen, H., and Mulligan, M.K. (2020). Facilitating Complex Trait Analysis via Reduced Complexity Crosses. *Trends in genetics : TIG* 36, 549-562.
25. Sigmon, J.S., Blanchard, M.W., Baric, R.S., Bell, T.A., Brennan, J., Brockmann, G.A., Burks, A.W., Calabrese, J.M., Caron, K.M., Cheney, R.E., *et al.* (2020). Content and Performance of the MiniMUGA Genotyping Array: A New Tool To Improve Rigor and Reproducibility in Mouse Research. *Genetics* 216, 905-930.
26. Keane, T.M., Goodstadt, L., Danecek, P., White, M.A., Wong, K., Yalcin, B., Heger, A., Agam, A., Slater, G., Goodson, M., *et al.* (2011). Mouse genomic variation and its effect on phenotypes and gene regulation. *Nature* 477, 289-294.
27. Mitra, I., Huang, B., Mousavi, N., Ma, N., Lamkin, M., Yanicky, R., Shleizer-Burko, S., Lohmueller, K.E., and Gymrek, M. (2020). Genome-wide patterns of de novo tandem repeat mutations and their contribution to autism spectrum disorders. *bioRxiv* 2020.03.04.974170 <https://doi.org/10.1101/2020.03.04.974170>.
28. Kong, A., Frigge, M.L., Masson, G., Besenbacher, S., Sulem, P., Magnusson, G., Gudjonsson, S.A., Sigurdsson, A., Jonasdottir, A., Jonasdottir, A., *et al.* (2012). Rate of de novo mutations and the importance of father's age to disease risk. *Nature* 488, 471-475.
29. Hannan, A.J. (2018). Tandem repeats mediating genetic plasticity in health and disease. *Nature reviews Genetics* 19, 286-298.
30. Fotsing, S.F., Margoliash, J., Wang, C., Saini, S., Yanicky, R., Shleizer-Burko, S., Goren, A., and Gymrek, M. (2019). The impact of short tandem repeat variation on gene expression. *Nature genetics* 51, 1652-1659.
31. Chaisson, M.J.P., Sanders, A.D., Zhao, X., Malhotra, A., Porubsky, D., Rausch, T., Gardner, E.J., Rodriguez, O.L., Guo, L., Collins, R.L., *et al.* (2019). Multi-platform discovery of haplotype-resolved structural variation in human genomes. *Nature communications* 10, 1784.
32. Hurles, M.E., Dermitzakis, E.T., and Tyler-Smith, C. (2008). The functional impact of structural variation in humans. *Trends in genetics : TIG* 24, 238-245.
33. Yalcin, B., Wong, K., Agam, A., Goodson, M., Keane, T.M., Gan, X., Nellaker, C., Goodstadt, L., Nicod, J., Bhomra, A., *et al.* (2011). Sequence-based characterization of structural variation in the mouse genome. *Nature* 477, 326-329.
34. Quinlan, A.R., Clark, R.A., Sokolova, S., Leibowitz, M.L., Zhang, Y., Hurles, M.E., Mell, J.C., and Hall, I.M. (2010). Genome-wide mapping and assembly of structural variant breakpoints in the mouse genome. *Genome research* 20, 623-635.
35. Simon, M.M., Greenaway, S., White, J.K., Fuchs, H., Gailus-Durner, V., Wells, S., Sorg, T., Wong, K., Bedu, E., Cartwright, E.J., *et al.* (2013). A comparative phenotypic and genomic analysis of C57BL/6J and C57BL/6N mouse strains. *Genome biology* 14, R82.
36. Doran, A.G., Wong, K., Flint, J., Adams, D.J., Hunter, K.W., and Keane, T.M. (2016). Deep genome sequencing and variation analysis of 13 inbred mouse strains defines candidate phenotypic alleles, private variation and homozygous truncating mutations. *Genome biology* 17, 167.
37. Beck, J.A., Lloyd, S., Hafezparast, M., Lennon-Pierce, M., Eppig, J.T., Festing, M.F., and Fisher, E.M. (2000). Genealogies of mouse inbred strains. *Nature genetics* 24, 23-25.
38. Charles River Laboratories. Available from: <https://www.criver.com/>
39. Jackson Laboratory. Available from: <https://www.jax.org/>
40. Sarsani, V.K., Raghupathy, N., Fiddes, I.T., Armstrong, J., Thibaud-Nissen, F., Zinder, O., Bolisetty, M., Howe, K., Hinerfeld, D., Ruan, X., *et al.* (2019). The Genome of C57BL/6J "Eve", the Mother of the Laboratory Mouse Genome Reference Strain. *G3* 9, 1795-1805.
41. Lippert, C., Casale, F.P., Rakitsch, B., and Stegle, O. (2014). LIMIX: genetic analysis of multiple traits. *003905* <https://doi.org/10.1101/003905>.

42. McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R., Thormann, A., Flicek, P., and Cunningham, F. (2016). The Ensembl Variant Effect Predictor. *Genome biology* 17, 122.
43. Chang, Y.F., Imam, J.S., and Wilkinson, M.F. (2007). The nonsense-mediated decay RNA surveillance pathway. *Annual review of biochemistry* 76, 51-74.
44. Yang, H., Wang, J.R., Didion, J.P., Buus, R.J., Bell, T.A., Welsh, C.E., Bonhomme, F., Yu, A.H., Nachman, M.W., Pialek, J., *et al.* (2011). Subspecific origin and haplotype diversity in the laboratory mouse. *Nature genetics* 43, 648-655.
45. Wang, J.R., de Villena, F.P., and McMillan, L. (2012). Comparative analysis and visualization of multiple collinear genomes. *BMC bioinformatics* 13 Suppl 3, S13.
46. Mulligan, M.K., Ponomarev, I., Boehm, S.L., 2nd, Owen, J.A., Levin, P.S., Berman, A.E., Blednov, Y.A., Crabbe, J.C., Williams, R.W., Miles, M.F., *et al.* (2008). Alcohol trait and transcriptional genomic analysis of C57BL/6 substrains. *Genes, brain, and behavior* 7, 677-689.
47. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature biotechnology* 37, 907-915.
48. Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31, 166-169.
49. Chiang, C., Layer, R.M., Faust, G.G., Lindberg, M.R., Rose, D.B., Garrison, E.P., Marth, G.T., Quinlan, A.R., and Hall, I.M. (2015). SpeedSeq: ultra-fast personal genome analysis and interpretation. *Nature methods* 12, 966-968.
50. Faust, G.G., and Hall, I.M. (2014). SAMBLASTER: fast duplicate marking and structural variant read extraction. *Bioinformatics* 30, 2503-2505.
51. Tarasov, A., Vilella, A.J., Cuppen, E., Nijman, I.J., and Prins, P. (2015). Sambamba: fast processing of NGS alignment formats. *Bioinformatics* 31, 2032-2034.
52. Garrison, E., and Marth, G.T. (2012). Haplotype-based variant detection from short-read sequencing. *arXiv:1207.3907 [q-bio.GN]* <https://arxiv.org/abs/1207.3907>
53. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., de Bakker, P.I., Daly, M.J., *et al.* (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *American journal of human genetics* 81, 559-575.
54. Willems, T., Zielinski, D., Yuan, J., Gordon, A., Gymrek, M., and Erlich, Y. (2017). Genome-wide profiling of heritable and de novo STR variations. *Nature methods* 14, 590-592.
55. Benson, G. (1999). Tandem repeats finder: a program to analyze DNA sequences. *Nucleic acids research* 27, 573-580.
56. Layer, R.M., Chiang, C., Quinlan, A.R., and Hall, I.M. (2014). LUMPY: a probabilistic framework for structural variant discovery. *Genome biology* 15, R84.
57. Abyzov, A., Urban, A.E., Snyder, M., and Gerstein, M. (2011). CNVnator: an approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. *Genome research* 21, 974-984.
58. Pedersen, B.S., and Quinlan, A.R. (2018). Mosdepth: quick coverage calculation for genomes and exomes. *Bioinformatics* 34, 867-868.