Patterns of *Mustelid gammaherpesvirus 1* (MusGHV-1) genital reactivation linked to stressors in adult European badgers (*Meles meles*)

Ming-shan Tsai^{1*}, Sarah Francois², Chris Newman^{1,3}, David W. Macdonald¹, Christina D. Buesching^{3,4}

Affiliation

1. Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, Recanati-Kaplan Centre, Abingdon Road, Tubney House, Tubney, Oxfordshire OX13 5QL, UK
2. Evolve.zoo, Department of Zoology, University of Oxford, Peter Medawar Building, South Park Road, Oxford, OX1 3SY, UK
3. Cook's Lake Farming Forestry and Wildlife Inc (Ecological Consultancy), Queens County, Nova Scotia, Canada
4. Department of Biology, Irving K. Barber Faculty of Science, The University of British Columbia, Okanagan, Kelowna, British Columbia, Canada

*Correspondence: cindy150051@gmail.com

Abstract

Herpesvirus infections are common and mostly asymptomatic in vertebrates, but can result in impaired reproduction. It is therefore important to understand infection patterns and associated risk factors, particularly the effects of different stressors. Here we use Mustelid gammaherpesvirus 1 (MusGHV-1) infection in European badgers (*Meles meles*) as a host-pathogen wildlife model to study the effects of a variety of demographic, social, physiological and environmental stressors on viral reactivation in the genital tract. We collected 251 genital swabs from 151 free-ranging individuals across 3 trapping seasons (spring, summer and autumn). We screened for MusGHV-1 using PCR and explored possible links between genital MusGHV-1 reactivation and stressors, and their interactions, using logistic regression. In adults, reactivation was more likely in males, especially those in poorer body condition during summer. In females, reactivation was more likely when living in social groups comprising a higher percentage of cubs, but counter to our predictions, recent lactation appeared not influential. In relation to age, reactivation was more common in individuals over 8 years old than among prime age adults, and among juveniles (<2 years old), especially females and individuals in better body condition, likely due to early puberty. Environmentally, reactivation was more prevalent in summer when food abundance is typically low. Our results evidence age effects on MusGHV-1 reactivation; in juveniles MusGHV-1 shedding in the genital tract is likely related to primary infection, while in adults, genital MusGHV-1 reactivation from latency was associated with aging, social and/or environmental stress.

Keywords: Stress, herpesvirus reactivation, gammaherpesvirus infection, sexually transmittable diseases, STI, wildlife disease epidemiology

39

40 Importance

41 The immuno-suppressive effects of elevated stress levels facilitate disease development, and can
42 ultimately cause host extinction at the population level, especially where diseases are transmitted
43 sexually. The impacts of stress on host-pathogen dynamics through disease, however, are still poorly
44 understood outside the laboratory or captive environments. Our study provides rare evidence from a
45 free-ranging wild mammal population that the infection dynamics of a common and sexually
46 transmittable gammaherpesvirus are linked to demographic, social, physiological and environmental
47 stress. We propose that the effects of stressors on STIs and viral reactivation are an important factor
48 to be taken into account in conservation efforts when working with vulnerable wildlife populations.

49

50 Introduction

51 Herpesvirus infection is common in vertebrates with most vertebrate herpesviruses belonging to 4
52 subfamilies, the *Alphaherpesvirinae*, *Betaherpesvirinae*, *Gammaherpesvirinae* and *Deltaherpesvirinae*
53 (1, 2). Herpesvirus species are generally host-specific, but cross-species transmission is more frequent
54 than previously assumed (3, 4). After primary infection, the herpesvirus enters a latent stage in the
55 host cell (e.g. lymphocytes in gammaherpesvirus infection), and can be reactivated repeatedly
56 throughout life by stress (5, 6), trauma (e.g. surgery:(7)) or primary co-infection with other pathogens
57 (8). Reactivation is a process of viral lytic infection, which involves virus replication within the host cell,
58 eventually destroying the cell and releasing infectious virions. Reactivation of herpesviruses usually
59 occurs in the epithelial cells of mucosa that function as portals for external contact (e.g., mouth, nose,
60 eyes and genital tract), thus facilitating transmission. Reactivation is, however, typically asymptomatic
61 or induces only mild disease, but can also promote development of severe diseases like cancer (9),
62 depending on strain pathogenicity (10) and co-infection with other pathogens causing
63 immunodeficiency (e.g. Human herpesvirus 8 and HIV (11)), and is associated with a higher risk of
64 contracting co-infection with additional pathogens with high virulence (e.g. *Chlamydia pecorum*
65 infection in koalas suffering from gammaherpesvirus reactivation (12)).

66

67 Chronic stress has proven a significant risk factor, causing immune system dysregulation. where
68 corticosteroids inhibit the pro-inflammatory cytokine responses, allowing the virus to (re-)activate and
69 undergo lytic proliferation unchecked (13). This link between elevated corticosteroid levels and
70 herpesvirus reactivation has been proven experimentally (horses: (14); captive reindeer: (15) and
71 through observation (e.g. humans: (6, 16); captive Grévy's zebras (*Equus grevyi*): (5), but has not been
72 investigated in free-living wildlife populations.

73

74 Here, we use European badgers (*Meles meles*) as a wildlife model to investigate how different risk
75 factors and stressors affect herpesvirus reactivation. Badgers are seasonally breeding mustelids that
76 are commonly infected with the *Mustelid gammaherpesvirus 1* (MusGHV-1: a large double-stranded

DNA virus belonging to the *Gammaherpesvirinae*, genus *Percavirus*), where prevalence of viral DNA in blood samples can reach up to 100% in the UK and in Ireland (17, 18), and 55% - 82.5% in genital swab samples (19, 20). Gammaherpesvirus reactivation can cause severe disease in humans (21, 22) and domestic animals (23–25), and has increasingly been associated with illness in wildlife species (26–29). In badgers, previous research has linked otherwise asymptomatic MusGHV-1 reactivation in genital tracts to impaired female reproductive performance (20) and indicates that during the main mating season, adult males are at particular risk of genital MusGHV-1 reactivation. Nevertheless, the impact of social, physiological and environmental stress on MusGHV-1 reactivation has thus far not been investigated. Badgers are subject to a variety of stressors: faecal corticoid levels indicate that badgers experience seasonal variation in stress levels (30, 31), which may be due to variation in food availability (30, 31) and can result in mortality (32, 33). Sociologically, higher social group density is associated with female reproductive suppression (34, 35), reduced body condition and fecundity (36), and increased bite wounding among male badgers (37). Furthermore, aging reduces tolerance to stress (38), specifically altering the balance of innate and acquired immunity in badgers (39), and increasing their risk of herpesvirus reactivation (20), as also observed in other carnivora species (40, 41), sometimes resulting in chronic and continuous herpesvirus reactivation (42). Therefore metrics of body condition, especially reduced body-condition as a consequence of recent lactation (30, 43), can indicate that the individual may be experiencing physiological stress (30).

To evaluate the impact of potential stressors on genital MusGHV-1 reactivation, we conducted population-wide molecular screening using genital swabs taken from a free-ranging badger population in the south of England across 3 seasons (spring, summer and autumn). We investigated whether environmental factors (i.e., season and social group size), host demographic parameters (i.e., sex, age and lactation), and host health (i.e., body condition) affect risk of genital MusGHV-1 reactivation.

Materials and methods

Field data and sample collection

Samples were collected from 151 individual live-trapped badgers in Wytham Woods, Oxfordshire, UK (51°46'26"N, 1°19'19"W; caught in May, September and November 2018 following the methodology described in Macdonald et al. (44); for details see Table 1). All trapping and animal sampling protocols were approved by the University of Oxford' Animal Welfare and Ethical Review Board. Trapping was conducted under Natural England license (currently 2019-38863, Badger Act 1992) and all animal handling procedures were carried out by qualified Personal Individual License (PIL) holders under Home Office license (current PPL 30/3379, Animals (Scientific Procedures) Act 1986). For each capture, we recorded sex, sett (i.e., communal den used by a badger social group) of capture, body condition score (BCS, categorized as 1= very thin to 5= very fat), and lactational status (determined by teat measurements of females in spring: (45)). Because each badger in Wytham is given an individual tattoo at first capture (usually as a cub (46)), exact age (in years) was known for most (243 of 251) animals in

the dataset. For the remaining 8 badgers first caught as adults, age was inferred by toothwear according to the method described Bright Ross et al. (47). We defined 4 age classes: i) juveniles < 2 years old (cubs and yearlings were combined to increase sample sizes, as there was no difference between Mus GHV-1 prevalence in cubs and yearlings: Fisher's exact test: p-value=0.7449); and - based on sex-steroid levels (48); ii) young adults: $2 \leq x < 5$ years old; iii) old adults: $5 \leq x < 8$ years old; iv) very old adults: ≥ 8 years old. The number of cubs and adults resident in each sett was estimated using minimum number alive (MNA) estimates (44, 47).

Sterile cotton tops with wooden shafts were used to swab the genital tracts of all females (cubs and adults) and all males (except for very small male cubs in spring for animal welfare reasons), and stored in 2 ml sterile microcentrifuge tubes. All samples were frozen and stored at -20°C immediately after sampling. Badgers were released at their site of capture on the same day, after full recovery from anaesthesia.

DNA extraction and purification

Each swab was reconstituted with 400µl sterile double distilled water and vortexed gently at room temperature for 10 minutes. A 200µl aliquot was taken from the reconstituted swab fluids, and viral DNA was extracted and purified using a commercial kit (DNeasy Blood and Tissue Kit, Qiagen) following manufacturer's instructions. Purified DNA was then eluted in 100µl of the provided buffer.

Screening for MusGHV-1 DNA using polymerase chain reaction (PCR) and sequencing

The purified DNA was screened using a MusGHV-1-specific primer pair designed by King et al. 2004 (18), targeting 281 base pairs of the partial DNAPol gene. For each reaction, a total of 20 µl PCR solution was mixed with 10µl HotStartTaq Master Mix (Qiagen, containing 1 unit of HotStartTaq DNA Polymerase, 12µM of $MgCl_2$ and 1.6µM of each dNTP), 0.5 µM of each primer, and 2µl CoralLoad gel loading dye and 5 µl DNA template. Amplification conditions were kept at 95°C for 5 mins to activate DNA polymerase, followed by 45 cycles of denaturation at 95°C for 45 seconds, primer annealing at 60°C for 45 seconds, and chain elongation at 72°C for 1 minute, followed by a final extension at 72°C for 10 minutes. Finally, the PCR products were loaded in 2% agarose gel to check the amplification results under UV light. Samples with positive results were then amplified again with substituted front primer (5' CCA AGC AGT GCA TAG GAG GT 3') to generate longer sequences (771 base pairs). PCR products were then purified and sent for genotyping using Sanger sequencing to confirm the identity of produced amplicon. Sequences returned were then aligned by Clustal W method (49) and analyzed for variation using MEGA X (10.1.7) (50). Representative sequences were selected and published on GenBank under accession number MT332100 and MT332101 assigned.

Statistical analysis

Statistical analyses were performed with the R and R Studio software (version 1.21335) (51). Prevalence of genital MusGHV-1 DNA was calculated by dividing the number of PCR positive cases by the total number of tested cases, and 5% upper and lower confidence intervals were calculated using the Wilson method (52). Logistic regression (glmer function, R package lme4) with badger identity (tattoo) number as a random effect was used to measure univariate effects of MusGHV-1 reactivation in genital tracts with season, sex, age, age class, BCS, number of residents per sett, and percentage of cubs per social group (calculated by sett), where we categorized these data on percentage of cubs per group into low and high using 30% as the dividing point according to distribution of the data (Figure S1). Effects of lactation were analyzed using Fisher's exact tests due to low sample sizes and presented using odds ratios. The final multivariable model was selected through the manual backwards selection method. Model residual diagnostics were conducted using R package DHARMA (version 0.3.3.0). Model fit was established using area-under-receiver-operating characteristics (AUC) (53). Kruskal-Wallis tests were used to compare genital MusGHV-1 positive and negative individuals of different BCS. Because juveniles (especially cubs) are generally thinner than adults, and thus have a different body condition distribution (Figure S2), we calculated a body condition index (BCI) as $\ln(\text{body weight})/\ln(\text{body length})$ (36)). We analysed the association of MusGHV-1 reactivation and individual BCS for juveniles and adults separately. Linear models (lm function, R package lme4) were used to assess the association of age and MusGHV-1 reactivation prevalence.

Results

The overall prevalence of genital MusGHV-1 reactivation was 35.9% (90/251, 95% CI: 30.2% - 42.0%), and prevalence was generally higher in summer (45%, 36/80) than in spring (34.4%, 33/96) and autumn (28%, 21/75) (Table 2; Figure 1).

Age effects on seasonal and sex-specific patterns of genital MusGHV-1 reactivation

There was strong evidence for an effect of age on genital MusGHV-1 where prevalence followed a U-shaped age curve, being lowest for badgers at the age of 5 or 6 years old (Figure 2, quadratic term, adjusted $R^2 = 0.683$, F-statistic= 12.83 on 2 and 9 DF, p-value= 0.002). When divided by age classes, prevalences in juveniles (46.6%, 34/73) and very old badgers (47.7%, 21/44) were higher than young (30.9%, 30/97) and old (13.5%, 5/37) adults (Table 2). However, no effect of sex (logistic regression analysis, p=0.679) was observed in the univariate analysis.

BCS effects on seasonal and sex-specific patterns of genital MusGHV-1 reactivation

Although adults with lower BCS had a higher probability of genital MusGHV-1 reactivation according to our univariate analysis (p-value=0.022, Table 2), when grouped by seasons and sex this relationship was only significant in adult males in summer (Kruskal-Wallis tests, p-value = 0.006) (Figure 3).

Group effects on seasonal and sex-specific patterns of genital MusGHV-1 reactivation

Genital MusGHV-1 reactivation was significantly more prevalent in females living in setts comprised by >30% cubs (logistic regression analysis, p -value = 0.002), but there was no evidence for the absolute number of adults, cubs nor the total number of badgers resident in each sett affecting reactivation rates (Table 2).

Effects of recent reproductive success on female genital MusGHV-1 prevalence

There was no evidence for recent lactation affecting the reactivation rate in sexually mature females in spring (Fisher's exact test, p -value=1) (Table 2).

Multivariable analysis of genital MusGHV-1 reactivation

Because the univariate analysis indicated that juveniles and adults appear to have different patterns of reactivation, we separated these data to generate multivariable models of juveniles ($n=73$) and adults ($n=178$). We included all variables initially for which there was any evidence for an effect in the univariate analyses (p -value < 0.10, Table 2). The most parsimonious models for juveniles (Table 3) and adults (Table 4) both had diagnostically acceptable (53) AUC area of 0.748 and 0.726 (Figure S3 and S4), respectively. In juveniles, multivariable analysis showed that all females, and all juveniles with higher BCS, were at particular risk of genital MusGHV-1 reactivation (Table 3). In adults, in contrast, males are at a higher risk of MusGHV-1 reactivation than are females, where all adults experience the highest reactivation risk in summer, and older (≥ 8 years) badgers, and female living in setts with a higher percentage (over 30%) of cubs are at particular risk (Table 4).

Genetic diversity of MusGHV-1 in the Wytham badger population

We sequenced 5 MusGHV-1 positive PCR products of a partial DNA polymerase gene. All sequences were trimmed to 694 base pairs and confirmed to be MusGHV-1 according to the NCBI online blasting service, returning 98.7% ($n=3$) and 100% ($n=2$) nucleotide identity to the published MusGHV-1 sequence isolated from a badger in Cornwall, England (Accession number: AF275657).

Discussion

Herpesvirus reactivation triggered by stress has been widely confirmed naturally and experimentally by corticosteroid injection in humans and domestic animals (5, 14). Linking stress and viral reactivation in wildlife, however, is particularly challenging due to the difficulties of monitoring individual stress levels in the field, and typically this relationship can only be confirmed experimentally by taking subjects into captivity at least temporarily (15). Using indicators that have been linked to stress hormone levels in previous studies can thus provide an informative way to study the relationship between stress and herpesvirus reactivation in free-ranging wildlife.

Faecal corticosteroid measurements from badgers in Ireland (30) evidence higher stress levels in summer likely associated with dry environmental conditions that result in lower earthworm availability

(i.e., the badgers' main food type (46)); similarly, in our own study population, summer drought is an established mortality factor due to starvation/ malnutrition (32, 33). This corresponds with our finding that, in all adults, seasonal MusGHV-1 reactivation rates were highest in summer, but females tended to have higher viral reactivation levels than males in spring – possibly due to reproductive stresses, while the reverse was true in autumn (Figure 1) (30). Interestingly, however, we found no correlation between BCS and MusGHV-1 reactivation across all seasons in females, implying that reduced body condition was not necessarily indicative of physiological stress. In fact, Bright Ross et al. (subm) found complex relationships between body-condition, survival and reproductive success in this same population, where although breeding females lose condition, they often end up being no thinner than non-breeding females as they were in much better condition in winter (i.e. before pregnancy/ lactation).

From the perspective of male rates of reactivation, males with higher testosterone levels tend to be thinner during spring and summer (54), but tend to mate more often (55). In our recent survey of Irish badger populations (20), the high rate (over 80%) of genital MusGHV-1 reactivation in adult males during the peak (postpartum) mating season, from mid-January to mid-February, implies not only a link to mating activity, but also a mechanism enhancing transmission. This corroborates another finding in the same study that males with more spermatozoa have a higher prevalence of genital MusGHV-1 reactivation (20); linking higher sexual activity to higher STI prevalence as reported also in many other studies (56–59). We were unable to explicitly test effects relating to badger mating behaviour in our study because we can not trap during late pregnancy and neonatal cub care, to avoid stressing mothers or depriving cubs of maternal care. Nevertheless, although reactivation rate in autumn, when food sources are most abundant, and badgers undergo a period of reproductive quiescence (60), and thus experience less implicit stress, was significantly lower compared only to summer, but not to spring. This suggests that other factor(s) (e.g. sex hormone cycles (61), oxidative stress (62), genital microbiome (63) or bacterial co-infections) might also be affecting reactivation rates, beyond the scope of our current study.

In terms of age class effects, the high genital reactivation rate detected in cubs and yearlings suggests that badgers contract MusGHV-1 early in life, before reaching sexual maturity. Although MusGHV-1 reactivates repeatedly throughout life, reactivation tends to be less frequent in young and old adults compared to adults in their prime, although rates increase in very old individuals. This matches patterns in humans where most people become infected with herpes during their childhood/ adolescence (e.g. 100% and 70% seroprevalence of EBV before age 14 in Hong Kong and the United Kingdom (64)), then typically experience viral latency during their prime, but can suffer from increasingly longer and more frequent herpesvirus reactivation that sometimes cause mild disease (e.g., shingles (65) in old age due to lowered immune response (66, 67)).

Since vertical transmission of MusGHV-1 through the placenta is unlikely (20), and the potential for infection from the vaginal tract during parturition is equally low due to low genital MusGHV-1 reactivation rate in pregnant females (20), we hypothesise that cubs contract primary infection through close contact with virus-shedding conspecifics (20). Thereafter genital reactivation in cubs may arise after primary acute infection through non-sexual routes and subsequent latency, as observed in the murine model where *Murine herpesvirus 4* (MuHV-4, also a gammaherpesvirus), inoculation in the nasal cavity results in acute infection in the respiratory tract and lungs and establishes latency in the spleen, but then reactivates in the vaginal tract 17-21 days after inoculation (61). MuHV-4 nasal cavity inoculation, however, does not result in reactivation in male genital tracts, and transmission is only possible from females to males. After sexual intercourse with virus-shedding female mice, the virus then replicates in the male penis for 3 weeks. Interestingly, also in badgers, female juveniles are at higher risk of MusGHV-1 reactivation in the genital tract than are males. Furthermore, juveniles in better body condition, regardless of sex, exhibit higher reactivation prevalence. Indeed, juvenile males in better body condition enter puberty earlier than thinner males (11 months compared to 22 – 28 months: (57)). Once juveniles enter puberty they will experience an increased risk of contracting MusGHV-1 through sexual contact and/ or that their latent infection is reactivated through mating resulting in viral shedding in the genital tract.

Our results also show that social group structure can affect prevalence of genital herpesvirus reactivation, particularly the proportion of cubs within a residential group. This trend was more apparent in females than in males. This may be because badger cubs generally carry higher pathogen burdens than adults (68, 69), and thus increase per capita immunity burden among all badgers resident in the respective sett (64).

Conclusion

Our study demonstrates - for the first time in the wild - the link between stress experienced by the host and latent virus reactivation. Amplified stress levels induced by human disturbance as well as food insecurity and more frequent catastrophic weather events arising from human induced rapid environmental change (HIREC) could therefore not only increase the risk of disease development, promotion of transmission within a population, but also negatively impact host reproductive fitness through latent virus reactivation in the reproductive tract (19). Careful monitoring of endemic latent virus infection as well as surveillance for possible newly emerging strains should be included when planning *in situ* and *ex situ* conservation programmes for endangered species (70).

Acknowledgments

The authors would like to thank Dr. Nadine Sugianto, Dr. Sil van Lieshout, Dr. Tanesha Allen, and Julius Bright-Ross for assistance with sample collection, fieldwork and MNA data provision. MST would like to thank the Ministry of Education in Taiwan and Lady Margaret Hall, University of Oxford, for

scholarship support. The authors also thank Paul Johnson and Ta-Chun Liu for providing advice on statistics. The authors claim no conflict of interests in the present work.

Author Contributions

Project conception: CBD,MST; samples collection: CBD,CN,MST ; laboratory work: MST ; data analysis: MST; writing and revision: MST,CBD,SF,DWM; All authors have read and approved the manuscript.

313 Table 1: Details of swab sampling

Age class	Age	Spring		Summer		Autumn		Total
		Female	Male	Female	Male	Female	Male	
Cub	0	15	15	10	4	9	9	62
Yearling	1	1	3	2	2		3	11
	2	7	7	7	9	5	6	41
Young adult	3	4	8	3	5	2	8	30
	4	2	5	4	5	2	8	26
Old adult	5	2	1	6	1	3	1	14
	6	3	1	3	1	3		11
	7		2		2		2	6
	NA	1	1	1	1		2	6
	8	4	4	4	3	1	3	19
Very old adult	9	2	1	2	1	1	1	8
	10	3	1	3		4		11
	11							0
	12							0
	13	1	1		1	1		4
	NA		1				1	2
	Total	45	51	45	35	31	44	251

Table 2: Overview of genital MusGHV-1 reactivation prevalence and univariate logistic regression analysis. Formula: MusGHV ~ Variate + (1|Tattoo); number of observations: 251; groups by tattoo number: 150

	Positive	Total	Prevalence	Prevalence 95% CI	Odds ratio (OR)	OR 95% CI	P-value
Sex							
Male	45	130	34.62%	27% - 43.1%	0.59	0.53 - 1.52	0.679
Female	45	121	37.19%	29.1% - 46.1%			
Season							
Spring	33	96	34.38%	25.6% - 44.3%	1.37	0.7 - 2.69	0.362
Summer	36	80	45.00%	34.6% - 55.9%	2.16	1.08 - 4.33	0.03
Autumn	21	75	28.00%	19.1% - 39%			
Age class							
Juvenile (<2 years old)	34	73	46.58%	35.6% - 57.9%	5.58	1.95 - 15.92	0.001
Young (2 - 4 years old)	30	97	30.93%	22.6% - 40.7%	2.87	1.02 - 8.08	0.046
Old (5 - 7 years old)	5	37	13.51%	5.9% - 28%			
Very old (>7 years old)	21	44	47.73%	33.8% - 62.1%	5.84	1.92 - 17.78	0.002
Age							
Age		243					<0.001
Age ²		243					<0.001
Body Condition ^a							
Body condition score		182					0.0416
Sett group size							
Total		251					0.814
Adult		251					0.434
Cub		251					0.086
Cub percentage per sett							
Low (<30%)	58	191	30.37%	24.3% - 37.2%			
High (>30%)	32	60	53.33%	40.9% - 65.4%	2.62	1.23 - 4.83	0.002
Lactational status ^{cd}							
Not Lactated	3	10	30.00%	10.8% - 60.3%			
Lactated	6	18	33.33%	16.3% - 56.3%	1.17	0.22 - 6.21	1

a: Only adults were included in this analysis
b: Only females captured in spring were included in this analysis
c: Fisher exact test

Table 3: Final general mixed effect model of multivariable logistic regression analysis for juveniles
Formula: MusGHV ~ Sex + Body condition + (1|Tattoo); number of observations: 72; groups by tattoo
number: 48

Group	Estimate	Standard error	z value	Adjusted OR	95% CI	p value
(Intercept)	-1.226	0.7851	-1.562			0.1184
Sex						
Female						
Male	-1.0519	0.5123	-2.054	0.35	0.13 - 0.95	0.04
Body Condition						
BCS	0.6245	0.2876	2.172	1.87	1.06 - 3.28	0.0299

Table 4: Final general mixed effect model of multivariable logistic regression analysis for adults
Formula: MusGHV ~ Sex + Season + AgeGroup + Cub percentage + Sex*Cub percentage + (1|Tattoo);
Number of observations: 178; Groups by tattoo number: 101

Group	Estimate	Standard error	z value	Adjusted OR	95% CI	p value
(Intercept)	-0.9952	0.4859	-2.048			0.04
Sex						
Female						
Male	0.8171	0.4126	1.98	2.26	1 - 5.1	0.048
Season						
Spring	-1.037	0.4255	-2.437	1.61	0.66 - 3.91	0.297
Summer	-2.2692	0.6333	-3.583	3.08	1.27 - 7.47	0.013
Autumn						
Age class						
Young	-1.1084	0.468	-2.368	0.35	0.15 - 0.82	0.015
Old	-2.4001	0.7052	-3.404	0.1	0.03 - 0.36	<0.001
Very old						
Cub percentage per sett						
Low (<30%)						
High (>30%)	1.8903	0.6968	2.713	6.62	1.69 - 25.95	0.006
Interaction						
Female:Cub percentage						
Male:Cub percentage	-2.3686	1.0492	-2.258	0.09	0.01 - 0.73	0.024

References

1. Roizman B, Baines J. 1991. The diversity and unity of herpesviridae. *Comp Immunol Microbiol Infect Dis* 14:63–79.
2. Pellett PE. 2014. Trunkloads of Viruses. *J Virol* 88:13520–13522.
3. Woźniakowski G, Samorek-Salamonowicz E. 2015. Animal herpesviruses and their zoonotic potential for cross-species infection. *Ann Agric Environ Med* 22:191–194.
4. Brito AF, Pinney JW. 2020. Intrahost speciations and host switches shaped the evolution of herpesviruses. *bioRxiv* 418111.
5. Seeber PA, Quintard B, Sicks F, Dehnhard M, Greenwood AD, Franz M. 2018. Environmental stressors may cause equine herpesvirus reactivation in captive Grévy's zebras (*Equus grevyi*). *PeerJ* 2018:1–18.
6. Stowe RP, Pierson DL, Barrett ADT. 2001. Elevated stress hormone levels relate to Epstein-Barr virus reactivation in astronauts. *Psychosom Med* 63:891–895.
7. Grinde B. 2013. Herpesviruses: latency and reactivation – viral strategies and host response. *J Oral Microbiol* 5:22766.
8. Donofrio G, Cavirani S, Van Santen V, Flammini CF. 2005. Potential secondary pathogenic role for bovine herpesvirus 4. *J Clin Microbiol* 43:3421–3426.
9. Barton E, Mandal P, Speck SH. 2011. Pathogenesis and Host Control of Gammaherpesviruses: Lessons from the Mouse. *Annu Rev Immunol* 29:351–397.
10. Goodman LB, Loregian A, Perkins GA, Nugent J, Buckles EL, Mercorelli B, Kydd JH, Palù G, Smith KC, Osterrieder N, Davis-Poynter N. 2007. A point mutation in a herpesvirus polymerase determines neuropathogenicity. *PLoS Pathog* 3:1583–1592.
11. Zhang P, Wang J, Zhang X, Wang X, Jiang L, Gu X. 2020. Identification of AIDS-Associated Kaposi Sarcoma: A Functional Genomics Approach. *Front Genet* 10:1–12.
12. Stalder K, Vaz PK, Gilkerson JR, Baker R, Whiteley P, Ficorilli N, Tatarczuch L, Portas T, Skogvold K, Anderson GA, Devlin JM. 2015. Prevalence and clinical significance of herpesvirus infection in populations of Australian marsupials. *PLoS One* 10:1–15.
13. Morey JN, Boggero IA, Scott AB, Segerstrom SC. 2015. Current directions in stress and human immune function. *Curr Opin Psychol* 5:13–17.
14. Barrandeguy M, Vissani A, Olguin C, Becerra L, Miño S, Pereda A, Oriol J, Thiry E. 2008. Experimental reactivation of equine herpesvirus-3 following corticosteroid treatment. *Equine Vet J* 40:593–595.
15. das Neves CG, Mørk T, Thiry J, Godfroid J, Rimstad E, Thiry E, Tryland M. 2009. Cervid herpesvirus 2 experimentally reactivated in reindeer can produce generalized viremia and abortion. *Virus Res* 145:321–328.
16. Glaser R, Pearl DK, Kiecolt-Glaser JK, Malarkey WB. 1994. Plasma cortisol levels and reactivation of latent Epstein-Barr virus in response to examination stress. *Psychoneuroendocrinology* 19:765–772.

- 373 17. Sin YW, Annavi G, Dugdale HL, Newman C, Burke T, MacDonald DW. 2014. Pathogen burden,
374 co-infection and major histocompatibility complex variability in the European badger (*Meles*
375 *meles*). *Mol Ecol* 23:5072–5088.
- 376 18. King DP, Mutukwa N, Lesellier S, Cheeseman C, Chambers MA, Banks M. 2004. Detection of
377 Mustelid Herpesvirus-1 Infected European Badgers (*Meles meles*) in the British Isles. *J Wildl*
378 *Dis* 40:99–102.
- 379 19. Kent A, Ehlers B, Mendum T, Newman C, Macdonald DW, Chambers M, Buesching CD. 2017.
380 Genital tract screening finds widespread infection with mustelid gammaherpesvirus 1 in the
381 European badger (*Meles meles*). *J Wildl Dis* 54:133–137.
- 382 20. Tsai MS, Fogarty U, Byrne AW, O'keeffe J, Newman C, Macdonald DW, Buesching CD. 2020.
383 Effects of mustelid gammaherpesvirus 1 (*Musghv-1*) reactivation in european badger (*meles*
384 *meles*) genital tracts on reproductive fitness. *Pathogens* 9:1–17.
- 385 21. Dittmer DP, Damania B. 2013. Kaposi sarcoma associated herpesvirus pathogenesis (KSHV) -
386 An update. *Curr Opin Virol* 3:238–244.
- 387 22. Young LS, Yap LF, Murray PG. 2016. Epstein-Barr virus: More than 50 years old and still
388 providing surprises. *Nat Rev Cancer* 16:789–802.
- 389 23. Chastant-Maillard S. 2015. Impact of bovine herpesvirus 4 (BoHV-4) on reproduction.
390 *Transbound Emerg Dis* 62:245–251.
- 391 24. McLuckie AJ, Barrs VR, Lindsay S, Aghazadeh M, Sangster C, Beatty JA. 2018. Molecular
392 diagnosis of *Felis catus* gammaherpesvirus 1 (FcaGHV1) infection in cats of known retrovirus
393 status with and without lymphoma. *Viruses* 10.
- 394 25. Marenzoni ML, Sforza M, Stefanetti V, Casagrande Proietti P, Brignone L, Del Sero A, Falcioni
395 F, Orvieto S, Tamantini C, Tiburzi A, Valentini S, Coletti M, Timoney PJ, Passamonti F. 2014.
396 Detection of Equid herpesvirus type 2 and 5 DNA in uterine flushings of mares with
397 reproductive disorders. *Vet Microbiol* 174:570–576.
- 398 26. Gagnon CA, Tremblay J, Larochelle D, Music N, Tremblay D. 2011. Identification of a novel
399 herpesvirus associated with cutaneous ulcers in a fisher (*Martes pennanti*). *J Vet Diagnostic*
400 *Investig* 23:986–990.
- 401 27. Tseng M, Fleetwood M, Reed A, Gill VA, Harris RK, Moeller RB, Lipscomb TP, Mazet JAK,
402 Goldstein T. 2012. Mustelid herpesvirus-2, a novel herpes infection in northern sea otters
403 (*Enhydra lutris kenyoni*). *J Wildl Dis* 48:181–185.
- 404 28. Nicolas de Francisco O, Esperón F, Juan-Sallés C, Ewbank AC, das Neves CG, Marco A, Neves E,
405 Anderson N, Sacristán C. 2020. Neoplasms and novel gammaherpesviruses in critically
406 endangered captive European minks (*Mustela lutreola*). *Transbound Emerg Dis* 1–13.
- 407 29. Abade dos Santos FA, Monteiro M, Pinto A, Carvalho CL, Peleteiro MC, Carvalho P, Mendonça
408 P, Carvalho T, Duarte MD. 2020. First description of a herpesvirus infection in genus *Lepus*.
409 *BioRxiv Prepr* <https://doi.org/http://dx.doi.org/10.1101/2020.01.21.913723>.

30. George SC, Smith TE, Mac Cana PSS, Coleman R, Montgomery WI. 2014. Physiological stress in the Eurasian badger (*Meles meles*): Effects of host, disease and environment. *Gen Comp Endocrinol* 200:54–60.
31. Virgós E, Mangas JG, Blanco-Aguilar JA, Garrote G, Almagro N, Viso RP. 2004. Food habits of European badgers (*Meles meles*) along an altitude gradient of Mediterranean environments: A field test of the earthworm specialization hypothesis. *Can J Zool* 82:41–51.
32. Macdonald DW, Newman C, Buesching CD, Nouvellet P. 2010. Are badgers “Under The Weather”? Direct and indirect impacts of climate variation on European badger (*Meles meles*) population dynamics. *Glob Chang Biol* 16:2913–2922.
33. Nouvellet P, Newman C, Buesching CD, Macdonald DW. 2013. A Multi-Metric Approach to Investigate the Effects of Weather Conditions on the Demographic of a Terrestrial Mammal, the European Badger (*Meles meles*). *PLoS One* 8.
34. Woodroffe R, MacDonald DW. 1995. Female/Female Competition in European Badgers *Meles meles*: Effects on Breeding Success. *J Anim Ecol* 64:12.
35. Creel SR, Creel NM. 1991. Energetics, reproductive suppression and obligate communal breeding in carnivores. *Behav Ecol Sociobiol* 28:263–270.
36. Macdonald DW, Newman C, Stewart PD, Domingo-Roura X, Johnson PJ. 2002. Density-dependent regulation of body mass and condition in badgers (*Meles meles*) from Wytham Woods. *Ecology* 83:2056–2061.
37. Macdonald DW, Harmsen BJ, Johnson PJ, Newman C. 2004. Increasing frequency of bite wounds with increasing population density in Eurasian badgers, *Meles meles*. *Anim Behav* 67:745–751.
38. Hunter RG, McEwen BS. 2013. Stress and anxiety across the lifespan: Structural plasticity and epigenetic regulation. *Epigenomics* 5:177–194.
39. Van Lieshout SHJ, Badás EP, Mason MWT, Newman C, Buesching CD, Macdonald DW, Dugdale HL. 2020. Social effects on age-related and sex-specific immune cell profiles in a wild mammal: Immune cell profiles in a wild mammal. *Biol Lett* 16:0–6.
40. Dall’Ara P, Labriola C, Sala E, Spada E, Magistrelli S, Lauzi S. 2019. Prevalence of serum antibody titres against feline panleukopenia, herpesvirus and calicivirus infections in stray cats of Milan, Italy. *Prev Vet Med* 167:32–38.
41. Troyer RM, Beatty JA, Stutzman-Rodriguez KR, Carver S, Lozano CC, Lee JS, Lappin MR, Riley SPD, Serieys LEK, Logan KA, Sweanor LL, Boyce WM, Vickers TW, McBride R, Crooks KR, Lewis JS, Cunningham MW, Rovnak J, Quackenbush SL, VandeWoude S. 2014. Novel Gammaherpesviruses in North American Domestic Cats, Bobcats, and Pumas: Identification, Prevalence, and Risk Factors. *J Virol* 88:3914–3924.
42. Sebastiano M, Chastel O, De Thoisy B, Eens M, Costantini D. 2016. Oxidative stress favours herpes virus infection in vertebrates: A meta-analysis. *Curr Zool* 62:325–332.

- 447 43. Woodroffe R, Macdonald DW. 1995. Costs of breeding status in the European badger, *Meles*
448 *meles*. *J Zool* 235:237–245.
- 449 44. Macdonald DW, Newman C, Nouvellet PM, Buesching CD. 2009. An Analysis of Eurasian
450 Badger (*Meles meles*) Population Dynamics: Implications for Regulatory Mechanisms. *J*
451 *Mammal* 90:1392–1403.
- 452 45. Dugdale HL, Davison D, Baker SE, Ellwood SA, Newman C, Buesching CD, Macdonald DW.
453 2011. Female teat size is a reliable indicator of annual breeding success in European badgers:
454 Genetic validation. *Mamm Biol* 76:716–721.
- 455 46. Macdonald DW, Feber RE, Newman C, Buesching CD. 2015. Badgers in the rural landscape -
456 conservation paragon or farmland pariah? Lessons from the Wytham Badger Project. *Wildl*
457 *Conserv Farml*, 2nd ed. 2:65–95.
- 458 47. Bright Ross JG, Newman C, Buesching CD, Macdonald DW. 2020. What lies beneath?
459 Population dynamics conceal pace-of-life and sex ratio variation, with implications for
460 resilience to environmental change. *Glob Chang Biol* 26:3307–3324.
- 461 48. Sugianto NA, Newman C, Macdonald DW, Buesching CD. 2020. Reproductive and Somatic
462 Senescence in the European Badger (*Meles meles*): Evidence from Lifetime Sex-Steroid
463 Profiles. *Zoology* 141:125803.
- 464 49. Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: Improving the sensitivity of
465 progressive multiple sequence alignment through sequence weighting, position-specific gap
466 penalties and weight matrix choice. *Nucleic Acids Res* 22:4673–4680.
- 467 50. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular evolutionary genetics
468 analysis across computing platforms. *Mol Biol Evol* 35:1547–1549.
- 469 51. R-Development-Core-Team. 2019. R: A language and environment for statistical computing. R
470 Foundation for Statistical Computing, Vienna, Austria.
- 471 52. Wilson EB. 1927. Probable Inference, the Law of Succession, and Statistical Inference. *J Am*
472 *Stat Assoc* 22:209.
- 473 53. Mandrekar JN. 2010. Receiver operating characteristic curve in diagnostic test assessment. *J*
474 *Thorac Oncol* 5:1315–1316.
- 475 54. Buesching CD, Heistermann M, Macdonald DW. 2009. Seasonal and inter-individual variation
476 in testosterone levels in badgers *meles meles*: Evidence for the existence of two
477 endocrinological phenotypes. *J Comp Physiol A Neuroethol Sensory, Neural, Behav Physiol*
478 195:865–871.
- 479 55. Dugdale HL, Griffiths A, Macdonald DW. 2011. Polygynandrous and repeated mounting
480 behaviour in European badgers, *Meles meles*. *Anim Behav* 82:1287–1297.
- 481 56. Ryder JJ, Pastok D, Hoare MJ, Bottery MJ, Boots M, Knell RK, Atkinson D, Hurst GDD. 2013.
482 Spatial variation in food supply, mating behavior, and sexually transmitted disease epidemics.
483 *Behav Ecol* 24:723–729.

57. Munday PE, Pritchard G, Harris JRW, Taylor-Robinson D. 1983. Prevalence of chlamydial infection in promiscuous women. *Sex Transm Infect* 59:103–104.
58. White J, Richard M, Massot M, Meylan S. 2011. Cloacal bacterial diversity increases with multiple mates: Evidence of sexual transmission in female common lizards. *PLoS One* 6.
59. Poiani A. 2000. Sexually Transmitted Diseases: A Possible Cost of Promiscuity in Birds? *Auk* 117:1061–1065.
60. Sugianto NA, Dehnhard M, Newman C, Macdonald DW, Buesching CD. 2021. A non-invasive method to assess the reproductive status of the European badger (*Meles meles*) from urinary sex-steroid metabolites. *Gen Comp Endocrinol* 301:113655.
61. François S, Vidick S, Sarlet M, Desmecht D, Drion P, Stevenson PG, Vanderplasschen A, Gillet L. 2013. Illumination of Murine Gammaherpesvirus-68 Cycle Reveals a Sexual Transmission Route from Females to Males in Laboratory Mice. *PLoS Pathog* 9.
62. Costantini D, Seeber PA, Soilemetzidou SE, Azab W, Bohner J, Buuveibaatar B, Cziráj G, East ML, Greunz EM, Kaczensky P, Lamglait B, Melzheimer J, Uiseb K, Ortega A, Osterrieder N, Sandgreen DM, Simon M, Walzer C, Greenwood AD. 2018. Physiological costs of infection: Herpesvirus replication is linked to blood oxidative stress in equids. *Sci Rep* 8:1–10.
63. Urbaniak C, Lorenzi H, Thissen J, Jaing C, Crucian B, Sams C, Pierson D, Venkateswaran K, Mehta S. 2020. The influence of spaceflight on the astronaut salivary microbiome and the search for a microbiome biomarker for viral reactivation. *Microbiome* 8:56.
64. Kangro HO, Osman HK, Lau YL, Heath RB, Yeung CY, Ng MH. 1994. Seroprevalence of antibodies to human herpesviruses in England and Hong Kong. *J Med Virol* 43:91–96.
65. Donahue JG. 1995. The incidence of herpes zoster. *Arch Intern Med* 155:1605–1609.
66. Stowe RP, Kozlova E V., Yetman DL, Walling DM, Goodwin JS, Glaser R. 2007. Chronic herpesvirus reactivation occurs in aging. *Exp Gerontol* 42:563–570.
67. Gouin JP, Hantsoo L, Kiecolt-Glaser JK. 2008. Immune dysregulation and chronic stress among older adults: A review. *Neuroimmunomodulation* 15:251–259.
68. Newman C, Macdonald DW, Anwar MA. 2001. Coccidiosis in the European badger, *Meles meles* in Wytham Woods: Infection and consequences for growth and survival. *Parasitology* 123:133–142.
69. Albery G, Newman C, Ross JB, Bansal S, Buesching C. 2020. Negative density-dependent parasitism in a group-living carnivore <https://doi.org/10.1101/2020.06.15.153726>.
70. Kelly TR, Karesh WB, Johnson CK, Gilardi KVK, Anthony SJ, Goldstein T, Olson SH, Machalaba C, PREDICT Consortium, Mazet JAK. 2017. One Health proof of concept: Bringing a transdisciplinary approach to surveillance for zoonotic viruses at the human-wild animal interface. *Prev Vet Med* 137:112–118.

Figure 1: Difference of genital MusGHV-1 prevalence between seasons (Logistic regression analysis)

Figure 2: Difference of genital MusGHV-1 prevalence between exact age (A) and age groups (B) (Logistic regression analysis)

Figure 3: Comparison of body condition score difference of adults between genital MusGHV-1 reactivation status. The p-values of each Kruskal-Wallis test results are showing above each box plot.

Figure S1: Data distribution of cub percentage (n=251). The cutoff point 30% (red dashed line) is used to divide the tail and the distribution in the left.

Figure S2: BCI data distribution by age groups showing distinct distribution of cub BCI comparing with other groups (A)

Figure S3: Diagnostic plot of the final multivariable logistic regression model of juvenile genital MusGHV-1 reactivation

DHARMA residual diagnostics

Figure S4: Diagnostic plot of the final multivariable logistic regression model of adult genital MusGHV-1 reactivation