1 QTL mapping: an innovative method for investigating the genetic 2 determinism of yeast-bacteria interactions in wine

3

4 Louise Bartle¹⁺, Emilien Peltier^{2,3,4}, Joanna Sundstrom^{1,5}, Krista Sumby^{1,5}, James G Mitchell⁶, Vladimir
5 Jiranek^{1,5}, Philippe Marullo^{2,3+}

6 ¹The University of Adelaide, Department of Wine Science, Urrbrae, SA, 5064, Australia

7 ²Université de Bordeaux, Unité de recherche Œnologie EA 4577, USC 1366 INRAe, Bordeaux INP;
8 ISVV, Villenave d'Ornon, France

9 ³Biolaffort, Bordeaux, France

10 ⁴Université de Strasbourg, CNRS, GMGM UMR 7156, Strasbourg, France

11 ⁵ Australian Research Council Training Centre for Innovative Wine Production, PMB1, Glen Osmond,
12 SA, 5064, Australia

13 ⁶Flinders University, College of Science and Engineering, Adelaide, SA, Australia

14 ⁺Corresponding Authors: Louise Bartle louise.bartle@adelaide.edu.au, Philippe Marullo
15 philippe.marullo@u-bordeaux.fr

16

17 **Keywords:** Yeast-bacteria interactions, malolactic fermentation, QTL, wine, *SSU1*

18

19 Abstract

20 The two most commonly used wine microorganisms, *Saccharomyces cerevisiae* yeast and
 21 *Oenococcus oeni* bacteria, are responsible for completion of alcoholic and malolactic fermentation
 22 (MLF), respectively. For successful co-inoculation, *S. cerevisiae* and *O. oeni* must be able to
 23 complete fermentation, however, this relies on compatibility between yeast and bacterial strains. For
 24 the first time, Quantitative Trait Loci (QTL) analysis was used to elucidate whether *S. cerevisiae*
 25 genetic makeup can play a role in the ability of *O. oeni* to complete MLF. Assessment of 67 progeny
 26 of an *S. cerevisiae* SBxGN cross, co-inoculated with a single *O. oeni* strain, SB3, revealed a major
 27 QTL linked to MLF completion by *O. oeni*. This QTL encompassed a well-known translocation, XV-t-
 28 XVI, that results in increased *SSU1* expression and is functionally linked with numerous phenotypes
 29 including lag phase duration and sulfite export and production. A reciprocal hemizyosity assay was
 30 performed to elucidate the effect of the gene *SSU1* in the SBxGN background. Our results instead
 31 revealed a strong effect of *SSU1* haploinsufficiency on *O. oeni*'s ability to complete malolactic
 32 fermentation during co-inoculation, and paves the way for the implementation of QTL mapping
 33 projects for deciphering the genetic bases of microbial interactions.

34

35 Key points:

36 For the first time QTL analysis has been used to study yeast-bacteria interactions.

37 A QTL encompassing a translocation, XV-t-XVI, was linked to MLF outcomes.

38 *S. cerevisiae* *SSU1* haploinsufficiency positively impacted MLF by *O. oeni*.

39

40 Introduction

41 Fermented beverages are the result of biotransformation of complex organo-chemical matrices by
42 microbial communities of moulds, yeast, bacteria and bacteriophage (Bokulich and Bamforth 2013;
43 Mounier et al. 2008; Renouf et al. 2007). Within these communities, the growth rate and metabolic
44 activities of each microbial species depends on (i) the biochemical composition of the medium, (ii) the
45 physicochemical conditions of the process (e.g. converting sugars to ethanol during juice
46 fermentation), (iii) the physiological state of the microbes and (iv) cell-to-cell contact and metabolite
47 interaction among microbial species.

48 Fermenting grape juice is a fast-changing environment that is especially interesting for studying how
49 the two most common wine microbes, the yeast *Saccharomyces cerevisiae* and lactic acid bacterium
50 *Oenococcus oeni*, coexist and interact. In an oenological context, microbial interactions may affect the
51 final profile of volatile compounds (Chasseriaud et al. 2018, Renault et al. 2015, 2016), that
52 subsequently can be detected as changes in wine sensorial complexity (Tempère et al. 2018). The
53 importance of microbial interactions in wine is evident from the wide number of studies focusing on
54 co-inoculated or sequential inoculation of *S. cerevisiae* and *O. oeni*, with the aim of decreasing overall
55 fermentation time while maintaining or increasing wine quality (Abrahamse and Bartowsky 2012;
56 Cañas et al. 2012, 2015; Knoll et al. 2012).

57 The mechanisms of yeast-lactic acid bacteria (LAB) interactions during juice fermentation have been
58 reviewed recently (Bartle et al. 2019). Broadly, microbial interactions may include cell-cell contact
59 (Nissen et al. 2003, 2004; Renault et al. 2013) or production of small metabolites (Renault et al. 2009;
60 Sadoudi et al. 2012) and macromolecules (Comitini et al. 2005; Jarosz et al. 2014) that can inhibit
61 and/or activate the growth and activity of interacting microbes. Understanding the molecular
62 mechanisms of yeast-LAB interactions is a challenging task due to the confounding effects of the
63 evolving complex growth environment and microbial metabolite production, but the benefits of such
64 work include optimisation of yeast-LAB co-inoculation strategies for implementation in wineries.

65 *S. cerevisiae* and *O. oeni* interactions can affect their ability to complete alcoholic (AF) and malolactic
66 fermentation (MLF), respectively (Bartle et al. 2019). Yeast may produce metabolic compounds that
67 can inhibit LAB growth, including ethanol (Capucho and San Romão 1994; Gao and Fleet 1995;
68 Guzzo et al. 2000), SO₂ (Osborne and Edwards 2006), short and medium-chain fatty acids (Alexandre
69 et al. 2004; Capucho and San Romão 1994), and antimicrobial peptides (Atanassova et al. 2003;
70 Mendoza et al. 2010; Nehme et al. 2010). Yeast and LAB also have the potential to interact physically
71 in the form of mixed species biofilms (Arroyo-López et al. 2012; Bartle et al. 2019; Furukawa et al.
72 2010) or through co-aggregation (Furukawa et al. 2011), though to date there have not been reports
73 of this occurring between *S. cerevisiae* and *O. oeni*.

74 In addition to chemical and physical interactions, *S. cerevisiae* gene expression is affected by co-
75 inoculation with *O. oeni* (Rossouw et al. 2012). Differential gene expression in *S. cerevisiae* in
76 response to co-inoculation with *O. oeni* included up-regulation of genes related to yeast stress
77 response and possible competition for sulfur compounds compared to *S. cerevisiae* alone (Rossouw

et al. 2012). Several studies have also reported strain-specific compatibility between yeast and LAB during co-inoculation (Abrahamse and Bartowsky 2012; Antalick et al. 2013; Arnink and Henick-Kling 2005; Comitini and Ciani 2007; Rossouw et al. 2012; Tristezza et al. 2016). Considering the strain specificity of compatibility outcomes, the intraspecific genetic variability of interacting species requires further investigation and analysis. To our knowledge, the identification of genetic variations to explain “strain compatibility” is a novel approach.

For *S. cerevisiae*, the genetic determinism of any complex trait can be investigated by mapping quantitative trait loci (QTLs) in a segregating progeny (Liti and Louis 2012). In the context of wine, this strategy has been used for elucidating the genetic basis of many traits of industrial interest (Peltier et al. 2019) including acetic acid production (Salinas et al. 2012), rate of nitrogen uptake (Brice et al. 2014; Jara et al. 2014), resistance to stuck fermentation (Marullo et al. 2019), resistance to low pH (Martí-Raga et al. 2017), and the production of aroma compounds (Eder et al. 2018; Huang et al. 2014; Roncoroni et al. 2011; Steyer et al. 2012). To date, QTL mapping has been performed for single pure cultures focusing on traits related to yeast fitness or effect on wine quality. However, this strategy may be applied to any trait resulting in measurable phenotypic variability. In the present work, we applied QTL mapping to delineate how *S. cerevisiae* genetic variability may affect the success of MLF in co-inoculated fermentations with a commercial strain of *O. oeni*.

95

96 **Materials and methods**

97 **Media**

98 Shiraz juice: Shiraz grapes (2017 vintage, Coombe vineyard, The University of Adelaide, Waite
99 Campus, Urrbrae, South Australia) were harvested, de-stemmed, crushed and left to macerate at 0
100 °C for 7 days to enable polyphenolic extraction. Shiraz must be pressed and the juice stored at -20 °C
101 until required. No SO₂ or antibacterial agents were added to the juice during pressing. Prior to
102 experimentation Shiraz juice was filtered (0.45 µm, catalogue # FHT45, Air-Met Scientific, Victoria,
103 Australia) to remove grape matter and solids. Initial measurements of total sugar were estimated by
104 refractometry and sugar reduced to 250 g L⁻¹ by addition of water. L-malic acid was increased to 2.5 g
105 L⁻¹ by addition of pure L-malic acid and pH was decreased to 3.5 by addition of tartaric acid, followed
106 by addition of 100 mg L⁻¹ diammonium phosphate. Finally, the juice was filter sterilised (0.2 µm).

107 Liquid de Man, Rogosa and Sharpe medium (MRS; catalogue # AM103, Amyl Media, Victoria,
108 Australia), supplemented with 20% apple juice (MRSAJ) was used for growing bacteria prior to
109 inoculation. MRS was autoclaved and sterile apple juice (0.2 µm filtered) added post sterilisation and
110 before use. MRSAJ supplemented with agar (2%) and cycloheximide (0.5%) following sterilisation of
111 the medium, was used for enumeration of bacteria.

112 All yeast strains were initially streaked for single colonies on YPD agar (2% glucose, 2% peptone, 1%
113 yeast extract, 2% agar) and grown at 28 °C, before growth of single isolates in YPD (2% glucose, 2%
114 peptone, 1% yeast extract) at 28 °C overnight. If required, Geneticin (G418, 100 µg mL⁻¹; catalogue #

G8168, Sigma-Aldrich, New South Wales, Australia) was added to YPD cultures to select for strains carrying the *KanMX* deletion cassette.

Strains and fermentations

Yeast strains

Strains used in this work are listed in Table 1. QTL analysis was performed using the SBxGN yeast background. SBxGN is the F1-hybrid of SB and GN strains, two diploid, fully homozygous strains derived from the wine starters Actiflore® BO213 and Zymaflore® VL1, respectively (Peltier et al. 2018b). The population used for QTL mapping consisted of 67 haploid progeny clones derived from the hybrid BN, an isogenic variant of SBxGN (Marullo et al. 2007a). These haploid meiotic progenies have been previously genotyped by whole genome sequencing (Martí-Raga et al. 2017). The effect of the gene *SSU1* was assayed using the reciprocal hemizygosity assay by deleting each parental copy of *SSU1* individually in the SBxGN F1-hybrid (Steinmetz et al. 2002a). The reciprocal hemizygous hybrids Δ G092 and Δ S092 were previously obtained as described by Zimmer and colleagues (2014).

Yeast cell concentration and viability

Yeast live cell concentrations were determined by flow cytometry. Yeast were diluted 100-times in sterile phosphate buffered saline (PBS), then stained with propidium iodide at a final concentration of 0.1 mg mL⁻¹. Samples were analysed using a Guava® easyCyte™ 12HT flow cytometer (Luminex, Yokohama, Japan; previously Millipore, Darmstadt, Germany). Yeast concentrations were adjusted to inoculate sterile Shiraz juice at a final rate of 5 x 10⁶ live cells mL⁻¹.

Bacteria

Freeze-dried SB3 (Laffort®, South Australia, Australia) was grown anaerobically in MRSAJ for four days at 30 °C in 20% CO₂. Twenty-four hours prior to inoculation, bacteria were centrifuged (2,236 x g), the supernatant discarded and the cell pellet washed in sterile Shiraz juice before overnight incubation in fresh sterile Shiraz juice at 30 °C. Bacteria were adjusted to an OD₆₀₀ of 0.55 immediately prior to inoculation. For QTL library fermentations, 200 µL of bacterial culture were added to each fermentation vessel manually through a silicone septum with a 21-gauge needle. For the hemizygote fermentations, 200 µL of bacterial culture were transferred from a 96-well deep well plate to each fermentation vessel using an automated fermentation system (developed after performing the QTL experiment).

Fermentations

Fermentations were conducted using an automated fermentation system built on an EVO Freedom workdeck (Tecan, Männedorf, Switzerland; Fig. S1: Online resource 2). The system enabled 384 concurrent fermentations at a volume of up to 25 mL. Full details of the system were described by Hranilovic et al. (2018) and can also be found on the University of Adelaide Biotechnology and

Fermentation Facility website (<https://sciences.adelaide.edu.au/agriculture-food-wine/research/biotechnology-and-fermentation-facility>).

Fermentation vessels were filled with 20 mL of sterile Shiraz juice and inoculated with yeast (5×10^6 live cells mL⁻¹) followed by LAB inoculation 24 hours later. Sampling occurred daily, and fermentations were homogenised by stirring prior to sampling. For the QTL mapping experiment, both parental strains (SB and GN), the hybrid BN and the 67 haploid progenies were fermented as pure cultures (in duplicate) or co-inoculated with SB3 (in triplicate). To test the effect of *SSU1*, hemizygous and wild type F1- hybrids were assessed in triplicate for both pure and co-inoculated fermentations with SB3.

Fermentation monitoring

Glucose and fructose consumption

Glucose and fructose concentrations were determined enzymatically using commercially available kits (catalogue # K-FRUGL, Megazyme, Bray, Ireland) following methods modified by Walker et al. (2014). Glucose and fructose consumption were used as determinants for progress of alcoholic fermentations, which were deemed complete when total glucose plus fructose concentration was < 4 g L⁻¹.

The amount of glucose/fructose consumed over time was modelled by local polynomial regression fitting with the R-loess function, setting the span parameter to 0.8. Five kinetics parameters were extracted from the model, which are described in Table 2.

L-malic acid concentration

L-malic acid was measured using an enzymatic test kit (catalogue # 4A165, Vintessential Laboratories, Victoria, Australia) with modifications so that a plate-reader/spectrophotometer (Infinite 200 PRO, Tecan, Männedorf, Switzerland) could be used to measure absorbance. Specifically, each well of a 96 well micro-titre plate was dosed with 70 µL buffer (0.1M gly-gly, 0.1M L-glutamate, pH 10), 14 µL nicotinamide adenine dinucleotide (40 mg mL⁻¹), 70 µL distilled water, 0.7 µL glutamate oxaloacetate transaminase (800 U mL⁻¹) and 5 µL of sample or one of the L-malic acid standards (ranging from 0 - 3.0 g L⁻¹). The plate was incubated at 22 °C for 3 minutes and the first absorbance was read at 340 nm; 7 µL of the 1:10 diluted L-malate dehydrogenase (12,000 U mL⁻¹) was added and mixed into each well; the plate was incubated at 22 °C for 15 minutes before the second absorbance was measured at 340 nm. L-malic acid in each sample was calculated from standard curves prepared with known L-malic acid concentrations. L-malic acid degradation was used as the determinant for MLF progress. L-malic acid consumed over time was modelled by local polynomial regression fitting with the R-loess function setting the span parameter to 0.75. MLF was deemed complete when L-malic acid concentration was < 0.1 g L⁻¹ and designated *tend-MLF* (Table 2).

L-malic acid end point parameters were determined for yeast alone and yeast-SB3 co-inoculation fermentations. These parameters were: percentage of L-malic acid consumed or produced by yeast

alone in relation to the starting L-malic acid concentration of 2.5 g L⁻¹ (*Pct_malic_AF*), and percentage of L-malic acid consumed by yeast and LAB in co-inoculated fermentations (*Pct_malic_co*).

To estimate the overall L-malic acid reduction by LAB when co-inoculated with yeast, the average concentration of L-malic acid for co-inoculated fermentations at the end of experimentation was subtracted from the average L-malic acid concentration for corresponding yeast-alone fermentations. This parameter was designated *Malic_acid_LAB_consumed*. The end of the experiment was defined as either reduction of L-malic acid to < 0.1 g L⁻¹ or approximately 11 days after the first yeast-LAB pair completed MLF.

A summary of all parameters assessed in this study can be found in Table 2.

Statistical analysis

All statistical analyses were performed using R versions 3.4.4 or higher (R Core Team 2017). Kendall correlation coefficient test was performed using R/stats package v3.6.2 (R Core Team 2017). The QTL mapping analysis was performed with the R/qtl package (Broman et al. 2003) by using the Haley-Knott regression model that provides a fast approximation of standard interval mapping (Haley and Knott 1992). A threshold corresponding to a 5% and 10% false discovery rate (FDR) was computed by performing 1000 permutations in order to assess the significance of the LOD score for QTL peaks (Churchill and Doerge 1994). The overall procedure was described by Peltier et al. (2018b) for multiple environments mapping.

Linear modelling was performed to evaluate the effect of allele, yeast background and translocation on MLF and AF parameters using the following formula:

$$y_{ij} = m + B_i + T_j + B_iT_j + \epsilon_{ij}$$

Where y_{ij} is the value for the background i ($i = 1,2$) with translocation j ($j = 1,2$), m is the overall mean, B_i is the yeast background effect, T_j is the translocation effect, B_iT_j is the interaction effect between yeast background and translocation, ϵ_{ij} is the residual error. Tukey post-hoc test ($\alpha = 0.05$) was used to elucidate differences between ANOVA test groups.

Results

Biometric assessment of MLF completion in the SBxGN progeny population

To identify QTLs influencing the completion of MLF, L-malic acid consumption by *O. oeni* was measured in co-inoculated Shiraz grape juice fermentations. Alone, *S. cerevisiae* strains from the BN progeny were able to consume a fraction of L-malic acid (Fig. 1a). The concentration of residual L-malic acid at the end of AF in yeast-alone fermentations ranged from 1.41 g L⁻¹ to 2.75 g L⁻¹, which

corresponded to between 44% consumption and 10% production of L-malic acid in respect to the starting concentration of 2.5 g L⁻¹ (Fig. 1b; Table S1: Online resource 1).

In the present study, the focus was the impact of yeast genotype on LAB MLF efficiency. Therefore, we measured L-malic acid consumption over time for fermentations co-inoculated with *S. cerevisiae* strains and LAB SB3 (Fig. 1c). As expected, L-malic acid consumption was much higher for many of the yeast-LAB co-inoculated fermentations. However, SB3 was only able to complete MLF in 39 of the 71 co-inoculated fermentations (Fig. 1d). Since LAB were only able to complete MLF when co-inoculated with some of the SBxGN progenies, this provided evidence of strong yeast-LAB interactions.

Though there were differences in residual L-malic acid across fermentations with different yeast strains, the ability of yeast to consume L-malic acid (as seen for yeast-alone fermentations, Fig. 1a) did not seem to impact MLF completion time by SB3 in co-inoculations. Kendall rank correlation coefficient revealed only a weak positive correlation (0.21, $p = 0.009$) between the amount of L-malic acid consumed by yeast and SB3 MLF completion time.

Fig. 1 a) L-malic acid concentration measured over the course of the experiment for yeast-alone fermentations for the population of 67 SBxGN yeast progeny. Values are the mean of duplicates. **b)** All yeast-alone strains ranked by percentage of L-malic acid consumption (positive %) or production (negative %), measured at the end of the experiment in relation to the starting L-malic acid concentration of 2.5 g L⁻¹. Percentages were calculated from the mean of duplicates. Colours indicate yeast parental strains: BN (orange), SB (blue) and GN (purple). All other yeast progeny are shown in green. **c)** MLF progress measured for yeast co-inoculated with SB3 LAB. Values are the mean of triplicates. The horizontal line at 0.1 g L⁻¹ indicates when MLF was deemed complete. **d)** All yeast strains in co-inoculations with SB3 LAB were ranked by the percentage of L-malic acid consumed, measured at the end of the experiment. Percentages were calculated from the mean of triplicates. Colours are the same as panel b

QTL mapping

To determine the concentration of L-malic acid consumed only by LAB at the end of the experiment (as defined in materials and methods), the average final L-malic acid concentration found in yeast-LAB co-inoculations was subtracted from the corresponding L-malic acid concentration in yeast-alone fermentations. This new parameter, the L-malic acid consumed by LAB (*Malic_acid_LAB_consumed*), provides a proxy for SB3 MLF efficiency after co-inoculation with different yeast strains. This new parameter has a nearly continuous distribution among the SBxGN progeny (Fig. 2a). Genetic regions linked to the variation of this trait were tracked by applying a linkage analysis. Despite the small number of progenies tested, two loci were linked to this phenotype at an FDR of 10% (Fig. 2b). One QTL peak, located on *S. cerevisiae* chromosome XVI, achieved a LOD score of 7.58 which is highly significant with respect to the threshold value of 4.58 that was estimated by 1000 permutations with an FDR of 5%. An additional peak, located on *S. cerevisiae* chromosome XV had a lower LOD score of 4.02. This LOD score reaches the threshold value of 4.00 which corresponds to an FDR of 10%.

The best marker of this linkage analysis was located at genomic position XVI_374156 and was therefore named XVI_374. Due to the density of markers surrounding XVI_374 (6 markers within 817 bp) and *SSU1* spanning the genomic region of chromosome XVI: 373793-375169, there was high confidence in the specificity of *SSU1* being the target of the QTL peak. The inheritance of this marker impacts the consumption of L-malic acid by the LAB SB3. Indeed, co-inoculated fermentations with yeast progenies carrying the SB allele of XVI_374 consumed more L-malic acid than those performed with yeast progenies carrying the GN allele (Fig. 2c). Interestingly, this phenotypic discrepancy is not due to the ability of yeast to consume L-malic acid. Additionally, in yeast-alone fermentations, the inheritance of XVI_374 from either SB or GN did not alter the percentage of L-malic acid consumed by the yeast (Fig. 2d). In contrast, most of the strains containing the yeast SB allele for this QTL allowed SB3 to complete MLF (Fig. 2c). Altogether, these data provide clear evidence that genetic regions of the *S. cerevisiae* genome have a direct impact on the metabolic activity of LAB during co-inoculation.

One peak detected did fall within the 10% FDR (Fig. 2b) and therefore warrants discussion. This peak was located on chromosome XV, at the marker XV_162503 which corresponds to the gene *PHM7*. Although *PHM7* is a gene of unknown function, in the SBxGN background a genetic linkage in this region was predictable since the left arm of chromosome XV is physically linked to chromosome XVI in the strain GN. This well-documented translocation event was previously named XV-t-XVI (Zimmer et al. 2014; Fig. 3) and segregates in a mendelian fashion in the SBxGN progeny. The XV-t-XVI translocation has been demonstrated to impact several fermentation-related phenotypes including lag phase duration, fermentation rate, and SO₂ production. The molecular basis of such phenotypes is due to modification of the promoter environment of the *SSU1* gene (Fig. 3) that encodes Ssu1p, a transmembrane sulfite efflux pump (Marullo et al. 2020; Peltier et al. 2018b; Zimmer et al. 2014). These previous findings suggest that *SSU1* would be a relevant candidate gene able to explain the yeast-LAB interaction observed in our experiment.

Fig. 2 a) Yeast ranked by the concentration of L-malic acid that was able to be consumed by SB3 during co-inoculation with each yeast strain. Values determined by subtracting the average residual L-malic acid concentration of the yeast-alone fermentations from the average residual L-malic acid concentration in the corresponding yeast-LAB co-inoculation fermentations. Colours indicate yeast parental strains: BN (orange), SB (blue) and GN (purple). All other yeast progeny are shown in grey. **b)** Genomic location of QTL peaks for the parameter *Malic_acid_LAB_consumed*. Threshold values are estimated from 1000 permutations and 5% FDR, indicated by the solid horizontal line. The dotted horizontal line indicates a LOD score threshold of 4. Significant (peak above threshold) and potential (peaks near a LOD score of 4) QTLs were found on chromosomes XV (left) and XVI (right). **c)** Distribution of yeast progenies with respect to the concentration of L-malic acid consumed by SB3 in co-inoculations with each yeast strain. Progenies are grouped by yeast background (SB, left; GN, right). **d)** Distribution of yeast progenies based on percentage of L-malic acid consumed (measured at the end of experimentation) for yeast alone (left panel) or when co-inoculated with SB3 (right panel). Progenies are grouped by yeast background

Fig. 3 Representation of the translocation located in SBxGN that results in increased *SSU1* gene expression due to reduced proximity between *SSU1* and promotor regions. SBxGN has an XV-t-XVI translocation that leads to a single copy of wild-type XV and XVI chromosomes (all black) and reciprocal XV and XVI translocated chromosomes (black and white). Hemizygous strains SΔG092 and GΔS092 with a single functional *SSU1* allele (orange) were generated to perform a reciprocal hemizygosity assay. Hemizygous strains were created by replacing a single copy of *SSU1* with a KanMX cassette (blue)

Functional study of a QTL closely related to *SSU1*

To determine the possible influence of *SSU1* on MLF outcome, we applied a reciprocal hemizygosity assay, enabling evaluation of the impact of the two parental alleles in an isogenic context, as proposed by Steinmetz et al. (2002b). To test this hypothesis, we used the hemizygous hybrids SΔG092 and GΔS092 in which one copy of the *SSU1* gene was replaced with the *KanMX4* cassette (Zimmer et al. 2014). Such strains are isogenic to the F1-hybrid SBxGN but carry only one functional copy of *SSU1*: SΔG092 carries the *SSU1* allele of GN while GΔS092 carries the *SSU1* allele of SB. The two hemizygous hybrids were compared to the SBxGN hybrid carrying both functional *SSU1* alleles.

Co-inoculated and yeast-alone fermentations were carried out in the same conditions applied for QTL mapping. In co-inoculated fermentation, all the malolactic fermentations were completed. However, the time taken for SB3 to reduce L-malic acid to $< 0.1 \text{ g L}^{-1}$ was significantly different depending on the yeast strain (Fig. 4a). Indeed, both hemizygous hybrids enabled shorter MLF completion time (by at least 48 hours) than the parental hybrid SBxGN, revealing a strong haploinsufficiency effect of the *SSU1* deletion. Surprisingly, MLF completion was not significantly impacted by inheritance of either the SB or GN allele (Fig. 4b). This result led to the conclusion that the *SSU1* gene was not the main cause of the SBxGN QTL on chromosome XVI that was based on L-malic acid consumption.

In addition to MLF, AF kinetics of isogenic hybrids co-inoculated with and without SB3 were also determined. The AF completion time (*tend-AF*) was significantly longer for SΔG092 compared to SBxGN and GΔS092 (Fig. 4c and d), but was unaffected by the presence of LAB. Additionally, in co-inoculated fermentations, SΔG092 (functional GN allele) had a significantly slower fermentation (Fig. 4c; ANOVA, Tukey post-hoc, $p < 0.05$, Table S2: Online resource 2). Differences in AF kinetics between *SSU1* hemizygous hybrids have been previously reported, where the GN allele correlated to reduced lag phase, increased cell viability (Zimmer et al. 2014) and longer fermentation completion

time (Peltier et al. 2018b) in connection with the SO₂ content of grape juice. For the present work, no sulfite was added to the grape juice, removing the influence of this antimicrobial agent. Therefore, the differences of fermentation kinetics observed for the SΔG092 may be in part due to yeast-LAB interactions. This unexpected result suggests that LAB co-inoculation with yeast may impact yeast fermentation fitness and allelic variations of *SSU1* in yeast could modulate such interactions.

Fig. 4 a) Time taken for SB3 to complete malolactic fermentation during co-inoculation with SBxGN, SΔG092 and GΔS092. Bar colour indicates the yeast *SSU1* genotype. Significant differences (ANOVA, Tukey post-hoc; $p < 0.05$) between yeast are indicated by *. **b)** L-malic acid consumption over time by SB3 when co-inoculated with SBxGN (triangle), SΔG092 (circle) or GΔS092 (square). Lines demonstrate fitting of local polynomial regression using the R-loess function, span = 0.75. **c)** Time to complete alcoholic fermentation for yeast alone (filled bars) vs yeast co-inoculated with SB3 (colourless bars). Significant differences (ANOVA, Tukey post-hoc; $p < 0.05$) are indicated by *. **d)** Total sugar consumption over time by SBxGN (triangle), SΔG092 (circle) and GΔS092 (square) alone (dark), or co-inoculated with SB3 (light). Lines demonstrate fitting of local polynomial regression using the R-loess function, span = 0.8. Values are the mean of triplicates and error bars are the standard deviation

Discussion

Implementation of QTL mapping for narrowing down genetic regions controlling microbial interactions

Previously, studies investigating yeast-LAB interactions during juice fermentation relied on AF and MLF kinetics and production of volatile and non-volatile compounds (Arnink and Henick-Kling 2005; Comitini and Ciani 2007; Mendoza et al. 2010; Nehme et al. 2008, 2010). Many combinations of yeast and LAB, either sequentially or co-inoculated in juice or wine, have revealed that yeast-LAB compatibility is strain specific (Comitini et al. 2005; Comitini and Ciani 2007; Muñoz et al. 2014). Considering the differences reported for different yeast and LAB strains, and production of different metabolites by yeast strains, there is no question of the influence that yeast genetic makeup has on co-inoculation outcomes. In addition to studies on strain combination and metabolite production, further work has revealed differential *S. cerevisiae* gene expression in response to co-inoculation with *O. oeni* (Rossouw et al. 2012). Though insightful, none of these works have identified specific genetic differences between yeast strains that may influence compatibility with *O. oeni*. Hence, the present work has laid a foundation for understanding how *S. cerevisiae* genetic makeup can impact MLF outcomes during co-inoculation with *O. oeni*. The application of a QTL mapping strategy was used to find specific influences of yeast genetic makeup on yeast-LAB interactions.

In the past, QTL mapping has been used for investigating plant-pathogen interactions unveiling links between plant genotype and pathogen resistance (Chen et al. 2010; Decroocq et al. 2005; Eun et al. 2016). In the current work, the same strategy was applied to the two main species involved in winemaking: *S. cerevisiae* and *O. oeni*, to delineate the influence of yeast genetic background in microbial interactions.

QTL mapping has been used numerous times for *S. cerevisiae* to determine the genotypic traits that influence yeast AF completion (Marullo et al. 2019), acetic acid production (Marullo et al. 2007b; Salinas et al. 2012), nitrogen uptake (Brice et al. 2014; Jara et al. 2014) and aroma compound production (Eder et al. 2018; Huang et al. 2014; Roncoroni et al. 2011; Steyer et al. 2012) which are important for fermentation progress and wine quality. However, to the best of our knowledge, QTL mapping has not been used to study yeast-bacteria interactions in wine. To achieve this task, we used the BN progeny clones (pop BN, Table 1) that have been previously used for investigating the genetic determinism of fermentation traits (Martí-Raga et al. 2017; Zimmer et al. 2014). Each progeny clone was co-inoculated (i.e. exposed) to LAB during fermentation and the percentage of L-malic acid consumed was measured. For the *S. cerevisiae* strains alone, L-malic acid at the end of fermentation ranged from 44% consumption to 10% production when compared to the starting concentration of 2.5 g L⁻¹ (Fig. 1b; Table S1: Online resource 1) and the ability of yeast to consume or produce this amount of L-malic acid is in agreement with previous findings (Delcourt et al. 1995; Peltier et al. 2018a; Yéramian et al. 2007). The continuous distribution of L-malic acid consumption or production

observed among the yeast progeny suggests that this trait is controlled by many genes. A study detailing those genes is currently under preparation (Peltier, pers. comm.).

Since *S. cerevisiae* strains can consume some of the L-malic acid present in grape must, the specific consumption of L-malic acid by LAB was determined by subtracting L-malic acid concentration in yeast-alone fermentations from values in co-inoculations. This computed trait (*Malic_acid_LAB_consumed*) was used to estimate the influence of yeast strain on *O. oeni* strain SB3's ability to complete MLF. The ability of SB3 to complete MLF in co-inoculated conditions showed a gaussian distribution, suggesting that several *S. cerevisiae* genes control this phenotype (Fig. 2a). Although the number of progenies tested was low for achieving QTL mapping (67 clones), we implemented a linkage analysis that revealed a major QTL (XVI_374) with a LOD score above an FDR threshold of 5%. The marker for this major QTL was located within *SSU1*, giving reason to perform follow-up work with *SSU1* hemizygote strains. Beside this major QTL, one other loci (XV_162) reached a lower FDR threshold of approximately 10%. The QTL XV_162 marker corresponds to *PHM7*, a gene of unknown function. However, this marker was physically linked with the major QTL XVI_374 due to the presence of the XV-t-XVI translocation that was previously identified in this genetic background (Zimmer *et al.* 2014). Therefore, our strategy was useful for narrowing down two unlinked genetic regions that impacted the success of MLF by SB3 in co-inoculated fermentations.

Putative role of *SSU1* gene in yeast-LAB interactions

Once identified by linkage analysis, QTLs can be dissected at the gene and single-nucleotide polymorphism (SNP) level to further understand the underlying molecular genetic mechanism of phenotypic traits (Peltier *et al.* 2019). However, this functional characterisation is a complex task that cannot always be achieved. In the present study, there was no clear demonstration of the impact of natural allelic variations linked to SB3 MLF completion in co-inoculated fermentations.

The QTLs identified in this study link two markers, XVI_374 and XV_162, to the phenotype investigated. These two QTLs belonging to chromosomes XV and XVI must be considered as a single locus since they are physically linked by a reciprocal translocation event, XV-t-XVI, in the parental strain GN. This translocation is detected in numerous wine related strains (Marullo *et al.* 2020; Treu *et al.* 2014; Zimmer *et al.* 2014) and can impact numerous yeast phenotypes including lag phase duration, cell viability (Zimmer *et al.* 2014), growth rate in the presence of SO₂ (Marullo *et al.* 2020) and alcoholic fermentation kinetics (Peltier *et al.* 2018b). These phenotypes have been attributed to the modification of the promoter sequence of the gene *SSU1*, that encodes Ssu1p, an intermembrane transporter responsible for *S. cerevisiae* sulfite efflux (Park and Bakalinsky 2000). In GN, the translocation results in decreased distance between the *ADH1* promoter region and *SSU1*, triggering increased and constitutive *SSU1* expression (Zimmer *et al.* 2014). Interestingly, two other gross chromosomal rearrangements related to the *SSU1* promoter have been described in the past (García-Ríos *et al.* 2019; Pérez-Ortín *et al.* 2002) and were associated with increased *SSU1* expression as

well as increased sulfite resistance (García-Ríos et al. 2019; Marullo et al. 2020). These chromosomal rearrangements have facilitated adaptation of wine yeast, enabling increased SO₂ efflux from the yeast cell in oenological conditions, and has been reviewed recently (Divol et al. 2012; García-Ríos and Guillamón 2019). In terms of co-inoculation, yeasts that efficiently export sulfite could negatively impact *O. oeni* growth and MLF progress, as SO₂ can inhibit the internal ATPase of this lactic acid bacterium (Carrete et al. 2002). This is supported by the linkage analysis since the GN allele had a deleterious effect on MLF completion (Fig. 2d). We tested this hypothesis by reciprocal hemizyosity assay, including fermentations with and without LAB co-inoculation and determining the percentage of L-malic acid consumed by LAB. Although MLF was completed in all the co-inoculated fermentations, a strong haploinsufficiency effect was observed between the native hybrid SBxGN and the two hemizygous hybrids (Fig. 4a). Thus, the lack of one copy of *SSU1* within yeast can enable considerably reduced duration of MLF by LAB, no matter which *SSU1* allele is deleted (Fig. 4a and b). This result suggests that the *SSU1* gene may play a role in the phenotype investigated. However, the comparison of hemizygous hybrids fails to confirm the inhibitor role of the *SSU1* GN allele specifically, compared to SB. This negative result may be due to a masking effect of *SSU1* haploinsufficiency, suggesting that the negative impact of the GN allele may require two functional *SSU1* alleles to be fully expressed.

Intriguingly, the AF kinetics of the hemizygous strain SΔG092, which carries the *SSU1* GN allele, was slightly delayed compared to GΔS092 and SBxGN (Fig. 4c and d). In addition, when SΔG092 was co-inoculated with LAB, this delay was longer (up to 20 hours) suggesting a possible inhibition of yeast AF by bacteria. This deleterious effect of the GN allele on fermentation kinetics has been previously reported (Peltier et al. 2019) and was attributed to the increase of external sulfite in the fermented medium, a direct effect of increased SO₂ efflux. However, in our conditions, sulfite concentration was below the detection threshold, and no SO₂ was added to the grape juice.

The inability to confirm the QTL using the reciprocal hemizyosity assay could be due to *SSU1* not being the main gene involved in MLF completion by LAB or AF activity. In fact, other genes genetically linked to the GN allele of *SSU1* could be involved in those phenotypes. Among them, *ADH1* may be an interesting candidate. *ADH1* encodes an alcohol dehydrogenase that is located on chromosome XV, but is also physically associated with the *SSU1* gene in the XV-t-XV translocation genotype. In the first hours after yeast inoculation, expression of *ADH1* is not fully active. Ethanol production is delayed and replaced by glycerol-pyruvic fermentation in order to cope with the depletion of NAD⁺ in the cytosol (Ribereau-Gayon et al. 2006). We hypothesise that the XV-t-XVI translocation could also modify the expression of *ADH1*, at least in the early stage of yeast culture. Decreased *ADH1* expression in the early phase of AF could lower alcohol dehydrogenase activity, increasing the accumulation of acetaldehyde. This phenomenon has been previously observed for different wine starters including VL1, the parental strain of GN (Cheriat et al. 2010). The biological implication of acetaldehyde utilisation by wine LAB in free or bound form is not yet fully understood, apart from the liberation of free SO₂ from the acetaldehyde-SO₂ complex (Liu and Pilone 2000). High levels (> 100 mg L⁻¹) of acetaldehyde may inhibit growth of LAB (El-Gendy et al. 1983). Thus, the

impact of the XVI_374 QTL observed in the SBxGN progeny might be due to other genetic polymorphisms related to the XV-t-XVI translocation, such as the modification of the *ADH1* promoter region. In future, additional experimental efforts will be required to further investigate these molecular causes.

In conclusion, for the first time, yeast genetic background was assessed for its role in yeast-LAB compatibility during fermentation. The impact of *SSU1* haploinsufficiency on LAB ability to complete MLF was clear, but further work is needed to understand the role of the XV-t-XVI translocation on MLF outcomes. The inclusion of more progeny and use of a different genetic background may help identify other genes that result in phenotypic differences in yeast-LAB co-inoculations. Further work may also reveal the underlying cause of the major QTL identified in this study, which was not able to be confirmed by the *SSU1* hemizygote strains. However, the influence of *SSU1* in this work does add to the understanding of the pleiotropic role of *SSU1*, since it has been reported to impact yeast AF, growth and SO₂ production, and now also has the potential to impact co-inoculation outcomes with LAB. This work begins to unravel the complexity of *S. cerevisiae* genetic differences that can lead to a phenotype that impacts *O. oeni* during co-inoculation. Understanding the delicate interplay between genotype and phenotype can create opportunities for wine yeast manufacturers to develop yeast that work effectively with LAB, without negatively impacting yeast AF performance.

Declarations

Funding

This work was supported by Australia's grape growers and winemakers through their investment body, Wine Australia, with matching funds from the Australian Government. LB was supported by joint scholarships from The University of Adelaide and Wine Australia (AGW Ph 1510). JS was supported by Wine Australia project funding (UA1707). JS, KS and VJ are supported by The Australian Research Council Training Centre for Innovative Wine Production (www.ARCwinecentre.org.au; project number IC170100008), which is funded by the Australian Government with additional support from Wine Australia and industry partners. The University of Adelaide is a member of the Wine Innovation Cluster in Adelaide (<http://www.thewaite.org/waite-partners/wine-innovation-cluster/>). PM and EP are supported by Biolaftort (Laffort® Research & Development subsidiary) for this project, in addition PM received a grant from Aquitaine Region (Sesam Project) for genome sequencing and QTL analysis.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Availability of data

Raw data may be supplied upon request, at the discretion of the corresponding authors.

Authors' contributions

All authors contributed to the study conception and design. LB carried out laboratory fermentation experiments; PM and EP provided yeast strains. Data analysis and interpretation was performed by LB, EP and PM. LB wrote the manuscript and EP, JS, KS, JGM, VJ and PM reviewed and revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Hlne Mesnage for her assistance in performing the QTL fermentation experiment. We also acknowledge Nick van Holst Pellekaan for capturing images of the automated fermentation platform.

Conflicts of interest

EP and PM are employed by Biolaffect. This does not alter the authors' adherence to all the journal policies on sharing data and materials.

References

- Abrahamse CE, Bartowsky EJ (2012) Timing of malolactic fermentation inoculation in Shiraz grape must and wine: influence on chemical composition. *World J Microbiol Biotechnol* 28: 255–265. doi: 10.1007/s11274-011-0814-3
- Alexandre H, Costello PJ, Remize F, Guzzo J, Guilloux-Benatier M (2004) *Saccharomyces cerevisiae*–*Oenococcus oeni* interactions in wine: current knowledge and perspectives. *Int J Food Microbiol* 93:141–154. doi: 10.1016/j.ijfoodmicro.2003.10.013
- Antalick G, Perello MC, de Revel G (2013) Co-inoculation with yeast and LAB under winery conditions: modification of the aromatic profile of merlot wines. *S Afr J Enol Vitic* 34:223–232. doi: 10.21548/34-2-1098
- Arnink K, Henick-Kling T (2005) Influence of *Saccharomyces cerevisiae* and *Oenococcus oeni* strains on successful malolactic conversion in wine. *Am J Enol Vitic* 56:228-237.
- Arroyo-López FN, Bautista-Gallego J, Domínguez-Manzano V, Romero-Gil F, Rodríguez-Gómez P, García-García A, Garrido-Fernández R, Jiménez-Díaz (2012) Formation of lactic acid bacteria–yeasts communities on the olive surface during Spanish-style Manzanilla fermentations. *Food Microbiol* 32:295-301. doi: 10.1016/j.fm.2012.07.003
- Atanassova M, Choiset Y, Dalgalarondo M, Chobert JM, Dousset X, Ivanova I, Haertlé T (2003) Isolation and partial biochemical characterization of a proteinaceous anti-bacteria and anti-yeast compound produced by *Lactobacillus paracasei* subsp. *paracasei* strain M3. *Int J Food Microbiol* 87:63-73. doi: 10.1016/S0168-1605(03)00054-0
- Bartle L, Sumbly K, Sundstrom J, Jiranek V (2019) The microbial challenge of winemaking: yeast-bacteria compatibility. *FEMS Yeast Res* 19:foz040. doi: 10.1093/femsyr/foz040
- Bokulich NA, Bamforth CW (2013) The microbiology of malting and brewing. *Microbiol Mol Biol Rev* 77:157-172. doi: 10.1128/MMBR.00060-12
- Brice C, Sanchez I, Bigey F, Legras JL, Blondin B (2014) A genetic approach of wine yeast fermentation capacity in nitrogen-starvation reveals the key role of nitrogen signaling. *BMC Genomics* 15:495. doi: 10.1186/1471-2164-15-495
- Broman KW, Wu H, Sen S, Churchill GA (2003) R/qtl: QTL mapping in experimental crosses. *Bioinformatics* 19:889-890. doi: 10.1093/bioinformatics/btg112
- Cañas PMI, Pérez-Martín F, Romero EG, Prieto SS, Herreros MDLLP (2012) Influence of inoculation time of an autochthonous selected malolactic bacterium on volatile and sensory profile of Tempranillo and Merlot wines. *Int J Food Microbiol* 156:245-254. doi: 10.1016/j.ijfoodmicro.2012.03.033
- Cañas PMI, Romero EG, Pérez-Martín F, Seseña S, Palop ML (2015) Sequential inoculation versus co-inoculation in Cabernet Franc wine fermentation. *Food Sci Technol Int* 21:203–212. doi: 10.1177/1082013214524585
- Capucho I, San Romão MV (1994) Effect of ethanol and fatty acids on malolactic activity of *Leuconostoc oenos*. *Appl Microbiol Biotechnol* 42:391–395. doi: 10.1007/BF00902747
- Carrete R, Vidal MT, Bordons A, Constantí M (2002) Inhibitory effect of sulfur dioxide and other stress compounds in wine on the ATPase activity of *Oenococcus oeni*. *FEMS Microbiol Lett* 211:155-159. doi: 10.1111/j.1574-6968.2002.tb11218.x
- Chasseriaud L, Coulon J, Marullo P, Albertin W, Bely M (2018) New oenological practice to promote non-*Saccharomyces* species of interest: Saturating grape juice with carbon dioxide. *Appl Microbiol Biotechnol* 102:3779-3791. doi: 10.1007/s00253-018-8861-4

566 Chen X, Niks RE, Hedley PE, Morris J, Druka A, Marcel TC, Vels A, Waugh R (2010) Differential gene
567 expression in nearly isogenic lines with QTL for partial resistance to *Puccinia hordei* in barley. BMC
568 Genomics 11:629. doi: 10.1186/1471-2164-11-629

569 Churchill GA, Doerge RW (1994) Empirical threshold values for quantitative trait mapping. Genetics
570 138:963-971.

571 Comitini F, Ciani M (2007) The inhibitory activity of wine yeast starters on malolactic bacteria. Ann
572 Microbiol 57:61-66. Doi: 10.1007/BF03175051

573 Comitini F, Ferretti R, Clementi F, Mannazzu I, Ciani M (2005) Interactions between *Saccharomyces*
574 *cerevisiae* and malolactic bacteria: preliminary characterization of a yeast proteinaceous compound
575 (s) active against *Oenococcus oeni*. J Appl Microbiol 99:105-111. doi: 10.1111/j.1365-
576 2672.2005.02579.x

577 Decroocq V, Foulongne M, Lambert P, Gall OL, Martin C, Pascal T, Schurdi-Levraud V, Kervella J
578 (2005) Analogues of virus resistance genes map to QTLs for resistance to sharka disease in *Prunus*
579 *davidiana*. Mol Genet Genomics 272:680-689. doi: 10.1007/s00438-004-1099-0

580 Delcourt F, Taillandier P, Vidal F, Strehaiano P (1995) Influence of pH, malic acid and glucose
581 concentrations on malic acid consumption by *Saccharomyces cerevisiae*. Appl Microbiol Biotechnol
582 43:321-324. doi: 10.1007/BF00172832

583 Divol B, Du Toit M, Duckitt E (2012) Surviving in the presence of sulphur dioxide: Strategies
584 developed by wine yeasts. Appl Microbiol Biotechnol 95:601-613. doi: 10.1007/s00253-012-4186-x

585 Eder M, Sanchez I, Brice C, Camarasa C, Legras JL, Dequin S (2018) QTL mapping of volatile
586 compound production in *Saccharomyces cerevisiae* during alcoholic fermentation. BMC Genomics
587 19:166. doi: 10.1186/s12864-018-4562-8

588 El-Gendy SM, Abdel-Galil H, Shahin Y, Hegazi FZ (1983) Acetoin and diacetyl production by homo-
589 and heterofermentative lactic acid bacteria. J Food Protect 46:420-425. doi: 10.4315/0362-028X-
590 46.5.420

591 Eun MH, Han JH, Yoon JB, Lee J (2016) QTL mapping of resistance to the Cucumber mosaic virus
592 P1 strain in pepper using a genotyping-by-sequencing analysis. Hortic Environ Biotechnol 57:589-
593 597. doi: 10.1007/s13580-016-0128-3

594 Furukawa S, Nojima N, Yoshida K, Hirayama S, Ogihara H, Morinaga Y (2011) The importance of
595 inter-species cell-cell co-aggregation between *Lactobacillus plantarum* ML11-11 and *Saccharomyces*
596 *cerevisiae* BY4741 in mixed-species biofilm formation. Biosci Biotechnol Biochem 75:1430-1434. doi:
597 10.1271/bbb.100817

598 Furukawa S, Yoshida K, Ogihara H, Yamasaki M, Morinaga Y (2010) Mixed-species biofilm formation
599 by direct cell-cell contact between brewing yeasts and lactic acid bacteria. Biosci Biotechnol Biochem.
600 74: 2316-2319. doi: 10.1271/bbb.100350

601 Gao C, Fleet GH (1995) Degradation of malic and tartaric acids by high density cell suspensions of
602 wine yeasts. Food Microbiol 12:65-71. doi: 10.1016/S0740-0020(95)80080-8

603 García-Ríos E, Guillamón JM (2019) Sulfur dioxide resistance in *Saccharomyces cerevisiae*: beyond
604 *SSU1*. Microb Cell 6:527-530. doi: 10.15698/mic2019.12.699

605 García-Ríos E, Nuévalos M, Barrio E, Puig S, Guillamón JM (2019) A new chromosomal
606 rearrangement improves the adaptation of wine yeasts to sulfite. Environ Microbiol 21:1771–1781.
607 doi: 10.1111/1462-2920.14586

608 Goto-Yamamoto N, Kitano K, Shiki K, Yoshida Y, Suzuki T, Iwata T, Yamane Y, Hara S (1998) *SSU1*-
609 R, a sulfite resistance gene of wine yeast, is an allele of *SSU1* with a different upstream sequence. J
610 Ferment Bioeng 86:427-433. doi: 10.1016/S0922-338X(98)80146-3

611 Guzzo J, Jobin MP, Delmas F, Fortier LC, Garmyn D, Tourdot-Maréchal R, Lee B, Diviès C (2000)
612 Regulation of stress response in *Oenococcus oeni* as a function of environmental changes and
613 growth phase. Int J Food Microbiol 55:27-31. doi: 10.1016/S0168-1605(00)00209-9

614 Haley CS, Knott SA (1992) A simple regression method for mapping quantitative trait loci in line
615 crosses using flanking markers. Heredity 69:315-324. Doi: 10.1038/hdy.1992.131

616 Hranilovic A, Gambetta JM, Schmidtke L, Boss PK, Grbin PR, Masneuf-Pomarede I, Bely M, Albertin
617 W, Jiranek V (2018) Oenological traits of *Lachancea thermotolerans* show signs of domestication and
618 allopatric differentiation. Sci Rep 8:1-13. doi: 10.1038/s41598-018-33105-7

619 Huang C, Roncoroni M, Gardner RC (2014) *MET2* affects production of hydrogen sulfide during wine
620 fermentation. Appl Microbiol Biotechnol 98:7125-7135. doi: 10.1007/s00253-014-5789-1

621 Jara M, Cubillos FA, García V, Salinas F, Aguilera O, Liti G, Martínez C (2014) Mapping genetic
622 variants underlying differences in the central nitrogen metabolism in fermenter yeasts. PLoS One
623 9:e86533. doi: 10.1371/journal.pone.0086533

624 Jarosz DF, Brown JC, Walker GA, Datta MS, Ung WL, Lancaster AK, Rotem A, Chang A, Newby GA,
625 Weitz DA, Bisson LF (2014) Cross-kingdom chemical communication drives a heritable, mutually
626 beneficial prion-based transformation of metabolism. Cell 158:1083-1093. doi:
627 10.1016/j.cell.2014.07.025

628 Knoll C, Fritsch S, Schnell S, Grossmann M, Krieger-Weber S, Du Toit M, Rauhut D (2012) Impact of
629 different malolactic fermentation inoculation scenarios on Riesling wine aroma. World J Microbiol
630 Biotechnol 28:1143-1153. doi: 10.1007/s11274-011-0917-x

631 Liti G, Louis EJ (2012) Advances in quantitative trait analysis in yeast. PLoS Genet 8:e1002912. doi:
632 10.1371/journal.pgen.1002912

633 Liu SQ, Pilone GJ (2000) An overview of formation and roles of acetaldehyde in winemaking with
634 emphasis on microbiological implications. Int J Food Sci Technol 35: 49-61. doi:10.1046/j.1365-
635 2621.2000.00341.x

636 Martí-Raga M, Peltier E, Mas A, Beltran G, Marullo P (2017) Genetic causes of phenotypic adaptation
637 to the second fermentation of sparkling wines in *Saccharomyces cerevisiae*. G3 Genes Genom Genet
638 7:399-412. doi: 10.1534/g3.116.037283

639 Marullo P, Aigle M, Bely M, Masneuf-Pomarede I, Durrens P, Dubourdieu D, Yvert G (2007b) Single
640 QTL mapping and nucleotide-level resolution of a physiologic trait in wine *Saccharomyces cerevisiae*
641 strains. FEMS Yeast Res 7:941-952. doi: 10.1111/j.1567-1364.2007.00252.x

642 Marullo P, Claisse O, Eder ML, Börlin M, Feghali N, Bernard M, Legras JL, Albertin W, Rosa AL,
643 Masneuf-Pomarede I (2020) *SSU1* checkup, a rapid tool for detecting chromosomal rearrangements
644 related to the *SSU1* promoter in *Saccharomyces cerevisiae*: an ecological and technological study on
645 wine yeast. Front Microbiol 11:1331. doi: 10.3389/fmicb.2020.01331

646 Marullo P, Durrens P, Peltier E, Bernard M, Mansour C, Dubourdieu D (2019) Natural allelic variations
647 of *Saccharomyces cerevisiae* impact stuck fermentation due to the combined effect of ethanol and
648 temperature; a QTL-mapping study. BMC Genomics 20:680. doi: 10.1186/s12864-019-5959-8

649 Marullo P, Yvert G, Bely M, Aigle M, Dubourdieu D (2007a) Efficient use of DNA molecular markers to
650 construct industrial yeast strains. FEMS Yeast Res 7:1295-1306. doi: 10.1111/j.1567-
651 1364.2007.00281.x

652 Mendoza LM, de Nadra MCM, Fariás ME (2010) Antagonistic interaction between yeasts and lactic
653 acid bacteria of oenological relevance: Partial characterization of inhibitory compounds produced by
654 yeasts. *Food Res Int* 43:1990-1998. doi: 10.1016/j.foodres.2010.05.017

655 Mounier J, Monnet C, Vallaes T, Arditi R, Sarthou AS, Hélias A, Irlinger F (2008) Microbial
656 interactions within a cheese microbial community. *Appl Environ Microbiol* 74:172-181. doi:
657 10.1128/AEM.01338-07

658 Muñoz V, Beccaria B, Abreo E (2014) Simultaneous and successive inoculations of yeasts and lactic
659 acid bacteria on the fermentation of an unsulfited Tannat grape must. *Braz J Microbiol* 45:59-66. doi:
660 10.1590/S1517-83822014000100009

661 Nehme N, Mathieu F, Taillandier P (2008) Quantitative study of interactions between *Saccharomyces*
662 *cerevisiae* and *Oenococcus oeni* strains. *J Ind Microbiol Biotechnol* 35:685-693. doi: 10.1007/s10295-
663 008-0328-7

664 Nehme N, Mathieu F, Taillandier P (2010) Impact of the co-culture of *Saccharomyces cerevisiae*–
665 *Oenococcus oeni* on malolactic fermentation and partial characterization of a yeast-derived inhibitory
666 peptidic fraction. *Food Microbiol* 27:150-157. doi: 10.1016/j.fm.2009.09.008

667 Nissen P, Nielsen D, Arneborg N (2003) Viable *Saccharomyces cerevisiae* cells at high
668 concentrations cause early growth arrest of non-*Saccharomyces* yeasts in mixed cultures by a cell–
669 cell contact-mediated mechanism. *Yeast* 20:331-341. doi: 10.1002/yea.965

670 Nissen P, Nielsen D, Arneborg N (2004) The relative glucose uptake abilities of non-*Saccharomyces*
671 yeasts play a role in their coexistence with *Saccharomyces cerevisiae* in mixed cultures. *Appl*
672 *Microbiol Biotechnol* 64:543-550. doi: 10.1007/s00253-003-1487-0

673 Osborne JP, Edwards CG (2006) Inhibition of malolactic fermentation by *Saccharomyces* during
674 alcoholic fermentation under low- and high-nitrogen conditions: a study in synthetic media. *Aust J*
675 *Grape Wine Res* 12:69-78. doi: 10.1111/j.1755-0238.2006.tb00045.x

676 Park H, Bakalinsky AT (2000) *SSU1* mediates sulphite efflux in *Saccharomyces cerevisiae*. *Yeast*
677 16:881-888. doi: 10.1002/1097-0061(200007)16:10<881::AID-YEA576>3.0.CO;2-3

678 Peltier E, Bernard M, Trujillo M, Prodhomme D, Barbe JC, Gibon Y, Marullo P (2018a) Wine yeast
679 phenomics: A standardized fermentation method for assessing quantitative traits of *Saccharomyces*
680 *cerevisiae* strains in enological conditions. *PLoS One* 13:e0190094. doi:
681 10.1371/journal.pone.0190094

682 Peltier E, Friedrich A, Schacherer J, Marullo P (2019) Quantitative trait nucleotides impacting the
683 technological performances of industrial *Saccharomyces cerevisiae* strains. *Front Genetics* 10:683.
684 doi: 10.3389/fgene.2019.00683

685 Peltier E, Sharma V, Raga MM, Roncoroni M, Bernard M, Jiranek V, Gibon Y, Marullo P (2018b)
686 Dissection of the molecular bases of genotype x environment interactions: a study of phenotypic
687 plasticity of *Saccharomyces cerevisiae* in grape juices. *BMC Genomics* 19:772. doi: 10.1186/s12864-
688 018-5145-4

689 Pérez-Ortín JE, Querol A, Puig S, Barrio E (2002) Molecular characterization of a chromosomal
690 rearrangement involved in the adaptive evolution of yeast strains. *Genome Res* 12:1533-1539. doi:
691 10.1101/gr.436602

692 R Core Team (2017) R: A language and environment for statistical computing. R Foundation for
693 Statistical Computing. <https://www.R-project.org/>. Accessed 4 September 2017

694 Rankine BC, Pocock KF (1970) Alkalimetric determination of sulphur dioxide in wine. *Aust Wine Brew*
695 *Spirit Rev* 88:40-44.

Renault P, Coulon J, de Revel G, Barbe JC, Bely M (2015) Increase of fruity aroma during mixed *T. delbrueckii*/*S. cerevisiae* wine fermentation is linked to specific esters enhancement. *Int J Food Microbiol* 207:40-48. doi: 10.1016/j.ijfoodmicro.2015.04.037

Renault P, Coulon J, Moine V, Thibon C, Bely M (2016) Enhanced 3-sulfanylhexas-1-ol production in sequential mixed fermentation with *Torulaspora delbrueckii*/*Saccharomyces cerevisiae* reveals a situation of synergistic interaction between two industrial strains. *Front Microbiol* 7:293. doi: 10.3389/fmicb.2016.00293

Renault P, Miot-Sertier C, Marullo P, Hernández-Orte P, Lagarrigue L, Lonvaud-Funel A, Bely M (2009) Genetic characterization and phenotypic variability in *Torulaspora delbrueckii* species: potential applications in the wine industry. *Int J Food Microbiol* 134:201-210. doi: 10.1016/j.ijfoodmicro.2009.06.008

Renault PE, Albertin W, Bely M (2013) An innovative tool reveals interaction mechanisms among yeast populations under oenological conditions. *Appl Microbiol Biotechnol* 97:4105-4119. doi: 10.1007/s00253-012-4660-5

Renouf V, Claisse O, Lonvaud-Funel A (2007) Inventory and monitoring of wine microbial consortia. *Appl Microbiol Biotechnol* 75:149-164. doi: 10.1007/s00253-006-0798-3

Ribéreau-Gayon P, Dubourdieu D, Donèche B, Lonvaud A (2006) Handbook of enology, Vol. 1: The microbiology of wine and vinifications, 2nd edn. Wiley, doi:10.1002/0470010363

Roncoroni M (2014) Quantitative trait loci mapping in winemaking yeast. Dissertation, University of Auckland

Roncoroni M, Santiago M, Hooks DO, Moroney S, Harsch MJ, Lee SA, Richards KD, Nicolau L, Gardner RC (2011) The yeast *IRC7* gene encodes a β -lyase responsible for production of the varietal thiol 4-mercapto-4-methylpentan-2-one in wine. *Food Microbiol* 28:926-935. doi: 10.1016/j.fm.2011.01.002

Rossouw D, Du Toit M, Bauer FF (2012) The impact of co-inoculation with *Oenococcus oeni* on the transcriptome of *Saccharomyces cerevisiae* and on the flavour-active metabolite profiles during fermentation in synthetic must. *Food Microbiol* 29:121–131. doi: 10.1016/j.fm.2011.09.006

Sadoudi M, Tourdot-Maréchal R, Rousseaux S, Steyer D, Gallardo-Chacón JJ, Ballester J, Vichi S, Guérin-Schneider R, Caixach J, Alexandre H (2012) Yeast–yeast interactions revealed by aromatic profile analysis of Sauvignon Blanc wine fermented by single or co-culture of non-*Saccharomyces* and *Saccharomyces* yeasts. *Food Microbiol* 32:243-253. doi: 10.1016/j.fm.2012.06.006

Salinas F, Cubillos FA, Soto D, Garcia V, Bergström A, Warringer J, Ganga MA, Louis EJ, Liti G, Martinez C (2012) The genetic basis of natural variation in oenological traits in *Saccharomyces cerevisiae*. *PloS One* 7:e49640. doi: 10.1371/journal.pone.0049640

Steinmetz LM, Scharfe C, Deutschbauer AM, Mokranjac D, Herman ZS, Jones T, Chu AM, Giaever G, Prokisch H, Oefner PJ, Davis RW (2002b) Systematic screen for human disease genes in yeast. *Nat Genet* 31:400-404. doi: 10.1038/ng929

Steinmetz LM, Sinha H, Richards DR, Spiegelman JL, Oefner PJ, McCusker JH, Davis RW (2002a) Dissecting the architecture of a quantitative trait locus in yeast. *Nature* 416:326-330. doi: 10.1038/416326a

Steyer D, Ambroset C, Brion C, Claudel P, Delobel P, Sanchez I, Erny C, Blondin B, Karst F, Legras JL (2012) QTL mapping of the production of wine aroma compounds by yeast. *BMC Genomics* 13:573. doi: 10.1186/1471-2164-13-573

739 Tempère S, Marchal A, Barbe JC, Bely M, Masneuf-Pomarede I, Marullo P, Albertin W (2018) The
740 complexity of wine: clarifying the role of microorganisms. *Appl Microbiol Biotechnol* 102:3995-4007.
741 doi: 10.1007/s00253-018-8914-8

742 Treu L, Toniolo C, Nadai C, Sardu A, Giacomini A, Corich V, Campanaro S (2014) The impact of
743 genomic variability on gene expression in environmental *Saccharomyces cerevisiae* strains. *Environ*
744 *Microbiol* 16:1378-1397. doi: 10.1111/1462-2920.12327

745 Tristezza M, di Feo L, Tufariello M, Grieco F, Capozzi V, Spano G, Mita G (2016) Simultaneous
746 inoculation of yeasts and lactic acid bacteria: Effects on fermentation dynamics and chemical
747 composition of Negroamaro wine. *LWT-Food Sci Technol* 66:406-412. doi: 10.1016/j.lwt.2015.10.064

748 Walker ME, Nguyen TD, Liccioli T, Schmid F, Kalatzis N, Sundstrom JF, Gardner JM Jiranek V (2014)
749 Genome-wide identification of the Fermentome; genes required for successful and timely completion
750 of wine-like fermentation by *Saccharomyces cerevisiae*. *BMC Genomics* 15:552. doi: 10.1186/1471-
751 2164-15-552

752 Weng J, Gu S, Wan X, Gao H, Guo T, Su N, Lei C, Zhang X, Cheng Z, Guo X, Wang J (2008)
753 Isolation and initial characterization of GW5, a major QTL associated with rice grain width and weight.
754 *Cell Res* 18:1199-1209. doi: 10.1038/cr.2008.307

755 Yéramian N, Chaya C, Suárez Lepe JA (2007) L-(-)-malic acid production by *Saccharomyces spp.*
756 during the alcoholic fermentation of wine. *J Agric Food Chem* 55:912-919. doi: 10.1021/jf061990w

757 Zimmer A, Durand C, Loira N, Durrens P, Sherman DJ, Marullo P (2014) QTL dissection of lag phase
758 in wine fermentation reveals a new translocation responsible for *Saccharomyces cerevisiae*
759 adaptation to sulfite. *PLoS One* 9:e86298. doi: 10.1371/journal.pone.0086298

760

761 **Table 1:** Yeast strains used in this study

Strain	Comment	Genotype	Origin	Culture collection: WDCM-791 (CRB ISVV, Bordeaux)
SB	Monosporic clone of Actiflore® BO213 (Laffort®, Bordeaux, France)	<i>HO/HO</i> , diploid	Peltier et al. (2018b)	CRBO L2001
GN	Monosporic clone of Zymaflore® VL1 (Laffort®, Bordeaux, France)	<i>HO/HO</i> , diploid	Peltier et al. (2018b)	CRBO L2002
SBxGN	F1 hybrid SBxGN	<i>HO/HO</i> , diploid	Peltier et al. (2018b)	CRBO L2003
BN	F1 hybrid hoSBxGN	<i>HO/ho::kanMx4</i> , diploid	Marullo et al. (2007b)	CRBO L2004
pop BN	67 progeny clones of BN (hoSBxGN). Labelled with prefix “CM” followed by an ID number	<i>ho::kanMx4</i> , haploids	Marullo et al. (2007b)	
SΔG092	Hemizygous hybrid isogenic to SBxGN	<i>ho/ho</i> , <i>YPL092^{SB}::kanMX4/ YPL092^{GN}</i> , diploid	Zimmer et al. (2014)	CRBO L2006
GΔS092	Hemizygous hybrid isogenic to SBxGN	<i>ho/ho</i> , <i>YPL092^{GN}::kanMX4/ YPL092^{SB}</i> , diploid	Zimmer et al. (2014)	CRBO L2007

762

763 **Table 2:** AF and MLF measures used to perform QTL mapping, BN progeny evaluation or statistical
 764 analysis for comparison of hemizygous strains with their corresponding SBxGN hybrids.
 765 Abbreviations, if assigned, are shown below:

AF measures	Abbreviation	MLF measures	Abbreviation
Time to complete AF (hours)	<i>tend-AF</i>	Residual L-malic acid concentration for yeast alone fermentations (g L ⁻¹)	Not assigned
Time to reach equivalent of 35% CO ₂ (175.53 g L ⁻¹ total sugar)	t35-AF	Percentage L-malic acid consumption for yeast-LAB co-inoculation fermentations	<i>Pct_malic_co</i>
Time to reach equivalent of 50% CO ₂ (143.62 g L ⁻¹ total sugar)	t50-AF	Estimated overall L-malic acid consumed by LAB (g L ⁻¹)	<i>Malic_acid_LAB_consumed</i>
Time to reach equivalent of 80% CO ₂ (79.79 g L ⁻¹ total sugar)	t80-AF	Time to complete MLF (hours)	<i>tend-MLF</i>
Slope between t50-AF and t80-AF	s50-80-AF		
Percentage L-malic acid consumption or production by yeast alone	<i>Pct_malic_AF</i>		

766