Decreasing arousal modulates perceptual decision-making

Sridhar R. Jagannathan^{1*}, Corinne A. Bareham^{1,2,3}, Tristan A. Bekinschtein^{1*}

¹Department of Psychology, University of Cambridge, Cambridge, United Kingdom

²Department of Clinical Neurosciences, University of Cambridge, Cambridge, United Kingdom

³School of Psychology, Victoria University of Wellington, Wellington, New Zealand

*Corresponding authors: srj34@cam.ac.uk, tb419@cam.ac.uk

ABSTRACT

The ability to make decisions based on external information, prior knowledge, evidence and context, is a crucial aspect of cognition and may determine the success and survival of an organism. Despite extensive work on decision making mechanisms and models, understanding the effects of arousal on its neural and cognitive processes remain limited. Here we characterise behavioural and neural dynamics of perceptual decision making in awake and low alertness periods in humans, and characterise the compensatory mechanisms as arousal decreases. Well-rested human participants, fluctuating between full wakefulness and low alertness, performed an auditory tone-localisation task and its behavioural dynamics was quantified with psychophysics, signal detection theory and drift-diffusion modelling, revealing slower reaction times, inattention to the left side of space, and a lower rate of evidence accumulation in periods of low alertness. Unconstrained multivariate pattern analysis (decoding) showed a ~260ms delayed onset driven by low alertness of the neural signatures differentiating between left and right decision, with a spatial reconfiguration from centro-parietal to lateral frontal regions 160-300ms. To understand the neural compensatory mechanisms with decreasing arousal, we connected the evidence-accumulation behavioural parameter to the neural activity, showing in the early period (150-300ms) a shift in the associated patterns from bilateral inferior frontal gyri in awake to right dorsolateral in low alertness. The second period of evidence accumulation shows posterior involvement in awake and, due to the delayed processing during drowsiness, a wide frontopolar, dorsolateral prefrontal cortices and right parietal during decreased arousal. The change in the neurobehavioural dynamics for both perceptual encoding and central accumulation cognitive processes define a clear reconfiguration of the brain networks' regions and dynamics needed for the implementation of decision making in both early perceptual and later central processes, revealing compensatory mechanisms of resilience of cognition when challenged by decreased arousal.

INTRODUCTION

The question of how decisions are made has shaped the world's systems of government, justice and social order (1). From a cognitive neuroscience perspective, the modelling of psychological and neural factors allows for the understanding of decision making via its underlying brain mechanisms, depending both from the specific constraints of the information provided from the external world and the internal state of the corresponding system. Studies on how the brain implements simple decisions have revealed several neurocognitive processes at the perceptual, central integration and motor implementation levels (2–4), but the modulatory effects of the internal milieu, homeostasis, arousal, alertness, and circadian influences on such processes have received less attention (5,6). Specifically, low alertness fluctuations on decision processes have been tackled by sleep deprivation and brain injury studies but hardly by normal variations in the stability of wakefulness (7). How does the brain compensate for the detrimental effects of decreasing arousal while making an optimal decision?

Perceptual decision making is successfully studied under the sequential sampling models (8,9), and referred to as a set of processes involving the processing of sensory information to make a behavioural choice from a set of available options (10). The various elements of this process (Figure s1) are: a) Perceptual stage: a sensory system that transforms physical stimulus intensities for the perceptual systems to deliver decision information; b) Central integration stage: a decision system that integrates and accumulates such decision-information variables and makes an optimal choice based on relative evidence; and c) Motor stage: a motor system that implements the appropriate motor plan and subsequent action. According to sequential sampling models, accurate perceptual decisions separate the noise from the signal by repeatedly sampling and integrating evidence until there is enough in favour of one of the decision choices. The preferred concept to understand and develop hypotheses is called decision variable, and it is an accumulation of priors, evidence, internal milieu and value into a quantity that is interpreted by the decision rule to produce a choice (9). This study aims to investigate how alertness modulates the decision variable in spatial auditory perception.

The neural underpinnings of perceptual decisions have been explored both in model-free and model-constrained manners in neuroimaging studies in rodents, monkeys and human observers. Signals from domain general regions such as the right insula, posterior left SFS/DLPFC, left TPJ, posterior parietal cortex (LIP) were related to processes of perceptual and central evidence accumulation (11–16). The above studies provide a general overview on the different regions of the brain involved in processes related to decision-making. To explore the question of how alertness modulates these decision-making related processes, we need to first define the specific aspect of wakefulness to be used as the experimental manipulation (17,18). Alertness related modulations can

76 be classed into ‘tonic’ fluctuations that span multiple trials and time-periods, and ‘phasic’ moment
 77 to moment modulations in arousal produced in response to an ongoing task. A few recent studies
 78 have shown, in humans and mice, that arousal, measured by brain-stem systems (pupil response),
 79 modulates individual decision making in moment to moment fluctuations of phasic arousal (19,20).
 80 Further to this, van Kempen and collaborators (21) showed that lower tonic and higher phasic
 81 arousal, as defined by the pupil measurements, predicted shorter reaction times and was associated
 82 with a centroparietal positivity in EEG space. However, these studies have used pupil responses;
 83 which is a physiological marker indirectly linked to neural arousal systems, and further, can also be
 84 influenced by other factors like visual contrast, spatial frequencies of the stimulus etc. (22). In
 85 addition to this, the above mentioned studies have not measured influences of tonic fluctuations in
 86 decreased arousal, which is difficult for pupillometry or eye tracker measures. Hence, we require a
 87 study that can directly investigate the effect of intrinsic low alertness (tonic) fluctuations on
 88 decision-making.

89
 90
 91 Before undertaking such an investigation, we need to resolve the problem of measuring arousal
 92 relatively independently and directly from the brain. For this, we use subclassification of stages of
 93 alertness pioneered by Hori and collaborators (23). This manual method of scoring is based on
 94 spectral power as a continuous measure and visually defined grapho-elements in the transition to
 95 sleep. We took a computational approach to this problem, validated and automated this procedure
 96 to avoid human error, and further cross-validated it to show robust outcomes (24) across different
 97 EEG datasets (64, 128 channel) with participants performing a variety of tasks (semantic
 98 decision-making, auditory masking). Recently this method was successfully applied to explore the
 99 effect of alertness on cognitive control (18).

100
 101 This sets up the stage for the primary questions that will be analysed in this study: a) In terms of
 102 behaviour, how does a perceptual decision making task get affected by changing levels of tonic
 103 alertness. b) How are such modulations in behaviour connected to different elements of the decision
 104 making process i.e. perceptual, central, motor stage (Figure s1) and how are these stages modulated
 105 by alertness. c) How are the specific temporal periods and spatial regions implementing the
 106 different elements of the decision making process modulated by alertness. d) Finally, how does the
 107 brain, when compromised by lower levels of alertness, adapt/reconfigure to perform the decision.

108
 109 To answer the above questions, we decided to use an experiment (auditory spatial localisation task)
 110 with a known behavioural asymmetry (25) and understand how neurobehavioural decision-making
 111 mechanisms are modulated by fluctuations in tonic alertness. First, we use a combination of
 112 multi-level modelling, psychophysics and signal detection theory to show how behaviour is
 113 modulated by variations in decreasing alertness. Second, we employ a hierarchical drift-diffusion
 114 model to tease apart and parameterise the different elements of the decision-making process and
 115 examine individually how alertness modulated them. Third, we use decoding at the temporo-spatial
 116 levels to understand how different elements of the decision-making processes (perceptual, evidence

accumulation, motor implementation) may be implemented in the brain, and how such processes are modulated by fluctuations in alertness. Fourth, we connect the distinct behavioural parameters (evidence accumulation rate) of the decision making process to the neural processes, thereby capturing how the brain reconfigures, spatial and temporally, to maintain behavioural performance when challenged by decreased alertness.

RESULTS

Behavioural evidence of decision making modulated by alertness

We organized the results from direct and model-free to theoretically constrained. Error proportion, reaction times and subjective midline crosses are described as direct measures to evaluate the effects of alertness, while signal detection and drift-diffusion models theoretically constrain the interpretation of its parameters to perceptual, central and motor processes sequentially occurring during spatial decision making. We further organized the brain analyses similarly, using multivariate decoding to widely characterise the spatiotemporal neural signatures of the decision and, constrained by the behavioural results, we map the neural dynamics of evidence accumulation in full wakefulness and low alertness. This approach allowed for both, neurobehavioural characterization and hypotheses-driven evaluation, of the effects of low arousal on perceptual decision-making.

Error-proportion modulated by alertness

We first used multilevel modelling to understand how the errors made by each participant in an auditory tone-localization task (Figure 1A) was influenced by the stimulus presented ('left' or 'right' auditory tone) and the state of the participant ('alert' or 'drowsy'). We defined 4 different multilevel (linear mixed) models where errors were modulated by various combinations of stimulus and alertness states (see methods section). The analysis of variance table of the winning model shows that both alertness ($F(1, 95.07) = 14.04, p < 0.0005$) and stimulus type ($F(1, 95.07) = 18.80, p < 0.00005$) have an effect on error-proportion. Further, there was a reliable interaction between alertness levels and stimulus type ($F(1, 95.07) = 6.88, p < 0.05$). Next, post-hoc analysis revealed a reliable difference between alert and drowsy conditions for left stimuli ($p < 0.00005$), and not for right stimuli ($p = 0.859$). These behavioural results (Figure 1B) replicate the findings of (25) in an independent study, task design, alertness measure (and lab), reporting an increase in location assignment errors on tones from the left side of the midline when people became drowsy.

Figure 1: Auditory spatial attention task: A) participants had to localize the direction of auditory tones coming from left and right side of the midline. B) proportion of errors committed in alert and drowsy periods across left and right stimuli. Multilevel modelling reveals that error rates depend on stimulus type (left,right) and state of participant (alert,drowsy). Post-hoc tests indicate that error proportion is reliably modulated by alertness but only for left stimuli. *** indicates $p < 0.00005$, ns indicates not significant (or not reliable), error bars indicate standard error of the mean. C) mean reaction times for individual participants. Reaction times are variable and slower under drowsy conditions. D) spatial bias value per participant (using psychometric fits), bias level shifts towards the left side (indicating more left errors) for most participants. Negative bias

values indicate shifts in the subjective midline towards the left and positive values indicate shift towards the right. E) group level psychometric fits indicate the shift in subjective midline (dotted lines), shaded regions are confidence interval bounds. F) signal detection analysis shows that only d' (sensitivity) seems modulated by alertness while criterion remains unaffected.

Reaction times are modulated by alertness

Second, we aimed to quantify the modulation in the response profiles (reaction times) of individual participants by alertness levels. A paired samples t-test with $t(31) = -7.78$, $p < 0.05$ revealed a reliable effect of alertness on reaction times. These behavioural results (Figure 1C) converge with the original findings from Hori (23) indicating slower reaction times under lower levels of alertness, and in agreement to all our previous work (7,18,25–27).

Subjective midline modulated by alertness

Third, we used psychophysics to quantify the modulation of subjective midline per participant by alertness levels. For this purpose we fit a cumulative normal function (see methods section) to the proportion of rightward responses (per participant) under each stimulus condition from -60° to 60° from the midline under both, alert and drowsy periods. The fit was weighted by the number of trials (responses) in each stimulus condition. The mean of the function referred to as 'bias' is the subjective midline or spatial bias (where participants have 0.5 chance of pressing left or right responses). Most participants (Figure 1D) had their bias point shifted to the left (as they became less alert: drowsy), indicating more left errors (as they overestimate the right side of space). A small proportion of participants had bias points shifted to the right. Overall, a paired samples t-test $t(20) = 3.33$, $p < 0.005$, revealed a reliable difference in bias points between alert and drowsy periods. Further, we also plotted (Figure 1E) the mean of the psychometric fits of individual participants to show that the overall subjective midline has shifted to the left at the group level.

Signal detection parameters modulated by alertness

Fourth, we used signal detection theory to understand the factors that modulate decision making under varying levels of alertness (Figure 1F). d' (sensitivity) was modulated by alertness with $t(31) = 6.23$, $p < 0.05$ however criterion (response bias) did not show reliable differences, $t(31) = -1.87$, $p > 0.05$. This suggests that internal representations in the brain in terms of sensory/perceptual and noise distributions are modulated by alertness levels. Further, the lack of reliable differences in the response bias also suggests that participants were not arbitrarily pressing right responses for uncertain stimuli.

To summarise, the behavioural results hint that the first two stages of perceptual, evidence accumulation process are affected by decreasing arousal and that the final stage of motor implementation (response bias/criterion) may not be affected.

Decreased alertness modulates sequential sampling model parameters of evidence accumulation of decision making

203

204 Next, we aimed to quantify the different elements of the decision-making process using
205 drift-diffusion modelling. The drift-diffusion model captures the optimal procedure involved in
206 performing a 2-alternative forced choice (2AFC) task under sequential sampling framework. It
207 assumes that the observer accumulates evidence for one or other alternative in every other time
208 step (28), until integrated evidence reaches a threshold to decision (Figure 2A). The localization of
209 tones to the left and right side of space is essentially a 2-choice task with the participant always
210 forced to make a decision on the direction of tone. The model was implemented with a hierarchical
211 Bayesian procedure using hierarchical drift diffusion model (HDDM) (See methods section). For the
212 HDDMs, we fit the response of each participant instead of accuracy. This procedure is referred to as
213 Stimulus-coding, allows for the testing of several decision making parameters and is critical to
214 uncover response bias (20).

215

216 We examined 8 different variants of the model (see methods section), allowing the parameters
217 drift-rate (v) or bias-point (z) to vary based on state (alert or drowsy) or stimulus (left or right)
218 (Figure 2B,C). The winning model was chosen as the one with lowest deviance information criterion
219 (DIC) providing a balance between model fit and complexity. The winning model (#8) was composed
220 of drift-rate (v) varied according to state (alert or drowsy) and stimulus (left or right), whilst
221 bias-point (z) varied according to stimulus (left or right) only, furthering the claim that the changes
222 in alertness place a higher burden in the evidence accumulation process (drift rate), and less on the
223 motor implementation of the task. The winning model was then analysed for differences in
224 posterior densities of parameters. For this purpose, we used a bayesian estimate which is
225 fundamentally more informative and avoids the arbitrary choices like significance level and specific
226 statistical tests used by the frequentist based methods (29).

227

228 We found in the winning model, the proportion of posterior overlap in the bias point (between left
229 and right stimuli) was 24.2% (Figure 2E). This indicates that the bias-point was not reliably different
230 between the different stimuli. Further, the proportion of posterior overlap (between left and right
231 stimuli) in the drift rate for alert trials was 13.8% this reduces to 3.0% for drowsy trials (Figure 2H).
232 This indicates that the drift-rate (evidence accumulation rate) was reliably different between left
233 and right stimuli on drowsy trials (3.0%) in comparison to the alert trials (13.8%).

234

235 To summarise, the behaviour modeling results hint that the first two stages of perceptual encoding
236 and central evidence accumulation process (indicated by drift rate) are affected by decreasing
237 arousal.

238

Figure 2: Evidence accumulation models: A) Drift-diffusion model accounts for the reaction time distributions of responses across left and right stimuli ('Stimulus Coding'). 'v(drift)' indicates evidence accumulation rate, 'a' indicates the boundary separation across left and right responses, 'z' indicates the bias point, usually $z = 0.5$ for unbiased responses. B,C) changes in response distributions explained by shift in 'z', shift in 'v' respectively from alert to drowsy trials. grey lines indicate unbiased condition. D) posterior densities of 'z' for left and right stimuli in the winning model #8. a.u. indicates arbitrary units. E) differences

in the posterior densities indicate no evidence of bias point difference across stimuli. F,G) posterior densities of 'v' for left and right stimuli across alert and drowsy periods in the winning model #8. H) differences in the posterior densities of changes in drift rate indicate strong evidence in favour of change in drift rate across stimuli in the drowsy period compared to alert periods.

Lower alertness delays the processing of spatial decision making

Next, we aimed to uncover how the neural patterns in the EEG data are modulated by alertness levels. Conventional event related potentials (ERP) analysis for this purpose, relies on a-priori identified spatial locations or temporal segments to measure the differences across conditions. For data driven characterisation of the neural echoes of our experiment we decided to use time resolved decoding techniques that do not rely on such a-priori definitions and, further these techniques due to its multivariate nature, perform much better in detecting differences across experimental conditions (30).

Spatial and temporal signatures involved in spatial localization across alert and drowsy periods

First, we were interested in identifying the neural signatures involved in the performance of this task during alert condition. For this purpose, decoding involved in identifying the stimulus (X - left or right tone) presented from the EEG data (Y). This process involves the identification of the W (classifier weights) that can produce the transformation, $Y_t = W_t X_t$ where 't' represents time (Figure 3A,B). The performance of the classifier (W) is evaluated by training and testing the data at each time-point (t) using area under the curve (AUC) as measure. In Figure 3C, the shaded region represents those periods reliably decoded ($p < 0.05$) (see methods for more details).

We found that when participants were alert the decoding of stimuli (Figure 3D) started at 160 ms after the stimulus and reliably lasted until 730 ms (cluster permutation, $p < 0.05$) with mean AUC of 0.57. The peak discriminatory power was at 280 ms (AUC = 0.61). The average AUC between 200-300 ms was 0.58 ± 0.007 , $p < 0.05$, between 300-400 ms was 0.56 ± 0.002 , $p < 0.05$. However when the participants became 'drowsy' the decoding of stimuli (Figure 3E) shifted to 420 ms after the stimulus was presented and lasted until 730 ms (reliable with cluster permutation, $p < 0.05$) with mean AUC of 0.54. The peak discriminatory power was at 590 ms (AUC = 0.56). The average AUC between 420-500 ms was $M = 0.53 \pm 0.002$, $p < 0.05$, between 510-590 ms was $M = 0.54 \pm 0.004$, $p < 0.05$. This suggest that the processes related to the discrimination between left and right stimuli under drowsy conditions may cease to be directly informative to the decision in the early processes of sensory encoding and would only start later, around ~400ms, to show neural differentiation. Further to this, the lower discriminatory power points to a potentially less efficient (lower decodability, longer time duration) process of central evidence accumulation, or to a different neural implementation of the processes during low alertness.

286

Figure 3: Temporal decoding (stimuli): A) decoding consists of identifying Y (responses or stimuli) from X (EEG data). The model thus consists of $Y_t = W_t X_t$, where 't' represents time and W_t represents transformation (classifier weights). B) classifier weights W_t are determined by the optimal separation of different classes (here left, right stimuli). C) decoding performance is assessed using the area under curve (AUC) where shaded regions represent reliably different time periods. D,E) AUC under alert and drowsy periods, where the classifier was trained to discriminate between targets of left and right stimuli. Shaded regions show time periods with reliable ($p < 0.05$) discriminatory power. The mean reaction times (RT) of participants are plotted below. F) comparison of coefficients of classifier patterns in the early time periods. The boxed region highlights topographical differences in the frontal, posterior parietal and central electrodes. G) comparison of coefficients of classifier patterns in the later time periods. The boxed region highlights topographical differences in the frontal and central electrodes. ** indicates reliable discriminatory in the corresponding period in temporal decoding.

Next, we plotted the coefficients of the classifier patterns (derived from classifier weights W) for further neurophysiological interpretation (31). We decided to compare the classifier patterns across early (< 300 ms), and later (> 400 ms) time periods. To compare patterns in the early time periods across 'alert' and 'drowsy' periods, we plotted the same for every 40 ms between 180 ms to 420 ms in Figure 3F. The green box indicates clear differences across alert and drowsy in the early time periods. For the alert periods, the pattern between 180-220 ms indicates a strong involvement of signal in the fronto-central electrodes, whereas in the corresponding time periods under drowsy, the pattern seems to have shifted to more frontal electrodes although its contribution may be minimal as it was not reliably decodable. Under alert periods, the pattern shifts to more posterior regions (centro-parietal electrodes in the right side of the scalp) between 220 ms to 260 ms whereas under drowsy the pattern stays in the frontal electrodes itself, still not reliably decodable in the unconstrained voltage decoding. Further from 260-300 ms the pattern shifts to more parietal and occipital sites under alert periods and is only weakly parietal in the drowsy periods. The comparison of the classifier patterns across alert and drowsy periods (in early time periods), reveals different topographies that point to a clear differential processing of information that would further map onto shifted perceptual and central evidence accumulation stages of cognitive processing (2) possible affecting both, perceptual encoding and central accumulation. To compare patterns in the later time periods across 'alert' and 'drowsy' periods, we plotted the same for every 40 ms between 420 ms to 660 ms in Figure 3G. The magenta box indicates clear differences across alert and drowsy in the later time periods. Particularly, the patterns in the alert periods have a lower frontal activity and higher posterior activity, whereas in the drowsy periods, the activity in both the frontal and posterior sites is much more localised although decodability is overall lower. Although only suggestive, this change in the decodability intensity and distribution suggest a delayed processing dynamics or a change in the way of processing as alertness decreases.

Spatio temporal clustering of classifier patterns in alert and drowsy periods

Figure 4: Classifier pattern differences: To identify differences between the classifier patterns in Figure 3 (F,G) we performed spatio-temporal clustering. A) Cluster#1 indicates early periods (160 - 320 ms after stimulus) where alert activity is higher than drowsy periods. B) Cluster#2 indicates later periods (440 - 620 ms after stimulus) where alert activity is higher than drowsy periods. In terms of spatial locations, the early and later clusters differ in right parietal sites, central, middle electrode sites. C) Cluster#3 indicates early periods (180 - 310 ms after stimulus) where drowsy activity is higher than alert periods. D) Cluster#4 indicates later

335 periods (390 - 620 ms after stimulus) where drowsy activity is higher than alert periods. In terms of spatial
336 locations, the early and later clusters differ in left frontal sites, middle electrode sites.

337
338

339 The descriptive analysis of the classifier patterns indicate differences between alert and drowsy
340 periods. To establish the spatial and temporal signatures of such differences we performed a cluster
341 permutation test and identified regions where activity patterns in alert periods are different from
342 drowsy periods. This analysis resulted in the identification for four clusters.

343

- 344 a) Cluster #1 (alert activity > drowsy activity): indicates early periods (160 ms to 320 ms),
345 wherein activity is concentrated in parietal, central, posterior electrode sites. These spatial
346 patterns are likely to be involved in early perceptual/evidence accumulation processes
347 during the alert periods, as the majority of responses only start occurring from 400 ms
348 onwards.
- 349 b) Cluster #2 (alert activity > drowsy activity): indicates later periods (440 ms to 620 ms),
350 wherein activity is concentrated in occipital, parietal, central, posterior electrode sites.
351 These spatial patterns are difficult to interpret as the reaction times overlap with the
352 corresponding time periods. Hence this decoding pattern could be an amalgamation between
353 central evidence accumulation and motor implementation processes.
- 354 c) Cluster #3 (drowsy activity > alert activity): indicates early periods (180 ms to 310 ms),
355 wherein activity is concentrated in frontal electrode sites. These spatial patterns are likely to
356 be involved in early perceptual/sensory encoding processes during the drowsy periods, as the
357 majority of responses only start occurring from 900 ms onwards. The temporal
358 neurodynamics show delayed patterns in drowsy compared to awake (3 DE and 4 AC).
- 359 d) Cluster #4 (drowsy activity > alert activity): indicates later periods (390 ms to 620 ms),
360 wherein activity is concentrated in left frontal electrode sites. These spatial patterns are
361 likely to be involved in central evidence accumulation related processes. However it is
362 difficult to interpret as the corresponding time periods during alert overlap with response
363 related activity.

364

365 To summarise, these results indicate differences in spatial patterns related to early
366 sensory/evidence accumulation related process (cluster #1,3) across alert and drowsy periods. The
367 next step is to identify and tease apart motor preparation related processes occurring during later
368 time periods.

369

370 **Spatial and temporal signatures of the motor implementation of the decision across alert** 371 **and drowsy periods**

372

373 In classic cognitive processing model frameworks (Figure s1), motor implementation occurs after
374 the central evidence processes, when information of the decision is routed to the premotor network.
375 We preprocessed the EEG data now locked to responses (see methods section) and decoded the

376 response hand itself (irrespective of the stimulus presented). These response related patterns thus
377 would depict information related to the motor implementation process irrespective of the decision
378 related processes.

Decoding response hand locked to responses

Decoding left vs right responses in alert periods

A

B

Decoding left vs right responses in drowsy periods

Spatio temporal clustering of response patterns across alert and drowsy periods

C

D

380 **Figure 5: Temporal decoding (responses):** To identify differences in motor preparation (response related),
381 we decoded the left and right button press events irrespective of stimuli presented. A,B) AUC under alert and
382 drowsy periods, where the classifier was trained to discriminate between targets of left and right button

presses. Shaded regions show time periods with reliable ($p < 0.05$) discriminatory power. The time point of 0 corresponds to the actual button press event. The ramp up of the AUC under both conditions starting from 210 ms (alert), 160ms (drowsy) before the actual response indicates the ability of the decoder to detect neural signatures related to responses. Spatio-temporal clustering of differences in classifier patterns produces two clusters with similar time periods in contrast to the clusters (with early and later time periods) produced by stimuli related classifier patterns. C) Cluster#1 indicates periods (-190 to 190 ms) where alert activity is higher than drowsy periods. D) Cluster#2 indicates periods (-290 to 190 ms) where drowsy activity is higher than alert periods. In terms of spatial locations, the clusters differ in frontal, central, middle, parietal electrode sites. The presence of a single cluster with similar time periods across alert and drowsy periods indicate that motor related processes start appearing at similar time periods and hence do not change with modulations in alertness levels.

The decoding performance: AUC (Figure 5A,B) starts ramping up from -210ms (alert), -160ms (drowsy) from the response time, and increases when the button is actually pressed at 0 sec, reaching the peak soon after. In both cases, awake and drowsy, the neural decoding results lasts till 190ms. These results are in line with motor implementation related processes commonly reported around 200 ms before the actual motor response, gradually ramping up to produce the response itself. It is also interesting to note that the AUC for alert is always higher than drowsy, indicating less variability in neural concerted activity, leading to better classifier performance without time span differences.

The classifier patterns related to responses are presented in Figure s5. To establish the spatial and temporal signatures of differences in motor related processes we performed a cluster permutation test and identified regions where activity patterns in alert periods are different from drowsy periods. This analysis resulted in the identification of two clusters.

- a) Cluster #1 (alert activity > drowsy activity): indicates periods (-190 ms to 190 ms), wherein activity is concentrated in premotor, parietal, posterior electrode sites.
- b) Cluster #2 (drowsy activity > alert activity): indicates periods (-290 ms to 190 ms), wherein activity is concentrated in premotor, frontal electrode sites.

The existence of a single cluster (in time) across both alert and drowsy periods indicate that motor related processes do not vary in time with modulations in alertness levels. Furthermore for both alert and drowsy periods the start of the decoding happens during similar time periods around -200ms and last till similar time periods ~190ms.

To summarise, these results indicate that motor preparation related processes start and end at similar time periods during both alert and drowsy trials, suggesting the differences in motor preparation process may not be critical to the changes in decision making in low arousal when compared to early sensory and evidence accumulation related processes.

The above analysis related to decoding of stimuli and responses indicate that sensory encoding, central evidence accumulation related processes are modulated by alertness. However the decoding

analysis in the later parts could be confounded by response related processes occurring during alert periods. Hence we devised the next step of the analysis wherein we use parameters derived from the drift diffusion modelling, to understand how different elements of the sensory encoding and evidence accumulation are represented in the neural domain. This analysis is determined by the drift diffusion model, based in a mathematical implementation of decision making theory.

Neural Signatures of evidence accumulation of decision making modulated by alertness

To take advantage of the information gained by the drift diffusion modelling we decided to capture the neural implementation of the evidence accumulation parameters in the decision making process, and how it might differ in wakefulness and low alertness periods. We specifically aimed at capturing the change in the neurobehavioural dynamics of the evidence accumulation process in the awake and drowsy periods. To establish this we developed a novel method based on the drift-diffusion analysis shown earlier. First, we used the winning model (#8) and computed trial-by-trial drift parameter (dv). Second, we used dv as a dependent variable and regressed the same against ERP data (see methods section).

It can be expressed as

$$dv \sim \beta_0 + \beta_1(ERP)$$

The above equation can be written in patsy form as below.

$$dv \sim ERP : C(state, Treatment('Alert')) : C(stim, Treatment('Right'))$$

Here, dv represents trial-by-trial drift-rate, ERP represents z-scored ERP data per trial per time point per electrode, state represents alertness levels ('alert' or 'drowsy'), stim represents stimulus types ('left' or 'right'). Further we compute the proportion of the overlap of the posterior distributions of the trace obtained for both left and right stimuli separately under alert and drowsy condition (Figure 6A). This analysis is repeated for each participant and we obtain regression patterns (similar to the classifier patterns) per participant.

Next, we plot the regression patterns in alert and drowsy periods. The patterns in the alert periods (Figure 6B) indicate that the topography focuses initially in the fronto-central electrodes (150 ms to 250 ms), which then shifts to centro-parietal electrodes (250 ms to 350 ms), similarly to the classifier analysis (decoding patterns from figures 3F and 3G). The patterns in the drowsy periods (Figure 6C) indicate that the topography initially focuses on the frontal electrodes (though weaker compared to alert periods) in the interval from 150 ms to 250 ms. Further the patterns shift to more central electrodes (again weaker when compared to alert periods) in the interval from 250 ms to 350 ms.

Figure 6: Regression patterns: A) trial by trial drift rate (dv) was regressed against EEG data (z-scored and averaged per 50 ms), the difference in the posterior of regression rate was quantified per electrode per time-point per participant. B,C) mean regression patterns in alert and drowsy periods projected in electrode space show a classic centro frontal to parietal progression from 150ms to 500ms. The green box highlights topographical differences in the frontal, middle, central parietal electrodes (in early time periods) and magenta box highlights the frontal, parietal electrode sites (in later time periods).

Spatio temporal clustering of regression patterns in alert and drowsy periods

Figure 7: Regression pattern differences: spatio-temporal clustering of differences in regression patterns shown in Figure 6 (B,C) between alert and drowsy periods. A) Cluster#1 indicates early periods (70 - 280 ms after stimulus) where alert activity is higher than drowsy periods. B) Cluster#2 indicates later periods (470 - 720 ms after stimulus) where alert activity is higher than drowsy periods. In terms of spatial locations, the early and later clusters differ in right parietal sites, central, middle electrode sites. C) Cluster#3 indicates early periods (70 - 330 ms after stimulus) where drowsy activity is higher than alert periods. D) Cluster#4 indicates

later periods (380 - 680 ms after stimulus) where drowsy activity is higher than alert periods. In terms of spatial locations, the early and later clusters differ in left frontal sites, middle electrode sites.

The descriptive analysis of the regression patterns indicate differences between alert and drowsy periods. To establish the spatial and temporal signatures of such differences we performed a cluster permutation test and identified regions where activity patterns in alert periods are different from drowsy periods. This analysis resulted in the identification of four clusters.

- a) Cluster #1 (alert activity > drowsy activity): indicates early periods (70 ms to 280 ms), where activity is concentrated in parietal and central electrode sites. These spatial patterns are likely to be involved in early perceptual and evidence accumulation processes.
- b) Cluster #2 (alert activity > drowsy activity): indicates later periods (470 ms to 720 ms), where activity is concentrated in occipital, parietal, posterior electrode sites. These spatial patterns are likely to be involved in central processes of evidence accumulation-related processes alone.
- c) Cluster #3 (drowsy activity > alert activity): indicates early periods (70 ms to 330 ms), wherein activity is concentrated in frontal/central electrode sites. These spatial patterns are likely to be involved in early perceptual/sensory encoding processes during the drowsy periods.
- d) Cluster #4 (drowsy activity > alert activity): indicates later periods (380 ms to 680 ms), wherein activity is concentrated in left frontal electrode sites. These spatial patterns are likely to be involved in central evidence accumulation-related processes alone.

These analyses indicate spatial/temporal regions where the alert and drowsy periods differ in terms of sensory encoding and central evidence accumulation related processes. Furthermore, we have also teased apart these differences from motor related processes by directly using a parameter (drift rate) from the hierarchical drift diffusion modelling.

Arousal modulates frontoparietal cortical neural patterns in perceptual decision making

Next we performed an exploratory analysis to identify the putative neural sources of these differences in electrodes (Figure 7), we used a source reconstruction procedure to project the classifier patterns of the different clusters back to their neural sources separately (see methods section). In the source space again we performed cluster permutation tests to identify where in the cortex the activity in alert periods is higher than in drowsy periods. We used the actual value of the source activity (instead of absolute values), as we are interested in the distance between the two different patterns and not in overall activity levels.

- a) Cluster #1: These regions are putative locations for early sensory, evidence accumulation related processes in alert periods. The analysis reveals regions (inferior frontal, superior temporal, orbitofrontal) previously associated with the ventral attention network (32,33) in both hemispheres, with emphasis on temporal cortices, in agreement with auditory processing of information.

- b) Cluster #2: These regions are putative locations for central evidence accumulation related processes in alert periods. The analysis reveals regions (inferior parietal, middle temporal, right lateral prefrontal) that are part of the dorsal attention network (32,33) and central processing of flexible information (34).
- c) Cluster #3: These regions are putative locations for early sensory related processes in drowsy periods. The analysis reveals regions (transverse temporal, superior temporal gyrus, temporoparietal, fronto-parietal - inferior frontal gyrus, middle frontal gyrus) regions in the right hemisphere also associated with the ventral attention network (32,33).
- d) Cluster #4: These regions are putative locations for central evidence accumulation related processes in drowsy periods. The analysis reveals regions (inferior parietal sulcus, superior parietal lobule, fronto-polar - in right hemisphere) that are part of the dorsal attention network (32,33). Additionally it also revealed fronto-parietal regions in the left hemisphere

To summarise, these results indicate that during alert periods: the early sensory/central accumulation related process (until 300ms from stimulus presentation) are located in the regions corresponding to the ventral attention network, further during later periods (after 300ms) evidence accumulation related process are located in the regions corresponding to dorsal attention network (which is specialised for spatial attention). For the drowsy periods: the early sensory encoding related process (until 300ms from stimulus presentation) are located in the regions corresponding to the fronto-parietal, ventral attention network, further during later periods (after 300ms) evidence accumulation related process are located in the regions corresponding to fronto-parietal, dorsal attention network. This indicates that fronto-parietal regions of the brain are further recruited for the late decision-making processes implemented in the drowsy periods, pointing to a cortical network reconfiguration in the evidence accumulation between normal and low alertness periods.

Projection of regression patterns in source space

Figure 8: Regression patterns projected into source space: To identify the putative regions/networks in the brain related to perceptual decision-making and its modulation by alertness we projected the regression patterns in Fig 6 to source space. A) regression patterns in cluster#1 (Fig 7A) in the early time period in electrode space are projected to source space and spatio-temporal clustering identifies differences in regions related to the ventral attention network. B) similarly regression patterns in cluster#2 (Fig 7B) reveals differences in dorsal attention networks. C) similarly regression patterns in cluster#3 (Fig 7C) reveals differences in fronto-parietal regions. D) similarly regression patterns in cluster#4 (Fig 7D) reveals differences in dorsal attention network and fronto-parietal regions.

568

569 DISCUSSION

570

571 In this study we were interested in how changes of arousal modulate perceptual decision making
572 and spatial attention. Informed by previous studies(25,26) we conducted a series of data-driven and
573 model-derived analyses to test different aspects of perceptual and cognitive processes underlying
574 behavioural and neural mechanisms of decision making.

575 We investigated how alertness modulates different neurocognitive components of perceptual
576 decision-making using a tone localisation task. We replicated the spatial bias induced by alertness
577 changes that has been shown in healthy participants, not dissimilar to that seen following brain
578 damage (unilateral spatial neglect). We characterised behavioural changes in decision making as
579 impacted by the fluctuations in arousal, revealing a decrease in performance due to lower sensitivity
580 to the stimuli but not due to response bias. We also showed a specific effect of alertness reducing
581 the rate of evidence accumulation in a drift diffusion model with little effects on the starting point
582 of the decision (bias). At the neural level we showed that the formation of brain patterns (relating to
583 stimuli presented) is delayed and decreased by lower levels of alertness, suggesting a
584 reconfiguration of the neural networks of the decision. We confirmed this suspicion by mapping the
585 behavioural model parameters of evidence accumulation (drift) into the neural evoked responses,
586 unveiling a more distributed neural patterns, in space and time, in drowsy evidence accumulation
587 and further recruitment of parietal and frontal cortices in low alertness as compared to awake. We
588 think this is evidence for a neural reconfiguration implementing compensatory mechanisms at play
589 in the decision dynamics when challenged by low alertness. For spatial attention theories we are
590 able to add further support to the interaction between ventral attention system (arousal related)
591 and dorsal attention system (spatial attention) when arousal is modulated, and to tease apart the
592 change in the predisposition of the system (bias-point, prior) from the modifications to the decision
593 process itself (drift-rate). For decision making theories, we have shown that low alertness in humans
594 can be used to study evidence accumulation mechanisms and added a new level of description by
595 characterising possible mechanisms of resilience of the neurocognitive processes when tonic
596 alertness decreases. Thus, we were able to show the behavioural flexibility of the system and
597 capture the temporal and spatial neural progression and reconfiguration caused by the change in
598 the internal context - describing its cognitive homeostasis.

599

600

601 **Behavioural changes in decision making show specific modulations of spatial** 602 **attention with decreased arousal**

603 We established, using multi-level modelling, that the errors for left-sided stimuli increased as
604 participants became drowsy. However the same did not occur for stimuli coming from the right side.
605 This evidence is convergent with Bareham et al(25) but with a different study design, analysis
606 methods and participant sample, furthering the conclusion that there is increased inattention or
607 neglect if stimuli comes from the left side when healthy people become drowsy. Advancing the
608 previous behavioural results, we fitted the proportion of rightward responses of individual
609 participants and identified the subjective midline (radial distance where a participant is equally
610 likely to press left or right response), finding that it shifted to the left side (indicating more right
611 responses) as individuals became drowsy (low alertness). This is in agreement with line-bisection
612 studies(35) showing healthy individuals bisect lines to the left of the veridical midline,
613 overestimating the right side of space. Additional studies(36) have also identified possible neural
614 origins of such bias ('pseudoneglect') and identify task independent activity in the ventral attention
615 network as one of the main modulating factors behind this shift. However these studies did not
616 examine whether trial to trial alertness of individuals was modulated. Several studies(33,37) have
617 shown that activity in the ventral attention network interacts with the arousal system (tonic
618 alertness), and alertness instability, as shown directly in this study, could be a cause for these
619 effects.

620 **Slower evidence accumulation when alertness decreases**

621 To characterise more general aspects of decision making modulated by alertness, we decided to
622 parse the different elements of the process. Using signal detection theory to identify the changes in
623 sensory signal and noise distributions due to alertness, we found that sensitivity (d') was reliably
624 -decreased- by alertness whereas criterion (response bias) was not(38). This suggests that the
625 sensory/perceptual representations have indeed been modulated by alertness, and that participants
626 were not arbitrarily pressing more rightward responses (in the face of uncertain stimuli) as alertness
627 decreases. To complement signal detection theory and further characterise how arousal modulates
628 decision making, we used the rich variability in reaction times produced by awake and drowsy trials
629 in a drift diffusion model, under the assumption that in each trial, noisy evidence is slowly
630 accumulated over time until criterion to make a decision is reached(28). We constructed several
631 models that allowed independent variation of parameters, such as starting point (bias) and drift rate
632 (evidence accumulation), based on stimulus type and alertness levels of the participant. The
633 statistical model with the higher evidence showed that drift rate was modulated by both alertness
634 levels and stimulus type, whereas the starting point was only affected by stimulus type (left or right
635 tone). Further analysis of the winning model also revealed that the evidence for modulation of the
636 starting point was weak while that for drift rate was stronger and reliably different for stimuli from
637 the left side compared to the right side. This shows that alertness indeed differentially modulates
638 evidence coming from the left side of space compared to the right side. Decision making studies in

the visual modality(39) have shown that drift rates are closely related to the strength of stimulus encoding, partially affected by attentional strength, and central evidence accumulation to a decision threshold. Further studies have also shown neural correlates of such parameters in the EEG(40,41) and its relationship with stimuli strength. Hence we consider the effect of alertness modulation on drift rate as acting specifically on spatial attention affecting the stimulus encoding, as well as in central processes of evidence accumulation, revealing a general mechanism where partially independent systems, arousal and decision making, interact.

646

647 **Alertness delays the neural signatures of perceptual decision making and alters the** 648 **dynamics of the neural networks**

We probed the neural signatures of decision making processes using time-multivariate decoding analysis in high-density EEG. The primary aim was to understand, unconstrained and model-free, how the brain decodes information related to correct decisions when the stimuli presented was ‘left’ or ‘right’ under alert and drowsy conditions. Under alert conditions the decoding started early, at 160 ms and lasted until 730 ms, however, under drowsy conditions the effects seemed delayed to 420 ms and lasted until 730 ms. This ~260 ms delay may indicate that, under lower arousal, the brain requires a longer time to decipher the direction of stimuli as suggested by slower reaction times and a change in the drift (evidence accumulation) of the diffusion model, or alternatively, that the neural code in the early window of processing is not captured by decoding evoked potentials locked to the stimulus presentation. Furthermore, the overall strength of decoding of the alert condition was systematically higher than the drowsy condition, suggesting that the brain processes responsible for decision making are less variable under the alert condition compared to the drowsy condition, or it may indicate less efficient decision processing during low alertness (see(42) for an account on sleep deprivation on cognition).

From a cognitive dynamics perspective the different neural time clusters point potentially to a first stage of information encoding and second of central evidence accumulation (43) that others have mapped to perceptual and central decision processes (3). If that is the case, the first neural signal cluster in drowsy (cluster#3 in Figure 4C), showing changes in the spatial pattern and a lack of reliable decoding may be interpreted as less efficient encoding and transfer of information in low alertness or may be using different network dynamics that are not captured by the direct decoding. The second cluster (cluster#4 in Figure 4D) more similar between states, may signal a lower efficiency of central evidence accumulation, as it happens around the time the central decision process occurs in cognitive control(13). Furthermore, the second window of processing may also show part of the motor implementation processes following the evidence accumulation triggering the motor plans, at least for the awake condition (cluster#2 in Figure 4B), where the responses overlap for several participants (Figure 3D). The overlapping processes from ~400ms in awake, a mixture of central evidence accumulation and motor implementation of the decision, depending on

the participant, are easily dissociated when drowsy, when the motor plan is implemented ~500ms later.

Next, we aimed to understand how such neural signatures are depicted at the sensor level. For this, we identified classifier patterns, which are neurophysiologically interpretable(31) from classifier weights. In the alert condition, the early pattern (180-220 ms) indicates activity over frontocentral electrodes, further shifting to more central and parietal sites later on. In the drowsy condition, the pattern initially starts at frontal sites (though not strong enough to be decoded) and shifts to central and parietal channels, later on, from 380 ms onwards. This indicates that the formation of central, parietal patterns (possibly related to the first stage of encoding, evidence accumulation) takes longer or starts later in drowsy periods compared to alert. Alternatively, the information decoded from the multivariate ERP patterns can be captured in awake but in drowsy the information changes code and it is not captured anymore in this early period. The spatial and temporal distribution of the decoding pattern resembles both a classic P2 or early P3, which has been dubbed as the build-to-threshold signal(44), and the centro-parietal positivity (CPP), a more specific signal of evidence accumulation(41,45) respectively. O'Connell et al showed that the CPP (an EEG evoked response) tracked evidence accumulation in perceptual decision making and was sensitive to perturbation, opening the window for the tracking of signals with sufficient temporal resolution to map the dynamics of decision making in humans(41) beyond what we have learnt in monkeys(14). Following this line of thought, our results highlight the neural dynamics of decision making when the external world remains unaltered, physical evidence held constant, but the internal milieu fluctuates exerting a modulatory influence on the cognitive decision making systems that caused them to reconfigure to solve the task at hand.

Evidence accumulation reveals slower and spatially extended frontoparietal network to resolve the decision when challenged by arousal

The decoding analysis is not guided by parameters that directly test mechanistic hypotheses. It is only based on correct responses from participants and hence it can't reveal the neural source of inattention to the left side of space, or differentiate between cognitive processes proposed by theory. further, the overlap of response periods in the later time window in alert trials, motivates us to use a parameter that can guide neural exploration with a direct readout of evidence accumulation of the decision, avoiding biases and motor related neural signatures. Here we developed a novel method using regression patterns that identified differences in drift rate across alert and drowsy periods into the neural dynamics.

We performed a cluster permutation test to identify regions in the brain where temporospatial regression patterns where alert and drowsy periods differ. The spatial topography of these four clusters resemble the clusters from the direct decoding results, providing strength to our preliminary interpretations from the decoding analysis (in the sensor space). To identify the spatial

712 location of these patterns we projected the data at the sensor level, and to cortical sources for the
713 four different clusters to see the cortical involvement.

714 We first look at clusters(#1,2) to understand how alert periods are active under this
715 tone-localisation task. These showed regions in the inferior frontal, superior temporal, orbito
716 frontal source regions that resemble the patterns of the ventral attention network with further
717 activity of auditory processing. Although we cannot directly differentiate between early perceptual
718 encoding and central accumulation with this decision, this first cluster most likely corresponds to
719 both in the alert periods. This part of the process of the spatial decision shows the early
720 involvement of these structures that have been linked, in several studies (46,47), to lesions in the
721 inferior frontal and superior temporal gyrus resulting in egocentric neglect (inattention to one side
722 of space from the viewer centric perspective). Furthermore, lower activity in the ventral attention
723 network was also reported for neglect patients suffering from deficits in arousal (33). This agrees
724 with the higher activity observed in the regression patterns for the alert periods compared to the
725 drowsy periods of lower arousal. The cluster#2, revealed regions in the inferior parietal, middle
726 temporal, right lateral prefrontal source regions that have been associated to the dorsal attention
727 network (46,47). This cluster most likely corresponds to the central evidence accumulation that is
728 directly involved in evaluation of the spatial attention information. To summarise, under alert
729 periods: when the tone is presented, a more ventral attention network gets recruited first, involved
730 in early sensory encoding and evidence accumulation, followed by the information processed
731 further a more dorsal attention network, where the central evidence accumulation primarily takes
732 place to reach the decision.

733 Next we look at clusters(#3,4) to understand the neural dynamics of evidence accumulation during
734 drowsy periods. Cluster#3, revealed regions in the transverse temporal, superior temporal gyrus,
735 temporoparietal junction, fronto-parietal regions (inferior frontal gyrus, middle frontal gyrus) in the
736 right hemisphere that can be considered part of the ventral attention network. Also here several
737 studies (48,49) have reported lesions in the right ventral frontal cortex in patients suffering from
738 both arousal related deficits and neglect to the right side of space. The highlighting of these regions
739 in our study indicates that they are different from the alert periods and their higher activity in the
740 regression patterns points to a potential source of spatial neglect (co-occurring with arousal
741 deficits), which is similar to that observed in patient studies. There is the need to further investigate
742 this possible dissociation in further studies, either with lesion metaanalysis or with virtual lesions in
743 normal volunteers. We think this cluster would correspond to a combination of sensory encoding,
744 evidence accumulation in the drowsy periods highlighting an early large cortical reconfiguration of
745 the evidence accumulation process with arousal decreases. Finally Cluster#4, revealed regions in the
746 inferior parietal sulcus, superior parietal lobule, fronto-polar regions that may be considered part of
747 the dorsal attention network in both hemispheres ascribed to spatial attention processes.
748 Additionally it also revealed further fronto-parietal regions involvement in the left hemisphere,
749 perhaps recruited to provide additional processing to reach criterion to compensate for arousal
750 related deficits.

Our account of a spatial neglect-like effect in participants under lower arousal is as follows: When participants are fully alert, the information is processed initially by the ventral attention network, followed by the dorsal attention network (specialised in spatial attention). Whereas when participants are drowsy, the ventral attention network although disproportionately affected in the right hemisphere (33,48,49), it is still involved in sensory encoding and early evidence accumulation processes. This reconfiguration in the ventral attention network is further propagated in the next stage, the dorsal attention network. This second part of the decision closely tallies with the proposal by Corbetta and colleagues to account for spatial neglect found in stroke patients suffering from lesions in right-hemispheric regions. But further, this network shows temporal and spatially extended recruitment of the parietal and frontopolar cortices to compensate for the direct effects of arousal in the resolution of the noisy evidence presented to reach a decision in the spatial task.

From a decision making perspective, we find that moving from fully alert to a lower arousal state the brain tries to recruit and expand evidence accumulation related process to fronto-parietal regions instead of the usual temporoparietal regions, consistent with an increase in cognitive demand proposed by the Multiple Demand system (50). Another line of evidence suggesting neural fluctuations influencing evidence accumulation comes from Summerfield and Wyart (51) where the phase of ongoing parietal delta oscillations impacted choice. We interpret that the neural reconfiguration occurring in the transition to lower states of consciousness (inattentive, mildly drowsy) brings slower rhythms and suppresses fast oscillatory events (disappearing alpha and beta), and that this changes the nature of the noise in the brain, forcing the cognitive system to exchange information differently. In this case the arousal changes may be impacting negatively the efficiency of the central accumulation of evidence and the system compensates by extending the neural processes in time and space.

As we lose consciousness, the neural system responsible for decision making adapts to the internal challenge of decreasing arousal and shows its resilience, exerting homeostatic regulations at the cognitive level to maintain performance. The attention and wakefulness fluctuations experienced in humans (and animals) are common during the day and they are not only dependent on the circadian and sleep-wake regulation pressure (52), but also on genetic, epigenetic, environmental and life history factors that shape the alertness aspects of attention as well as the alertness aspect of arousal (53,54).

Concluding remarks

In this study we investigated the effects of arousal fluctuations on decision making and proposed that despite the internally generated pressure due to the conscious state change, the decision making system is flexible enough to maintain performance by reallocating neural resources. We found that the decision becomes more liberal (in terms of performance) as alertness decreases, showing lower sensitivity with a less steep psychophysics curve. We further show that an evidence accumulation parameter (drift rate) explains the change in the patterns of responses with decreased alertness. At the neural level we demonstrate a change in the manner of processing of the decision

789 both by direct decoding of the neural dynamics, and when guided by the evidence accumulation
790 behaviour in each trial.

791 Transitions of consciousness in the near-awake to light-decrease of alertness is emerging as a model
792 for internally caused interactions (18,26,27,55,56). With lesion and pharmacological challenges,
793 neuropsychology and cognitive neuroscience have tried to define the necessary and sufficient brain
794 networks, areas and dynamics of the brain to implement cognition, revealing compensation,
795 reconfiguration and plasticity (57–59). Semi-causal effects of the internal interference exerted by
796 the relative independence of the arousal system on the cognitive process, paves the way for the use
797 of wakefulness and arousal challenges in a principled manner, adding new tools for cognitive brain
798 research. Cognitive neuroscience uses models, correlational and causal methods to reach consensus
799 about the underlying mechanism of thought (60). Here, we have followed a theoretically motivated
800 question about perceptual decision making systems, using behavioural modelling to understand the
801 system, but at the same time directly modulated the neural networks with arousal to uncover the
802 mechanisms of brain function.

803

804

805 **EXPERIMENTAL PROCEDURES**

806

807 **Stimuli and Protocol**

808 Forty-one healthy participants (no auditory, neurological or psychiatric abnormalities) participated
809 in this study. Data from nine participants was discarded due to a) technical problems with
810 headphone amplifier (8) b) Not following task instructions (1). Thus only data from 32 participants
811 (24.46 ± 3.72 years old, 14 males) was considered for further analysis. All participants were
812 self-reported to be right-handed and the same was also established by using Edinburgh Handedness
813 Scale (61). Each participant had a handedness score of above 0 (right-handed) with mean $80.26 \pm$
814 23.59 . Only easy sleepers (as per self report) were recruited and further they were administered with
815 the Epworth Sleepiness scale (62) on the day of the experiment. 29 participants had a sleepiness
816 score ≥ 7 (classified as easy sleepers) and 3 of them had a sleepiness score ≥ 4 . All participants
817 were asked not to consume any stimulant like Coffee/Tea before the experiment that day. The study
818 and the experimental protocol was approved by the cambridge psychology research ethics
819 committee and written informed consent was provided by all participants. A monetary
820 compensation of £30 was provided for participation in the study.

821

822 Each participant underwent two experimental sessions a) Alert b) Drowsy.

823

824 **Alert session:** participants were presented with 124 complex harmonic tones (guitar chord) that fell
825 on the left or right side of their veridical midline (0°) ranging from -59.31° to $+59.31^\circ$. These tones
826 were recorded using in-ear microphones in free-field (25). Six tones from -59.31° to -39.26° were

presented two times each; twelve tones from -35.24° to -1.86° were presented four times each. A similar pattern was repeated on the right side with twelve tones from 1.86° to 35.24° presented four times each, six tones from 39.26° to 59.31° presented two times each. The tones in the midline (0°) were presented four times, resulting in a total of 124 tones. The order of tones presented was randomized per participant. Further, participants were instructed to keep their eyes closed and respond (as quickly and as accurately as possible) with a button press (using left/right thumb) indicating the location of the tone (left or right). Each trial began after a random interval of 2-3 seconds and if the participant did not respond for 5 seconds, the next trial was started. The participants were also instructed to stay awake throughout this session.

Drowsy session: participants were presented with 740 complex harmonic tones (as above) that fell on the left or right of their veridical midline (0°) again ranging from -59.31° to $+59.31^{\circ}$. Six tones from -59.31° to -39.26° were presented twenty times each; twelve tones from -35.24° to -1.86° were presented twenty times each. Similar pattern was repeated on the right side with twelve tones from 1.86° to 35.24° being presented twenty times each, six tones from 39.26° to 59.31° presented twenty times each. The tone in the midline (0°) was presented twenty times, resulting in a total of 740 tones. The order of tones was again randomized per participant as in the alert session. participants were again instructed to keep their eyes closed and respond (as quickly and as accurately as possible) with a button press (by left/right thumb) indicating the direction of the tone (left or right). Each trial began after a random interval of 4-5 seconds and if the participant did not respond for 5 seconds, the next trial was started. In this session, the participants were allowed to fall asleep (and become drowsy) and were gently awoken if they didn't respond to more than 3 trials consecutively.

Before the start of the experiment, the participants were allowed a practise session to familiarise with the task.

Preprocessing

EEG data was acquired with 129 Ag/AgCl electrodes (Electrical Geodesics Inc) using Cz as the reference electrode. The impedances of all electrodes were kept below 100 K Ω (to ensure higher signal to noise ratio) and data was acquired at a sampling rate of 500 Hz. EEG data was pre-processed with custom made scripts in MATLAB (MathWorks Inc. Natick, MA, USA) using EEGLAB toolbox (63). The preprocessing steps are as follows: First, the peripheral electrodes that covered the regions of forehead, cheeks and neck were removed to reduce artifacts related to eye and muscle movements, thus retaining only 92 channels that covered the scalp. Second, the data was bandpass filtered with zero phase shift between 1 and 40 Hz using hamming windowed-sinc FIR filter and further resampled to 250 Hz. Third, pre-trial and post-trial epochs per trial were created. For the pre-trial epochs, the data was epoched from -4000ms to 0ms prior to the onset of the stimuli. The pre-trial epochs were created only in the drowsy session and not in the alert session (details below). For the post-trial epochs, the data was epoched from -200ms to 800ms to the onset of the stimuli for both the alert and drowsy sessions. Fourth, the trials that exceeded the amplitude

threshold of ± 250 μV were removed in a semi-automatic fashion. Fifth, the bad channels were detected in a two-step fashion: a) channels are considered bad (zero activity) if channel variance is below 0.5. b) The normalized power spectrum of the remaining channels was computed and any channel that exceeded the mean power spectrum by ± 3 standard deviations was marked bad. Sixth, to remove further artifacts related to eye-blink and muscle movement, independent component analysis (ICA) was performed on the channels not marked as bad in the previous step. ICA components that correspond to artifacts were rejected by manual inspection. Seventh, the bad channels were now interpolated using spherical interpolation. Eighth, the bad trials were detected again using an amplitude threshold of ± 250 μV and bad electrodes (those exceeding the threshold) in such trials were interpolated in a trial-by-trial fashion. Ninth, the post-trial epochs were re-referenced to the average of all channels (whereas the pre-trial epochs were maintained with the recorded reference - Cz).

879

880 Alertness levels

The preliminary step in both behavioural and neural analysis is to classify periods of data into 'alert' and 'drowsy'. The data from the pre-trial epochs were used to classify each trial into alert or drowsy. For the alert session, all pre-trial periods were considered to be awake as participants were explicitly instructed to stay awake (and none of the participants failed to respond to any of the trials in the alert session). In the drowsy session, the participants were allowed to fall asleep and hence some trials would be considered as alert and some would be drowsy. For each trial in the drowsy session, pre-trial epochs were analysed using the micro-measures algorithm (24). Each trial was classed as 'alert', 'drowsy(mild)', 'drowsy(severe)'. Further only trials that were classified as 'drowsy(mild)' were used as drowsy trials. The other trials were excluded because usually participants don't respond when under 'drowsy(severe)' trials. We similarly excluded the 'alert' trials in the drowsy session to instead compare the drowsy trials of the drowsy session to all of the trials in the alert session.

893

894 Behavioral analysis

895

896 Error proportion

In order to understand how the rate of errors differs across different stimuli (tones to the left or right side of midline) and how it is modulated by change in alertness levels we performed the following analysis. First, we computed the proportion of errors made by each participant under each alertness level ('awake', 'drowsy') and under each stimulus type ('left' or 'right' tone). If the total number of trials for any participant under any condition is less than 5, then the corresponding error proportion (for that condition) is ignored in the further analysis. We defined 4 different multilevel models to understand the modulation of error proportion by state of participant ('alert', 'drowsy') and stimulus ('left', 'right'). We used multilevel models, as different participants had different levels of alertness (differing number of alert, drowsy trials). In the null model, the error proportion depends only on the mean per participant (fixed effect) and the participant id (random effect). In

the second model (state model), the error proportion depends only on the state of the participant ('alert' or 'drowsy' as fixed effect) and individual participant (participant id is used as random effect). In the third model (stimulus model), the error proportion depends only on the stimulus ('left' or 'right' as fixed effect) and individual participant (participant id is used as random effect). In the fourth model (state-stimulus model), the error proportion depends on a combination of state of participant ('alert' or 'drowsy') and the stimulus ('left' or 'right'), both used as fixed effects and individual participants (participant id is used as random effect). These 4 models were fitted using the 'lmer' function ('lmerTest' package) in R (64) and the winning model is identified as the one with the highest log-likelihood by comparing it with the null model and performing a likelihood ratio chi-square test (χ^2). Finally the top two winning models were compared against each other using 'anova' function in R (65), to validate whether the winning model (if it is more complex) is actually better than the losing model (if it is simpler). The state-stimulus model emerged as the winning model.

The different models along with their log-likelihood values are shown below:

Model comparison:

Model	Parameters	Log-likelihood	Pr(> χ^2)
Null	Fixed: mean, Random: participant id	61.24	-
State	Fixed: state, Random: participant id	66.62	<0.001
Stimulus	Fixed: stimulus, Random: participant id	68.45	<0.001
State-Stimulus	Fixed: state*stimulus, Random: participant id	77.98	<0.001

Type III Analysis of Variance Table with Satterthwaite's method of the winning model (State-Stimulus):

Model elements	Sum Sq	Mean Sq	NumDF	DenDF	F value	Pr(>F)
State	0.21023	0.21023	1	95.07	14.0465	0.0003061
Stimulus	0.28140	0.28140	1	95.07	18.8018	3.607e-05
State:Stimulus	0.10312	0.10312	1	95.07	6.8897	0.0101019

The state-stimulus (winning model) was further analysed with the 'anova' function and it was found that there was a reliable main effect of both state ($p<0.001$) and stimulus ($p<0.0001$) on error proportion and also the interaction between state and stimulus also had a reliable effect ($p<0.05$) on

error proportion. Estimated marginal means were computed using the ‘emmeans’ package in R to perform post-hoc tests using tukey adjustment for multiple comparisons.

Subjective midline shifts

The change in subjective midline was quantified by fitting psychometric functions to the responses produced by each participant under alert and drowsy conditions.

The proportion of rightward responses for each participant under each stimulus condition from -60° to $+60^\circ$ was fitted with a cumulative normal function using the generalized linear model ‘glm’ function in R. The link function used in the model was ‘probit’ which quickly asymptotes to the axes compared to the ‘logit’ function. The fit also constrained in such a way the number of trials under each stimulus angle weighed the shape of the function. The mean of such a cumulative normal function (the point where the curve crosses 0.5 in the y-axis) is referred to as the subjective midline (‘bias’). The subjective midline is the stimulus (angle) at which the participant performs at chance (0.5), which in an ideal world would be closer to the veridical midline (0°). This fitting procedure was performed for each participant separately under each condition (alert, drowsy). Examples of fits for two participants are shown in the supplemental information (Figure s1). The standard deviation of the cumulative function (slope or steepness) represents the ‘sensitivity’ of the system. In general, large variations in the bias point tend to reduce the sensitivity of the system. Further we also ignored participants (leaving 11 participants) in the drowsy condition that had a bias point of more than 60° (as the overall stimulus angle can vary only between -60° to $+60^\circ$) and standard deviation of more than 30° .

Drift-diffusion modelling

The different elements of the decision-making process can be teased apart by using the drift-diffusion model. The primary parameters of this model include

- i) drift-rate - ‘v’ -- rate of evidence accumulation
- ii) bias point - ‘z’ -- starting point of the decision making process
- iii) boundary separation distance - ‘a’ -- distance between the two decision boundaries
- iv) non-decision time - ‘Ter’ -- for accounting other processes like stimulus encoding (before the start of evidence accumulation process), response execution (after the end of evidence accumulation).

We implemented the drift diffusion process using hierarchical drift diffusion model (HDDM) (66) (version 0.7.1) that allows for a hierarchical Bayesian procedure to estimate the model parameters. The principal reason for using such hierarchical procedures is because different participants fall asleep in different ways (differing number of alert and drowsy trials). Using a hierarchical bayesian procedure allows for robust estimation of model parameters under such limited conditions of trials available (67). We used the response of each participant (‘left’ or ‘right’ button press) instead of

975 accuracy ('correct' or 'incorrect') to fit the HDDM. Such a procedure is referred to as
 976 Stimulus-coding and allows for robust estimation of 'z' without being biased by accuracy. To
 977 identify the best parameters, we identified 8 different models where each model allows the
 978 parameters ('z','v') to vary depending on state ('alert' or 'drowsy') or stimulus ('left' or 'right'). In
 979 each model, 15000 samples from the posterior distribution were estimated using Markov chain
 980 Monte Carlo methods. 5000 samples were further discarded as burn-in to discard the effect of initial
 981 values on the posterior distribution estimation. The winning model was chosen as the one with the
 982 lowest deviance information criterion (DIC). DIC allows for computing model accuracy while
 983 penalising for model complexity (68). Further the winning model was also checked for model
 984 convergence using the Gelman-Rubin statistic. The convergence statistic was computed for 5
 985 different model runs and was found to be closer to 0.99 (values closer to 1 but less than 1.2 indicate
 986 convergence) (68,69). Posterior predictive fits for a few sample participants are shown in the
 987 supplemental information (Figure s2). The DIC scores for different models are given below

Model	Drift-rate (v)	Bias-point (z)	DIC score
1	'state'	'state'	22516.09
2	'state'	'stim'	22474.67
3	'state'	'state','stim'	22273.48
4	'stim'	'state'	24057.42
5	'stim'	'stim'	23901.97
6	'stim'	'state','stim'	22110.07
7	'state','stim'	'state'	21713.90
8	'state','stim'	'stim'	21700.95

991
 992
 993
 994 **Neural analysis**
 995
 996 **Decoding**
 997 We used multivariate pattern analysis (MVPA) techniques to analyse the divergent pattern in EEG
 998 data. Decoding is a form of MVPA in which we analyse patterns of brain activity in order to predict
 999 the experimental condition (or stimuli) under which that pattern was generated. Traditional ERP
 1000 (Event related potentials) analyses rely on using a-priori identified spatial locations or temporal
 1001 segments in the data to measure the differences across conditions. However decoding techniques do
 1002 not rely on a-priori definitions and perform much better in detecting differences across
 1003 experimental conditions (30).

1004

1005 **Temporal decoding:**

1006 Temporal decoding involves using EEG data (X) composed of size: [Electrodes x Time points x

1007 Trials] to predict the experimental condition (Y). The experimental condition predicted here was

1008 the direction of the stimuli presented e.g. left or right stimuli. The first step in the decoding analysis

1009 consists of fitting an estimator (w) to a subset of the data (X) called (X_{train}) to predict a subset of

1010 the experimental condition (Y) called (Y_{train}). The second step involves using this trained

1011 estimator on another subset of the data (X) called (X_{test}) to predict subset of the experimental

1012 condition (Y) called (Y_{test}). The third step involves evaluating the performance of this estimator

1013 using a measure (e.g. accuracy) by comparing the prediction ($\hat{Y}_{test}$) with the actual label (Y_{test}).

1014

1015 **Estimator construction:**

1016 First, the EEG data is subjected to a standard scaler (using `StandardScaler()` from scikit-learn version

1017 0.22) that removes the mean of the data and scales it by its variance. Second, we used logistic

1018 regression to estimate the model parameters for maximally separating the different experimental

1019 conditions. Third, we implemented temporal decoding by using the sliding estimator (using

1020 `SlidingEstimator()` from scikit-learn) to fit the logistic regression model per time-point.

1021

1022 **Cross-validation:**

1023 The EEG data was first downsampled to 100 Hz and further binary classification was performed

1024 between two conditions ('left' and 'right' stimuli) separately across alert and drowsy conditions per

1025 individual participant. As the target of the classification was stimuli being presented we only

1026 considered the trials wherein the participant made the correct decision. The participant was

1027 considered for classification only if they had at least 25 trials under each condition. Further 5-fold

1028 cross validation was performed such that 4 folds were used for training and the last fold was used as

1029 a testing set.

1030

1031 **Validation measure:**

1032 The classifier performance was evaluated using Area Under the Curve (AUC) of the

1033 receiver-operating characteristic (ROC). It is implemented using 'roc_auc' in the sliding estimator

1034 function in scikit-learn. When AUC is about 0.5 the classifier performs at chance, while the AUC

1035 score of 1 has a very good separability across classes. We computed the AUC-ROC score per

1036 participant as the average of the score across all the cross-validation folds.

1037

1038 **Group statistics:**

1039 In order to identify the reliability of the AUC score at the group level, we performed a cluster

1040 permutation test (participants x timepoints) using MNE version 0.19.2 (`spatio_temporal_`

1041 `cluster_1samp_test`) (70). Thus producing p-values per time point at the group level, from which

1042 time points wherein we can infer those regions wherein AUC is reliably different from chance (0.5)

1043 at the group level.

Coefficients of patterns:

The parameters of the decoding model are not neurophysiologically interpretable in a straightforward way (31). Hence the parameters of the backward model (decoding) need to be transformed to produce the forward model. This is done by obtaining the coefficients of the estimator model per participant using the ‘get_coef’ function from MNE(‘_patterns’). For performing group statistics in electrode space, we used the same cluster permutation based approach as described earlier.

1051

Source reconstruction of patterns:

The coefficients created above can be projected into the source space to infer the brain regions involved in the pattern activity. Source reconstruction was achieved using a combination of Freesurfer (71) and MNE. First, the surface was reconstructed using ‘recon-all’ (Freesurfer) using the default ICBM152 template for the structural magnetic resonance image (MRIs). Next, the Boundary element model (BEM) was created using ‘make_watershed_bem’ from MNE. Further, scalp surfaces for the different element boundaries were created using ‘make_scalp_surface’ from MNE. Second, we performed the registration of the scalp surface with the default EEG channel locations (with fiducials as markers) manually using ‘coregistration’ from MNE. Third, forward solution was computed using ‘make_bem_model’ from MNE with conductivity = [0.3, 0.006, 0.3]. Fourth, to test if the source reconstruction of the electrode data is accurate we projected the ERP data of a sample participant into source space and analysed data from different regions of interest to confirm its validity as shown in the supplemental information (Figure s4). The classifier patterns of each participant were then projected into source space in the following manner. First, we computed the noise covariance using the baseline data from -0.2 to 0 ms. Second, we used the forward solution (8196 vertices) and the noise covariance to create an inverse operator using ‘minimum_norm.make_inverse_operator’ from MNE (loose=0.2). Third, used the individual classifier pattern per participant and applied the inverse operator on it (method = MNE, SNR = 3, $\lambda^2 = 1/(\text{SNR})^2$) to produce the source reconstruction of the classifier patterns per participant. For performing group statistics in the source space we used the same cluster permutation based approach as described earlier.

1073

Decoding locked to responses of participants:

To tease apart the process related to evidence accumulation from motor implementation, we decided to decode the response hand of the participant. The EEG data for the response decoding was created by epoching the data from -800ms to 200ms prior to the onset of the response from the participant. The response-locked trials were preprocessed in a similar fashion to the stimulus-locked trials (details in the section of Preprocessing). The alertness level for the corresponding response-locked trial was obtained from the labels of the stimulus-locked trial. The temporal decoding and other methods used are similar to the stimulus-locked trials, except that the target of decoding is the response hand being pressed (left or right thumb).

1083

1084 **Neuro-behavioural analysis**

1085

1086 **Regression with drift diffusion model**

1087 To identify the correlates of the evidence accumulation process, we used the model parameters
 1088 generated by the drift diffusion model (winning model #8). First, the ERP data (post trial epochs)
 1089 were z-scored per electrode per trial. Second, the ERP data was baseline corrected with pre-trial
 1090 data from -200ms to 0ms. Third, the ERP data was averaged every 50ms per electrode per trial to
 1091 create 20 time points (-200ms to 800ms) per electrode per trial. Fourth, the ERP data were entered
 1092 into a regression with trial by trial estimation of the drift rate using the HDDMRegressor from the
 1093 HDDM toolbox (66). The drift rate was allowed to vary per trial based on the ERP data per state
 1094 ('alert' or 'drowsy') per stimuli ('left' or 'right'). Fifth, the traces were computed per condition (state
 1095 and stimuli combination). Sixth, the differences in drift rate (between left and right stimuli) per
 1096 time point per electrode were computed in both alert and drowsy condition. This difference
 1097 represents the discriminability of the electrode in identifying the left and right stimuli at that time
 1098 point. Thus this analysis yielded differences in drift rate per electrode, per time point, per
 1099 condition, per participant. The differences can then be considered similar to the classifier patterns
 1100 and can be analysed both in electrode and source spaces. Further we also computed group level
 1101 differences using the cluster permutation techniques described earlier.

1102

1103 **Reproducing figures**

1104 The code for reproducing the main figures, raw datasets and other data accompanying this paper
 1105 will further be released upon acceptance or during the review of the paper.

1106

1107

1108 **AUTHOR CONTRIBUTIONS**

1109 Conceptualization: S.R.J and T.A.B.

1110 Methodology: S.R.J.

1111 Software: S.R.J.

1112 Investigation: S.R.J.

1113 Formal Analysis: S.R.J.

1114 Resources: S.R.J, C.A.B. and T.A.B.

1115 Data Curation: S.R.J and C.A.B.

1116 Writing – Original Draft: S.R.J.

1117 Writing – Review & Editing: S.R.J, C.A.B, and T.A.B.

1118 Visualization: S.R.J.

1119 Supervision: C.A.B. and T.A.B.

1120 Project Administration: S.R.J.

1121 Funding Acquisition: S.R.J and T.A.B.

1122

1123 ACKNOWLEDGMENTS

1124

1125 This research was funded by the Gates Cambridge Scholarship awarded to S.R.J and the Wellcome
1126 Trust Biomedical Research Fellowship WT093811MA awarded to T.A.B. as well as small
1127 departmental funding to S.R.J. and T.A.B. We thank Annamaria Laudini and Dritan Nikolla for
1128 assistance with data collection, Andrés Canales-Johnson, Will Harrison, Valdas Noreika and other
1129 members of the Consciousness and Cognition Lab in Cambridge for their valuable comments and
1130 support. We also thank lavazza for the continuous support with delicious and stimulating coffee.

1131

1132

SUPPLEMENTAL INFORMATION

Different stages of a cognitive task

1133

1134

1135

1136 **Figure s1:** Different stages of a cognitive task as identified by (2). In this example task, the stimulus period is
1137 represented by a green box and lasts from 0 till 180 ms and response occurs at approximately 400 ms. The
1138 different stages of processing are: a) perceptual stage (orange box) representing encoding of stimuli to
1139 decision making areas of the brain. b) central integration stage (red line) that represents accumulation of
1140 evidence towards a decision threshold. c) motor stage (violet box) represents the implementation of the
1141 decision from the previous stage. The distribution of reaction times for this example task are also plotted as
1142 histograms (yellow).

1143

1144

Subjective midline by sigmoid fits

1145

1146 **Figure s2:** Example fits of two different participants are shown. For each participant under each condition
 1147 ('alert', 'drowsy'), the proportion of rightward responses were fitted to the stimulus angle varying from -60°
 1148 to +60°. The 'glm' fit was constrained in such a way that the number of responses under each angle was used
 1149 as weights. The size of the dots represent the normalised (per condition) number of trials under each angle. A)
 1150 bias point (dotted line) shifts towards the left side as the participant becomes drowsy. In other words, as the
 1151 participant becomes drowsy they overestimate the right side of space. B) bias point (dotted line) shifts
 1152 towards the right side as the participant becomes drowsy. In other words, as the participant becomes drowsy
 1153 they overestimate the left side of space

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165 Posterior predictive fits from the best model

Posterior predictive distributions from the best model

1166

1167 **Figure s3:** Posterior predictive fits for the best model (#8) in hddm for some sample participants are shown.
 1168 The x-axis in each plot represents the reaction times in seconds, the positive values are times for a 'left'
 1169 button presses, whereas those on the negative x-axis are for 'right' button presses. The histograms (in red)
 1170 represent distributions of the observed data, whereas the blue lines represent model prediction. A) Posterior
 1171 predictive distributions when the participant is 'alert' and the stimulus presented was 'left' tone. B) Posterior
 1172 predictive distributions when the participant is 'alert' and the stimulus presented was 'right' tone. C)
 1173 Posterior predictive distributions when the participant is 'drowsy' and the stimulus presented was 'left' tone.

1174 D) Posterior predictive distributions when the participant is 'drowsy' and the stimulus presented was 'right'
1175 tone.

1176

1177 **Event related potential (ERP) values at significant electrodes from classifier patterns**

1178

ERP values at electrodes identified from clustering of classifier patterns

1179 **Figure s4:** spatio-temporal clustering of the classifier patterns across alert and drowsy periods reveals
1180 significant clusters at different time periods (from Figure 4). Here the significant electrodes are taken and
1181 their corresponding ERP values are plotted. The time course of the ERP values look similar to the time course
1182 of the classifier patterns itself.

1183

1184 **Classifier patterns for temporal decoding of responses**

1185 **Figure s5: Temporal decoding (responses):** A,B) Comparison of coefficients of classifier patterns in alert
1186 and drowsy periods. The patterns look fairly similar except the changes in intensity levels. The amplitude of
1187 the patterns are also several levels of magnitude higher (alert: 60 uV, drowsy: 30 uV) than temporal decoding
1188 of stimuli (both alert, drowsy: 2.4 uV Fig 3F,G).

1187 The classifier patterns for alert when response-locked (Figure s5), reveals scalp patterns related to
1188 motor implementation that seem similar to the later period of decoding when stimulus-locked (0.3
1189 onwards in Alert in Figure 3F,G), (-0.175 to -0.125 sec in Alert in Figure s5B). One possible
1190 interpretation is that these neural patterns may represent the same processes from 0.3 sec onwards
1191 in Figure 3F, motor implementation, as seen in the decoding for the response-locked. However, the
1192 classifier patterns for drowsy (Figure s5B) related to motor implementation, despite being spatial
1193 similar to stimulus-related decoding (0.42 to 0.46 sec in Drowsy in Figure 3G), (-0.175 to -0.125 sec
1194 in Drowsy in Figure s5B), these are separated by 300-700ms, with 0.42 sec drowsy decoding to
1195 actual reaction times (mean) only occurring at 1.17 sec. Hence, the decoding (of stimulus) under

drowsy conditions seems to start later, and this process is clearly different from motor related processes and shows an evidence accumulation related pattern that most likely develops later in a motor related process. This is different from the alert condition, wherein the classifier performance was initially driven by perceptual and central evidence accumulation processes (from 0.18 sec to 0.3 sec) and further by motor implementation related processes, possibly from 0.3 sec onwards.

Validation of source reconstruction

To verify the accuracy of the spatial localization procedure, we used the following methodology. First, we used EEG data from a sample participant (in the drowsy periods, as it has a larger number of trials) and computed noise covariance from 0.2 sec to 0 sec (pre stimulus period) using the 'shrunk' method in mne. Next the computed covariance was regularised. Second, the forward solution was computed using the transformation file, source space, beamformer solution as described in the methods section. Third, we computed the inverse operator using the noise covariance and the forward solution. Fourth, we applied the inverse operator on the raw data with parameters of $\text{snr} = 12$, $\text{lambda2} = 1.0 / \text{snr}^2$, $\text{method} = \text{'MNE'}$. For validation we used the 'MNE' method instead of 'dSPM' used in the decoding analysis, as we wanted to have a positive and negative amplitude in the source signals. Further we only picked the source signals whose orientations were normal to the surface. Fifth, we used labels from the left hemisphere in the source space and extracted signals in the regions of auditory, occipital, middle temporal, and frontal pole using the mode 'mean_flip'. Sixth, we extracted mean time courses inside each ROI and plotted the same in Figure s3. We identified N100 (peak negative signal 100 ms followed by stimulus presentation) only in the time course from the auditory region, whereas the other ROIs displayed weaker N100 amplitudes as shown below. This validates the accuracy of source localization as N100 is ERP amplitude is known to be localised in the auditory region.

source localization in left hemisphere rois in a sample participant

1221 **Figure s6:** source localization performed in a sample participant clearly reveals N100 presence in auditory
1222 regions at 100 ms after stimulus presentation. However occipital regions do not reveal the presence of N100
1223 thereby validating the source localization methods used.

1224

1225 REFERENCES

- 1226 1. Buchanan L, O'Connell A. A brief history of decision making. *Harv Bus Rev.* 2006
1227 Jan;84(1):32–41, 132.
- 1228 2. Sigman M, Dehaene S. Parsing a cognitive task: a characterization of the mind's bottleneck.
1229 *PLoS Biol.* 2005 Feb;3(2):e37.
- 1230 3. Sigman M, Dehaene S. Brain mechanisms of serial and parallel processing during dual-task
1231 performance. *J Neurosci.* 2008 Jul 23;28(30):7585–98.
- 1232 4. O'Connell RG, Shadlen MN, Wong-Lin K, Kelly SP. Bridging Neural and Computational
1233 Viewpoints on Perceptual Decision-Making. *Trends Neurosci.* 2018 Nov;41(11):838–52.
- 1234 5. Hull JT, Wright KP Jr, Czeisler CA. The influence of subjective alertness and motivation on
1235 human performance independent of circadian and homeostatic regulation. *J Biol Rhythms.* 2003
1236 Aug;18(4):329–38.
- 1237 6. Knowles JB. Sleep, Sleepiness and Performance Timothy H. Monk (Ed.) John Wiley & Sons, 1991
1238 [Internet]. Vol. 14, *Journal of Organizational Behavior.* 1993. p. 703–5. Available from:
1239 <http://dx.doi.org/10.1002/job.4030140710>
- 1240 7. Goupil L, Bekinschtein TA. Cognitive processing during the transition to sleep. *Arch Ital Biol.*
1241 2012 Jun;150(2-3):140–54.
- 1242 8. Link SW, Heath RA. A sequential theory of psychological discrimination. *Psychometrika.* 1975
1243 Mar;40(1):77–105.
- 1244 9. Gold JJ, Shadlen MN. The neural basis of decision making. *Annu Rev Neurosci.* 2007;30:535–74.
- 1245 10. Heekeren HR, Marrett S, Ungerleider LG. The neural systems that mediate human perceptual
1246 decision making. *Nat Rev Neurosci.* 2008 Jun;9(6):467–79.
- 1247 11. Heekeren HR, Marrett S, Bandettini PA, Ungerleider LG. A general mechanism for perceptual
1248 decision-making in the human brain. *Nature.* 2004 Oct 14;431(7010):859–62.
- 1249 12. Heekeren HR, Marrett S, Ruff DA, Bandettini PA, Ungerleider LG. Involvement of human left
1250 dorsolateral prefrontal cortex in perceptual decision making is independent of response
1251 modality. *Proc Natl Acad Sci U S A.* 2006 Jun 27;103(26):10023–8.
- 1252 13. Ho TC, Brown S, Serences JT. Domain general mechanisms of perceptual decision making in
1253 human cortex. *J Neurosci.* 2009 Jul 8;29(27):8675–87.

- 1254 14. Shadlen MN, Newsome WT. Neural Basis of a Perceptual Decision in the Parietal Cortex (Area
1255 LIP) of the Rhesus Monkey [Internet]. Vol. 86, Journal of Neurophysiology. 2001. p. 1916–36.
1256 Available from: <http://dx.doi.org/10.1152/jn.2001.86.4.1916>
- 1257 15. Shadlen MN, Newsome WT. Motion perception: seeing and deciding. Proc Natl Acad Sci U S A.
1258 1996 Jan 23;93(2):628–33.
- 1259 16. Roitman JD, Shadlen MN. Response of neurons in the lateral intraparietal area during a
1260 combined visual discrimination reaction time task. J Neurosci. 2002 Nov 1;22(21):9475–89.
- 1261 17. Bekinschtein T, Cologan V, Dahmen B, Golombek D. You are only coming through in waves:
wakefulness variability and assessment in patients with impaired consciousness [Internet].
1262 Progress in Brain Research. 2009. p. 171–89. Available from:
1263 [http://dx.doi.org/10.1016/s0079-6123\(09\)17712-9](http://dx.doi.org/10.1016/s0079-6123(09)17712-9)
- 1264 18. Canales-Johnson A, Beerendonk L, Blain S, Kitaoka S, Ezquerro-Nassar A, Nuiten S, et al.
1265 Decreased Alertness Reconfigures Cognitive Control Networks. J Neurosci. 2020 Sep
1266 9;40(37):7142–54.
- 1267 19. McGinley MJ, Vinck M, Reimer J, Batista-Brito R, Zagha E, Cadwell CR, et al. Waking State:
1268 Rapid Variations Modulate Neural and Behavioral Responses. Neuron. 2015 Sep
1269 23;87(6):1143–61.
- 1270 20. de Gee JW, Colizoli O, Kloosterman NA, Knapen T, Nieuwenhuis S, Donner TH. Dynamic
modulation of decision biases by brainstem arousal systems. Elife [Internet]. 2017 Apr 11;6.
1271 Available from: <http://dx.doi.org/10.7554/eLife.23232>
- 1272 21. van Kempen J, Loughnane GM, Newman DP, Kelly SP, Thiele A, O’Connell RG, et al. Behavioural
1273 and neural signatures of perceptual decision-making are modulated by pupil-linked arousal.
1274 Elife [Internet]. 2019 Mar 18;8. Available from: <http://dx.doi.org/10.7554/eLife.42541>
- 1275 22. Wang C-A, Baird T, Huang J, Coutinho JD, Brien DC, Munoz DP. Arousal Effects on Pupil Size,
1276 Heart Rate, and Skin Conductance in an Emotional Face Task [Internet]. Vol. 9, Frontiers in
1277 Neurology. 2018. Available from: <http://dx.doi.org/10.3389/fneur.2018.01029>
- 1278 23. Hori T, Hayashi M, Morikawa T. Topographical EEG changes and the hypnagogic experience
1279 [Internet]. Sleep onset: Normal and abnormal processes. 1994. p. 237–53. Available from:
1280 <http://dx.doi.org/10.1037/10166-014>
- 1281 24. Jagannathan SR, Ezquerro-Nassar A, Jachs B, Pustovaya OV, Bareham CA, Bekinschtein TA.
1282 Tracking wakefulness as it fades: Micro-measures of alertness. Neuroimage. 2018 Aug
1283 1;176:138–51.
- 1284 25. Bareham CA, Manly T, Pustovaya OV, Scott SK, Bekinschtein TA. Losing the left side of the
1285 world: rightward shift in human spatial attention with sleep onset. Sci Rep. 2014 May
1286 28;4:5092.
- 1287 26. Noreika V, Canales-Johnson A, Johnson A, Arnatkevičiūtė A, Koh J, Chennu S, et al.

- 1288 Wakefulness fluctuations elicit behavioural and neural reconfiguration of awareness [Internet].
1289 2020 [cited 2020 Sep 7]. p. 155705. Available from:
1290 <https://www.biorxiv.org/content/10.1101/155705v3>
- 1291 27. Comsa IM, Bekinschtein TA, Chennu S. Transient Topographical Dynamics of the
1292 Electroencephalogram Predict Brain Connectivity and Behavioural Responsiveness During
1293 Drowsiness. *Brain Topogr.* 2019 Mar;32(2):315–31.
- 1294 28. Ratcliff R, Smith PL, Brown SD, McKoon G. Diffusion Decision Model: Current Issues and
1295 History. *Trends Cogn Sci.* 2016 Apr;20(4):260–81.
- 1296 29. Kruschke JK. Bayesian Estimation Supersedes the t Test [Internet]. *PsycEXTRA Dataset.* 2012.
1297 Available from: <http://dx.doi.org/10.1037/e502412013-055>
- 1298 30. Fahrenfort JJ, van Driel J, van Gaal S, Olivers CNL. From ERPs to MVPA Using the Amsterdam
1299 Decoding and Modeling Toolbox (ADAM). *Front Neurosci.* 2018 Jul 3;12:368.
- 1300 31. Haufe S, Meinecke F, Görgen K, Dähne S, Haynes J-D, Blankertz B, et al. On the interpretation
1301 of weight vectors of linear models in multivariate neuroimaging. *Neuroimage.* 2014 Feb
1302 15;87:96–110.
- 1303 32. Yeo BTT, Thomas Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, et al. The
1304 organization of the human cerebral cortex estimated by intrinsic functional connectivity
1305 [Internet]. Vol. 106, *Journal of Neurophysiology.* 2011. p. 1125–65. Available from:
1306 <http://dx.doi.org/10.1152/jn.00338.2011>
- 1307 33. Corbetta M, Shulman GL. Spatial neglect and attention networks. *Annu Rev Neurosci.*
1308 2011;34:569–99.
- 1309 34. Fedorenko E, Duncan J, Kanwisher N. Broad domain generality in focal regions of frontal and
1310 parietal cortex. *Proc Natl Acad Sci U S A.* 2013 Oct 8;110(41):16616–21.
- 1311 35. Jewell G, McCourt ME. Pseudoneglect: a review and meta-analysis of performance factors in
1312 line bisection tasks. *Neuropsychologia.* 2000;38(1):93–110.
- 1313 36. Benwell CSY, Harvey M, Thut G. On the neural origin of pseudoneglect: EEG-correlates of shifts
1314 in line bisection performance with manipulation of line length. *Neuroimage.* 2014 Feb
1315 1;86:370–80.
- 1316 37. Robertson IH, Mattingley JB, Rorden C, Driver J. Phasic alerting of neglect patients overcomes
1317 their spatial deficit in visual awareness. *Nature.* 1998 Sep 10;395(6698):169–72.
- 1318 38. Deaton M, Tobias JS, Wilkinson RT. The Effect of Sleep Deprivation on Signal Detection
1319 Parameters [Internet]. Vol. 23, *Quarterly Journal of Experimental Psychology.* 1971. p. 449–52.
1320 Available from: <http://dx.doi.org/10.1080/14640747108400257>
- 1321 39. Smith PL, Ratcliff R. An integrated theory of attention and decision making in visual signal

- 1322 detection. Psychol Rev. 2009 Apr;116(2):283–317.
- 1323 40. Nunez MD, Vandekerckhove J, Srinivasan R. How attention influences perceptual decision
1324 making: Single-trial EEG correlates of drift-diffusion model parameters. J Math Psychol. 2017
1325 Feb;76(Pt B):117–30.
- 1326 41. O’Connell RG, Dockree PM, Kelly SP. A supramodal accumulation-to-bound signal that
1327 determines perceptual decisions in humans. Nat Neurosci. 2012 Dec;15(12):1729–35.
- 1328 42. Killgore WDS. Effects of sleep deprivation on cognition. Prog Brain Res. 2010;185:105–29.
- 1329 43. Kelly SP, O’Connell RG. The neural processes underlying perceptual decision making in
1330 humans: Recent progress and future directions. Journal of Physiology-Paris. 2015 Feb
1331 1;109(1):27–37.
- 1332 44. Twomey DM, Murphy PR, Kelly SP, O’Connell RG. The classic P300 encodes a
1333 build-to-threshold decision variable. Eur J Neurosci. 2015 Jul;42(1):1636–43.
- 1334 45. Loughnane GM, Newman DP, Bellgrove MA, Lalor EC, Kelly SP, O’Connell RG. Target Selection
1335 Signals Influence Perceptual Decisions by Modulating the Onset and Rate of Evidence
1336 Accumulation [Internet]. Vol. 26, Current Biology. 2016. p. 496–502. Available from:
1337 <http://dx.doi.org/10.1016/j.cub.2015.12.049>
- 1338 46. Shulman GL, Pope DLW, Astafiev SV, McAvoy MP, Snyder AZ, Corbetta M. Right hemisphere
1339 dominance during spatial selective attention and target detection occurs outside the dorsal
frontoparietal network. J Neurosci. 2010 Mar 10;30(10):3640–51.
- 1340 47. Dietz MJ, Friston KJ, Mattingley JB, Roepstorff A, Garrido MI. Effective connectivity reveals
1341 right-hemisphere dominance in audiospatial perception: implications for models of spatial
1342 neglect. J Neurosci. 2014 Apr 2;34(14):5003–11.
- 1343 48. Heilman KM, Bowers D, Valenstein E, Watson RT. Hemispace and Hemispatial Neglect. In:
1344 Jeannerod M, editor. Advances in Psychology. North-Holland; 1987. p. 115–50.
- 1345 49. Corbetta M, Kincade MJ, Lewis C, Snyder AZ, Sapir A. Neural basis and recovery of spatial
1346 attention deficits in spatial neglect. Nat Neurosci. 2005 Nov;8(11):1603–10.
- 1347 50. Duncan J. The multiple-demand (MD) system of the primate brain: mental programs for
1348 intelligent behaviour. Trends Cogn Sci. 2010 Apr;14(4):172–9.
- 1349 51. Wyart V, de Gardelle V, Scholl J, Summerfield C. Rhythmic fluctuations in evidence
1350 accumulation during decision making in the human brain. Neuron. 2012 Nov 21;76(4):847–58.
- 1351 52. Borbély AA, Daan S, Wirz-Justice A, Deboer T. The two-process model of sleep regulation: a
1352 reappraisal. J Sleep Res. 2016 Apr;25(2):131–43.
- 1353 53. Bekinschtein TA, Dehaene S, Rohaut B, Tadel F, Cohen L, Naccache L. Neural signature of the
1354 conscious processing of auditory regularities. Proc Natl Acad Sci U S A. 2009 Feb

- 1355 3;106(5):1672–7.
- 1356 54. Mitchell KJ. *Innate: How the Wiring of Our Brains Shapes Who We Are*. Princeton University
1357 Press; 2020. 312 p.
- 1358 55. Tagliazucchi E, Laufs H. Decoding wakefulness levels from typical fMRI resting-state data
1359 reveals reliable drifts between wakefulness and sleep. *Neuron*. 2014 May 7;82(3):695–708.
- 1360 56. Song C, Tagliazucchi E. Linking the nature and functions of sleep: insights from multimodal
imaging of the sleeping brain [Internet]. Vol. 15, *Current Opinion in Physiology*. 2020. p. 29–36.
1361 Available from: <http://dx.doi.org/10.1016/j.cophys.2019.11.012>
- 1362 57. Adolphs R. Human Lesion Studies in the 21st Century. *Neuron*. 2016 Jun 15;90(6):1151–3.
- 1363 58. Valero-Cabré A, Amengual JL, Stengel C, Pascual-Leone A, Coubard OA. Transcranial magnetic
stimulation in basic and clinical neuroscience: A comprehensive review of fundamental
principles and novel insights [Internet]. Vol. 83, *Neuroscience & Biobehavioral Reviews*. 2017.
1364 p. 381–404. Available from: <http://dx.doi.org/10.1016/j.neubiorev.2017.10.006>
- 1365 59. Yeung AWK, Tzvetkov NT, Atanasov AG. When Neuroscience Meets Pharmacology: A
1366 Neuropharmacology Literature Analysis [Internet]. Vol. 12, *Frontiers in Neuroscience*. 2018.
1367 Available from: <http://dx.doi.org/10.3389/fnins.2018.00852>
- 1368 60. Krakauer JW, Ghazanfar AA, Gomez-Marin A, MacIver MA, Poeppel D. Neuroscience Needs
1369 Behavior: Correcting a Reductionist Bias. *Neuron*. 2017 Feb 8;93(3):480–90.
- 1370 61. Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory.
1371 *Neuropsychologia*. 1971 Mar;9(1):97–113.
- 1372 62. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale.
Sleep. 1991 Dec;14(6):540–5.
- 1373 63. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG
1374 dynamics including independent component analysis. *J Neurosci Methods*. 2004 Mar
1375 15;134(1):9–21.
- 1376 64. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed Effects
1377 Models [Internet]. Vol. 82, *Journal of Statistical Software*. 2017. Available from:
1378 <http://dx.doi.org/10.18637/jss.v082.i13>
- 1379 65. Fox J, Weisberg S. *An R Companion to Applied Regression*. SAGE Publications; 2018. 608 p.
- 1380 66. Wiecki TV, Sofer I, Frank MJ. HDDM: Hierarchical Bayesian estimation of the Drift-Diffusion
1381 Model in Python. *Front Neuroinform*. 2013 Aug 2;7:14.
- 1382 67. Zhang J, Rittman T, Nombela C, Fois A, Coyle-Gilchrist I, Barker RA, et al. Different decision
1383 deficits impair response inhibition in progressive supranuclear palsy and Parkinson’s disease.
1384 *Brain*. 2016 Jan;139(Pt 1):161–73.

- 1385 68. Spiegelhalter DJ, Best NG, Carlin BP, van der Linde A. Bayesian measures of model complexity
1386 and fit [Internet]. Vol. 64, Journal of the Royal Statistical Society: Series B (Statistical
1387 Methodology). 2002. p. 583–639. Available from: <http://dx.doi.org/10.1111/1467-9868.00353>
- 1388 69. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. Bayesian Data Analysis, Third
1389 Edition. CRC Press; 2013. 675 p.
- 1390 70. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MEG and EEG
1391 data analysis with MNE-Python. Front Neurosci. 2013 Dec 26;7:267.
- 1392 71. Fischl B. FreeSurfer. Neuroimage. 2012 Aug 15;62(2):774–81.