1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

The calcium binding protein S100β marks hedgehog-responsive resident vascular stem cells within vascular lesions

[¶]Mariana Di Luca¹, [¶]Emma Fitzpatrick¹, [¶]Denise Burtenshaw¹, [¶]Weimin Liu², Jay-Christian Helt², Roya Hakimjavadi¹, Eoin Corcoran¹, Yusof Gusti¹, Daniel Sheridan¹, Susan Harman¹, Catriona Lally³, Eileen M. Redmond^{*2} and Paul A. Cahill^{*1}

¹ Dublin City University, Vascular Biology & Therapeutics Group, School of Biotechnology (Dublin, Ireland); ² University of Rochester, Dept Surgery (Rochester, New York, United States) ³Trinity College Dublin, Trinity Centre for Bioengineering, Trinity Biomedical Sciences Institute, (Dublin, Ireland).

[¶] Contributed equally

* Both senior authors

Address for correspondence

Paul A. Cahill., BSc., PhD
Professor, School of Biotechnology
Chair of Vascular Biology & Therapeutics, School of Biotechnology
Lonsdale Bldg, Office X2-09, Faculty of Science and Health
Dublin City University, Glasnevin, Dublin 9. Ireland
E: paul.cahill@dcu.ie, Ph: 353-1-700-8466, Fax: 353-1-700-5412

Abstract

A hallmark of subclinical atherosclerosis is the accumulation of vascular smooth muscle cell (SMC)-like cells leading to intimal thickening. While medial SMCs contribute, the participation of hedgehog responsive resident vascular stem cells (vSCs) to lesion formation remains unclear. Using transgenic eGFP mice and genetic lineage tracing of S100 β vSCs *in vivo*, we identified S100 β /Sca1 cells derived from a S100 β non-SMC parent population within lesions that co-localise with smooth muscle α -actin (SMA) cells following iatrogenic flow restriction, an effect attenuated following hedgehog inhibition with the smoothened inhibitor, cyclopamine. *In vitro*, S100 β /Sca1 cells isolated from atheroprone regions of the mouse aorta expressed hedgehog signalling components, acquired the di-methylation of histone 3 lysine 4 (H3K4me2) stable SMC epigenetic mark at the Myh11 locus and underwent myogenic differentiation in response to recombinant sonic hedgehog (SHh). Both S100 β and PTCH1 cells were present in human vessels while S100 β cells were enriched in arteriosclerotic lesions. Recombinant SHh promoted myogenic differentiation of human induced pluripotent stem cell-derived S100 β neuroectoderm progenitors *in vitro*. We conclude that hedgehog responsive S100 β vSCs contribute to lesion formation and support targeting hedgehog signalling to treat subclinical arteriosclerosis.

Keywords: Arteriosclerosis, intimal thickening, resident vascular stem cells, S100 β calcium binding proteins, stem cell antigen-1, lineage tracing, hedgehog signalling, human induced pluripotent stem cells (HiPSCs).

53 Introduction

54 Atherosclerosis is a chronic progressive inflammatory disease that leads to myocardial
55 infarction and stroke¹. Pathologic observations in humans vessels confirm that early
56 'transitional' lesions enriched with smooth muscle (SMC)-like cells called adaptive lesions are
57 routinely present in atherosclerotic-prone regions of arteries before pathologic intimal
58 thickening, lipid retention, and the appearance of a developed plaque². Endothelial cell (EC)
59 dysfunction due to disturbed blood flow is classically associated with the development of
60 atheroprone lesions³, while the embryological origin (e.g., neuroectoderm versus mesoderm)
61 of SMCs may also influence disease localisation and progression⁴. These early lesions are
62 routinely modelled in mouse carotid arteries following flow restriction⁵ and can further
63 rapidly develop into advanced atherosclerotic plaques in ApoE knockout on a Western diet^{6,7}.

64 Lineage tracing and single cell RNA sequence analysis (scRNA-seq) have provided compelling
65 evidence in various animal models for the involvement of a rare population of (i)
66 differentiated myosin heavy chain 11 (Myh11) medial SMC that are stem cell antigen-1
67 positive (Sca1⁺)^{8,9} and (ii) various adventitial and medial progenitors in progressing intimal
68 thickening¹⁰⁻¹³. While some adventitial progenitors are derived from a parent Myh11 SMC
69 population¹⁴, non-SMC derived SRY-related HMG-box 10 (Sox10) adventitial/medial resident
70 multipotent vascular stem cells (vSC), glioma-associated oncogene homolog 1 (Gli) and Nestin
71 progenitor cells have also been implicated^{13,15}. Despite these insights, the putative role of
72 resident vSCs that do not originate from medial Myh11⁺ SMCs in promoting intimal
73 thickening remains controversial.

74 S100 β , a member of the S100 multigenic family expressing small (9kDa and 14kDa) Ca²⁺-
75 binding proteins of the EF-hand type, is primarily localised to astrocytes and glial cells under
76 normal physiological conditions and is positively associated with age and various
77 neurodegenerative and neuroinflammatory diseases¹⁶. S100 β promotes intimal thickening
78 through interaction with the receptor for advanced glycation end-products (RAGE)¹⁷ and
79 induces migration and infiltration of inflammatory cells^{18,19}. S100 β plays a crucial role in
80 maintaining an intermediate state of Sca1 progenitor cells following injury-induced neointimal
81 thickening through a RAGE and stromal cell-derived factor-1 α /CXCR4 signalling mechanism
82²⁰.

Hedgehog signalling (Hh) is essential for normal embryonic development and plays a pivotal role in the maintenance of adult progenitor/stem cells in tissue repair after injury²¹. Hh signalling is initiated upon binding of a ligand to the canonical receptors, PTCH1 and PTCH2 (Patched 1 and 2)²², which leads to disinhibition of SMO (Smoothened) and activation of a complex signalling cascade that regulates the GLI (glioma-associated oncogene homolog 1) family zinc finger transcription factors (GLI1, GLI2, and GLI3)²³. GLI activators induce the transcription of Hh target genes primarily involved in cell proliferation, cell survival, and cell fate specification, including PTCH and GLI^{24,25}. HHIP (hedgehog interacting protein) modulates hedgehog signalling activity by binding and inhibiting the action of hedgehog proteins²⁶. Genome wide association studies (GWAS) in coronary artery disease patients have identified variants at the 14q32 HHIP1 (Hedgehog interacting protein-like 1) that encode a paralog of HHIP²⁷ to increase Hh signalling and regulate SMC proliferation and migration *in vitro*²⁸. In atherosclerotic mice, Hh has a permissive role through increased lipid uptake by macrophages²⁹, while global Hhpl1 deletion reduces plaque burden²⁸. Hh signalling, especially SHh and signal peptide CUB domain and EGF like domain containing 2 (Scube2), are overexpressed in injured arteries during lesion formation after flow restriction³⁰⁻³², autogenous vein grafts³³, and hypoxic pulmonary arterial SMCs³⁴. Adventitial Sca1⁺ cells that co-localise with SHh and Ptch1³⁵ significantly contribute to intimal thickening *in vivo*^{31,36,37} while Hh inhibition with cyclopamine or local perivascular depletion of Ptch1 attenuates intimal thickening following iatrogenic flow-restriction *in vivo*^{38,39}.

Given the pivotal role of Hh in the maintenance of adult progenitor/stem cells, tissue repair and atherosclerosis^{21,28}, we sought to determine the role of Hh responsive S100 β /Sca1 vSCs and their myogenic progeny in contributing to intimal thickening and the progression of adaptive lesions *in vivo* and, furthermore, whether Hh signalling directs murine S100 β vSCs and hiPSC-derived S100 β neuroectoderm stem cells to undergo proliferation and myogenic differentiation *in vitro*.

Results

Perivascular Sca1 and S100 β stem vSCs are present in normal adult vessels

The presence of perivascular Sca1 and S100 β vSCs was first confirmed in whole mounted vessels from Sca1-eGFP and S100 β -Cre-tdT mice. A discrete number of Sca1-eGFP and S100 β -tdT cells were present on collagen fibres within the adventitial layer of the thoracic aorta, when compared to wild-type (WT) controls [Suppl Figure 1a-d]. The cells were located predominantly at the aortic arch and at bifurcations with intercostal vessels, and their presence diminished along the descending and abdominal aorta [Suppl Figure 1b, c]. When the media was removed, Sca1-eGFP cells were clearly present in adventitial layer of the thoracic aorta [Suppl Figure 1d] and in perivascular regions of the carotid artery [Suppl Figure 1d].

Sca1 and S100 β cells are present in intimal lesions from murine carotid arteries following iatrogenic flow restriction.

The contribution of Sca1 cells to intimal medial thickening following iatrogenic flow restriction was first assessed in Sca1-eGFP transgenic mice following partial ligation of the left carotid artery (LCA). Vessels were harvested on day 3, 7 and 14 days post-ligation and paraformaldehyde fixed and paraffin-embedded cross sections were evaluated for eGFP (indicative of Sca1 expression) by confocal microscopy. Morphometric analysis confirmed that adventitial and intimal volumes increased significantly over time following ligation, without any significant change to medial volume, concomitant with a significant reduction in luminal volume after 7 and 14 days [Figure 1a].

The number of cells within the adventitial, medial and intimal layers of the LCA did not significantly change 3d post ligation [Figure 1b-d]. No Sca1-eGFP cells were visible within the medial layer of sham vessels which was populated primarily by α -smooth muscle actin (SMA) SMCs [Figure 1e]. While a small but significant number of Sca1-eGFP cells were present within the adventitial and intimal layers of sham vessels, the fraction of Sca1-eGFP cells significantly increased within the intimal layer 3d post ligation [Figure 1e, f]. Immunohistochemical analysis using anti-eGFP confirmed the presence of intimal Sca1-eGFP cells in ligated vessels [Figure 1e] and further demonstrated that intimal Sca1-GFP cells co-express PECAM-1/CD31 and endothelial cell nitric oxide (eNOS) [Figure 1e]. Moreover, there was also a notable

increase in the expression of the pro-apoptotic protein, Bax within the intimal, adventitial and medial layer of the LCA and the medial layer of the RCA 3d post ligation [Figure 1g,h], consistent with disturbed flow- and strain -induced apoptosis⁴⁰.

While the majority of Sca1-eGFP cells were present in the adventitial and intimal layers of sham operated vessels, a small fraction were also visible within the medial layer after 7, 14 and 21d [Figure 2a, b and g and Suppl Figure 2a, b]. Upon injury, there was a notable appearance of Sca1-eGFP cells within the expanding intimal layer of ligated vessels after 7, 14 and 21d [Figure 2a, b, d, g] that coincided with an increase in the number of cells [Figure 2c] and the volume of the neointimal layer [Figure 1a]. Cumulative analysis revealed that there was a significant increase in the fraction of Sca1-eGFP cells within neointimal layers after 7 and 14 and 21d, respectively, when compared to sham vessels or contralateral RCA [Figure 2d, h]. The majority of Sca1 cells within these lesions co-localised with SMA⁺ cells [Figure 2e, f, Suppl Figure 3a].

The contribution of S100 β vSC to intimal thickening following iatrogenic flow restriction was also assessed in S100 β -eGFP transgenic mice. While the majority of S100 β -eGFP cells were present in the adventitial layer of sham operated vessels and the contralateral RCA, a small fraction were also present within the medial layer after 21d [Suppl Figure 2a, c, Figure 2g, h]. In contrast, no intimal S100 β -eGFP cells were present in sham vessels or the contralateral RCA [Suppl Figure 2a c-e]. Upon injury, the fraction of S100 β -eGFP cells within lesions significantly increased when compared to sham vessels and the contralateral RCA [Suppl Figure 2d, e and Figure 2g,h]. Importantly, the number of double positive Sca1/S100 β cells increased within the neointimal layers of ligated vessels compared to the sham [Suppl Figure 2d, e] and contralateral RCA control [Figure 2g, h]. Similarly, the number of double positive S100 β /SMA cells increased within lesions compared to the contralateral RCA [Figure 2i, j]. Taken together, these data suggest double positive Sca1/S100 β cells populate vascular lesions following iatrogenic injury *in vivo*.

Lineage tracing analysis of marked perivascular S100 β cells iatrogenic flow restriction.

In order to determine the source of intimal double positive Sca1/S100 β cells following flow restriction, lineage tracing analysis was performed using transgenic S100 β -eGFP-CreERT2-Rosa-26-tdTomato reporter mice [Suppl Figure 4a]. As cre-mediated recombination results in

a permanent heritable change in a cell's genome, these cells and their progeny remain tdTomato (tdT) positive. The animals were treated with tamoxifen (Tm) for 7 days to induce nuclear translocation of CreER(Tm) and subsequent recombination to indelibly mark S100 β cells with red fluorescent tdT seven days before ligation was performed [Suppl Figure 4b]. Only S100 β cells present during the period of Tm treatment and their subsequent progeny following injury are marked with tdT. The tissue specificity and recombination efficiency of the Tm-induced Cre activity was confirmed in bone-marrow smears and neuronal tissue from S100 β -eGFP-CreERT2-Rosa-26-tdTomato mice. Treatment with Tm resulted in Cre-mediated recombination and expression of S100 β -tdT in >90% of neuronal cells of the carotid sinus nerve (Hering's nerve) without any S100 β -tdT cells observed in the bone marrow smears [Suppl Figure 4c]. The neuronal staining was confirmed by immunohistochemical analysis of S100 β expression using both anti-S100 β and anti-tdT antibodies (data not shown).

Morphometric analysis of the S100 β -eGFP-creER2-Rosa26-tdT mice confirmed significant intimal thickening in transgenic mice following ligation injury when compared to the sham LCA and contralateral RCA [Figure 3a-c]. Treatment of S100 β -eGFP-creER2-Rosa26-tdT transgenic mice with tamoxifen indelibly marked S100 β cells (S100 β -tdT) within the adventitial layer of the sham LCA prior to flow restriction [Figure 3d]. No cells are marked when these double transgenic mice were treated with the vehicle control (corn oil) [Figure 3d]. Importantly, no S100 β -tdT cells were observed in the intimal (EC) or medial (SMC) layers of these sham vessels following tamoxifen treatment [Figure 3d]. However, following ligation, there was a striking increase in the number of S100 β -tdT marked cells within the medial and intimal layers of the ligated LCA, compared to sham controls [Figure 3e, f]. In order to rule out the possibility that Tm treatment continues to label a significant number of cells for weeks after last Tm treatment⁴¹, we confirmed that the fraction of S100 β -tdT marked cells remained significantly higher in the neointima of LCA four weeks after the last Tm injection when compared to the RCA [Suppl Figure 4d, e]. Moreover, the fraction of S100 β -tdT marked cells within the LCA adventitial, medial and intimal layers of ligated vessels after one and four weeks from the last Tm injection was of a similar magnitude [Suppl Figure 4f].

When S100 β -tdT marked cells were evaluated for co-expression of Sca1 and SMA by immunohistochemistry using anti-Sca1, anti-SMA and anti-tdT antibodies, S100 β -tdT marked cells co-localised with Sca1 cells in the adventitial and medial layers (but not the intimal layer)

of the control contralateral RCA before injury [Suppl Figure 3d, Figure 3g]. There was a notable increase in the number of double stained Sca1/S100 β -tdT cells in the media and neointima of the LCA compared to the contralateral RCA [Figure 3g] and the sham LCA control [Suppl Figure 3b] in double stained SMA/S100 β -tdT cells within the neointima and media compared to the sham control [Suppl Figure 3c]. Cumulative analysis confirmed both a significant increase in the fraction of S100 β -tdT cells [Figure 3h] and the fraction of double stained S100 β /Sca1 cells [Figure 3i] within the LCA medial and intimal layers compared to the contralateral RCA control. Moreover, the majority of Sca1 cells within ligated vessels were S100 β -tdT cells [Figure 3j]. Since intimal and medial cells are not marked with tdT prior to injury, these data suggest that these cells are derived from a non-SMC S100 β parent population following flow restriction. Moreover, the majority of intimal Sca1 cells are derived from an S100 β parent population.

Hedgehog signalling components control intimal thickening following iatrogenic flow restriction

The effect of systemic administration of the Hedgehog inhibitor cyclopamine (10 mg/kg, IP, every other day) on neointimal thickening and the accumulation of Sca1-eGFP cells was evaluated. 2-hydroxypropyl- β -cyclodextrin (H β CD) was used as a vehicle control. Vessels were harvested 7 and 14 d post-ligation and morphometric analysis was performed. Whole vessel *Gli1* and *Gli2* mRNA levels were significantly elevated in ligated vessels treated with H β CD when compared to sham controls (*Gli1*: 6.1 ± 1.3 after 3 d; *Gli2*: 8.1 ± 0.9 fold after 14d, $p \leq 0.05$ $n=4$), an effect that was significantly reduced following cyclopamine treatment (*Gli2*: 3.2 ± 0.3 fold, $n=4$, $p \leq 0.05$). Immunohistochemical analysis revealed that the Hh receptor, Ptch1, was up-regulated in LCA following injury concomitant with enhanced *Gli2* expression in LCA when compared to sham vessels [Figure 4a]. *Gli2* co-localised with Sca1-eGFP expressing cells and was markedly increased in ligated vessels treated with vehicle (H β CD) when compared to sham-operated vessels, especially in the neointima and media of the LCA, an effect that was attenuated following Hh inhibition [Figure 4a]. Cyclopamine treatment significantly reduced the fraction of Sca1-eGFP expressing cells [Figure 4b, c] concomitant with reduced intimal thickening 7 d and 14 d-post ligation [Figure 4d-g]. Collectively, these data suggest that intimal thickening is attenuated by systemic administration of the

smoothened inhibitor, cyclopamine, concomitant with a reduction in the accumulation of Gli2/Sca1 Hh-responsive cells within the intimal and medial layers.

Isolation and characterisation of resident S100 β /Sca1 vascular stem cells from murine aorta.

The expression of SMC differentiation markers genes, *Myh11* and *Cnn1* was enriched in medial segments from both atheroprone aortic arch (AA) and atheros resistant thoracic/abdominal aorta (TA) regions of the mouse aorta, when compared to NE-4C cells and immortalised MOVAS mSMCs in culture [Figure 5a]. The level of S100 β and Sox10 expression, both markers of resident vascular stem cells^{12,42}, was then assessed in AA and TA regions, with or without the adventitia [Figure 5b, c]. There was significant enrichment of both S100 β and Sox10 in the AA when compared to the TA. Moreover, the levels were significantly enriched within the adventitia since its removal resulted in a dramatic reduction in the expression of both transcripts. Nevertheless, medial expression of both genes was still observed in the AA region [Figure 5b, c].

Resident vascular stem cells (vSCs) were subsequently isolated from mouse aortic arch (AA) and thoracic aorta (TA) by enzymatic digestion and sequential plating, and were characterised by flow cytometry, immunocytochemistry and gene expression for both stem cell markers and SMC differentiation markers. Flow cytometric analysis confirmed that the cells were Sca1 and S100 β positive when compared to Sca1 and S100 β positive control cell lines (i.e., C3H 10T1/2 and NE-4C, respectively) [Figure 5d]. The vSCs did not enrich for *Myh11* or *Cnn1* transcripts when compared to fresh aortic tissue [Figure 5e, f] or express these proteins [Figure 5g,h] but exhibited greater telomere length as a measure of stemness when compared to freshly isolated aortic SMC and to the immortalised MOVAS SMCs in culture [Figure 5i].

Immunocytochemical analysis revealed that vSCs from both AA and TA regions were Sca1 positive and expressed neuroectodermal markers S100 β , Sox10, Sox17, and Nestin but were negative for SMC differentiation markers *Cnn1* and *Myh11* [Figure 5g, h]. The vSCs were enriched for neuroectodermal marker genes *S100 β* , *Sox10* and *Nestin* [Suppl Figure 5a-c], but not for mesoderm markers *Kdr*, *Pax1* and *Tbx6* [Suppl Figure 5d-f], when compared to SMCs and NE-4 cells in culture. Collectively, these data indicate that neuroectodermal S100 β /Sca1

cells are primarily located within the adventitial perivascular region of atheroprone regions of the mouse aorta and can be isolated and expanded *in vitro* to retain these phenotypic markers.

Hedgehog signalling promotes myogenic differentiation of resident murine S100 β /Sca1 vascular stem cells *in vitro*.

The functional effect of sonic hedgehog (SHh) signalling on the growth and myogenic differentiation of vSCs was evaluated *in vitro* using multipotent Sca1/S100 β cells isolated from mouse aorta. Whole vessel Gli1 mRNA levels increased over time following ligation [Suppl Figure 6c]. Treatment of vSCs isolated from AA and TA with recombinant SHh (rSHh, 0.5 μ g/ml, 24h) promoted Hh signalling by increasing Hh target gene mRNA levels (*Gli 1*, *Gli 2* and *Ptch1*), an effect blocked following pre-treatment with cyclopamine [Figure 5j, k and Suppl Figure 6a, b]. Concurrent treatment of cells with rSHh for 7 days resulted in a significant increase in SMC differentiation marker gene expression (*Cnn1* and *Myh11*), an effect attenuated by pre-treatment of cells with cyclopamine [Figure 5j, k]. SHh treatment also significantly increased the growth of AA vSCs after 7 days, an effect attenuated by pre-treatment with cyclopamine [Suppl Figure 6d].

The level of enrichment of the SMC specific epigenetic histone mark, di-methylation of lysine 4 on histone H3 (H3K4me2) and the repressor mark, tri-methylation of lysine 27 on histone H3 (H3K27me3) at the *Myh11* locus was assessed using chromatin immunoprecipitation (ChIP) assays before and after rSHh treatment. Freshly isolated differentiated aortic medial SMC and murine embryonic stem cells (mESCs) were used as positive/negative controls for each mark and were markedly enriched for H3K4me2 and H3K27me3 at the *Myh11* promoter, respectively [Figure 5l-n]. In contrast to medial SMCs, AA vSCs were enriched for H3K27me3 but lacked enrichment of H3K4me2 at this locus. Treatment with recombinant SHh (rSHh, 0.5 μ g/ml, 7d) increased the enrichment levels of the H3K4me2 mark while concurrently decreasing the enrichment of the H3K27me3 mark, an effect attenuated by cyclopamine [Figure 5l-n]. Finally, SHh treatment significantly increased the fraction of Cnn1⁺ and Myh11⁺ cells in AA vSCs, an effect attenuated by pre-treatment with cyclopamine [Figure 5o,p].

S100 β cells populate human arteriosclerotic lesions

The expression of SMA, S100 β and the hedgehog receptor, PTCH1, was determined by immunofluorescence staining in normal and arteriosclerotic human aortic sections. The expression of SMA was abundant in the medial layers of both normal and arteriosclerotic vessels while reduced within the neointima [Figure 6a]. The expression of S100 β was confined to the adventitial layer of normal vessels but was significantly enhanced within the adventitial and intimal layers of arteriosclerotic sections [Figure 6a]. The expression of PTCH1 was confined to the adventitial layers of normal vessels but was also sparsely expressed in the medial and intimal layers of arteriosclerotic sections [Figure 6a].

Hedgehog signalling promotes myogenic differentiation of human induced pluripotent stem cell (HiPSC)-derived neuroectodermal S100 β vascular stem cells *in vitro*.

To address whether human neuroectoderm S100 β progenitors (NEPs) respond to hSHh by undergoing myogenic differentiation, S100 β NEPs were isolated from human induced pluripotent stem cells (HiPSC) using a chemically defined protocol⁴³. Human iPSCs enriched for Oct4 were treated with neuroectodermal inductive stimuli (FGF-2 and the selective inhibitor of ALK5, SB413642) for 7 days [Figure 6b] and expressed S100 β but not SMC differentiation markers CNN1 and MYH11 [Figure 6b]. Treatment of NEPs with myogenic media supplemented with TGF- β 1/PDGF-BB significantly increased the fraction of CNN1 and MYH11 cells after a further 12 d in culture, while concomitantly decreasing the number of S100 β cells [Figure 6c]. The levels of S100 β , CNN1 and MYH11 transcripts mirrored the changes in protein, where NEP-derived SMCs became significantly enriched for CNN1 and MYH11 following TGF- β 1/PDGF-BB treatment as they diminish their expression of S100 β [Figure 6d]. Treatment of NEPs with rSHh (1.0 μ g/ml) for 12 d resulted in a significant increase in CNN1 and MYH11 mRNA levels, an effect attenuated by cyclopamine [Figure 6e, f], concomitant with an increase in the number of CNN1 positive cells [Figure 6g]. SHh treatment also significantly increased the growth of S100 β NEPs after 12 days in culture, an effect attenuated by pre-treatment with cyclopamine [Suppl Figure 6e].

Discussion

In this study, we used transgenic mice and genetic lineage tracing approaches to delineate the involvement of Hh-responsive S100 β resident vSCs in progressing neointima formation following flow restriction. Our findings indicate that intimal thickening involves significant accumulation of Hh-responsive S100 β -derived progeny. Elucidation of this new cellular S100 β source for generating SMC-like cells following injury expands our current understanding of vascular pathology and may represent an important therapeutic strategy for combating subclinical atherosclerosis.

SMC-like cells within neointimal lesions reportedly are monoclonal/oligoclonal in origin arising from the expansion of a pre-existing resident clonal population of cells⁴⁴⁻⁴⁷. Despite extensive research, their origin has remained controversial⁴⁸. As a general rule, SMCs are purported to switch from a contractile phenotype to a less mature synthetic phenotype following vascular injury resulting in the loss of expression of SMC differentiation contractile markers, increased proliferation, and the synthesis and release of pro-inflammatory cytokines, chemotaxis-associated molecules, and growth factors⁴⁹. However, whether these clonal progeny within lesions are derived primarily from differentiated medial SMCs^{8,9,49}, endothelial cells^{10,50}, and/or resident vSCs¹⁰⁻¹³ remains unclear. Over the last decade, vascular SMC lineage tracing studies have provided compelling evidence of a significant role for pre-existing medial SMCs in vascular pathologies^{8,51-53}. Single cell RNA sequence analysis (scRNA-seq) further identified a rare Sca1⁺ Myh11-Cre marked SMC subpopulation that gives rise to functionally distinct neointimal cells during lesion progression in mice^{8,9}. However, although the Myh11-CreERT2 transgene was originally deemed specific for vascular SMC⁵⁴, data indicating expression of this and other 'SMC differentiation genes' in non-SMC populations has recently emerged⁵⁵⁻⁵⁷ raising the possibility that other resident cells may be inadvertently marked and contribute to lesion formation. Myh11-CreER SMC lineage tracing studies have also demonstrated that the fraction of SMCs originating from pre-existing SMCs may be considerably smaller after severe transmural injury in mice further suggesting a contributory role for cells other than medial SMCs⁵⁸.

In this study, we provide genetic evidence that vascular lesions contain an abundance of dual labelled Hh responsive S100 β /Sca1 vSCs derived from a non-SMC perivascular and/or medial S100 β parent population. Notwithstanding the fact that Sca1 cells are present in bone-

marrow⁵⁹, bone-marrow cells were not marked using the S100 β -CreERT2-tdT. It is therefore most likely that Sca1/S100 β cells marked with S100 β -CreERT2-tdT within lesions were locally derived⁴⁷. S100 β cells predominate within the adventitial layer of healthy vessels (murine and human) since removal of the adventitia dramatically reduced aortic S100 β transcript levels in mice. Nevertheless, a small fraction of S100 β /Sca1⁺ cells are present in medial layer of normal vessels and could be successfully isolated from aortic medial segments following enzymatic digestion and sequential plating *in vitro*. Indeed, previous studies using *in situ* immunohistochemical analysis and scRNAseq also confirms the presence of a sparse Sox10/S100 β cell population within the media, in addition to the adventitia^{9,60,61}. However, these enrichment levels were minimal compared to an S100 β ⁺ neuronal cell and vascular adventitial 'fibroblast-like cell' phenotypes within the vessel wall, supporting the contention that a perivascular adventitial S100 β cell may be the predominant source of S100 β within the vessel wall. Similarly, a discrete population of immature medial cells expressing CD146 that emerge during embryonic development may also contribute to neointimal formation in response to injury⁵⁸. Whether these CD146 immature SMC progenitors are also S100 β /Sca1 remains unclear although S100 β vSCs derived from rat and murine aortic explants express CD146^{12,42}. In the future, it will be important to determine whether medial S100 β vSCs differ from CD146⁵⁸ and Myh11-CreER Sca1 cells⁹ in their propensity to modulate different phenotypes and undergo clonal expansion using various injury models.

Adventitial progenitors have long been implicated in the progression of neointimal lesions following various vascular injuries^{66,67}. However, in the absence of Sca1 or c-kit⁺ stem cell lineage tracing experiments *in vivo*, the hypothesis of a regenerative function for adventitial Sca1 or c-kit⁺ vSCs in lesion formation has been difficult to prove until recently. While perivascular cells must first traverse the internal and external elastic lamina to accumulate within the intima following injury⁶⁴, disruption of the elastic lamellae and flow-dependent enlargement of fenestrae can facilitate such migration⁶⁵⁻⁶⁷. Cell fate mapping studies have now clarified the role for adventitial PDGFR α Sca1 progenitors in lesion progression following severe injury¹¹ while c-kit progenitors minimally facilitate endothelial regeneration or SMC differentiation to SMCs⁶⁸. As Sca1-CreER marked cells were not present in vascular lesions following wire-induced injury¹¹, the origin of Sca1 neointimal cells following flow restriction in the current study is unlikely to be from a Sca1 parent population despite expressing Sca1

(albeit using a different injury model). Indeed, the percentage of perivascular S100 β -CreER marked cells that were Sca1⁺ in normal vessels was < 50% indicating that a Sca1^{neg}/S100 β ^{pos} population might be responsible. In this context, S100 β has been shown to maintain an intermediate state of Sca1 progenitor cells following lesion formation²⁰. Alternatively, this residual Sca1^{pos}/S100 β ^{neg} population of adventitial Sca1 cells might represent Sca1 cells derived from differentiated Myh11-CreER SMCs that expand in number following injury and contribute to adventitial remodelling¹⁴ and lesion formation⁹. While Sca1/S100 β cells were predominantly found in the developing intimal layer after 7, 14 and 21 days, these cells were also present in the medial layer following injury. In agreement with previous studies demonstrating significant apoptosis of medial SMCs following flow restriction⁶⁹, the appearance of medial Sca1/S100 β cells marked with S100 β -CreERT2-tdT following early expression of the pro-apoptotic protein Bax within the media is consistent with regeneration of medial SMCs following injury.

Previous lineage tracing studies have also implicated endothelial cells (EC) that undergo endothelial-to-mesenchymal transition (EndMT) to SMC-like cells as contributing to neointimal formation during vein graft remodelling^{10,50}. EndMT promotes neointimal hyperplasia and induces atherogenic differentiation of EC⁷⁰. In the current study, EC were not marked with S100 β -CreERT2-tdT prior to flow restriction suggesting that the S100 β -CreERT2-tdT marked cells within lesions do not originate from an EC parent population. These apparent differences may result from the use of different models of vascular injury where vein graft and atherosclerosis models involve different biomechanical stimuli and/or a different inflammatory environments compared to the flow restriction model^{50,71}. While EC did not contribute to neointimal formation in developed lesions, a potential role for adventitial Sca1 cells in regenerating the endothelium following ligation was observed. As flow restriction leading to alterations in shear stress and cyclic strain is strongly associated with increased endothelial cell apoptosis *in vitro*⁷² and *in vivo*⁷³, significant Bax expression at the adventitial surface was observed after flow restriction. Bcl-2/Bax ratios are an important rheostat to determine the incidence of apoptosis in general and increases in Bax expression have been widely reported and validated using the TUNNEL assay to indicate apoptosis of endothelial cells *in vitro* in response to low shear stress (LSS)⁷⁴ and *in vivo* following carotid artery ligation in wildtype mice⁷⁵ or following flow perturbation in ApoE deficient mice⁷⁶. The increase in

Bax expression concomitant with an increase in the number of Sca1 cells that maintain a CD31 and eNOS endothelial cell phenotype at the adluminal surface is consistent with the appearance of new endothelial cells in the injured vessel that are derived from the proliferation of adluminal Sca1 cells. Indeed, recent cell fate mapping studies using Sca1-Cre-ERT2-tdT mice have since confirmed this observation by demonstrating expansion of Sca1-CreERT marked cells in the intimal layer that maintain CDH5 endothelial cell fate following injury¹¹.

There are many potential reasons for the different outcomes of lineage tracing studies that addressed the origin of lesional SMC-like cells including technical considerations around fixation, tissue preservation, brightness and contrast settings for image acquisition, autofluorescent material such as lipofuscin and single cell resolution⁷⁷. Another important reason may involve the use of tamoxifen (Tm) in time-sensitive Tm-dependent Cre-LoxP models. Tm-induced Cre recombinase has been shown to indelibly mark cells up to 4 weeks after the last Tm injection thereby potentially confounding the interpretation of many previous studies⁴¹. In this context, we found that the fraction of S100 β -CreER (tdT) marked cells within the various compartments of ligated vessels four weeks after the last Tm injection was similar to that observed after one week, indicating that, at least in our hands, medial SMC that might undergo de-differentiation and acquire the S100 β promoter are not being inadvertently marked after injury.

Several GWAS studies have implicated HHIP1 at the chromosome 14q32 CAD locus^{27,78} and identified it as a secreted Hh signaling protein that modulates atherosclerosis-relevant vascular phenotypes²⁸. Here we provide direct evidence that inhibition of Hh signalling controls neointimal thickening following flow restriction by attenuating the volume and accumulation of Sca1 cells within lesions. Furthermore, Hh signalling directly promotes Sca1/S100 β cells from mouse and S100 β neuroectoderm stem cells derived from human induced pluripotent stem cells (HiPSC) to undergo growth and myogenic differentiation to SMC-like cells *in vitro*. The possibility that Hh responsive S100 β /Sca1 cells isolated from murine AA and TA regions are derived from a Myh11 medial SMC population was ruled out by examining the H3K4me2 epigenetic profile of S100 β /Sca1 cells at the Myh11 locus, which is preserved during SMC phenotypic modulation *in vitro* and *in vivo* following injury⁷⁹. Hh responsive S100 β /Sca1 vSCs do not have this SMC histone mark and only become enriched

following myogenic differentiation to SMC-like cells, an effect attenuated by the Hh inhibitor, cyclopamine. The accumulation of Sca1/S100 β cells within lesions may be due, in part, to asymmetric division of Sca1/S100 β progenitor stem cells (from the adventitia and/or media) and subsequent proliferation and migration of their partially differentiated myogenic progeny to the intima. Indeed, SHh promotes Sca1/S100 β growth and the proliferation of modulated, partially differentiated SMCs *in vitro*³². Moreover, previous studies have shown that it is a Sca1 adventitial population in particular that proliferate following injury using immunohistochemical co-staining with Ki-67³¹. This is not surprising as adventitial Sca1 stem cells co-localise with SHh and its receptor, Ptch1³⁵ while several previous studies support a role for Hh signalling in vascular lesion formation in mice^{29,38,39}. It appears the Yes-Associated Protein (YAP) is required for incorporation and expansion of Sca1-derived neointimal cells following injury¹¹ where the Hh pathway reportedly acts upstream of the Hippo pathway to regulate stem cell maintenance⁸⁰. Indeed, our data suggest S100 β cells are also present in normal vessels within the adventitial layer where Hh components reside^{81,82}. In arteriosclerotic lesions, S100 β expression is exaggerated within the neointimal layer with sparse staining for PTCH1 present. As S100 β NEP-derived SMCs are considered crucial to the generation of SMC in vascular lesions within atheroprone regions of the murine vasculature^{4,43}, it is tempting to speculate that Hh responsive S100 β -derived SMC-like progeny may also play a pivotal role in lesion formation in humans. Future studies that address the interactions between Hippo pathway and Hh may yield therapeutic strategies aimed at interfering with YAP activity through inhibition of hedgehog pathway, which downregulates the YAP protein^{83,84}.

In conclusion, the identification of a population of S100 β /Sca1 cells within vascular lesions that are derived from a non-SMC S100 β parent population is novel and highlights these cells as an important source for intimal thickening following injury. Furthermore, the role of SHh in mediating this process expands our understanding of the mechanisms contributing to the progression of vascular lesions. While Hh signalling has previously been implicated in the development of vascular lesions through modulation of a differentiated SMC phenotype²⁸, our study further suggests that Hh signalling may also control myogenic differentiation and accumulation S100 β /Sca1 derived SMC-like progeny within vascular lesions. While hedgehog signalling appears critical, further research is required to investigate the precise mechanisms

473 driving migration, proliferation and differentiation of stem cells during lesion formation. It is
 474 possible that SMC-like cells within lesions comprise several different stages of the myogenic,
 475 osteogenic and myeloid differentiation of various stem cells and hence display a variety of
 476 divergent phenotypes. The elucidation of these mechanisms will provide vital information
 477 needed for the development of more efficient and novel therapies for vascular proliferative
 478 diseases. Our results provide a potential strategy for the treatment of atherosclerosis by
 479 combating cell migration and asymmetric differentiation of a perivascular-derived S100 β
 480 parent vSC population.

481

Methods

Mice Breeding and Genotyping

All procedures were approved by the University of Rochester Animal Care Committee in accordance with the guidelines of the National Institutes of Health for the Care and Use of Laboratory Animals. Sca1-eGFP transgenic mice were obtained from JAX labs; Stock #012643, strain name B6.Cg.Tg(Ly6a-EGFP)G5Dzk/j. These transgenic mice have an enhanced green fluorescent protein (eGFP) under the control of murine lymphocyte antigen 6 complex, locus A (Ly6a) promoter. Hemizygous Ly6a-GFP mice are viable, fertile, normal in size and do not display any gross physical or behavioural abnormalities⁸⁵.

S100 β -EGFP/Cre/ERT2 transgenic mice (JAX Labs, stock #014160, strain name B6;DBA-Tg(S100 β -EGFP/cre/ERT2)22Amc/j) express the eGFP^{CreER^{T2}} (Enhanced Green Fluorescent Protein and tamoxifen inducible cre recombinase/ESR1) fusion gene under the direction of the mouse S100 β promoter. Ai9 mice (Jax Labs, stock #007909, strain name B6.Cg-Gt(ROSA)26Sor^{tm9(CAG-tdTomato)Hze}/J) express robust tdTomato fluorescence following Cre-mediated LoxP recombination. For lineage tracing experiments S100 β -eGFP/Cre/ERT2–tdTomato double transgenic mice of both genders were generated by crossing S100 β -eGFP/Cre-ERT2 mice with Ai9 reporter mice. The tdTomato transgene expression pattern corresponds to genomic marked S100 β cells, and the eGFP transgene expression pattern corresponds to constitutive expression of S100 β . Mice were genotyped using genomic DNA prepared from tail samples. All male and female mice were included in the study and were 8–10 weeks old.

Tamoxifen-induced genetic lineage tracing

S100 β -CreER-Rosa26tdT mice (average weight 20g, 6–8 wks old) were injected IP with tamoxifen (Tm) dissolved in corn oil at 75 mg/kg for 5 consecutive days. Carotid artery ligation (partial, or complete), or sham operation, was performed at 1 wk or 4 wks after the last injection of Tm. At the indicated time post-ligation or sham operation, anesthetized mice were perfusion fixed and carotid arteries harvested for analysis.

Carotid Artery Ligation

Ligation of the left common carotid artery was performed essentially as described previously³⁸ one and four weeks after Tm-induced cre-recombination, in male and female S100 β -eGFP/Cre/ERT2-dTomato double transgenic mice. Prior to surgery mice received a single dose of Buprenorphine SR (sustained release) analgesia (0.5-1.0 mg/kg SQ) (and every 72 hrs thereafter as needed). The animal was clipped and the surgical site prepped using betadine solution and alcohol. A midline cervical incision was made. For partial carotid artery ligation, with the aid of a dissecting microscope, the left external and internal carotid arterial branches were isolated and ligated with 6-0 silk suture reducing left carotid blood flow to flow via the patent occipital artery. The neck incision (2 layers, muscle and skin) was sutured closed. Partial ligation of the left carotid artery in this manner resulted in a decrease (~80%) in blood flow, leaving an intact endothelial monolayer. Buprenorphine was administered at least once post-op 6-12 hrs. For complete ligation, the left common carotid was isolated and ligated just beneath the bifurcation with 6-0 silk suture. In sham surgery group, carotid was similarly manipulated but not ligated. The neck incision (2 layers, muscle and skin) was sutured closed and the animal allowed recover under observation. After 3, 7, 14 and 21 days, mice were perfused with 4% PFA via the left ventricle for 10 min, and carotid arteries were harvested for analysis.

Cyclopamine treatment

Sca1-eGFP mice were treated with the smoothened inhibitor, cyclopamine, or the vehicle 2-hydropropyl- β -cyclodextrin (H β CD) (Sigma-Aldrich) alone as a control, (essentially as described previously)⁸⁶. Mice were injected (250 μ l max volume) intraperitoneally (IP) 1 day before ligation, then every other day after ligation at a dose of 10 mg/kg Cyclopamine (Sigma-Aldrich) dissolved in a solution of 45% (w/v) H β CD. There was no effect of cyclopamine treatment on mouse body weight, compared to the H β CD vehicle control group (data not shown).

Histomorphometry

At specified times post-ligation, mice were anesthetized (ketamine/xylazine) and perfusion fixed with 4% paraformaldehyde in sodium phosphate buffer (pH 7.0). Fixed carotids were embedded in paraffin for sectioning. Starting at the carotid bifurcation landmark (single

lumen) a series of cross-sections (10 x 5 μ m) were made, every 200 μ m through 2 mm length of carotid artery. Cross-sections were de-paraffinized, rehydrated in graded alcohols and stained with Verhoeff-Van Gieson stain for elastic laminae and imaged using a Nikon TE300 microscope equipped with a Spot RT digital camera (Diagnostic Instruments). Digitized images were analyzed using SPOT Advanced imaging software. Assuming a circular structure *in vivo*, the circumference of the lumen was used to calculate the lumen area, the intimal area was defined by the luminal surface and internal elastic lamina (IEL), the medial area was defined by the IEL and external elastic lamina (EEL), and the adventitial area was the area between the EEL and the outer edge, essentially as described by us previously³⁸

Haematoxylin and Eosin Staining of Tissue

For human artery cross sections, slides were placed in ice-cold acetone and held at -20°C for 15 min. They were then washed 3 x PBS before immersion in Harris Haematoxylin at room temperature for 8 min. Slides were processed by sequential washing – in distilled water for 5 min; 1 % acid alcohol for 30 sec; 0.1 % sodium bicarbonate for 1 min and 95 % ethanol. Slides were then washed in Eosin for 1 min, 75 % ethanol for 3 min, 95 % ethanol for 3 min and 100 % ethanol for 3 min before being submerged in Histoclear for 3 min and permanently mounted using 3 drops of DPX, cleaned with 100 % ethanol before analysis.

For murine vessels, paraffin rehydration was conducted at room temperature by immersing the slides in xylene for 20 minutes. The slides were then immersed in the following gradients of ethanol for 5 minutes respectively: 100 %, 90 %, 70 % and 50 % ethanol. The slides were then rinsed in distilled H₂O₂ before washing in 1x PBS for 10 minutes. The slides were stored in 1x PBS until ready when they were immersed in Harris Haematoxylin at room temperature for 8 min.

S100 β ⁺ cells were identified as either “genomically marked” red tdTomato expressing cells or “currently expressing” green fluorescent protein (eGFP) cells visualized in deparaffinized S100 β -eGFP/Cre/ERT2-dTomato mouse carotid cross sections mounted with Sigma Fluoroshield with DAPI, using an FV1000 Olympus or a Nikon A1R HD laser scanning confocal microscope. Numbers of red or green fluorescent cells and DAPI nuclei (blue) in carotid cross section images from different experimental groups were either analyzed by Fiji ImageJ software in each vessel compartment using a grid system.

Whole mount histochemical analysis

Briefly, arteries harvested from transgenic mice were washed in PBS before transfer to 10 % neutral buffered formalin and incubated at 4 °C overnight. The tissue was rinsed 3 x PBS before mounting on histology slides using DAPI containing fluoroshield (Sigma). The whole tissue was then imaged using a Nikon Eclipse TS100 inverted microscope, EXFO X-Cite™ 120 fluorescence system and assessed using SPOT imaging software.

Immunofluorescent Staining of tissues.

Immunostaining essentially as previously described³⁸. Carotid artery cryosections were air-dried for 1 h at room temperature, followed by incubation with blocking buffer (5% donkey serum, 0.1% Triton X-100 in PBS) for 30 minutes at room temperature and then incubated with primary antibody overnight at 4°C in antibody solution (2.5% BSA, 0.3M Glycine and 1% Tween in DPBS). Murine and human arterial sections were stained with primary antibodies [Supplementary Table I]. Isotype IgG control and secondary antibody only controls were performed. For antigen retrieval, slides were brought to a boil in 10 mM sodium citrate (pH 6.0) then maintained at a sub-boiling temperature for 10 minutes. Slides were cooled on the bench-top for 30 minutes then washed in deionized water (3 x 5 min) each before being washed in PBS (3 x 5 min). The antigen retrieval protocol diminishes endogenous eGFP and tdT tomato transgene signals. Therefore, those sections were co-stained with anti-eGFP antibody and anti-Td tomato antibody [Supplementary Table I].

For immunofluorescence staining, 5 consecutive images were obtained and processed using ImageJ software™ to analyze the collected images. Mouse carotid artery cross sections were mounted with fluoroshield with DAPI and analyzed using an FV1000 Olympus or a Nikon A1R HD laser scanning confocal microscope. Images were merged using the Image color-merge channels function. Merged signals and split channels were used to delineate the signals at single-cell resolution. Settings were fixed at the beginning of both acquisition and analysis steps and were unchanged. Brightness and contrast were lightly adjusted after merging.

Immunocytofluorescent staining of cells

Cells seeded onto UV sterilized coverslips were fixed with 3.7 % formaldehyde, (15 min, RT). If cells required permeabilization for the detection of intracellular antigens, cells were

incubated in 0.025 % Triton X-100 PBS (room temp, 15 min). All coverslips were blocked (1 hr, RT) using 5 % BSA, 0.3 M Glycine, 1 % Tween PBS solution (1 hr, RT). Cells were incubated with primary antibodies overnight at 4°C [Supplemental Table II], then washed twice with PBS to remove any unbound primary antibody before being incubated (1 hr, RT) with the recommended concentration of fluorochrome-conjugated secondary antibodies diluted in blocking buffer [Supplementary Table II]. Following 2x wash in PBS, cell nuclei were stained using DAPI: PBS (dilution 1:1000) (15 min, RT). For each primary and secondary antibody used, a secondary control and an IgG isotype control was performed to assess nonspecific binding. An Olympus CK30 microscope and FCell™ software was used to capture images. Images were analyzed using ImageJ software as described above. Settings were fixed at the beginning of both acquisition and analysis steps and were unchanged. Brightness and contrast were lightly adjusted after merging.

Cell culture and harvesting

Murine aortic SMCs (MOVAS (ATCC® CRL-2797™) were cultured in Dulbecco's Modified Eagles growth medium (DMEM) supplemented with Fetal Bovine Serum (10%), L-glutamine (2 mM, Gibco) and Penicillin-Streptomycin (100U/mL, Gibco). Murine neuroectodermal stem cells (mNE-4Cs, ATCC® CRL-2925™) were grown in Eagle's Minimum Essential Medium supplemented with Fetal Bovine Serum (10%), L-glutamine (2 mM, Gibco) and Penicillin-Streptomycin (100U/mL, Gibco) using Poly-L-Lysine coated plates (15 µg/ml). Murine embryonic C3H 10T1/2 cells (ATCC® CRL-226™) were grown in Eagle's Basal medium supplemented with heat-inactivated fetal bovine serum to a final concentration of 10% and 2mM L-glutamine. Murine embryonic stem cells (mESCs) ES-D3 [D3] (ATCC® CRL-1934™) were maintained in culture in Mouse ES Cell Basal Medium (ATCC SCRR-2011) supplemented with 2-mercaptoethanol (0.1 mM) and 15% ES-Cell Qualified FBS (ATCC SCRR-30-2020) and a 1,000 U/mL mouse leukemia inhibitory factor (LIF) (Millipore Cat. No. ESG1107). Murine vSCs from AA and TA regions of the mouse aorta were grown in DMEM media supplemented with chick embryo extract (2%), FBS embryonic qualified (1%, ATCC), B-27 Supplement, serum free, N-2 Supplement (Cell Therapy Systems), recombinant mouse bFGF (0.02 µg/mL), 2-Mercaptoethanol (50nM), retinoic acid (100 nM), Penicillin-Streptomycin (100U/mL, Gibco) (MM).

Isolation of S100β/Sca1 murine resident vascular stem cells (vSCs)

Using an optimised enzymatic cell dissociation technique protocol vSCs from both atheroprone (aortic arch - neuroectodermal, NE), and athero-resistant (thoracic aorta-paraxial mesodermal, PM) regions of the murine aorta were isolated using sequential seeding on adherent and non-adherent plates. Mouse thoracic aortas (4 at a time) were harvested and placed in cold Hank's solution for adipose tissue removal and lumen rinsing. The adventitia was enzymatically removed by incubation of the vessels in collagenase solution of MEM α containing GlutaMAXTM (2 ml), collagenase type 1A (0.7 mg/ml), soybean trypsin inhibitor (50 mg/ml), and bovine serum albumin (1 mg/ml) for approx. 10-20 min at 37°C. Once the adventitia became loose, it was carefully removed as an intact layer using forceps under a dissecting microscope. The aorta was divided into aortic arch (AA) and thoracic aorta (TA) regions and cut into 1 mm pieces and digested with Elastase type III (Sigma) at 37°C. Dispersed cells were centrifuged and washed twice in warm maintenance medium (MM) (DMEM supplemented with 2% chick embryo extract, 1% FBS, 0.02 μ g/ml, bFGF basic protein, B-27 Supplement, N-2 supplement, 1% Penicillin-Streptomycin, 50 nM 2-Mercaptoethanol and 100 nM retinoic acid) before seeding (1st seeding) on a 6-well non-adherent plate in MM. At 48 h, suspension cells were transferred (2nd seeding) to a fresh well of a CELLstartTM pre-coated 6-well plate in MM. Additional MM was added to the remaining cells in the non-adherent surface. Cells were incubated at 37°C, 5% CO₂ for 1 week with minimal disturbance. The stem cells exhibited a uniform neural-like morphology in low density culture adopting a dendritic-like tree shape and retaining their morphological characteristics at low density throughout repeated passage. Cells were fed with MM every 2-3 days and passaged every 3-4 days or when ~70% confluent.

Generation of human neuroectoderm progenitors (NEPs) from induced pluripotent stem cells (HiPSCs)

Human induced pluripotent stem cells (HiPSC) were obtained from HipSci (Cambridge, UK) and cultured per the manufacturer's instructions. Briefly, human dermal fibroblasts were isolated from excess skin of patients undergoing plastic surgery and reprogrammed as previously described⁸⁷. HiPSCs were grown on plates pre-coated using Vitronectin (10 μ g/ml) and maintained in complete Essential eight medium replacing 95% of the medium daily. A ROCK inhibitor (10 μ M) was added to optimize colony growth. Cells were sub-cultured with 0.05 mM EDTA for 4-8 min at room temperature until colonies displayed bright halos around

the edges and small holes appeared throughout. Cells were frozen down in knock-out serum solution with 10% DMSO. HiPSCs at 70% confluency were routinely split at a ratio of 1:10 and cells were used between passage 20-40.

For neuroectoderm differentiation, cells were grown in chemically defined neuroectodermal medium (CDM) containing IMDM, F12 nutrient mix, chemically defined lipid concentrate, transferrin, monothioglycerol, penicillin-streptomycin and 0.5 g of PVA, and insulin, as previously described⁴³. Neuroectoderm differentiation was initiated in CDM supplemented with FGF2 (12 ng/ml) and SB431542 (10 μ M) for 7 days. Neuroectodermal (NE) progenitors were maintained in CDM-PVA supplemented with FGF2 (12 ng/ml) and SB431542 (10 μ M) and sub-cultured using TrypLE™ for 3–5 min at 37°C in a 5% CO₂ incubator. NE progenitors were used between passage 1-10.

To initiate myogenic SMC differentiation, NEPs were subcultured in SMC differentiation medium containing CDM + PDGF-BB (10 ng/ml, PeproTech) + TGF- β 1 (2 ng/ml, PeproTech) for 12 days. Neuroectoderm-derived SMCs (NE-SMC) were maintained in CDM supplemented with 10% FBS. After 12 d of PDGF-BB and TGF- β 1 treatment, all cells appeared spindle-shaped. Complete medium was then changed to DMEM 10% FBS, 1% Penicillin streptomycin for expansion. Cells were frozen down using 90% (volume/volume) FBS + 10% (vol/vol) DMSO. The NE-SMC were used between passage 3-20.

Human tissue

Tissue sections from paired normal human aorta and arteriosclerotic vessels were obtained from Amsbio LLC, tissue repository.

Chromatin Immunoprecipitation

ChIP was performed on cultured cells as previously described with slight modifications⁷⁹. Cells at 70% confluency were fixed with 1% paraformaldehyde (10 min, RT). Cross-link was stopped by addition of 125 mM glycine for 10 min. The cross-linked chromatin was sonicated to shear chromatin into fragments of 200–600 base pairs. The sheared chromatin was immunoprecipitated with 2 μ g tri-methyl-histone H3 (Lys27) and di-methyl-histone H3 (Lys4), while negative control was incubated with mouse IgG and input DNA without antibody using the CHIP—IT Express HT Kit from Active Motif (Cat no: 53018) according to the

manufacturer's instructions [Supplementary Table III]. A chromatin IP DNA Purification Kit (Cat No 58002 Active Motif) was used to purify the samples after CHIP before PCR was performed using Myh11 primers [5' -CCC TCC CTT TGC TAA ACA CA - 3' and 5' - CCA GAT CCT GGG TCC TTA CA – 3]. Sensimix SYBR® no-ROX Bioline Kit (QT650) was used to perform Real Time One-Step PCR according to the manufacturers' instructions. The antibodies used for ChIP are outlined in Supplemental Table I

Quantitative PCR

Total RNA was prepared from cultured cells using the ReliaPrep™ RNA Cell Miniprep System kit from Promega according to the manufacturer's protocol. Two micrograms of RNA was used for reverse transcription with Rotor-Gene SYBR Green RT-PCR (QIAGEN) or The SensiMix™ SYBR® No-ROX (BioLine) protocols for Real TimeOne-Step RT-PCR using the Real Time Rotor-GeneRG-3000™ light cyclers from Corbett Research using primers listed in Supplementary Table IV.

Flow cytometry and analysis

Isolated cells were incubated with I fluorochrome-conjugated antibodies and control IgG's or (ii) the appropriate primary antibody with fluorochrome-conjugated secondary antibodies at a final concentration of 1µg per sample and incubated at 4°C for 30 minutes [Supplementary Table V]. Fluorescence labeled cells analysis was performed using a BD FACS Aria flow cytometry system (BD Biosciences), and data were analyzed with FlowJo™ software (Tree Star, Ashland, Ore) and De Novo software FCS Express 4 Flow Cytometry (Pasadena, CA)

Telomere Assay

The average telomere length in cells was measured using a ScienCell's Absolute Mouse Telomere Length Quantification qPCR Assay Kit (AMTLQ) according to the manufacturer's instructions. The telomere primer set recognizes and amplifies telomere sequences and is designed to directly measure the average telomere length of murine cells.

QUANTIFICATION AND STATISTICAL ANALYSIS

All data were determined from multiple individual biological samples and presented as mean values ± standard error of the mean (SEM). All in vitro experiments were performed in

720 triplicate and repeated three times unless otherwise stated. All mice were randomly assigned
 721 to groups (including both male and female), and the analysis was performed blind by two
 722 groups. For statistical comparisons, all data was checked for normal gaussian distribution
 723 before parametric and non-parametric tests were performed. An unpaired two-sided
 724 Student's t-test was performed using GraphPad Prism software v9™ for comparing differences
 725 between two groups, and ANOVA test for over multiple comparisons. Significance was
 726 accepted when $p \leq 0.05$.

Data Availability:

The data, materials and reagents generated or analysed that support the findings of this study are available from the corresponding authors (PAC and EMR) on reasonable request.

Acknowledgments:

We thank Diana Scott for histological tissue processing, and Drs. Linda Callahan and Paivi Jordan for confocal microscope expertise. This research was funded in part by Science Foundation Ireland grant SFI-11/PI/1128, Health Research Board (HRB) of Ireland grant HRA-POR-2015-1315, the European Union's INTERREG VA Programme, managed by the Special EU Programmes Body (SEUPB) to PAC, and NIH R21AA020365 and R21AA023213 to EMR, Irish Research Council (IRC) GOIPG/2014/43 (M.DiL) and GOIPG/2016/1529 (E.C), King Abdulla Scholarship IR15208 ID1066336148 to Y.G.

Author Contributions:

E.F., W.L., J-C.H, performed the animal experiments and M.DiL and SH performed the blind analysis of the images. E.F performed the human arteriosclerotic tissue analysis. E.F., M.DiL., R.H., D.B., E.C., performed the murine and rat cell culture experiments, D.B performed the HiPSCs cell work. EC performed the telomere assay and R.H.J, MdiL and Y.G., performed the ChIP analysis. E.M.R., C.L., and P.A.C drafted the manuscript, revised the manuscript and discussed the results. E.M.R., and P.A.C., designed and coordinated the experiments, interpreted the data, revised and confirmed the paper. All authors reviewed and confirmed the manuscript.

Competing Interests:

The authors declare no financial and non-financial conflicts of interest.

References

1. Bennett, M. R., Sinha, S. & Owens, G. K. Vascular Smooth Muscle Cells in Atherosclerosis. *Circ. Res.* **118**, 692–702 (2016).
2. Sakamoto, A. *et al.* Pathologic intimal thickening: Are we any closer to understand early transitional plaques that lead to symptomatic disease? *Atherosclerosis* **274**, 227–229 (2018).
3. Halcox, J. P. J. *et al.* Endothelial Function Predicts Progression of Carotid Intima-Media Thickness. *Circulation* **119**, 1005–1012 (2009).
4. Sinha, S., Iyer, D. & Granata, A. Embryonic origins of human vascular smooth muscle cells: implications for in vitro modeling and clinical application. *Cell. Mol. Life Sci.* **71**, 2271–88 (2014).
5. Korshunov, V. A. & Berk, B. C. Flow-induced vascular remodeling in the mouse: a model for carotid intima-media thickening. *Arterioscler. Thromb. Vasc. Biol.* **23**, 2185–91 (2003).
6. Liu, W., Redmond, E. M., Morrow, D. & Cullen, J. P. Differential effects of daily-moderate versus weekend-binge alcohol consumption on atherosclerotic plaque development in mice. *Atherosclerosis* **219**, 448–54 (2011).
7. Nam, D. *et al.* Partial carotid ligation is a model of acutely induced disturbed flow, leading to rapid endothelial dysfunction and atherosclerosis. *Am. J. Physiol. Heart Circ. Physiol.* **297**, H1535–43 (2009).
8. Chappell, J. *et al.* Extensive Proliferation of a Subset of Differentiated, yet Plastic, Medial Vascular Smooth Muscle Cells Contributes to Neointimal Formation in Mouse Injury and Atherosclerosis Models. *Circ. Res.* **119**, 1313–1323 (2016).
9. Dobnikar, L. *et al.* Disease-relevant transcriptional signatures identified in individual smooth muscle cells from healthy mouse vessels. *Nat. Commun.* **9**, 4567 (2018).
10. Yuan, F. *et al.* Contribution of Vascular Cells to Neointimal Formation. *PLoS One* **12**, e0168914 (2017).
11. Tang, J. *et al.* Arterial Sca1+ Vascular Stem Cells Generate De Novo Smooth Muscle for Artery Repair and Regeneration. *Cell Stem Cell* **26**, 81–96.e4 (2020).
12. Tang, Z. *et al.* Differentiation of multipotent vascular stem cells contributes to vascular diseases. *Nat. Commun.* **3**, 875 (2012).
13. Kramann, R. *et al.* Adventitial MSC-like Cells Are Progenitors of Vascular Smooth Muscle Cells and Drive Vascular Calcification in Chronic Kidney Disease. *Cell Stem Cell* **19**, 628–642 (2016).
14. Majesky, M. W. *et al.* Differentiated Smooth Muscle Cells Generate a Subpopulation of Resident Vascular Progenitor Cells in the Adventitia Regulated by Klf4. *Circ. Res.* **120**, 296–311 (2017).
15. Wan, M. *et al.* Injury-activated transforming growth factor β controls mobilization of mesenchymal stem cells for tissue remodeling. *Stem Cells* **30**, 2498–511 (2012).
16. Donato, R. *et al.* S100B's double life: intracellular regulator and extracellular signal. *Biochim. Biophys. Acta* **1793**, 1008–22 (2009).
17. Cao, T. *et al.* S100B promotes injury-induced vascular remodeling through modulating smooth muscle phenotype. *Biochim. Biophys. Acta - Mol. Basis Dis.* **1863**, 2772–2782 (2017).
18. Sagheddu, R. *et al.* Targeting RAGE as a potential therapeutic approach to Duchenne muscular dystrophy. *Hum. Mol. Genet.* **27**, 3734–3746 (2018).
19. Zhou, S., Zhu, W., Zhang, Y., Pan, S. & Bao, J. S100B promotes microglia M1 polarization and migration to aggravate cerebral ischemia. *Inflamm. Res.* **67**, 937–949 (2018).

- 802 20. Wu, Y. *et al.* S100B is required for maintaining an intermediate state with double-
803 positive Sca-1+ progenitor and vascular smooth muscle cells during neointimal
804 formation. *Stem Cell Res. Ther.* **10**, 294 (2019).
- 805 21. Petrova, R. & Joyner, A. L. Roles for Hedgehog signaling in adult organ homeostasis and
806 repair. *Development (Cambridge)* vol. 141 3445–3457 (2014).
- 807 22. Marigo, V., Davey, R. A., Zuo, Y., Cunningham, J. M. & Tabin, C. J. Biochemical evidence
808 that patched is the hedgehog receptor. *Nature* **384**, 176–179 (1996).
- 809 23. Briscoe, J. & Théron, P. P. The mechanisms of Hedgehog signalling and its roles in
810 development and disease. *Nat. Rev. Mol. Cell Biol.* **14**, 416–429 (2013).
- 811 24. Marigo, V. & Tabin, C. J. Regulation of patched by Sonic hedgehog in the developing
812 neural tube. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 9346–9351 (1996).
- 813 25. Marigo, V., Johnson, R. L., Vortkamp, A. & Tabin, C. J. Sonic hedgehog differentially
814 regulates expression of GLI and GLI3 during limb development. *Dev. Biol.* **180**, 273–283
815 (1996).
- 816 26. Chuang, P. T. & McMahon, A. P. Vertebrate hedgehog signalling modulated by
817 induction of a hedgehog-binding protein. *Nature* **397**, 617–621 (1999).
- 818 27. Schunkert, H. *et al.* Large-scale association analysis identifies 13 new susceptibility loci
819 for coronary artery disease. *Nat. Genet.* **43**, 333–8 (2011).
- 820 28. Aravani, D. *et al.* *HHIPL1*, a Gene at the 14q32 Coronary Artery Disease Locus,
821 Positively Regulates Hedgehog Signaling and Promotes Atherosclerosis. *Circulation* **140**,
822 500–513 (2019).
- 823 29. Beckers, L. *et al.* Disruption of hedgehog signalling in ApoE^{-/-} mice reduces plasma
824 lipid levels, but increases atherosclerosis due to enhanced lipid uptake by
825 macrophages. *J. Pathol.* **212**, 420–8 (2007).
- 826 30. Ali, H. *et al.* Localization and characterization of a novel secreted protein, SCUBE2, in
827 the development and progression of atherosclerosis. *Kobe J. Med. Sci.* **59**, E122-31
828 (2013).
- 829 31. Dutzmann, J. *et al.* Sonic hedgehog-dependent activation of adventitial fibroblasts
830 promotes neointima formation. *Cardiovasc. Res.* **113**, 1653–1663 (2017).
- 831 32. Morrow, D. *et al.* Sonic Hedgehog induces Notch target gene expression in vascular
832 smooth muscle cells via VEGF-A. *Arterioscler. Thromb. Vasc. Biol.* **29**, 1112–8 (2009).
- 833 33. Li, F. *et al.* Sonic hedgehog signaling induces vascular smooth muscle cell proliferation
834 via induction of the G1 cyclin-retinoblastoma axis. *Arterioscler. Thromb. Vasc. Biol.* **30**,
835 1787–94 (2010).
- 836 34. Wang, G. *et al.* Activation of the sonic hedgehog signaling controls human pulmonary
837 arterial smooth muscle cell proliferation in response to hypoxia. *Biochim. Biophys. Acta*
838 **1803**, 1359–67 (2010).
- 839 35. Passman, J. N. *et al.* A sonic hedgehog signaling domain in the arterial adventitia
840 supports resident Sca1+ smooth muscle progenitor cells. *Proc. Natl. Acad. Sci. U. S. A.*
841 **105**, 9349–54 (2008).
- 842 36. Shikatani, E. A. *et al.* c-Myb Regulates Proliferation and Differentiation of Adventitial
843 Sca1+ Vascular Smooth Muscle Cell Progenitors by Transactivation of Myocardin.
844 *Arterioscler. Thromb. Vasc. Biol.* **36**, 1367–76 (2016).
- 845 37. Tsai, T.-N. *et al.* Contribution of stem cells to neointimal formation of decellularized
846 vessel grafts in a novel mouse model. *Am. J. Pathol.* **181**, 362–73 (2012).
- 847 38. Fitzpatrick, E. *et al.* Alcohol Reduces Arterial Remodeling by Inhibiting Sonic Hedgehog-
848 Stimulated Stem Cell Antigen-1 Positive Progenitor Stem Cell Expansion. *Alcohol. Clin.*
849 *Exp. Res.* **41**, 2051–2065 (2017).

- 850 39. Redmond, E. M. *et al.* Inhibition of patched-1 prevents injury-induced neointimal
851 hyperplasia. *Arterioscler. Thromb. Vasc. Biol.* **33**, 1960–4 (2013).
- 852 40. Fang, Y., Wu, D. & Birukov, K. G. Mechanosensing and mechanoregulation of
853 endothelial cell functions. *Compr. Physiol.* **9**, 873–904 (2019).
- 854 41. Reinert, R. B. *et al.* Tamoxifen-Induced Cre-loxP Recombination Is Prolonged in
855 Pancreatic Islets of Adult Mice. *PLoS One* **7**, e33529 (2012).
- 856 42. Kennedy, E. *et al.* Embryonic rat vascular smooth muscle cells revisited - a model for
857 neonatal, neointimal SMC or differentiated vascular stem cells? *Vasc. Cell* **6**, 6 (2014).
- 858 43. Cheung, C., Bernardo, A. S., Trotter, M. W. B., Pedersen, R. A. & Sinha, S. Generation of
859 human vascular smooth muscle subtypes provides insight into embryological origin-
860 dependent disease susceptibility. *Nat. Biotechnol.* **30**, 165–73 (2012).
- 861 44. Murry, C. E., Gipaya, C. T., Bartosek, T., Benditt, E. P. & Schwartz, S. M. Monoclonality
862 of smooth muscle cells in human atherosclerosis. *Am. J. Pathol.* **151**, 697–705 (1997).
- 863 45. Basatemur, G. L., Jørgensen, H. F., Clarke, M. C. H., Bennett, M. R. & Mallat, Z. Vascular
864 smooth muscle cells in atherosclerosis. *Nat. Rev. Cardiol.* **16**, 727–744 (2019).
- 865 46. Gomez, D. & Owens, G. K. Reconciling Smooth Muscle Cell Oligoclonality and
866 Proliferative Capacity in Experimental Atherosclerosis. *Circ. Res.* **119**, 1262–1264
867 (2016).
- 868 47. Hoglund, V. J., Dong, X. R. & Majesky, M. W. Neointima formation: a local affair.
869 *Arterioscler. Thromb. Vasc. Biol.* **30**, 1877–9 (2010).
- 870 48. Nguyen, A. T. *et al.* Smooth muscle cell plasticity: fact or fiction? *Circ. Res.* **112**, 17–22
871 (2013).
- 872 49. Campbell, J. H. & Campbell, G. R. Smooth Muscle Phenotypic Modulation—A Personal
873 Experience. *Arterioscler. Thromb. Vasc. Biol.* **32**, 1784–1789 (2012).
- 874 50. Cooley, B. C. *et al.* TGF- Signaling Mediates Endothelial-to-Mesenchymal Transition
875 (EndMT) During Vein Graft Remodeling. *Sci. Transl. Med.* **6**, 227ra34–227ra34 (2014).
- 876 51. Nemenoff, R. A. *et al.* SDF-1 α induction in mature smooth muscle cells by inactivation
877 of PTEN is a critical mediator of exacerbated injury-induced neointima formation.
878 *Arterioscler. Thromb. Vasc. Biol.* **31**, 1300–8 (2011).
- 879 52. Herring, B., Hoggatt, A. M., Burlak, C. & Offermanns, S. Previously differentiated medial
880 vascular smooth muscle cells contribute to neointima formation following vascular
881 injury. *Vasc. Cell* **6**, 21 (2014).
- 882 53. Shankman, L. S. *et al.* KLF4-dependent phenotypic modulation of smooth muscle cells
883 has a key role in atherosclerotic plaque pathogenesis. *Nat. Med.* **21**, 628–637 (2015).
- 884 54. Owens, G. K. Molecular control of vascular smooth muscle cell differentiation and
885 phenotypic plasticity. *Novartis Found. Symp.* **283**, 174–91; discussion 191–3, 238–41
886 (2007).
- 887 55. Chakraborty, R. *et al.* Promoters to Study Vascular Smooth Muscle. *Arterioscler.*
888 *Thromb. Vasc. Biol.* **39**, 603–612 (2019).
- 889 56. Murgai, M. *et al.* KLF4-dependent perivascular cell plasticity mediates pre-metastatic
890 niche formation and metastasis. *Nat. Med.* **23**, 1176–1190 (2017).
- 891 57. Sheikh, A. Q., Lighthouse, J. K. & Greif, D. M. Recapitulation of Developing Artery
892 Muscularization in Pulmonary Hypertension. *Cell Rep.* **6**, 809–817 (2014).
- 893 58. Roostalu, U. *et al.* Distinct Cellular Mechanisms Underlie Smooth Muscle Turnover in
894 Vascular Development and Repair. *Circ. Res.* **122**, 267–281 (2018).
- 895 59. Tobin, S. W. *et al.* Dual roles for bone marrow-derived Sca-1 cells in cardiac function.
896 *FASEB J.* **31**, 2905–2915 (2017).
- 897 60. Tang, Z., Wang, A., Wang, D. & Li, S. Smooth muscle cells: to be or not to be? Response

- 898 to Nguyen et Al. *Circ. Res.* **112**, 23–6 (2013).
- 899 61. Kalluri, A. S. *et al.* Single-Cell Analysis of the Normal Mouse Aorta Reveals Functionally
900 Distinct Endothelial Cell Populations. *Circulation* **140**, 147–163 (2019).
- 901 62. Psaltis, P. J. *et al.* Characterization of a resident population of adventitial macrophage
902 progenitor cells in postnatal vasculature. *Circ. Res.* **115**, 364–375 (2014).
- 903 63. Hu, Y. *et al.* Abundant progenitor cells in the adventitia contribute to atherosclerosis of
904 vein grafts in ApoE-deficient mice. *J. Clin. Invest.* **113**, 1258–65 (2004).
- 905 64. Wang, Y.-L., Liu, L.-Z., He, Z.-H., Ding, K.-H. & Xue, F. Phenotypic transformation and
906 migration of adventitial cells following angioplasty. *Exp. Ther. Med.* **4**, 26–32 (2012).
- 907 65. Campbell, G. J. & Roach, M. R. Fenestrations in the internal elastic lamina at
908 bifurcations of human cerebral arteries. *Stroke* **12**, 489–96.
- 909 66. Wong, L. C. Y. & Langille, B. L. Developmental Remodeling of the Internal Elastic Lamina
910 of Rabbit Arteries. *Circ. Res.* **78**, 799–805 (1996).
- 911 67. Guo, Z.-Y., Yan, Z.-Q., Bai, L., Zhang, M.-L. & Jiang, Z.-L. Flow shear stress affects
912 macromolecular accumulation through modulation of internal elastic lamina fenestrae.
913 *Clin. Biomech.* **23**, S104–S111 (2008).
- 914 68. Chen, Q. *et al.* Genetic lineage tracing analysis of c-kit⁺ stem/progenitor cells revealed
915 a contribution to vascular injury-induced neointimal lesions. *J. Mol. Cell. Cardiol.* **121**,
916 277–286 (2018).
- 917 69. Skelly, C. L., Chandiwal, A., Vosicky, J. E., Weichselbaum, R. R. & Roizman, B. Attenuated
918 herpes simplex virus 1 blocks arterial apoptosis and intimal hyperplasia induced by
919 balloon angioplasty and reduced blood flow. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 12474–
920 12478 (2007).
- 921 70. Moonen, J.-R. A. J. *et al.* Endothelial-to-mesenchymal transition contributes to fibro-
922 proliferative vascular disease and is modulated by fluid shear stress. *Cardiovasc. Res.*
923 **108**, 377–86 (2015).
- 924 71. Souilhol, C., Harmsen, M. C., Evans, P. C. & Krenning, G. Endothelial–mesenchymal
925 transition in atherosclerosis. *Cardiovasc. Res.* **114**, 565–577 (2018).
- 926 72. Pan, L. *et al.* Shear stress induces human aortic endothelial cell apoptosis via
927 interleukin-1 receptor-associated kinase 2-induced endoplasmic reticulum stress.
928 *Mol. Med. Rep.* **16**, 7205–7212 (2017).
- 929 73. Heo, K.-S. *et al.* PKC ζ mediates disturbed flow-induced endothelial apoptosis via p53
930 SUMOylation. *J. Cell Biol.* **193**, 867–884 (2011).
- 931 74. Dong, G. *et al.* Low shear stress-induced autophagy alleviates cell apoptosis in HUVECs.
932 *Mol. Med. Rep.* **15**, 3076–3082 (2017).
- 933 75. Zhang, Y. *et al.* Protective effect of irisin on atherosclerosis via suppressing oxidized low
934 density lipoprotein induced vascular inflammation and endothelial dysfunction. *PLoS*
935 *One* **11**, (2016).
- 936 76. Franck, G. *et al.* Flow Perturbation Mediates Neutrophil Recruitment and Potentiates
937 Endothelial Injury via TLR2 in Mice: Implications for Superficial Erosion. *Circ. Res.* **121**,
938 31–42 (2017).
- 939 77. Bentzon, J. F. & Majesky, M. W. Lineage tracking of origin and fate of smooth muscle
940 cells in atherosclerosis. *Cardiovasc. Res.* **114**, 492–500 (2018).
- 941 78. van der Harst, P. & Verweij, N. Identification of 64 Novel Genetic Loci Provides an
942 Expanded View on the Genetic Architecture of Coronary Artery Disease. *Circ. Res.* **122**,
943 433–443 (2018).
- 944 79. Gomez, D., Shankman, L. S., Nguyen, A. T. & Owens, G. K. Detection of histone
945 modifications at specific gene loci in single cells in histological sections. *Nat. Methods*

946 **10**, 171–177 (2013).
947 80. Fernandez-L, A. *et al.* YAP1 is amplified and up-regulated in hedgehog-associated
948 medulloblastomas and mediates Sonic hedgehog-driven neural precursor proliferation.
949 *Genes Dev.* **23**, 2729–41 (2009).
950 81. Queiroz, K. C. S. *et al.* Dichotomy in Hedgehog signaling between human healthy vessel
951 and atherosclerotic plaques. *Mol. Med.* **18**, 1122–7 (2012).
952 82. Dunaeva, M., van Oosterhoud, C. & Waltenberger, J. Expression of Hedgehog signaling
953 molecules in human atherosclerotic lesions: An autopsy study. *Int. J. Cardiol.* **201**, 462–
954 464 (2015).
955 83. Felley-Bosco, E. & Stahel, R. Hippo/YAP pathway for targeted therapy. *Transl. lung*
956 *cancer Res.* **3**, 75–83 (2014).
957 84. Cotton, J. L. *et al.* YAP/TAZ and Hedgehog Coordinate Growth and Patterning in
958 Gastrointestinal Mesenchyme. *Dev. Cell* **43**, 35–47.e4 (2017).
959 85. Ma, X., Robin, C., Ottersbach, K. & Dzierzak, E. The Ly6A Sca1 GFP Transgene is
960 Expressed in all Adult Mouse Hematopoietic Stem Cells. *Stem Cells* **20**, 514–521 (2002).
961 86. Van Den Brink, G. R. *et al.* Sonic hedgehog regulates gastric gland morphogenesis in
962 man and mouse. *Gastroenterology* **121**, 317–328 (2001).
963 87. Vallier, L. *et al.* Signaling Pathways Controlling Pluripotency and Early Cell Fate
964 Decisions of Human Induced Pluripotent Stem Cells. *Stem Cells* **27**, 2655–2666 (2009).
965

Figure Legends:

Figure 1. Expression of Sca1-eGFP and S100 β -eGFP cells following iatrogenic flow restriction after 3 days.

a. Morphometric analysis of adventitial, medial, intimal and luminal volumes following partial ligation of the left carotid artery compared to sham controls. Data are the mean $\pm$ SEM of 4 animals, #p<0.05 vs sham.

b-d. The number of DAPI nuclei/hpf in cross-sections of (b) adventitia, (c) media and (d) intima following partial ligation of the left carotid artery (LCA) after 3d compared to untouched (control) and sham. Data are the mean $\pm$ SEM of 3-5 representative images per animals, n=3, #p<0.05.

e. Verhoeff-Van Gieson stained sections and corresponding confocal fluorescence images of DAPI nuclei (blue), Sca1-eGFP (green) and immunofluorescence of α -actin (SMA) (red), anti-eGFP (green), anti-CD31 (red) and anti-eNOS (red) expression in ligated LCA vessels expression in sham-operated and ligated LCA vessels. Arrows highlight Sca1-eGFP expressing cells in the adventitia (A), media (M) and neointima (NI) and CD31 and eNOS in intima. Scale bar = 20 μ m.

f. The fraction of Sca1 cells/high power field (hpf) in the adventitia, media, and intima of sham and ligated vessels after 3 days. Data are the mean $\pm$ SEM of 3- 5 sections, #p $\leq$ 0.05 from a minimum of 4 animals per experimental group.

g. Representative immunofluorescence images for the pro-apoptotic marker, Bax, (red) in ligated left carotid artery (LCA) vessels and contralateral right carotid artery (RCA) after 3d. IgG was used as a control.

h. The number of Bax cells/hpf in the adventitia, media, and intima of sham and ligated vessels after 3d. Arrows highlight Bax expressing cells within the adventitia (A), media (M), and neointima (NI). Data are the mean $\pm$ SEM of 3-6 sections per experimental group from 3 animals. #p $\leq$ 0.05. Scale bar = 20 μ m.

Figure 2. Expression of Sca1-eGFP and S100 β -eGFP cells following iatrogenic flow restriction after 7, 14 and 21 days.

a,b. Representative Verhoeff-Van Gieson stained sections and corresponding confocal fluorescence images of DAPI nuclei (blue) and Sca1-eGFP (green) expression in LCA after **(a)** 7 days and **(b)** 14 days. Scale bar = 20 μ m.

c. The number of DAPI nuclei/hpf in cross-sections of adventitia, media and intima following partial ligation of the LCA after 7 and 14 days compared to sham controls. Data are the mean $\pm$ SEM of 3-5 representative images per animal, n=4.

d. The fraction of Sca1 cells/high power field (hpf) in the adventitia, media, and intima of sham and ligated vessels after 7 days and 14 days. Data are the mean $\pm$ SEM of 10-15 sections from 4 animals per experimental group.

e. Representative confocal fluorescence images of DAPI nuclei (blue), immunofluorescence of anti-eGFP (green) expression and anti-SMA (Red) in the LCA 14 days post-ligation in Sca1-eGFP mice. Data are representative of 5 images per animal.

f. Cumulative analysis of Sca1⁺ cells, SMA⁺ cells, double stained Sca1/SMA positive cells and the percentage of Sca1⁺ cells that were SMA⁺ in the adventitial, medial and intimal layer of RCA and LCA vessels after 21 days, respectively. Data are the mean $\pm$ SEM, n=15-20 sections from 5 animals.

g. Representative confocal fluorescence images of DAPI nuclei (blue), immunofluorescence of anti-eGFP (green) expression and anti-Sca1 (far-red white) in sham and ligated LCA after 21 days in S100 β -eGFP mice. Scale bar = 20 μ m.

h. Cumulative analysis of S100 β ⁺ cells, Sca1⁺ cells, double stained Sca1/S100 β positive cells and the percentage of S100 β ⁺ cells that were Sca1⁺ in the adventitial, medial and intimal layer of RCA and LCA vessels, respectively after 21 days in S100 β -eGFP mice. Data are the mean $\pm$ SEM, n=15-20 sections from 5 animals/group.

i. Representative confocal fluorescence images of DAPI nuclei (blue), immunofluorescence of anti-eGFP (green) expression and anti-SMA (far red, white) in the RCA and LCA 21 days post-

ligation in S100 β -eGFP mice. Data are from a minimum of 4 animals per experimental group.
Scale bar = 20 μ m.

j. Cumulative analysis of S100 β ⁺ cells, SMA⁺ cells, double stained S100 β /SMA positive cells and the percentage of S100 β ⁺ cells that were SMA⁺ in the adventitial, medial and intimal layer of RCA and LCA vessels, respectively after 21 days in S100 β -eGFP mice. Data are the mean $\pm$ SEM, n=15-20 sections from 4 animals/group.

Figure 3. Lineage tracing analysis of S100 β -Cre(ERT2)-tdT marked cells following iatrogenic flow restriction for 21 days.

a-c. Verhoeff-Van Gieson stained sections and morphometric analysis of adventitial, medial, intimal and luminal volumes of the **(b)** contralateral RCA control compared to the **(c)** LCA following ligation for 21d. Data are the mean $\pm$ SEM of 10 images, #p<0.05 vs sham controls.

d-f. Representative confocal fluorescence images of DAPI nuclei (blue) and S100 β -tdT (red) cells in mice treated with corn oil (control) or tamoxifen for 7 days prior to washout for 1 week in **(d)** sham and **(e, f)** ligated LCA. Data are minimum of 4 animals per experimental group. Scale bar = 20 μ m. Arrows highlight S100 β -tdT (red) marked cells.

g. Representative confocal fluorescence images of DAPI nuclei (blue), immunofluorescence of anti-tdT (red) and anti-Sca1 (far red, white) expression in the contralateral RCA and LCA 21 days post ligation. Scale bar = 20 μ m.

h-j. Cumulative analysis of the fraction of S100 β -tdT⁺ cells in the **(h)** RCA and **(i)** LCA and **(j)** Sca1/S100 β double stained cells in the adventitial, medial and intimal layer of RCA and LCA vessels, respectively after 21 days in S100 β -CreERT2-tdT mice.

Figure 4. Inhibition of Hh signalling attenuates intimal thickening and Sca1 cell expansion following iatrogenic flow restriction.

a. Representative immunofluorescence images of Hh receptor patched 1 (Ptch1) (red), Hh target gene Gli2 (red), Sca1-eGFP (green) expression and merged (eGFP/Gli2/DAPI) in carotid

artery cross sections from sham-operated, ligated vehicle control (H β CD), and ligated + cyclopamine (10 mg/kg, IP) treated mice after 14 days. Scale bar = top panel: 50 μ m, all other panels: 20 μ m.

b, c. Cumulative data showing the fraction of Sca1-eGFP cells within the adventitia, media and intima from sham-operated, ligated vehicle control (H β CD), and ligated + cyclopamine (Cyl) Sca1-eGFP mice after **(b)** 7 days and **(c)** 14 days. Data are the mean $\pm$ SEM of five sections per experimental group, #p $\leq$ 0.05 vs ligated H β CD controls and are representative of a minimum of 4 animals per experimental group.

d, e. Representative H&E staining of carotid artery cross sections from sham, ligated control (H β CD vehicle), and ligated + cyclopamine (Cyl) treated animals after 7 d **(d)** and **(e)** 14 d. Scale bar = 50 μ m.

f,g. Morphometric analysis of vessel compartment volumes for ligated control (H β CD), and ligated + cyclopamine (Cyl) treated animals after **(f)** 7d and **(g)** 14 d. Data mean $\pm$ SEM of 7-12 sections analysed from 4 animals, #p $\leq$ 0.05 vs ligated control (H β CD).

Figure 5. Resident S100 β /Sca1 vascular stem cells from atheroprone and atheroresistant regions of the mouse aorta *in vitro*.

a. Relative levels of *Myh11* and *Cnn1* in AA and TA regions of the mouse aorta. Data are expressed as the Log2 fold change in mRNA levels relative to neural stem cells (NE-4C) in culture and are the mean $\pm$ SEM of three aortic specimens, #p $\leq$ 0.05 vs NE-4C cells. The housekeeping gene *Gapdh* was used as a control.

b, c. The level of enrichment of mRNA for neuroectodermal markers **(b)** S100 β and **(c)** Sox10 within atheroprone AA (aortic arch) and atheroresistant TA (thoracic/descending aorta) regions of the mouse aorta in the absence or presence of the adventitial (Adv) layer. The housekeeping gene *hypoxanthine guanine phosphoribosyltransferase (hprt)* was used as a control. Data are expressed as the Log2 fold change in mRNA levels relative to murine aortic SMCs in culture and are the mean $\pm$ SEM of three aortic specimens, #p $\leq$ 0.05 vs MOVAS SMCs.

d. Flow cytometry analysis of vSCs isolated from AA and TA regions of the mouse aorta and grown in maintenance media for S100 β (dark green) and Sca1 (light green) expression. NE-4C and CH3-10T1/2 cells were used as positive controls, respectively. Filled dark and light green curves represent negative control samples.

e,f. Relative levels of **(e)** *Myh11* and **(f)** *Cnn1* in vSCs isolated from AA and TA regions of the mouse aorta, compared to AA and TA aortic tissue. Data are expressed as the Log2 fold change in mRNA levels relative to neural stem cells (NE-4C) in culture and are the mean $\pm$ SEM of three independent cultures and three aortic specimens, #p $\leq$ 0.05 vs NE-4C cells.

g,h. Representative immunocytochemical analysis of stem cell markers S100 β , Nestin, Sox10, Sox17, Sca1 and SMC differentiation markers, *Cnn1* and *Myh11* in vSCs isolated and grown in maintenance media (MM) from **(g)** AA and **(h)** TA regions of the mouse aorta. Scale bar 20 μ m.

i. Absolute telomere length of vSCs compared to cultured MOVAS SMC and fresh aortic tissue. The telomere primer set recognizes and amplifies telomere sequences. The single copy reference (SCR) primer set recognizes and amplifies a 100 bp-long region on mouse chromosome 10, and serves as reference for data normalization. Data are expressed as absolute telomere length (kb) and are the mean $\pm$ SEM of three plates and aortic specimens, #p $\leq$ 0.05 vs MOVAS SMCs.

j,k. Relative levels of *Gli1*, *Myh11* and *Cnn1* within vSCs isolated from **(j)** AA and **(k)** TA regions of the mouse aorta in the absence or presence of rSHh (0.5 μ g/ml) with or without the smoothened inhibitor, cyclopamine (10 μ M). Data are expressed as the Log2 fold change in mRNA levels relative to vSCs alone (control) and are the mean $\pm$ SEM of three representative wells from two independent experiments, #p $\leq$ 0.05 versus rSHh alone.

l-m. ChiP analysis of the ratio of **(l)** H3K4me2:H3K27me3 and **(m)** H3K27me3:H3K4me2 enrichment at the *Myh11* locus in AA vSCs in the absence or presence of rSHh (0.5 μ g/ml) with or without cyclopamine (10 μ M) for 7 d.

n. Fold enrichment of H3K4me2 at the *Myh11* locus in AA vSCs in the absence or presence of rSHh (0.5 μ g/ml) with or without cyclopamine (15 μ M) for 7 d. Fresh aortic tissue and mouse ECSs was used as positive and negative controls, respectively Data are the mean $\pm$ SEM, n=3, #p $\leq$ 0.05 versus aorta/ESC, *p $\leq$ 0.05 vs vSC, §p $\leq$ 0.05 vs rSHh.

o. Representative immunocytofluorescence staining and **p.** the fraction of Cnn1 and Myh11 positive cells in the absence or presence of rSHh (0.5 µg/ml) with or without cyclopamine (15µM) for 7 days. Representative images shown, Scale bar = 50 µm. Cumulative data is the mean ± SEM, n=3, #p≤ 0.05 versus control, *p≤ 0.05 vs rSHh.

Figure 6. Human neuroectoderm progenitors express S100β cells and are present in vascular lesions.

a. Representative phase contrast photomicrographs, hematoxylin and eosin (H&E) staining, DAPI staining (blue) and immunocytochemical staining for smooth muscle cell α-actin (SMA) (red), S100β (red) and PTCH1 as indicated in the adventitial (A), medial (M), and intimal (I) sections of normal human and arteriosclerotic arteries. Scale bar=50µm. Data are representative of 3 sections from 3 patient cohorts.

b. Phase contrast images, fluorescence staining of DAPI nuclei (blue) and immunocytochemical analysis of the expression of OCT4 (green), S100β (green) in NEPs and CNN1 (green) and MYH11 (green) in NEP-derived SMCs. All cells were generated as described previously⁴³. Scale bar = Right panels: 50µm, Left panels: 20µm. Data are representative of five images per experimental group from three independent cultures.

c. The fraction of S100β, CNN1⁺ and MYH11⁺ cells in HiPSCs, NEPs and NEP-SMCs. Data are mean ± SEM, n=3, #p≤ 0.05 versus HiPSCs.

d. Relative levels of S100β, CNN1 and MYH11 in HiPSCs, NEPs and NEP-SMCs, respectively. Data are expressed as the Log2 fold change in mRNA levels relative to HiPSCs in culture and are the mean ± SEM, n=3, #p≤ 0.05 versus HiPSCs.

e,f. Relative levels of **(e)** CNN1 and **(f)** MYH11 in NEPs in the absence or presence of hSHh (0.5µg/ml) with or without cyclopamine (10µM) for 12 d. Data are representative of three independent cultures, #p≤ 0.05 versus NEPs, *p≤ 0.05 vs hSHh.

g. Immunocytochemical analysis of CNN1 (green) expression and DAPI (blue) staining in NEPs in the absence or presence of recombinant hSHh (0.5 µg/ml) for 12 d. Data are representative of five images per experimental group from two independent cultures.