1 **PilB from *Streptococcus sanguinis* is a bimodular type IV pilin with a direct**
2 **role in adhesion**

3

4 Claire Raynaud^{1,¶}, Devon Sheppard^{1,¶}, Jamie-Lee Berry¹, Ishwori Gurung¹, Vladimir
5 Pelicic^{1,2,*}

6

7 ¹MRC Centre for Molecular Bacteriology and Infection, Imperial College London,
8 London, United Kingdom

9

10 ²Laboratoire de Chimie Bactérienne, Aix-Marseille Université-CNRS (UMR 7283),
11 Institut de Microbiologie de la Méditerranée, Marseille, France

12

13 [¶]These authors contributed equally

14

15 *Corresponding author

16 E-mail: vladimir.pelicic@inserm.fr

17 **ABSTRACT**

18 Type IV pili (T4P) are functionally versatile filamentous nanomachines, nearly
 19 ubiquitous in prokaryotes. They are predominantly polymers of one major pilin, but
 20 also contain minor pilins whose functions are often poorly defined, and likely to be
 21 diverse. Here, we show that the minor pilin PilB from the T4P of *S. sanguinis* displays
 22 an unusual bimodular 3D structure, with a bulky von Willebrand factor A-like (vWA)
 23 module "grafted" onto a small pilin module via a short unstructured loop. Structural
 24 modelling suggests that PilB is only compatible with a localisation at the tip of T4P.
 25 By performing a detailed functional analysis, we found that (i) the vWA module
 26 contains a canonical metal ion-dependent adhesion site (MIDAS), preferentially
 27 binding Mg^{2+} and Mn^{2+} , (ii) abolishing metal-binding has no impact on the structure of
 28 PilB or piliation, (iii) metal-binding is important for *S. sanguinis* T4P-mediated
 29 twitching motility and adhesion to eukaryotic cells, and (iv) the vWA module shows
 30 an intrinsic binding ability to several host proteins. These findings reveal an elegant,
 31 yet simple, evolutionary tinkering strategy to increase T4P functional versatility, by
 32 grafting an adhesive module onto a pilin for presentation by the filaments. This
 33 strategy appears to have been extensively used by bacteria, in which modular pilins
 34 are widespread and exhibit an astonishing variety of architectures.

35 INTRODUCTION

36 Type IV pili (T4P) are functionally versatile filaments widespread in prokaryotes,
 37 implicated in a variety of functions such as adhesion, twitching motility, DNA uptake
 38 *etc*¹. T4P are helical polymers consisting of type IV pilins, usually one major pilin and
 39 several minor (low abundance) ones, assembled by distinctive multi-protein
 40 machineries. These defining features are shared by a superfamily of filamentous
 41 nanomachines known as type IV filaments (T4F)¹, which are ubiquitous in
 42 prokaryotes².

43 T4P have been intensively studied for decades in diderm bacteria because
 44 they play a central role in pathogenesis in several important human pathogens³. The
 45 following global picture of T4P biology has emerged from these studies. The pilus
 46 subunits, type IV pilins, are characterised by a short N-terminal sequence motif
 47 known as class III signal peptide, which consists of a hydrophilic leader peptide
 48 ending with a small residue (Gly or Ala), followed by a tract of 21 predominantly
 49 hydrophobic residues⁴. This tract constitutes the N-terminal segment (α 1N) of an α -
 50 helix (α 1) of ~50 residues, which is the universally conserved structural feature in
 51 type IV pilins. Usually, the α 1N helix protrudes from a globular head most often
 52 consisting of a β -sheet composed of several antiparallel β -strands, which gives pilins
 53 their characteristic "lollipop" shape⁴. The hydrophilic leader peptide is then processed
 54 by a dedicated prepilin peptidase⁵ upon pilin translocation across the cytoplasmic
 55 membrane (CM) by the general secretory pathway^{6,7}. Processed pilins remain
 56 embedded in the CM via their α 1N, generating a pool of subunits ready for
 57 polymerisation. Filament assembly, which occurs from tip to base, is mediated at the
 58 CM by a complex multi-protein machinery (10-20 components)¹, centred on an
 59 integral membrane platform protein and a cytoplasmic extension ATPase⁸. Recent
 60 cryo-EM structures have revealed that T4P are right-handed helical polymers in
 61 which pilins are held together by extensive interactions between their α 1N helices,
 62 which are partially melted and run approximately parallel to each other within the

63 filament core^{9,10}. One of the properties of T4P key for their functional versatility is
 64 their ability to retract, which has been best characterised for T4aP (where "a"
 65 denotes the subtype). In T4aP, retraction results from rapid filament depolymerisation
 66 powered by the cytoplasmic retraction ATPase PilT¹¹, which generates important
 67 tensile forces^{12,13}.

68 Studying T4P in monoderm bacteria represents a promising alternative
 69 research avenue¹⁴. *Streptococcus sanguinis*, a commensal of the oral cavity that
 70 commonly causes life-threatening infective endocarditis (IE), has emerged as a
 71 monoderm model for deciphering T4P biology¹⁵. Our comprehensive functional
 72 analysis of *S. sanguinis* T4P¹⁶ revealed that they are canonical T4aP. Indeed,
 73 filaments are (i) assembled by a multi-protein machinery similar to diderm T4aP
 74 species, but simpler with only ten components, (ii) retracted by a PilT ATPase,
 75 generating tensile forces similar to diderm species, and (iii) powering intense
 76 twitching motility, leading to spreading zones around bacteria growing on plates,
 77 visible by the naked eye. Subsequently, we performed a global biochemical and
 78 structural analysis of *S. sanguinis* T4P¹⁷, showing that (i) they are hetero-polymers
 79 composed of two major pilins, PilE1 and PilE2, rather than one as normally seen, (ii)
 80 the major pilins display classical type IV pilin 3D structure, and (iii) the filaments
 81 contain a low abundance of three minor pilins (PilA, PilB, and PilC), which are
 82 required for piliation.

83 The present study was prompted by a perplexing observation, *i.e.*, that the
 84 minor pilin PilB harbours a protein domain that has been extensively studied in
 85 several eukaryotic proteins where it mediates adhesion to a variety of protein
 86 ligands¹⁸. This suggested that PilB might be an adhesin, promoting T4P-mediated
 87 adhesion of *S. sanguinis* to host cells and proteins. Therefore, since both the
 88 molecular mechanisms of T4P-mediated adhesion and the exact role of minor pilins
 89 in T4P biology remain incompletely understood¹, we decided to perform a

90 structure/function analysis of PilB, which is reported here. This uncovered a
91 widespread strategy for minor pilins to enhance the functional properties of T4P.

RESULTS

PilB displays a modular pilin architecture

PilB, one of the three minor pilins in *S. sanguinis* T4P¹⁷, exhibits a canonical N-terminal class III signal peptide, the defining feature of type IV pilins⁴. This sequence motif consists of a seven-residue leader peptide composed predominantly of hydrophilic and neutral amino acids (aa), ending with a conserved Gly (Fig. 1A). This leader peptide, which is processed by the prepilin peptidase PilD¹⁷, is followed by a stretch of 21 predominantly hydrophobic aa, except for a negatively charged Glu in position 5 (Fig. 1A). Processed PilB is unusually large for a pilin, with a predicted molecular mass of 50.5 kDa (Fig. 1B). For comparison, the two major pilins of *S. sanguinis* T4P, PilE1 and PilE2¹⁶, have typical pilin sizes of 14.7 and 14.1 kDa, respectively (Fig. 1B). The larger size of PilB is due to the presence of a C-terminal domain (Fig. 1B) readily detectable by a bioinformatic analysis, which belongs to the von Willebrand factor A-like domain superfamily (InterPro entry IPR036465). We will refer to this domain as vWA. The prototypical vWA domain is found in the von Willebrand factor (vWF), a human blood protein required for haemostasis¹⁹, the physiological process that prevents/stops bleeding. vWA domains, which are found in more than 300,000 proteins in the three domains of life, have been extensively studied in eukaryotic proteins where they mediate adhesion to a variety of protein ligands¹⁸. They have been much less studied in bacteria. Of note, the PilB vWA domain is predicted to contain a metal coordination site known as MIDAS, for metal ion-dependent adhesion site²⁰ (Fig. 1A), which was found to be important for ligand-binding in several eukaryotic vWA-containing proteins²¹.

The above-described architecture is unusual for type IV pilins for two reasons. First, in contrast to classical pilins that consist only of a pilin module⁴, defined by a short N-terminal IPR012902 motif within the class III signal peptide (Fig. 1B), PilB apparently has an additional module. Second, the extra C-terminal module in PilB

corresponds to a well-defined functional domain not specific to T4P biology, vWA, which is often associated with adhesion to protein ligands^{18,21}. This has not been previously reported in T4P. This is what we call a modular architecture, and why we refer to PilB as a modular pilin.

Taken together, these findings suggest that PilB is a modular pilin in which a functional module has been grafted during evolution onto a pilin moiety in order to promote T4P-mediated adhesion of *S. sanguinis* to protein ligands.

Crystal structure of PilB reveals a bimodular pilin in which a small type IV pilin module is linked to a bulky vWA module via a short loop

High-resolution structural information is required to confirm that PilB is composed of two modules, but also to understand how modular pilins are polymerised in the filaments and how they modulate T4P functionality. We therefore endeavoured to solve the 3D structure of PilB by X-ray crystallography. To facilitate protein purification, we used a synthetic *pilB* gene codon-optimised for expression in *Escherichia coli*, and produced a recombinant protein in which the N-terminal 35 aa of PilB (encompassing the hydrophobic α 1N) (Fig. 1A) were replaced by a hexahistidine tag (6His)¹⁷. This is a commonly used approach in the field since the truncation of α 1N has minimal structural impact on the rest of the protein²². The resulting 48.4 kDa 6His-PilB protein was soluble and could be purified using a combination of affinity and gel-filtration chromatography. The protein readily crystallised in multiple conditions, and after optimising the best diffracting crystals, we collected a complete dataset on crystals forming in the space group *P*6₁ (Table 1). After phase determination, done using crystals produced in the presence of selenomethionine (SeMet), we solved the 2.26 Å structure of native 6His-PilB. As can be seen in Fig. 2A, this structure reveals a clear bimodular architecture with a small pilin moiety (highlighted in blue) linked to a bulky vWA moiety (in red) by a short nine-residue unstructured loop (grey).

While a bioinformatic analysis could only predict that the extreme N-terminus of PilB corresponds to an IPR012902 class III signal peptide motif, our structure reveals that the first 180 residues of processed PilB clearly display a type IV pilin fold⁴ and thus indeed correspond to a pilin module (Fig. 2B). The pilin module exhibits a long N-terminal α -helix packed against, not one β -sheet as usual, but two consecutive β -sheets consisting of six and three β -strands respectively, which together form the globular head of the pilin. The 432918 topology, *i.e.*, the order of the β -strands in the first β -sheet (Fig. 2B), is unusual since the β -strands are not contiguous along the protein sequence. Moreover, the last portion of this β -sheet forms a Ψ -loop²³ in which two antiparallel strands (β 8 and β 9) are linked via β 1 in between, connected to both of them by hydrogen bonds. This motif occurs rarely in proteins²³.

As for the vWA module (Fig. 2C), the structure strengthens the predictions of the bioinformatic analysis. The vWA moiety of PilB adopts a canonical vWA fold^{20,24}, with a central β -sheet (composed of five parallel and one antiparallel β -strands) surrounded on both sides by a series of α -helices. Consequently, the vWA module of PilB shows high structural similarity to many vWA-containing proteins with which it shares little sequence identity. For example, the vWA module of PilB is very similar to the third vWA domain of human vWF²⁵ (Fig. S1), with a root mean square deviation (RMSD) of 1.72 Å when the two structures are superposed. As in eukaryotic vWA-containing proteins^{20,24}, PilB exhibits a MIDAS located on top of the central β -sheet (Fig. 2C). However, in contrast to these proteins, the MIDAS in PilB is flanked by two protruding "arms", which is reminiscent of the RrgA adhesin from *Streptococcus pneumoniae*²⁶. The first arm is mainly unstructured, while the second folds into a four-stranded β -sheet (Fig. 2C). The MIDAS motif in PilB, which is formed by residues conserved in vWA-containing proteins, non-contiguous in the sequence (Fig. 1A) but in close proximity in the 3D structure (Fig. 2C), is functional since it coordinates a metal ion in the crystal. We have modelled the metal as Mg⁺² because of its abundance in the protein expression medium and the high affinity of PilB for it

(see below). The Ser₂₀₆, Ser₂₀₈, Thr₂₉₁ and Asp₃₁₉ residues in the MIDAS motif²⁰ of PilB form direct hydrogen bonds with the metal through oxygen atoms (Fig. 2C), while two additional coordination sites are provided by water molecules.

An important biological implication of the PilB structure is that modular pilins, despite their large size, are likely to be polymerised into T4P in the same way as classical pilins⁴, *i.e.*, via their N-terminal pilin module. We therefore tested by structural modelling whether PilB could pack into filaments. First, we produced a full-length 3D structural model of PilB including the missing α 1N (Fig. S2), which was truncated in the recombinant protein that we purified. Since a portion of α 1N in major pilins is melted during filament assembly, as observed in several T4aP cryo-EM structures^{9,10}, the α 1N of PilB was modelled with a melted segment. This is consistent with the presence of the helix-breaking Gly residue in position 21 of α 1N (Fig. 1A). Then, we fitted this full-length PilB into a previously generated model of *S. sanguinis* T4P, a right-handed helical heteropolymer where major pilins PilE1/PilE2 are held together by interactions between their α 1N helices (Fig. 3A), which was based on the cryo-EM structure of *Neisseria meningitidis* T4P⁹. Despite its unusual modular structure, PilB can be readily modelled into T4P, its pilin module establishing extensive hydrophobic interactions via its α 1N with the α 1N of neighbouring major pilins (Fig. 3A). This suggests that PilB will assemble into filaments in the same way as classical pilins^{9,10}. However, PilB can only be accommodated at the tip of the filaments because the bulky vWA module sits on top of the pilin module in the PilB structure, and essentially prevents other pilin subunits from being modelled above it (Fig. 3B). Accordingly, when PilB is modelled in the body of the filament (Fig. S3A), it exhibits important steric clashes with neighbouring major pilins (Fig. S3B).

Together, these structural findings show that PilB is a bimodular protein composed of two fused but clearly distinct structural modules. The pilin module adopts a canonical type IV pilin fold⁴, which explains how modular pilins are polymerised into T4P, most probably at their tip. The second module, which is linked

to the end of the pilin module via a short unstructured loop, adopts a vWA fold^{20,24} with a clearly defined MIDAS that coordinates a metal. Since the vWA motif in many eukaryotic proteins is involved in adhesion to protein ligands^{18,21}, our structure strengthens our working hypothesis that PilB might be an adhesin.

Functional analysis of the MIDAS in PilB reveals that metal binding, although structurally dispensable, is important for T4P functionality

Our PilB structure revealed that Mg^{2+} , despite not being added during crystallisation, is bound by the MIDAS. In eukaryotic proteins, the MIDAS sometimes coordinates Mn^{2+} as well^{20,24}. We therefore tested the metal binding specificity of the MIDAS in PilB using ThermoFluor. This fluorescent-based method, which measures changes in thermal denaturation temperature, is a commonly used approach for detecting and quantifying protein-ligand interactions²⁷. We determined the affinity of purified PilB for the Mg^{2+} , Mn^{2+} , and Ca^{2+} divalent cations (Fig. 4A). While no binding was detected to Ca^{2+} , we found that PilB binds Mg^{2+} and Mn^{2+} efficiently in the micromolar range, with estimated K_d of 70 and 54 μM , respectively. To confirm that metal binding involves the MIDAS motif, we produced the PilB_{D319A} protein in which the key MIDAS residue Asp₃₁₉ (Fig. 2C) was changed into an Ala by site-directed mutagenesis. Binding assays performed with PilB_{D319A} showed that changing this one residue abolishes the metal-binding ability of PilB for both Mg^{2+} and Mn^{2+} (Fig. 4B). These findings show that the MIDAS in PilB is functional and preferentially binds Mg^{2+} and Mn^{2+} .

Next, to determine whether metal presence/absence might impact the 3D structure of PilB, we solved the structure of PilB_{D319A} by X-ray crystallography. The PilB_{D319A} protein readily crystallised in the same condition as the wild-type (WT) protein, and we collected a complete dataset on crystals diffracting to a resolution of 3 Å (Table 1). The structure of PilB_{D319A} (Fig. 5A), which was solved by molecular replacement, shows that, no metal is occupying the mutated MIDAS pocket on top of the central β -sheet (Fig. 5B), which is consistent with results of metal binding assays.

When the structures of PilB and PilB_{D319A} were compared, we found that they are essentially identical, superposing onto each other (Fig. 5C) with an RMSD of merely 0.48 Å, including the two arms flanking the MIDAS pocket. This shows that metal-binding by the MIDAS has no detectable structural impact on PilB.

Next, we explored whether MIDAS-mediated metal-binding by PilB is important for piliation and/or T4P-powered twitching motility, both of which were previously shown to be abolished in a $\Delta pilB$ mutant¹⁶. We therefore constructed an unmarked *S. sanguinis* mutant in which the endogenous *pilB* gene was altered by site-directed mutagenesis to produce PilB_{D319A} with an inactive MIDAS. We first tested whether the *pilB*_{D319A} mutant retains the ability to assemble T4P using filament purification¹⁶. As can be seen in Fig. 6A, in which purified T4P were separated by SDS-PAGE and stained with Coomassie blue, the *pilB*_{D319A} mutant is piliated. This is evidenced by the presence of the two bands corresponding to major pilins Pile1 and Pile2, which are absent in a non-piliated $\Delta pilD$ control (Fig. 6A). Moreover, the amount of T4P that can be purified from the *pilB*_{D319A} mutant and WT strain appear comparable. We then tested whether the pili in the *pilB*_{D319A} mutant are able to mediate twitching motility¹⁶. For the WT strain, twitching motility is evidenced by spreading zones around bacteria grown on agar (Fig. 6B). Spreading zones were absent for the *pilB*_{D319A} mutant, which therefore exhibits no detectable twitching motility (Fig. 6B). This shows that the MIDAS-mediated metal-binding ability of PilB, while dispensable for piliation, is important for T4P-mediated twitching motility.

Together, these findings show that the MIDAS in PilB is a functional metal-binding site, dispensable for piliation and protein folding, but essential for T4P functionality.

T4P-mediated adhesion to eukaryotic cells requires PilB, which specifically binds several human proteins

Since vWA is involved in adhesion in many eukaryotic proteins^{18,21}, our original hypothesis was that PilB might mediate *S. sanguinis* adhesion to host cells and/or proteins, which we aimed to test next. First, we determined whether *S. sanguinis* T4P might be involved in its well-known ability to adhere to host cells²⁸. After testing a few eukaryotic cell lines, we opted for CHO cells because the WT strain adheres very efficiently to them. When CHO cells were infected by *S. sanguinis* at a multiplicity of infection (MOI) of 10, 31.6 ± 9.1 % of the bacterial inoculum adhered to the cells. In contrast, a non-piliated $\Delta pilD$ mutant showed a significantly reduced adhesion, with an 18-fold decrease relative to the WT (Fig. 7). Next, we tested our original assumption that PilB might be an adhesin, by quantifying the adhesion of the *pilB*_{D319A} mutant. As can be seen in Fig. 7, although the *pilB*_{D319A} mutant is piliated, its adhesion to CHO cells is dramatically impaired, with a 33-fold decrease when compared to the WT. These findings show that *S. sanguinis* T4P are multi-functional filaments also important for adhesion to eukaryotic cells, and that PilB plays an important role.

Since the vWA domain in multiple eukaryotic proteins has been shown to mediate cell-extracellular matrix (ECM) interactions²¹, we reasoned that PilB might recognise similar ligands because it exhibits a canonical vWA module (Fig. S1). We tested this hypothesis by performing binding assays with purified PilB using enzyme-linked immunosorbent assay (ELISA). In brief, we coated 96-well plates with selected putative ligands, added serial dilutions of purified 6His-PilB, and detected binding using an anti-6His antibody. We tested binding to fibrinogen and the ECM proteins fibronectin, elastin, and laminin. While PilB exhibits no binding to BSA that was used as a negative control (Fig. 8A), dose-dependent binding to fibronectin and fibrinogen was observed, but not to the other ECM proteins that were tested (elastin and laminin). Specific binding to fibronectin and fibrinogen was in the high nanomolar range, with calculated K_d of 865 and 494 nM, respectively (Fig. 8A). Under these *in vitro* experimental conditions metal-coordination by the MIDAS is dispensable for

binding to fibronectin and fibrinogen, as demonstrated by binding assays using purified PilB_{D319A} protein, which showed that PilB_{D319A} binds these ligands as well as PilB (Fig. S4). Finally, to confirm the prediction that binding of PilB to the above ligands is mediated by its vWA module, we then produced and purified the PilB_{vWA} protein corresponding only to the vWA module (see Fig. 1A). We found that PilB_{vWA} could also bind to fibronectin and fibrinogen (Fig. 8B), with calculated K_d of 337 and 997 nM, respectively, which were comparable to PilB. These findings confirm that the adhesive ability of PilB is due to its vWA module.

Taken together, these findings show that *S. sanguinis* T4P are multi-functional filaments mediating adhesion to eukaryotic cells, and that PilB is a *bona fide* adhesin using its vWA module to bind several human protein ligands, which it shares with eukaryotic vWA-containing proteins.

Pilins with modular architectures are widespread in bacteria

While PilB orthologs are ubiquitous in *S. sanguinis*, which also produces a second modular pilin, PilC¹⁷, where the extra module belongs to the concanavalin A-like lectin/glucanase domain superfamily (IPR013320) (Fig. 1B), we wondered how widespread and how diverse modular pilins might be. We therefore searched the InterPro database²⁹ for all the proteins with an N-terminal IPR012902 domain, which also contain an extra domain not specific to T4P biology. This showed that modular pilins are (i) widespread with more than 1,200 proteins displaying such architecture (Supplementary data 1), (ii) present both in monoderm and diderm species, and (iii) diverse, with as many as 264 different architectures detected. Although a bimodular architecture is the most prevalent, there are modular pilins with multiple additional domains, the most extreme case being an 860-residue protein from *Candidatus* Falkowbacteria, with 12 copies of the IPR013211 motif of unknown function (Supplementary data 1). A closer inspection of the 15 most frequent modular pilin architectures offers a glimpse of their diversity (Fig. 9). While in many of these

proteins the extra domain has no clear function (IPR007001, IPR011871, IPR026906, PF05345, IPR006860, IPR003961, IPR021556), for others a function can be predicted. These functions include (i) binding to carbohydrates via PF13385 (that overlaps with the IPR013320 lectin domain superfamily), PF13620 (carbohydrate-binding-like fold), or IPR011658 (PA14 carbohydrate-binding domain), (ii) peptidase activity via IPR030392, and (iii) binding to proteins via IPR002035 (that overlaps with the IPR036465 vWA domain superfamily). These findings suggest that the rather simple modular design strategy – during which a functional module has been grafted during evolution onto a pilin moiety – appears to have been used often during evolution both by monoderm and diderm bacteria and is expected to increase the functional versatility of T4P.

DISCUSSION

T4F are an important research topic because of their virtual ubiquity in prokaryotes and their ability to mediate several key biological processes¹. Furthermore, the molecular mechanisms of most T4F-mediated functions and the exact role of minor pilins remain incompletely understood. Therefore, in this report we focused on T4aP – the prototypical T4F¹ – in the recently established monoderm model *S. sanguinis*¹⁵, and performed a structure/function analysis of the unusual minor pilin PilB, which we predicted might play a role in T4P-mediated adhesion. This led to several notable findings discussed below, and confirmed predictions that the study of T4P in monoderms has the potential to shine new light on these filaments^{14,15}.

The first important finding in this study is that PilB defines a new class of widespread and extremely diverse type IV pilins – the modular pilins – in which an N-terminal pilin module is fused via a short linker to distinct modules that mediate clearly defined functions. Modular pilins are likely to be tip-exposed in the filaments because of their peculiar architecture. While previous 3D structures of a few large minor pilins suggest that they are modular pilins, their second modules have no clear function and do not correspond to protein domains identifiable by available bioinformatic tools. In CofB from enterotoxigenic *E. coli* (ETEC) T4bP, there are two additional structural domains linked to the C-terminus of the pilin module by a flexible linker, a β -repeat domain followed by a β -sandwich domain³⁰. CofB, which forms a trimer predicted to be exposed at the tip of ETEC T4bP³¹, appears to be an adapter for a secreted protein CofJ³², having a direct role in adhesion. In ComZ from *Thermus thermophilus* T4aP, the additional structural domain is a large β -solenoid inserted not at the end of the pilin module but into the β -sheet³³. The role of ComZ is not known but it might be involved in binding extracellular DNA to promote its uptake during transformation³³. This modular architecture is not restricted to T4P as it is also observed for a minor pilin from another T4F, GspK from type II secretion systems (T2SS)³⁴. GspK, in which the additional structural domain is an α -domain of unknown

function inserted into the β -sheet of the pilin module, has also been proposed to be at the tip of T2SS pseudopili, together with two other non-modular minor pilins (GspI and GspJ) with which it interacts to form a heterotrimer³⁴. These examples suggest that we have here probably underestimated the global distribution of modular pilins, which are likely to be much more widespread because in many of them the additional modules are not yet defined by protein signatures in the databases. However, what is clear from our global analysis is that the functions associated with these modular pilins are potentially extremely diverse. Although a "common theme" appears to promote the interaction of T4F with a variety of ligands – including proteins (via vWA in PilB, and the β -repeat/ β -sandwich module in CofB), carbohydrates (via a variety of lectin domains including the concanavalin A-like lectin/glucanase domain in PilC), or DNA (the putative role of the β -solenoid module in ComZ) – other functions are possible. This is suggested by the modular architectures IPR012902-IPR030392 or IPR012902-IPR011493 in which the second module is a predicted peptidase belonging to S74 and M26 families, respectively.

The functional characterisation of the vWA module in PilB including its MIDAS, showing that it is a *bona fide* adhesin, is another significant achievement of this study. First, the vWA domain, which is ubiquitous in the three domains of life and has been extensively studied in eukaryotes^{18,21}, has been much less studied in bacteria. Second, T4P-mediated adhesion remains among the most poorly understood T4P functions¹. Our functional analysis of the vWA module in PilB significantly extends what was known for prokaryotic vWA-containing proteins and highlights important similarities and differences with extensively studied prokaryotic vWA-containing proteins. Our 3D structure shows that the vWA module in PilB exhibits striking similarity to the vWA domain in eukaryotic vWA-containing proteins^{20,24}, with a canonical MIDAS coordinating a metal. The main difference is that the MIDAS in PilB is flanked by two protruding arms, similarly to what has been described for RrgA from *S. pneumoniae*²⁶. Interestingly, RrgA is a subunit with intrinsic adhesive properties³⁵

of sortase-assembled pili in monoderms³⁶, which are unrelated to T4P. The parallel between RrgA and PilB denotes a case of convergent evolution in which two unrelated types of pili have independently evolved a similar strategy to mediate adhesion. Testing metal binding by the MIDAS in PilB, which was previously done only for eukaryotic vWA-containing proteins³⁷, highlight important similarities. MIDAS show no significant binding to Ca²⁺, and a slight preference for Mn²⁺ over Mg²⁺, although the difference in affinity is much smaller than in eukaryotic proteins³⁷. Metal binding can be abolished by altering the MIDAS motif, which has no impact on PilB structure^{37,38}. Abolishing metal binding has no detectable effect on piliation, which is analogous to what has been reported for vWA-containing adhesins of sortase-assembled pili^{39,40}, but it impairs T4P-mediated twitching motility. It is unclear at this stage whether the lack of motility of the *pilB*_{D319A} mutant is due to a reduced T4P adhesion to the agar, which would be consistent with PilB role in adhesion, or to impaired filament retraction, which powers movement¹¹. We also provide evidence that the vWA module of PilB binds several human protein ligands, which it shares with eukaryotic vWA-containing proteins such as integrins and/or vWF^{19,21}. However, unlike in these proteins where binding is often impaired when the MIDAS is inactivated²⁰, a notable difference is that binding to fibronectin and fibrinogen is unaffected in a PilB_{D319A} mutant. This either suggests that the MIDAS is not implicated in binding these specific ligands, which has been described for vWF binding to collagen²⁵, or that our *in vitro* binding assays are not sensitive enough to detect subtle but significant differences in binding.

The finding that PilB plays a key role in adhesion of *S. sanguinis* to host cells and structures, via its vWA module, has implications for the pathogenesis of this species in particular, and for our understanding of T4P-mediated adhesion in general. Our findings are consistent with the possibility that PilB-mediated adhesion to host proteins might play a role in IE⁴¹, a life-threatening infection often caused by *S. sanguinis*. Indeed, during IE, bacteria that have gained access to the bloodstream

adhere to pre-existing sites of valvular damage where sub-endothelial ECM proteins are exposed, and a blood clot is present, containing large amounts of platelets, fibrinogen/fibrin, and fibronectin⁴². Our finding that PilB adheres directly to two of these proteins, but additional ligands cannot be excluded, suggests that PilB might be important at this early stage in IE, which could be tested in future studies. Our findings, which arguably make PilB the best characterised T4P adhesin alongside PilC/PilY1 found in diderm T4aP⁴³⁻⁴⁷, have general implications for our understanding of T4P-mediated adhesion. The vWA module in PilB, which is most likely exposed at the pilus tip, is ideally placed to maximise bacterial adhesion to host protein receptors. T4P spring-like properties (gonococcal T4P can be stretched 3 times their length, which is a reversible process⁴⁸) are expected to help bacteria that are bound via a tip-located adhesin, such as PilB, to withstand adverse forces in their particular environment, *e.g.*, blood flow in a heart valve. This is likely to apply to other modular pilins as well, which harbour different modules predicted to function in adhesion. The parallel with the best characterised T4P adhesin PilC/PilY1 is obvious. This protein, which is not a pilin, is an adhesin that has been proposed to be presented at the T4P tip⁴⁴ via its interaction with a tip-located complex of four widely conserved minor pilins⁴⁹. All PilC/PilY1 have in common a C-terminal IPR008707 β -propeller domain while their N-termini are different⁵⁰. Since this is analogous to the situation with modular pilins, we wondered whether it could be an indication of a modular design for PilC/PilY1. This indeed seems to be the case since a search of the InterPro database²⁹ for all the proteins with an IPR008707 domain shows that 68 different PilC/PilY1 modular architectures are detected (Supplementary data 2). Strikingly, many of the N-terminal modules in PilC/PilY1 are shared with modular pilins. These observations suggest that the same tinkering strategy has been used both by pilins and PilC/PilY1 to increase the functional versatility of T4P. In both instances, a "carrier" module for presentation at the tip of the filaments (either a pilin, or an

IPR008707 domain that interacts with a tip-located complex of minor pilins) has been fused to variety of "effector" modules, directly involved in a variety of functions.

In conclusion, by performing a detailed structure/function of the minor pilin PilB from *S. sanguinis*, we have shed light on several aspects of T4P biology. Our findings are not only of relevance for *S. sanguinis*, most notably for colonisation of its human host, they have general implications for T4P/T4F by uncovering a prevalent strategy used by these widespread filamentous nanomachines to promote their well-known exceptional functional versatility¹. The resulting conceptual framework paves the way for further investigations, which will indubitably improve our understanding of these fascinating filaments.

MATERIALS AND METHODS

Strains and growth conditions

Strains and plasmids used in this study are listed in Table S1. For cloning, we used *E. coli* DH5 α . For protein purification we used *E. coli* BL21(DE3) or *E. coli* BL21 B834(DE3). *E. coli* was grown in liquid or solid Lysogenic Broth (LB) (Difco) containing, when required, 100 μ g/ml spectinomycin or 50 μ g/ml kanamycin (both from Sigma). For purification of protein labelled with seleno-methionine (SeMet), bacteria were grown in chemically defined medium (CDM) supplemented with 20 mg/ml SeMet (Sigma). Chemically competent *E. coli* cells were prepared as described⁵¹. DNA manipulations were done using standard molecular biology techniques⁵². All PCR were done using high-fidelity DNA polymerases (Agilent). Primers used in this study are listed in Table S2. The pET-28b (Novagen) derivative, pET28-*pilB*₃₆₋₄₆₁ for expressing 6His-PilB₃₆₋₄₆₁ was described previously¹⁷. In this plasmid, the portion of a synthetic *pilB* gene codon-optimised for expression in *E. coli*, encoding the soluble portion of PilB, was fused to a non-cleavable N-terminal 6His tag. Similarly, we constructed pET28-*pilB*₁₉₂₋₄₆₁ for expressing PilB_{VWA}. To construct pET28-*pilB*_{D319A} for expressing 6His-PilB_{D319A}, we introduced a missense mutation in pET28-*pilB*₃₆₋₄₆₁ using QuickChange site-directed mutagenesis (Agilent).

The WT *S. sanguinis* 2908 strain and deletion mutants (Δ *pilD*, Δ *pilB*) were described previously¹⁶. *S. sanguinis* was grown on plates containing Todd Hewitt (TH) broth (Difco) and 1% agar (Difco), incubated at 37°C in anaerobic jars (Oxoid) under anaerobic conditions generated using Anaerogen sachets (Oxoid). Liquid cultures were grown statically under aerobic conditions in THT, *i.e.*, TH broth containing 0.05% tween 80 (Merck) to limit bacterial clumping. When required, 500 μ g/ml kanamycin was used for selection and 15 mM *p*-Cl-Phe (Sigma) for counterselection⁵³. *S. sanguinis* genomic DNA was prepared from overnight (O/N) liquid cultures using the kit XIT Genomic DNA from Gram-Positive Bacteria (G-

Biosciences). Strain 2908, which is naturally competent, was transformed as described^{16,53}. The unmarked *S. sanguinis pilB_{D319A}* mutant was constructed using a previously described two-step, cloning-independent, gene editing strategy⁵³. In brief, in the first step, we replaced the gene in the WT by a promoterless *pheS*aphA-3* double cassette, which confers sensitivity to *p*-Cl-Phe and resistance to kanamycin. To do this, we fused by splicing PCR the upstream and downstream regions flanking *pilB* to *pheS*aphA-3*, directly transformed the PCR product into the WT, and selected allelic exchange mutants on kanamycin plates. Allelic exchange was confirmed by PCR. In the second step, we replaced the *pheS*aphA-3* double cassette in this primary mutant by allelic exchange, with an unmarked *pilB_{D319A}* mutation. To do this, we first constructed the missense mutation by site-directed mutagenesis, using as a template a pCR8/GW/TOPO (Invitrogen) derivative in which the WT gene was cloned. Then, the PCR product was directly transformed into the primary mutant, with plating on *p*-Cl-Phe-containing plates. Markerless allelic exchange mutants, which are the only one sensitive to kanamycin, were identified by re-streaking colonies on TH plates with and without antibiotic.

Protein purification

To purify native PilB, PilB_{D319A} and PilB_{VWA} proteins, the corresponding pET-28b derivatives were transformed in *E. coli* BL21(DE3). Transformants were grown O/N at 37°C in liquid LB with kanamycin. The next day, this culture was diluted (1/500) in 1 l of the same medium and grown at 37°C to an OD₆₀₀ of 0.4-0.6. The temperature was then set to 16°C, the culture allowed to cool for 30 min, before protein expression was induced O/N by adding 0.5 mM IPTG (Merck). The next day, cells were harvested by centrifugation at 8,000 *g* for 20 min and subjected to one -80°C freeze/thaw cycle in binding buffer (50 mM HEPES pH 7.4, 200 mM NaCl, 15 mM imidazole), to which we added SIGMAFAST EDTA-free protease inhibitor cocktail (Sigma). Cells were disrupted by repeated cycles of sonication, *i.e.*, pulses of 5 sec

on and 5 sec off during 3-5 min, until the cell suspension was visibly less viscous. The cell lysate was then centrifuged for 30 min at 17,000 g to remove cell debris. The clarified lysate was first affinity-purified on an ÄKTA Purifier using His-Trap HP columns (GE Healthcare) and eluted with elution buffer (50 mM HEPES pH 7.4, 200 mM NaCl, 300 mM imidazole). Affinity-purified proteins were further purified, and simultaneously buffer-exchanged into (50 mM HEPES pH 7.4, 200 mM NaCl), by gel-filtration chromatography on an ÄKTA Purifier using a Superdex 75 10/300 GL column (GE Healthcare). Protein concentration was quantified spectrophotometrically using a NanoDrop.

To purify SeMet-labelled PilB for phasing, the corresponding pET-28b derivative was transformed in *E. coli* BL21 B834(DE3). Transformants were grown at 37°C in liquid LB with kanamycin, until OD₆₀₀ reached 0.6-0.7. Next, the cells were pelleted at 8,000 g for 5 min, and washed twice with 2 ml of CDM, which contains no Met. The pellet was then washed with 2 ml of CDM supplemented with 20 mg/ml L-Met (Sigma) and used to inoculate, at 1/200 dilution, 20 ml of CDM supplemented with 20 mg/ml Met. This culture was grown at 37°C for 16-18 h. Cells were pelleted and washed three times with CDM. Then, the pellet was re-suspended in 20 ml of CDM supplemented with 20 mg/ml SeMet, which was used to inoculate 1 l of CDM supplemented with SeMet. Cells were grown at 37°C until OD₆₀₀ reached 0.5-0.7. The temperature was then set to 16°C, the culture allowed to cool for 30 min, before protein expression was induced O/N by adding 1 mM IPTG (Merck) and 4 ml of 36 % glucose (w/v). Two and half hours later, we again added 4 ml of 36 % glucose to the culture. The next day, SeMet-labelled PilB was purified as above.

Crystallisation and structure determination

Purified proteins in 50 mM HEPES pH 7.4, 200 mM NaCl were concentrated to 50 mg/ml and tested for crystallisation using sitting-drop vapor diffusion, with 100 nl drops of protein solution and mother liquor. We tested a range of commercially

available kits (Molecular Dimensions, Hampton Research and Rigaku Reagents), which yielded a number of hits, mainly in high salt conditions. Crystallisation conditions were optimised to yield larger and better diffracting crystals. The PilB crystals used for the high-resolution structure determination were obtained when the purified protein was mixed 1:1 with crystallisation liquor (0.1 M Bis-tris propane pH 7, 3 M NaCl). The PilB_{D319A} crystals were obtained with crystallisation liquor (0.1 M Bis-tris pH 6.5, 3 M NaCl). Crystals were cryoprotected with 30% glycerol in crystallisation liquor, and flash-frozen in liquid nitrogen. All data was collected and processed using the Diamond Light Source beamline i03, and integrated in *P6₁* using the 3dii pipeline in *xia2*⁵⁴. Initial molecular replacement was performed with Phaser⁵⁵ on the 2.26 Å resolution PilB dataset using a low-resolution partial model produced from the SeMet data using autoSHARP⁵⁶. Manual building in Coot⁵⁷ was performed on the high-resolution dataset, and the full model was then used for molecular replacement in the low-resolution datasets. All structures were produced using Coot and phenix.refine⁵⁸ and validated using MolProbity⁵⁹.

Assaying metal-binding by purified PilB

The metal binding specificity of PilB was tested using ThermoFluor, a fluorescent-based method measuring changes in thermal denaturation temperature²⁷. Assays were done in a 96-well plate (Applied Biosystems) format. In the wells, we added to a final volume of 40 µl (i) 0-1 mM range of concentrations of MgCl₂, MnCl₂ and CaCl₂, (ii) 20 µM purified PilB or PilB_{D319A}, and (iii) 1/5,000 dilution of SYBR Orange (Thermo Fisher Scientific). Plates were then analysed using a temperature gradient, from 25 to 99 °C, on a StepOnePlus real-time qPCR machine (Applied Biosystems). The data were exported MATLAB and analysed in GraphPad. Analyses were performed with Prism (GraphPad Software). K_d were calculated using non-linear regression fits, applying saturation binding equation (One site - Total and non-specific binding) using Ca²⁺ as non-specific binding control.

Assaying protein ligand-binding by purified PilB

Binding of PilB, PilB_{VWA}, PilB_{D319A} to a variety of eukaryotic proteins was tested by ELISA as follows. Putative ligand proteins (elastin from human skin, fibrinogen from human plasma, laminin from human placenta, fibronectin from human plasma) (all from Sigma) were resuspended in carbonate-bicarbonate buffer (Sigma) at 5 µg/ml. Fifty µl was dispatched into the wells of MaxiSorp plates, and adsorbed O/N at 4°C. Wells were washed three times with PBS (Gibco) and blocked during 1 h with 3 % BSA (Probumin) or 1% gelatin (Sigma) in PBS. After washing with PBST (PBS containing 0.05 % tween 20), serial twofold dilutions of PilB (from 40 to 0.625 µg/ml) were added to the wells and incubated for 2 h at 37°C. After five washes with PBST, we added 50 µl anti-6His RTM antibody (Abcam) at 1/500 dilution in PBS, and incubated for 1 h at RT. After five washes with PBST, we added 50 µl Amersham ECL anti-rabbit IgG HRP-linked whole antibody (GE Healthcare) at 1/500 dilution in PBS, and incubated for 1 h at RT. After five washes with PBST, we added 100 µl/well of TMB solution (Thermo Scientific) and incubated the plates during 20 min at RT in the dark. Finally, we stopped the reaction by adding 100 µl/well of 0.18 M sulfuric acid, before reading the plates at 450 nm using a plate reader. Analyses were performed with Prism (GraphPad Software). K_d were calculated using non-linear regression fits, applying saturation binding equation (One site - Total and non-specific binding) using BSA or gelatin as non-specific binding control.

Assaying piliation of *S. sanguinis*

S. sanguinis T4P were purified as described^{16,17}. In brief, bacteria grown in THT until the OD₆₀₀ reached 1-1.5, at which point OD were normalised, a were pelleted and re-suspended in pilus buffer (20 mM Tris, pH 7.5, 50 mM NaCl). This suspension was vortexed for 2 min at full speed to shear T4P. After removing bacterial cells by two centrifugation steps and filtration through a 0.22 µm pore size syringe filter

(Millipore), pili were pelleted by ultracentrifugation. Pili were resuspended in pilus buffer, separated by SDS-PAGE, before gels were stained with Bio-Safe Coomassie (Bio-Rad).

Assaying twitching motility of *S. sanguinis*

Twitching motility was assessed on agar plates as described¹⁶. In brief, bacteria grown O/N were streaked as straight lines on freshly poured TH plates containing 1% Eiken agar (Eiken Chemicals). Plates were grown for several days at 37°C in anaerobic condition under high humidity, which is necessary for twitching. Plates were then photographed using an Epson Perfection V700 photo scanner.

Assaying adhesion of *S. sanguinis* to eukaryotic cells

We tested adhesion of *S. sanguinis* to CHO cells (Public Health England) as follows. Cells were replicated in flasks in DMEM medium (Gibco) containing 1 x MEM non-essential aa mix (Gibco) and 5 % fetal bovine serum (Gibco) and seeded at 100,000 cells/cm² in 24-well plates, which were incubated O/N at 37°C in the presence of 5% CO₂. The next day, cell monolayers were gently rinsed with PBS, and infected at an MOI of 10 with bacteria grown in TH. In brief, bacteria were grown for a few hours to OD₆₀₀ 0.5 units, adjusted at the same OD, pelleted by centrifugation at 1,100 g during 10 min, and resuspended in PBS. Bacteria in the inoculum were quantified by performing CFU counts on TH plates. After 1h of infection at 37°C, cell monolayers were gently rinsed four times with PBS, before cells with adherent bacteria were scraped in distilled water. Adherent bacteria were then quantified by performing CFU counts. Statistical analyses were performed with Prism. Comparisons were done by one-way ANOVA, followed by Dunnett's multiple comparison tests. An adjusted *P* value < 0.05 was considered significant (* *P*<0.05, ** *P*<0.01, *** *P*<0.001, **** *P*<0.0001).

Bioinformatics

Protein sequences were routinely analysed using DNA Strider⁶⁰. Prediction of protein domains, their global distribution and associated architectures was done by using InterProScan²⁹ to interrogate the InterPro database. This database was also used to download all the protein entries discussed in this paper. Molecular visualisation of protein 3D structures was done using PyMOL (Schrödinger). The PDBsum Generate⁶¹ server was used to provide at-a-glance overviews – secondary structure, topology diagram, protein motifs, schematic diagram of metal-protein interactions – of the 3D structures determined during this work. The DALI⁶² server was used for comparing protein structures in 3D. Protein 3D structures were downloaded from the RCSB PDB server. The 3d-SS⁶³ server was used to superpose 3D protein structures.

The cryo-EM structure of *N. meningitidis* T4P (PDB 5KUA)⁹ was used to model, using SWISS-MODEL⁶⁴, the N-terminal helices of PilE1, PilE2 and PilB within the filaments. Coot and PyMOL were then used to place the full-length structures within the T4P model.

DATA AVAILABILITY

The 3D structures determined during this study have been deposited in the PDB, under accession codes 7B7P for PilB, and 7BA2 for PilB_{D319A}. All the datasets generated during this study are included in this paper and its Supplementary information. Source data are provided. The InterPro database (<http://www.ebi.ac.uk/interpro>) was used to identify modular pilin architectures. The DALI server (<http://ekhidna2.biocenter.helsinki.fi/dali>) was used for comparing protein 3D structures. The PDBsum Generate server (<https://www.ebi.ac.uk/thornton-srv/databases/pdbsum/Generate.html>) was used to generate analytical overviews of the 3D structures we have determined. The 3d-SS server (<http://cluster.physics.iisc.ernet.in/3dss>) was used for 3D superposition of protein structures. The RCSB PDB server (<https://www.rcsb.org>) was used to download x3D structures of proteins.

REFERENCES

1. Berry, J.L. & Pelicic, V. Exceptionally widespread nano-machines composed of type IV pilins: the prokaryotic Swiss Army knives. *FEMS Microbiology Reviews* **39**, 134-154 (2015).
2. Denise, R., Abby, S.S. & Rocha, E.P.C. Diversification of the type IV filament superfamily into machines for adhesion, protein secretion, DNA uptake, and motility. *PLoS Biology* **17**, e3000390 (2019).
3. Pelicic, V. Type IV pili: *e pluribus unum*? *Molecular Microbiology* **68**, 827-837 (2008).
4. Giltner, C.L., Nguyen, Y. & Burrows, L.L. Type IV pilin proteins: versatile molecular modules. *Microbiology and Molecular Biology Reviews* **76**, 740-72 (2012).
5. LaPointe, C.F. & Taylor, R.K. The type 4 prepilin peptidases comprise a novel family of aspartic acid proteases. *Journal of Biological Chemistry* **275**, 1502-10 (2000).
6. Francetic, O., Buddelmeijer, N., Lewenza, S., Kumamoto, C.A. & Pugsley, A.P. Signal recognition particle-dependent inner membrane targeting of the PulG pseudopilin component of a type II secretion system. *Journal of Bacteriology* **189**, 1783-93 (2007).
7. Arts, J., van Boxtel, R., Filloux, A., Tommassen, J. & Koster, M. Export of the pseudopilin XcpT of the *Pseudomonas aeruginosa* type II secretion system via the signal recognition particle-Sec pathway. *Journal of Bacteriology* **189**, 2069-76 (2007).
8. Chang, Y.W. *et al.* Architecture of the type IVa pilus machine. *Science* **351**, aad2001 (2016).
9. Kolappan, S. *et al.* Structure of the *Neisseria meningitidis* type IV pilus. *Nature Communications* **7**, 13015 (2016).
10. Wang, F. *et al.* Cryoelectron microscopy reconstructions of the *Pseudomonas aeruginosa* and *Neisseria gonorrhoeae* type IV pili at sub-nanometer resolution. *Structure* **25**, 1423-1435 (2017).

11. Merz, A.J., So, M. & Sheetz, M.P. Pilus retraction powers bacterial twitching motility. *Nature* **407**, 98-102 (2000).
12. Maier, B. *et al.* Single pilus motor forces exceed 100 pN. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 16012-7 (2002).
13. Biais, N., Ladoux, B., Higashi, D., So, M. & Sheetz, M. Cooperative retraction of bundled type IV pili enables nanonewton force generation. *PLoS Biology* **6**, e87 (2008).
14. Melville, S. & Craig, L. Type IV pili in Gram-positive bacteria. *Microbiology and Molecular Biology Reviews* **77**, 323-41 (2013).
15. Pelicic, V. Monoderm bacteria: the new frontier for type IV pilus biology. *Molecular Microbiology* **112**, 1674-1683 (2019).
16. Gurung, I. *et al.* Functional analysis of an unusual type IV pilus in the Gram-positive *Streptococcus sanguinis*. *Molecular Microbiology* **99**, 380-392 (2016).
17. Berry, J.L. *et al.* Global biochemical and structural analysis of the type IV pilus from the Gram-positive bacterium *Streptococcus sanguinis*. *Journal of Biological Chemistry* **294**, 6796-6808 (2019).
18. Whittaker, C.A. & Hynes, R.O. Distribution and evolution of von Willebrand/integrin A domains: widely dispersed domains with roles in cell adhesion and elsewhere. *Molecular Biology of the Cell* **13**, 3369-87 (2002).
19. Sadler, J.E. Biochemistry and genetics of von Willebrand factor. *Annual Review of Biochemistry* **67**, 395-424 (1998).
20. Lee, J.O., Rieu, P., Arnaout, M.A. & Liddington, R. Crystal structure of the A domain from the alpha subunit of integrin CR3 (CD11b/CD18). *Cell* **80**, 631-8 (1995).
21. Dickeson, S.K. & Santoro, S.A. Ligand recognition by the I domain-containing integrins. *Cellular and Molecular Life Sciences* **54**, 556-66 (1998).
22. Craig, L. *et al.* Type IV pilin structure and assembly. X-ray and EM analyses of *Vibrio cholerae* toxin-coregulated pilus and *Pseudomonas aeruginosa* PAK pilin. *Molecular Cell* **11**, 1139-50 (2003).

23. Hutchinson, E.G. & Thornton, J.M. HERA—a program to draw schematic diagrams of protein secondary structures. *Proteins* **8**, 203-12 (1990).
24. Qu, A. & Leahy, D.J. Crystal structure of the I-domain from the CD11a/CD18 (LFA-1, α L β 2) integrin. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 10277-81 (1995).
25. Romijn, R.A. *et al.* Identification of the collagen-binding site of the von Willebrand factor A3-domain. *Journal of Biological Chemistry* **276**, 9985-91 (2001).
26. Izoré, T. *et al.* Structural basis of host cell recognition by the pilus adhesin from *Streptococcus pneumoniae*. *Structure* **18**, 106-15 (2010).
27. Niesen, F.H., Berglund, H. & Vedadi, M. The use of differential scanning fluorimetry to detect ligand interactions that promote protein stability. *Nature Protocols* **2**, 2212-21 (2007).
28. Kreth, J., Giacaman, R.A., Raghavan, R. & Merritt, J. The road less traveled - defining molecular commensalism with *Streptococcus sanguinis*. *Molecular Oral Microbiology* **32**, 181-196 (2017).
29. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**, 1236-40 (2014).
30. Kolappan, S., Ng, D., Yang, G., Harn, T. & Craig, L. Crystal structure of the minor pilin CofB, the initiator of CFA/III pilus assembly in enterotoxigenic *Escherichia coli*. *Journal of Biological Chemistry* **290**, 25805-18 (2015).
31. Kawahara, K. *et al.* Homo-trimeric structure of the type IVb minor pilin CofB suggests mechanism of CFA/III pilus assembly in human enterotoxigenic *Escherichia coli*. *Journal of Molecular Biology* **428**, 1209-1226 (2016).
32. Oki, H. *et al.* Interplay of a secreted protein with type IVb pilus for efficient enterotoxigenic *Escherichia coli* colonization. *Proceedings of the National Academy of Sciences of the United States of America* (2018).

33. Salleh, M.Z. *et al.* Structure and properties of a natural competence-associated pilin suggest a unique pilus tip-associated DNA receptor. *mBio* **10**, e00614-19 (2019).
34. Korotkov, K.V. & Hol, W.G. Structure of the GspK-GspI-GspJ complex from the enterotoxigenic *Escherichia coli* type 2 secretion system. *Nature Structural and Molecular Biology* **15**, 462-8 (2008).
35. Hillerigmann, M. *et al.* Pneumococcal pili are composed of protofilaments exposing adhesive clusters of RrgA. *PLoS Pathogens* **4**, e1000026 (2008).
36. Telford, J.L., Barocchi, M.A., Margarit, I., Rappuoli, R. & Grandi, G. Pili in Gram-positive pathogens. *Nature Reviews in Microbiology* **4**, 509-19 (2006).
37. Baldwin, E.T. *et al.* Cation binding to the integrin CD11b I domain and activation model assessment. *Structure* **6**, 923-35 (1998).
38. Qu, A. & Leahy, D.J. The role of the divalent cation in the structure of the I domain from the CD11a/CD18 integrin. *Structure* **4**, 931-42 (1996).
39. Konto-Ghiorgi, Y. *et al.* Dual role for pilus in adherence to epithelial cells and biofilm formation in *Streptococcus agalactiae*. *PLoS Pathogens* **5**, e1000422 (2009).
40. Nielsen, H.V. *et al.* The metal ion-dependent adhesion site motif of the *Enterococcus faecalis* EbpA pilin mediates pilus function in catheter-associated urinary tract infection. *MBio* **3**, e00177-12 (2012).
41. Que, Y.A. & Moreillon, P. Infective endocarditis. *Nature Reviews Cardiology* **8**, 322-36 (2011).
42. Werdan, K. *et al.* Mechanisms of infective endocarditis: pathogen-host interaction and risk states. *Nature Reviews Cardiology* **11**, 35-50 (2014).
43. Nassif, X. *et al.* Roles of pilin and PilC in adhesion of *Neisseria meningitidis* to human epithelial and endothelial cells. *Proceedings of the National Academy of Sciences of the United States of America* **91**, 3769-3773 (1994).
44. Rudel, T., Scheuerpflug, I. & Meyer, T.F. *Neisseria* PilC protein identified as a type-4 pilus-tip located adhesin. *Nature* **373**, 357-359 (1995).

45. Heiniger, R.W., Winther-Larsen, H.C., Pickles, R.J., Koomey, M. & Wolfgang, M.C. Infection of human mucosal tissue by *Pseudomonas aeruginosa* requires sequential and mutually dependent virulence factors and a novel pilus-associated adhesin. *Cellular Microbiology* **12**, 1158-73 (2010).
46. Johnson, M.D. *et al.* *Pseudomonas aeruginosa* PilY1 binds integrin in an RGD- and calcium-dependent manner. *PLoS One* **6**, e29629 (2011).
47. Porsch, E.A. *et al.* Calcium binding properties of the *Kingella kingae* PilC1 and PilC2 proteins have differential effects on type IV pilus-mediated adherence and twitching motility. *Journal of Bacteriology* **195**, 886-95 (2013).
48. Biais, N., Higashi, D.L., Brujic, J., So, M. & Sheetz, M.P. Force-dependent polymorphism in type IV pili reveals hidden epitopes. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 11358-63 (2010).
49. Treuner-Lange, A. *et al.* PilY1 and minor pilins form a complex priming the type IVa pilus in *Myxococcus xanthus*. *Nature Communications* **11**, 5054 (2020).
50. Orans, J. *et al.* Crystal structure analysis reveals *Pseudomonas* PilY1 as an essential calcium-dependent regulator of bacterial surface motility. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 1065-70 (2010).
51. Inoue, H., Nojima, H. & Okayama, H. High efficiency transformation of *Escherichia coli* with plasmids. *Gene* **96**, 23-8 (1990).
52. Sambrook, J. & Russell, D.W. *Molecular cloning. A laboratory manual*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).
53. Gurung, I., Berry, J.L., Hall, A.M.J. & Pelicic, V. Cloning-independent markerless gene editing in *Streptococcus sanguinis*: novel insights in type IV pilus biology. *Nucleic Acids Research* **45**, e40 (2017).
54. Winter, G., Lobley, C.M. & Prince, S.M. Decision making in *xia2*. *Acta Crystallographica Section D Biological Crystallography* **69**, 1260-73 (2013).

55. McCoy, A.J. *et al.* Phaser crystallographic software. *Journal of Applied Crystallography* **40**, 658-674 (2007).
56. Vonrhein, C., Blanc, E., Roversi, P. & Bricogne, G. Automated structure solution with autoSHARP. *Methods in Molecular Biology* **364**, 215-30 (2007).
57. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot. *Acta Crystallographica Section D Biological Crystallography* **66**, 486-501 (2010).
58. Afonine, P.V. *et al.* Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallographica Section D Biological Crystallography* **68**, 352-67 (2012).
59. Chen, V.B. *et al.* MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallographica Section D Biological Crystallography* **66**, 12-21 (2010).
60. Marck, C. 'DNA Strider': a 'C' program for the fast analysis of DNA and protein sequences on the Apple Macintosh family of computers. *Nucleic Acids Research* **16**, 1829-36 (1988).
61. Laskowski, R.A., Jablonska, J., Pravda, L., Varekova, R.S. & Thornton, J.M. PDBsum: Structural summaries of PDB entries. *Protein Science* **27**, 129-134 (2018).
62. Holm, L. & Laakso, L.M. Dali server update. *Nucleic Acids Research* **44**, W351-5 (2016).
63. Sumathi, K., Ananthalakshmi, P., Roshan, M.N. & Sekar, K. 3dSS: 3D structural superposition. *Nucleic Acids Research* **34**, W128-32 (2006).
64. Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Research* **46**, W296-W303 (2018).

ACKNOWLEDGEMENTS

This work was supported by funding from the MRC (MR/P022197/1) to V. P. We acknowledge the use of the crystallisation facility at Imperial College London, which was supported by BBSRC (BB/D524840/1) and the Wellcome Trust (202926/Z/16/Z). We thank Angelika Gründling (Imperial College London), Sophie Helaine (Harvard Medical School), Christoph Tang (University of Oxford), and Romé Voulhoux (Laboratoire de Chimie Bactérienne) for critical reading of the manuscript.

AUTHOR CONTRIBUTIONS

V. P. was responsible for conception and supervision of the work, and the computational studies. C. R., D. S., J. L. B., and I. G. performed the experimental studies. All authors contributed to writing of the manuscript

816 **COMPETING INTERESTS**

817 The authors declare no competing interests.

818 **Table 1. Crystal structures data collection and refinement statistics.**

Proteins	PilB (7B7P)	PilB_{D319A} (7BA2)
Maximum resolution (Å)	2.26	3.00
Space group	<i>P6₁</i>	<i>P6₁</i>
Unit cell parameters		
a, b, c (Å)	124.38, 124.38, 140.74	120.95, 120.95, 151.25
α, β, γ (°)	90, 90, 120	90, 90, 120
Number of observations	329,504	208,399
Number of unique observations	57,372	25,101
R _{merge} (%)	6.2	16.6
I/σI	13.4	7.6
CC ½	0.999	0.998
Resolution range used for refinement	56.88 – 2.26	75.63 – 3.00
Completeness (%)	99.4	93.8
R factor (%)	20.5	21.2
Free R factor (%)	23.7	27.9
Ramachandran favoured (%)	93.4	83.6
Ramachandran allowed (%)	6.4	12.0
Ramachandran outliers (%)	0.2	4.4
RMSD from ideal values		
bond length (Å)	0.008	0.005
bond angles (°)	0.974	1.043

819

LEGENDS TO FIGURES

Fig. 1. Bioinformatic analysis of PilB. **A)** Relevant features of PilB. The sequence is from *S. sanguinis* 2908. The N-terminal class III signal peptide, the defining feature of type IV pilins, is boxed. The 7-aa long leader peptide contains mostly hydrophilic (shaded in orange) and neutral (no shading) residues, and it ends with a conserved Gly. This leader peptide is processed by the prepilin peptidase PilD, which is indicated by the vertical arrow, generating a protein of 454 residues (50.5 kDa). The processed protein starts with a tract of 21 predominantly hydrophobic residues (shaded in blue), which invariably form an extended α -helix that is the main assembly interface within filaments. The C-terminal vWA module (IPR002035) in PilB is boxed, with the conserved residues forming the MIDAS highlighted in yellow. Arrowheads indicate the proteins that were produced and purified in this study, consisting of either two modules (black arrowhead) or just the vWA module (red arrowhead). **B)** Modular architectures of PilB and PilC minor pilins compared to the major pilins PilE1/PilE2. The proteins, from *S. sanguinis* 2908, have been drawn to scale. The black rounded rectangles correspond to the IPR012902 motif that is part of the class III signal peptide. The C-terminal domains in PilB and PilC are highlighted by coloured rounded rectangles, vWA domain in PilB (red) and lectin domain in PilC (yellow).

Fig. 2. Crystal structure of PilB. **A)** Orthogonal cartoon views of the 6His-PilB structure in which the two distinct modules have been highlighted in blue (pilin module) and red (vWA module), while the short loop connecting them is in grey. The orange sphere represents a magnesium ion. **B)** Left, close-up cartoon view of the pilin module coloured in rainbow spectrum from blue (N-terminus) to red (C-terminus). Right, topology diagram of the pilin module structure. **C)** Left, close-up cartoon view of the vWA module in which the β -strands composing the central β -

sheet are highlighted in red, while the surrounding α -helices are highlighted in yellow. The connecting loops are in grey, except for the two "arms" on top of the structure (coloured in orange and blue), which surround the MIDAS. Right, diagram of the magnesium coordination by the conserved MIDAS residues in the vWA module of PilB. Coordinating oxygen atoms are shown with dashed lines corresponding to hydrogen bonds.

Fig. 3. 3D model of PilB in *S. sanguinis* T4P. **A)** Packing of PilB (red) into *S. sanguinis* T4P, which is a right-handed helical heteropolymer of two major pilins PilE1 (blue) and PilE2 (grey). **B)** View of the T4P tip capped by PilB, or not.

Fig. 4. Metal binding by PilB. Purified PilB was incubated with increasing concentrations of divalent ions (Ca^{2+} , Mg^{2+} , Mn^{2+}) and binding was quantified by ThermoFluor. **A)** Metal binding by PilB. **B)** Metal binding by PilB_{D319A}, with an inactive MIDAS module. Results are the average $\pm$ standard deviations from 3 independent experiments.

Fig. 5. 3D crystal structure of PilB_{D319A}. **A)** Close-up cartoon view of the vWA module in PilB_{D319A}. **B)** Comparison of electron density maps in the MIDAS pocket for the PilB_{D319A} (upper panel) and PilB (lower panel) structures. **C)** Superposition of the vWA modules of PilB (grey) and PilB_{D319A} (orange). The two structures superpose with an RMSD of 0.48 Å.

Fig. 6. Phenotypic characterisation of a *S. sanguinis* mutant expressing PilB_{D319A} with an inactive MIDAS. **A)** Piliation was quantified by purifying T4P from cultures adjusted to the same OD₆₀₀, using a shearing/ultracentrifugation procedure. Purified T4P (identical volumes were loaded in each lane) were separated by SDS-PAGE and stained with Coomassie blue. A molecular weight marker (MW) was run in

the first lane. Molecular weights are indicated in kDa. **B)** Twitching motility was assessed by a macroscopic motility assay. Bacteria were streaked on plates, which were incubated several days at 37°C in a humid atmosphere and then photographed. Twitching motility is characterised by spreading zones around colonies.

Fig. 7. Adhesion of *S. sanguinis* to eukaryotic cells : testing the importance of T4P and the role of the MIDAS motif in PilB. Bacteria were incubated with CHO cells (MOI 10) for 1 h. After removing non-adherent bacteria by several washes, bacteria adhering to cells were enumerated by performing CFU counts. Results are expressed as adhesion relative to WT (set to 1), and are the average $\pm$ standard deviations from five independent experiments. For statistical analysis, one-way ANOVA followed by Dunnett's multiple comparison tests were performed (**** $P < 0.0001$).

Fig. 8. Dose-dependent binding of PilB to various protein ligands. Increasing concentrations of purified PilB was added to constant concentrations of immobilised putative ligands, and binding was quantified by ELISA. BSA served as negative control. Results are the average $\pm$ standard deviations from at least three independent experiments. **A)** Binding of PilB to fibrinogen, fibronectin, elastin, and laminin. **B)** Binding of PilB_{vWA}, consisting only of the vWA module, to fibrinogen and fibronectin.

Fig. 9. Global distribution of modular pilins. The fifteen most widespread modular pilin architectures in the InterPro database are presented. The numbers in parenthesis represent the number of proteins displaying that architecture. The representative proteins depicted, drawn to scale, are from the following species. 1, *Candidatus* Magasanikbacteria (UniProtKB/TrEMBL protein A0A0G0IU57). 2, *Photorhabdus luminescens* (A0A022PI42). 3, *Desulfuribacillus stibiiarsenatis*

903 (A0A1E5L295). 4, *Candidatus* Falkowbacteria (A0A1J4TDE2). 5, *Candidatus*
 904 Wolfebacteria (A0A0G1WFE5). 6, Clostridiales bacterium (A0A101H8M7). 7,
 905 *Corynebacterium glutamicum* (A0A1Q6BQB1). 8, *Thermosulfidibacter takaii*
 906 (A0A0S3QUH2). 9, *Candidatus* Gracilibacteria (A0A1J5F7A7). 10, *Candidatus*
 907 Saccharibacteria (A0A1Q3NLQ9). 11, Planctomycetes bacterium (A0A1G2ZHU9).
 908 12, *Actinoplanes awajinensis* subsp. *mycoplanecinus* (A0A101J7V4). 13,
 909 *Actinoplanes derwentensis* (A0A1H2D7E9). 14, Desulfobacterales bacterium
 910 (A0A1V1WSE4). 15, Parcubacteria group bacterium (A0A2D6FLV5).

A **B**

A **B**

pilus tip without PilB

pilus tip with PilB

A

PilB

BPilB_{D319A}

A

PilB_{D319A}

C

PilB
PilB_{D319A}

B

A

MW WT

 $\Delta pilD$ *pilB*_{D319A}**B**

WT

 $\Delta pilD$ *pilB*_{D319A}

A **B**