Nephrotoxicity of the BRAF-kinase inhibitor Vemurafenib is driven by off-target Ferrochelatase inhibition

Yuntao Bai¹, Ji Young Kim¹, Laura A. Jayne¹, Megha Gandhi¹, Kevin M. Huang¹, Josie A. Silvaroli¹, Veronika Sander², Jason Prosek³, Kenar D. Jhaveri⁴, Sharyn D. Baker¹, Alex Sparreboom¹, Amandeep Bajwa⁵, Navjot Singh Pabla^{1*}

¹Division of Pharmaceutics and Pharmacology, College of Pharmacy & Comprehensive Cancer Center, Ohio State University, Columbus, OH, USA. ²Department of Molecular Medicine and Pathology, University of Auckland, Auckland, New Zealand. ³The Ohio State University Wexner Medical Center, Columbus, OH, USA. ⁴Division of Kidney Diseases and Hypertension, Donald and Barbara Zucker School of Medicine at Hofstra-Northwell, Northwell Health, Great Neck, New York, USA. ⁵Transplant Research Institute, James D. Eason Transplant Institute, Department of Surgery, College of Medicine, The University of Tennessee Health Science Center, Memphis, TN, USA.

Running Title: Off-target mechanisms associated with vemurafenib nephrotoxicity.

Correspondence should be addressed to: Navjot Pabla, Division of Pharmaceutics and Pharmacology, College of Pharmacy and Cancer Center, 460 W 12th Ave, Columbus, OH 43221, USA. Phone: 614-292-1063. E-mail: pabla.2@osu.edu

Abstract

A multitude of disease and therapy related factors drive the frequent development of renal disorders in cancer patients. Along with chemotherapy, the newer targeted therapeutics can also cause renal dysfunction through on and off-target mechanisms. Interestingly, among the small-molecule inhibitors approved for the treatment of cancers that harbor BRAF-kinase activating mutations, vemurafenib can trigger tubular damage and acute kidney injury (AKI). To investigate the underlying mechanisms, here, we have developed cell culture and mouse models of vemurafenib nephrotoxicity. Our studies show that at clinically relevant concentrations vemurafenib induces cell-death in transformed and primary murine and human renal tubular epithelial cells (RTEC). In mice, two weeks of daily vemurafenib treatment causes moderate AKI with histopathological characteristics of RTEC injury. Importantly, RTEC-specific BRAF gene deletion did not influence renal function under normal conditions or alter the severity of vemurafenib-associated renal impairment. Instead, we found that inhibition of ferrochelatase (FECH), an enzyme involved in heme biosynthesis contributes to vemurafenib nephrotoxicity. FECH overexpression protected RTECs and conversely FECH knockdown increased the sensitivity to vemurafenib nephrotoxicity. Collectively, these studies suggest that vemurafenib-associated RTEC dysfunction and nephrotoxicity is BRAF-independent and caused in part by off-target FECH inhibition.

Keywords

Onconeurology, Acute kidney injury, BRAF kinase, Protein kinase inhibitors, Ferrochelatase, and Renal tubular epithelial cells.

Translational Statement

BRAF is the most frequently mutated protein kinase and a critical oncogenic driver in human cancers. In melanoma and other cancers with BRAF activating mutations, BRAF targeted small-molecule therapeutics such as vemurafenib, and dabrafenib have shown remarkable clinical benefits. However, recent clinical studies have shown that a significant number of patients that receive vemurafenib develop AKI through mechanisms that remain unknown. The present study describes the development of novel experimental models of vemurafenib nephrotoxicity and reveals the underlying off-target mechanisms that contribute to renal injury.

Introduction

A broad array of factors increase the risk of renal dysfunction in cancer patients^{1–3}. While certain malignancies can directly affect kidney function⁴, cancer therapy can also trigger fluid and electrolyte disorders⁵, acute kidney injury (AKI)⁶, and chronic kidney disease (CKD)⁷. Renal disorders jeopardize the continuation of cancer therapy and contribute to the development of debilitating short and long-term adverse sequelae. Along with chemotherapeutics, targeted antibodies⁸ and small-molecule inhibitors⁹ are also associated with renal impairment. Thus, one of the challenges in the emerging field of onconeurology^{10–13} is to determine the mechanisms associated with these toxicities in order to develop mitigating strategies.

Dysregulation of the mitogen-activated protein kinase (MAPK) pathway is a major driver of oncogenesis¹⁴. The Raf (rapidly accelerated fibrosarcoma) family of serine/threonine kinases act as a conduit between upstream Ras signaling and downstream MAPK kinase activation, relaying signaling cues from the extracellular environment and directing cell proliferation, differentiation, migration and survival¹⁵. Mammals possess three RAF proteins: RAF1 (CRAF), ARAF and BRAF, which play essential and distinct physiological roles. Importantly, somatic activating mutations in BRAF are frequent in hairy-cell leukemia (100%), melanoma (50%–60%), and thyroid cancer (40%–60%)^{15,16}. The recent development of orally bioavailable BRAF-inhibitors, namely vemurafenib^{17,18} and dabrafenib¹⁹ have brought exceptional clinical benefits, especially in melanoma patients.

Dermatological toxicities are the major side effects related to vemurafenib treatment²⁰. While renal toxicities were not observed in the clinical trials²¹, a growing body of literature suggests that a significant percentage of patients treated with vemurafenib can develop AKI^{22–24,24–26}. Acute tubular necrosis, electrolyte disorders, and subnephrotic-range proteinuria have been reported in a subset of vemurafenib-treated patients^{22,25}. These studies have shown that

patients treated with vemurafenib as compared to dabrafenib have a higher incidence of AKI and analysis of kidney biopsies have established tubular injury as the major histopathological lesion²².

Due to paucity of experimental models, the mechanisms underlying vemurafenib nephrotoxicity remain unclear. Furthermore, it is unknown if BRAF is essential for renal tubular function. It is also unknown whether vemurafenib nephrotoxicity is BRAF-dependent. To address these key questions, here we have developed a renal tubule-specific BRAF knockout mice and established *in vitro* and *in vivo* models of vemurafenib nephrotoxicity. Our studies suggest that BRAF knockout in renal tubules does not trigger renal impairment and vemurafenib-associated RTEC dysfunction and AKI is BRAF-independent.

Results

Vemurafenib induces cell-death in cultured RTECs. We initially sought to determine if vemurafenib could cause direct toxicity in cultured RTECs. To this end, we treated transformed tubular epithelial cells of murine (BUMPT) and human origin (HK-2) with vemurafenib and tested its effect on cellular survival. In these experiments, we also included cisplatin, a well-studied nephrotoxic drug²⁷, as well as BRAF inhibitor dabrafenib, and the multi-kinase and CRAF-inhibitor sorafenib²⁸. Similar to cisplatin, vemurafenib treatment reduced cellular viability as measured by trypan blue staining (**Fig. 1A-B**) in both BUMPT and HK-2 cells ($IC_{50} \sim 50 \mu M$). Notably, pharmacokinetic studies in humans have shown that plasma levels of vemurafenib can range from 50-100 μM ²¹. Confirmatory experiments with MTT assays showed that at 50 μM concentration, vemurafenib can cause ~50% reduction in cellular viability within 48 hours (**Suppl. Fig.1A-B**), with a parallel increase in caspase activation (**Fig. 1C-D**).

Intriguingly, kinase assays showed that at 50 μ M concentration vemurafenib, dabrafenib, and sorafenib inhibited BRAF-kinase activity to similar levels (**Suppl. Fig.1C-D**), however, under these conditions only vemurafenib triggered RTEC cell-death. We next questioned whether BRAF kinase is essential for RTEC survival and a causal factor in vemurafenib-induced cell-death. To address this, we carried out RNAi-mediated knockdown and CRISPR/Cas9-mediated BRAF knockout in BUMPT and HK-2 cells respectively (**Fig. 1E-F**). BRAF knockdown or knockout had no impact on cellular viability under normal conditions and neither did it influence vemurafenib-induced cell-death (**Fig. 1E-F and Suppl. Fig. 2**). These results suggest that vemurafenib can trigger cell-death in cultured RTECs, seemingly through a BRAF-independent mechanism.

Establishment of a mouse model of vemurafenib nephrotoxicity. To establish a murine model of vemurafenib nephrotoxicity, we initially performed a pharmacokinetic study in C57B6/J mice. In concordance with previous work²⁹, we found that at 20 mg/kg dose, plasma drug (~50 μ M) concentrations (**Suppl. Fig. 3**) were similar to those observed in humans²¹. We then determined the effect of 20 mg/kg twice-daily vemurafenib administration on kidney function (**Fig. 2A**). We noticed a significant increase in blood urea nitrogen (**Fig. 2B**) and serum creatinine (**Fig. 2C**) levels after 2 weeks of b.i.d. dosing. Histological examination (**Fig. 2D-E**) revealed tubular necrosis and TUNEL staining (**Fig. 2 D&F**) showed significant tubular epithelial cell-death. In C57B6/J mice, several injury³⁰, repair³¹ and inflammatory genes^{32–35} are upregulated during ischemia, cisplatin, and rhabdomyolysis associated AKI³⁶. We found a similar increase in injury, repair, and inflammation-related genes during vemurafenib nephrotoxicity (**Fig. 2G and Suppl. Fig. 4**). Similar to humans^{22,25}, we found that treatment discontinuation in mice reverses vemurafenib-associated renal impairment (day 20 versus 22).

Vemurafenib nephrotoxicity is BRAF-independent. To evaluate the RTEC-specific role of BRAF kinase, we generated conditional knockout mice (BRAF^{PT-/-}) by crossing the BRAF

floxed³⁷ mice with the Ggt1-Cre mice³⁸. We found that BRAF deficiency (**Figure 3A**) did not influence normal renal function (**Suppl. Fig. 5**). Moreover, when the control and BRAF^{PT-/-} littermates were challenged with vemurafenib, the extent of renal damage as evaluated by blood urea nitrogen, serum creatinine, as well as histological damage and injury biomarker analysis was found to be similar in both groups (**Figure 3B-F**). To corroborate the *in vivo* findings, we isolated primary RTECs from BRAF floxed mice and carried out *in vitro* Cre-mediated deletion (**Suppl. Fig. 6A**). Vemurafenib treatment induced cell-death in primary RTECs and BRAF gene-deletion did not influence cell-death under these conditions (**Suppl. Fig. 6B**). These findings support the notion that BRAF-kinase inhibition is unlikely to be the underlying cause of vemurafenib-associated AKI.

Identification of molecular targets associated with vemurafenib toxicity. Small-molecule kinase inhibitors can target the ATP-binding pocket; however, the high conservation of the ATP binding site within the kinase families poses a significant challenge in developing highly specific kinase inhibitors³⁹. Previous studies^{40,41} have systematically profiled the promiscuity of kinase inhibitors including vemurafenib. To ascertain if inhibition of these kinases contribute to vemurafenib cytotoxicity in RTECs, we used a chemical genetics approach⁴². BUMPT cells were transfected with plasmids encoding wild-type or inhibitor-resistant kinase genes and their effect on cellular survival was monitored (**Suppl. Fig. 7**). Surprisingly, inhibitor-resistant kinase overexpression did not rescue cytotoxicity in BUMPT cells (**Suppl. Fig. 7**), raising the possible role of a non-kinase related mechanism.

Interestingly, chemical proteomics studies^{43,44} have shown that vemurafenib but not dabrafenib can inhibit ferrochelatase (FECH). FECH is a mitochondrial protein that catalyzes the insertion of ferrous iron into protoporphyrin IX, and is the terminal enzyme involved in heme biosynthesis⁴⁵. To ascertain if FECH inhibition causes vemurafenib-associated RTEC cell-death, we initially quantified FECH activity in BUMPT and HK-2 cells. We found that vemurafenib

significantly inhibited FECH activity in RTEC cell-lines (**Fig. 4A-B**). Importantly, wild-type FECH overexpression protected BUMPT and HK-2 from vemurafenib associated cell-death (**Fig. 4C-G**). Overexpression of an inactive FECH mutant (FECH^{mut}) did not influence vemurafenib associated cell-death. These results indicate that FECH inhibition might contribute to RTEC cell-death under in vitro conditions.

In vivo FECH knockdown hastens vemurafenib nephrotoxicity. Immunofluorescence studies showed that FECH is highly expressed in the cortical tubular epithelial cells (**Suppl. Fig. 8A**). While FECH protein levels remained unaltered, enzymatic assays showed a progressive decline in FECH activity during vemurafenib nephrotoxicity (**Suppl. Fig. 8B-C**). Reduction in FECH activity was vemurafenib-specific and not a generalized consequence of AKI (**Suppl. Fig. 8D**). Given that a progressive decline in FECH activity preceded the development of AKI at 2 weeks, we questioned if FECH knockdown would influence vemurafenib nephrotoxicity. Using hydrodynamic siRNA injection approach^{46,47}, we identified a specific siRNA that reduced FECH protein expression by ~70% (**Fig. 5A-B**). FECH knockdown did not influence normal renal function, however strikingly, mice with FECH knockdown developed vemurafenib nephrotoxicity within three days in contrast to 2 weeks in control mice (**Fig. 5C-F**). This increased sensitivity was vemurafenib-specific because no difference in the extent of renal impairment was observed when the mice were challenged with cisplatin (**Suppl. Fig. 9**). These results suggest that FECH knockdown can remarkably hasten the development of vemurafenib-associated AKI.

Accelerated development of vemurafenib nephrotoxicity in FECH mutant mice. In humans, mutations associated with reduced ferrochelatase activity can cause erythropoietic protoporphyria (EPP), a disease characterized by cutaneous photosensitivity and liver damage. In the mice with homozygous (fch/fch) mutations that reduce the FECH activity to 2.7-6.3%, photosensitivity and hepatic dysfunction is observed⁴⁵. On the other hand, the heterozygotes (+/fch) mice have 45-65% of normal FECH activity and do not display any skin or liver damage.

Congruent with a previous study⁴⁵, we found that the heterozygous mutant mice had approximately 50% reduction in renal FECH activity (**Fig. 6A**). Next, we studied the consequence of reduced FECH activity on the development of vemurafenib nephrotoxicity. In comparison to control littermates, the heterozygous mice developed vemurafenib nephrotoxicity in an accelerated manner. At 7 days of vemurafenib treatment, when the control mice had no apparent renal impairment, BUN, serum creatinine, histological, and gene expression analysis (**Fig. 6 B-E**) showed a precipitous decline in renal function in the heterozygous mice. Given that vemurafenib inhibits FECH activity in RTECs and genetic reduction of FECH activity accelerates the development of AKI, we propose that inhibition of renal FECH function contributes to vemurafenib nephrotoxicity.

CRISPR/Cas9 mediated FECH knockout triggers cell-death in HK-2 cells. We questioned if genetic FECH inhibition would be sufficient to trigger cell-death in cultured renal tubular epithelial cells. To address this, we utilized a doxycycline-inducible CRISPR/Cas9 system, wherein we generated stable HK-2 cells that express either a control or FECH-targeted sgRNA, along with doxycycline-inducible Cas9. As shown in **Fig. 7A-B**, doxycycline treatment resulted in decline in FECH protein expression in the stable cells expressing the FECH targeted sgRNA. When we further monitored these cells for 24-72 hours post doxycycline induction, we noticed a clear decline in cellular viability in the FECH knockout cells as measured by trypan blue staining (**Fig. 7C**). At 72 hours, MTT and caspase activity assays confirmed reduced viability and increased cell-death in the FECH knockout cells (**Fig. 7D-E**). The observation that genetic FECH deficiency impairs RTEC viability supports the hypothesis that FECH inhibition could trigger RTEC cell-death and dysfunction.

Discussion

In the current study, we have established cell culture and murine models of vemurafenib nephrotoxicity that have provided important mechanistic insights. Our studies reveal that vemurafenib triggers RTEC dysfunction and nephrotoxicity through a BRAF-independent and ferrochelatase-dependent manner.

Similar to most protein kinase inhibitors, vemurafenib is mainly excreted via feces (94%), with urinary excretion (1%) playing an insignificant role^{21,48}. Based on our *in vitro* and *in vivo* studies, we propose that vemurafenib causes direct toxicity to tubular epithelial cells. Several lines of evidence suggest that this toxicity is independent of BRAF-kinase inhibition. Firstly, both vemurafenib and dabrafenib treatment resulted in BRAF inhibition, however only vemurafenib triggered RTEC cell-death. Secondly, BRAF deletion in murine and human RTECs did not influence cell survival under normal conditions. BRAF deletion or knockdown also did not influence vemurafenib-induced cell-death and AKI. This also eliminates the possibility that paradoxical MAPK activation initiated by inhibitor-induced wild type BRAF dimerization⁴⁹ might contribute to nephrotoxicity. Thirdly, chemical genetic studies with transfection of vemurafenib-resistant BRAF gatekeeper mutants did not confer protection from vemurafenib-induced cell-death. While genetic compensation remains a possibility, collectively our results suggest that BRAF-kinase is not an essential gene in RTECs and vemurafenib nephrotoxicity is BRAF-independent. Though it is important to note that AKI is a multifaceted disease² that results from a complex interplay between epithelial^{31,50,51}, immune³³, and endothelial^{52,53} cells. Thus, notwithstanding our studies with RTECs, further work is required to discern the role of immune and endothelial cells in vemurafenib nephrotoxicity.

We propose that vemurafenib inhibits ferrochelatase, which likely contributes to RTEC cell-death and AKI. Ferrochelatase is the terminal enzyme of the heme biosynthesis pathway⁴⁵. FECH is a mitochondrial protein that catalyzes the insertion of the reduced form of iron (Fe^{2+}) into protoporphyrin. In human, mutations associated with reduced FECH activity are associated

with erythropoietic protoporphyria (EPP) that is characterized by cutaneous photosensitivity and progressive liver damage⁴⁵. Proteomic profiling of leukemia cells previously revealed that vemurafenib but not dabrafenib is a potent FECH inhibitor^{43,44}. It was thus speculated that FECH inhibition might contribute to photosensitivity and skin toxicities associated with vemurafenib treatment. Our in vivo studies show that vemurafenib treatment results in a progressive decline in FECH activity and development of AKI two weeks post treatment initiation. Importantly, RNAi mediated knockdown or FECH loss-of-function mutations, cause a striking acceleration in the development and onset of vemurafenib-associated AKI (3-7 days versus 15 days). Furthermore, FECH overexpression in cultured RTECs provided significant protection from vemurafenib-induced cell-death.

In contrast with targeted anti-angiogenic therapeutics⁵⁴ where the entire class of drugs cause hypertension due to 'on-target' effects, not all BRAF inhibitors seem to cause AKI²². This is supported by the observation that patients treated with vemurafenib have a higher incidence of AKI, with a recent retrospective study suggesting that nearly 60% patient develop AKI within the first three months of treatment^{22,25}. We propose that this might be due to the unique property of vemurafenib to inhibit FECH. In support, BRAF gene deletion in RTECs did not result in renal impairment in vivo. While it is currently remains unknown if BRAF is essential in other renal cell types, it is interesting that BRAF inhibition has been suggested as a therapeutic option to prevent podocyte injury^{55,56}. As of yet, it is difficult to predict which patients are more likely to develop vemurafenib nephrotoxicity. It would be interesting to examine if genetic polymorphisms or physiological conditions associated with altered renal FECH activity play a contributing role.

Our study raises several questions that warrant future investigations. While we show that vemurafenib mediated FECH inhibition contributes to RTEC cell-death, the underlying mechanisms remain unknown. Altered iron homeostasis is a known feature of AKI⁵⁷⁻⁵⁹; however, the role of FECH in RTEC function and AKI have not been previously explored. It is

conceivable that vemurafenib-mediated FECH inhibition might result in accumulation of by-products of heme biosynthesis⁴⁴, which might cause direct toxicity to RTECs through increased oxidative stress. Furthermore, the reduction in heme biosynthesis could also result in mitochondrial dysfunction and RTEC cell-death. Additionally, while genetic FECH knockout triggers RTEC cell-death, it is unlikely that FECH inhibition in RTECs is solely responsible for vemurafenib nephrotoxicity. FECH-dependent and independent mechanisms in RTECs and other renal cells including endothelial and immune cells might play a role in this toxicity.

Due to the availability of BRAF inhibitors such as dabrafenib and the moderate nature of vemurafenib nephrotoxicity, currently there is no dire need to develop therapeutic strategies to prevent vemurafenib-associated AKI. However, several approved, experimental, and investigational kinase inhibitors have recently been demonstrated to inhibit FECH activity, through mechanisms that remain elusive⁴⁴. For instance, an ALK (anaplastic lymphoma kinase) inhibitor alcatinib can inhibit FECH⁴³ and is linked with renal dysfunction⁶⁰, although the causality remains to be investigated. Since, FECH inhibition is not limited to vemurafenib, but has emerged as a relatively common feature of several protein kinase inhibitors, future genetic and pharmacological studies are required to understand the role of this enzyme in renal function and AKI.

In summary, our work suggests that vemurafenib-associated AKI is an off target toxicity that can be partly attributed to FECH inhibition in RTECs. By identifying the underlying mechanisms, our study reveals that drugs with FECH-inhibiting ability might cause RTEC dysfunction and nephrotoxicity.

Methods

Animal Experiments. Mice were housed and handled in accordance with the Institutional Animal Care and Use Committee of the Ohio State University and the University of Tennessee Health Science Center. Details regarding animal procedures are provided in Supplementary methods.

Cell culture. BUMPT, HK-2, and primary murine RTECs were cultured according to previously described methods^{61,62}. Further details of experimental procedures are provided in Supplementary methods.

Statistical Analysis. Data are presented as mean with SD. $P < 0.05$ was considered statistically significant. Further details are provided in supplementary methods.

Disclosure Statement

No conflicts of interest, financial or otherwise, are declared by the authors.

Acknowledgements

We thank Dr. Yogesh Scindia for critical reading of the manuscript. We thank Dr. Shuiying Hu for assistance with pharmacokinetic analysis. This work was supported by funds from the OSU Cancer Center (to N.S.P.) and NIDDK Grant R01 DK-117183 (to A.B.). Y.B. was supported by postdoctoral fellowships from the Pelotonia foundation and the American Heart Association.

Author Contributions. Y.B., J.Y.K., L.A.J., M.G., J.A.S, V.S., K.M.H, S.D.B., A.S., A.B., and N.S.P acquired and analyzed most of the data. Y.B., V.S., J.S., K.D.J and N.S.P conceived the study and performed data analysis. N.S.P approved the final version of the manuscript.

Figure Legends

Figure 1: *Vemurafenib induces cell death in murine and human tubular epithelial cell lines.* Tubular epithelial cell lines of murine (BUMPT) and human (HK-2) origin were treated with vehicle, cisplatin or kinase inhibitors including vemurafenib followed by assessment of cell viability and cell death. **(A-B)** Dose response experiments (0-100 μ M) and trypan blue based cellular viability assays at 48 hours post treatment showed that vemurafenib can induce cell death in RTEC cell lines with IC50 values of approximately 50 μ M. Survival data was normalized to vehicle group and are presented as mean ($n=5$ biologically independent samples), from one out of three independent experiments, all producing similar results. **(C-D)** BUMPT and HK-2 cells were treated with vehicle or indicated drugs at 50 μ M concentration, followed by measurement of caspase activity at 48 hours. The results show that similar to cisplatin, vemurafenib can reduce RTEC viability. Data are presented as individual data points ($n=8$ biologically independent samples from three independent experiments). **(E)** RNAi mediated Braf knockdown in BUMPT cells did not influence vemurafenib associated cell death (50 μ M for 48 hours) as assessed by trypan blue based viability assay. Data are presented as individual data points ($n=8$ biologically independent samples from three independent experiments). A representative immunoblot (right panel) shows the successful knockdown of Braf gene. **(F)** CRISPR/Cas9 mediated Braf knockout in HK-2 cells did not influence vemurafenib associated cell death (50 μ M for 48 hours) as assessed by trypan blue based viability assay. Data are presented as individual data points ($n=8$ biologically independent samples from three independent experiments). A representative immunoblot (right panel) shows the successful knockout of Braf gene. In all the graphs ($n=5-8$ biologically independent samples),

experimental values are presented as mean $\pm$ s.d. The height of error bar = 1 s.d. and $p < 0.05$ was indicated as statistically significant. One-way ANOVA followed by Dunnett's was carried out and statistical significance is indicated by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 2: Development of a mouse model of vemurafenib nephrotoxicity. (A) Schematic representation of dosing strategy. Age-matched, 8-12 weeks old male C57BL/6J mice were treated with either vehicle or 20 mg/kg vemurafenib (p.o, b.i.d.) for 20 days followed by cessation of drug administration for 2 days and subsequent endpoint analysis of renal function. (A) Analysis of blood urea nitrogen levels showed that vemurafenib can induce AKI after 2 weeks of continuous treatment and cessation of drug administration reversed the increase in BUN levels. (C) Vemurafenib treatment also resulted in increased serum creatinine levels indicating significant renal impairment (D-E) Histological analysis of renal tissues showed that vemurafenib treated mice had clear tubular epithelial injury and cell death. Representative H&E staining depicting renal tubular damage (indicated by an asterisk) linked with vemurafenib-associated AKI. (D & F) TUNEL staining of renal tissues revealed significant tubular epithelial cell death in the vemurafenib treated mice. (G) Renal NGAL gene expression analysis further confirmed significant renal damage in vemurafenib treated mice. In all the bar graphs (n=7 biologically independent samples), experimental values are presented as mean $\pm$ s.d. The height of error bar = 1 s.d. and $p < 0.05$ was indicated as statistically significant. Student's t-test was carried out and statistical significance is indicated by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Scale bar (D): 100 μ m.

Figure 3: Vemurafenib nephrotoxicity is not influenced by RTEC-specific *Braf* gene deletion. To generate mice with renal tubule-specific *Braf* knockout, *Ggt1-Cre* mice were crossed with *Braf*-floxed mice. (A-B) Representative genotyping and immunoblots showing

successful knockout in the renal tissues. Littermate control and *Braf* conditional knockout mice (indicated by $Braf^{PT-/-}$) were used to study the role of *Braf* in vemurafenib nephrotoxicity. Age-matched, 8-12 weeks old male littermate mice were treated with either vehicle or 20 mg/kg vemurafenib (p.o, b.i.d.) for 20 days followed by subsequent endpoint analysis of renal function. Blood urea nitrogen (C), serum creatinine (D), and histological analysis (E) were performed to examine renal function and damage. The extent of functional renal impairment and damage was similar between the control and *Braf* deficient mice. (F) TUNEL staining also showed similar amount of renal epithelial cell death in the control and knockout tissues. (G) Renal NGAL gene expression analysis. In all the bar graphs (n=8 biologically independent samples), experimental values are presented as mean $\pm$ s.d. The height of error bar = 1 s.d. and $p < 0.05$ was indicated as statistically significant. One-way ANOVA followed by Tukey's multiple-comparison test was carried out and statistical significance is indicated by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 4: Ferrochelatase inhibition contributes to Vemurafenib mediated RTEC cell death.

Tubular epithelial cell lines of murine (BUMPT) and human (HK-2) origin were treated with vehicle, cisplatin or kinase inhibitors including vemurafenib at 50 μ M concentration, followed by assessment of ferrochelatase activity at 24 hours. (A-B) Ferrochelatase activity was inhibited by vemurafenib in BUMPT and HK-2 cells. (B) Representative immunoblot showing overexpression of FLAG-tagged wild type and mutant FECH. Blots are representative of three independent experiments. (D-G) Empty vector, wild type FECH, or FECH mutant (M98K) was overexpressed in BUMPT (transient transfection) and HK-2 (lentiviral transduction) cells followed by treatment with either vehicle or 50 μ M vemurafenib for 48 hours. Trypan blue based survival assays and caspase assays showed that wild type FECH overexpression can protect BUMPT and HK-2 cells from vemurafenib-associated cell death. In all the bar graphs (n=5-6 biologically

independent samples) from one out of three independent experiments, all producing similar results, and experimental values are presented as mean $\pm$ s.d. The height of error bar = 1 s.d. and $p < 0.05$ was indicated as statistically significant. One-way ANOVA followed by Tukey's multiple-comparison test was carried out and statistical significance is indicated by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 5: *In vivo* siRNA mediated FECH knockdown hastens the development of vemurafenib nephrotoxicity. Age-matched male (8-12 weeks) C57BL/6 mice were administered with three once-daily intravenous injections of control (non-specific) or FECH targeting siRNAs (25 μ g in 0.5 ml of PBS). In one group (mock) 0.5 ml of PBS was injected. One day later mice were treated with either vehicle or 20 mg/kg vemurafenib (p.o, b.i.d.) for 3 days followed by endpoint analysis of renal function. (A) Representative blots and (B) densitometric analysis show that the targeted siRNA was able to knock-down FECH proteins levels by approximately 75%. Blots are representative of three independent experiments, all producing similar results. Blood urea nitrogen (C), serum creatinine (D), and histological analysis (E) Renal NGAL gene expression analysis (F) showed that the FECH knockdown mice developed vemurafenib nephrotoxicity within 3 days of treatment, while the control group demonstrated no obvious renal injury or damage. In all the bar graphs (n=6-10 biologically independent samples), experimental values are presented as mean $\pm$ s.d. The height of error bar = 1 s.d. and $p < 0.05$ was indicated as statistically significant. One-way ANOVA followed by Tukey's multiple-comparison test was carried out and statistical significance is indicated by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 6: Accelerated development of vemurafenib nephrotoxicity in FECH mutant mice.

(A) Renal tissues from wild type and heterozygous (Fch/+) mutant mice (littermates) were used

to evaluate FECH activity using an enzymatic assay. The result show an approximately 50% reduction in FECH activity in the heterozygous mice. The wild type and heterozygous mice were then challenged with either vehicle or 20 mg/kg vemurafenib (p.o, b.i.d.) for 7 days followed by endpoint analysis of renal function. Blood urea nitrogen (**B**), serum creatinine (**C**), and histological analysis (**D**) Renal NGAL gene expression analysis (**E**) showed that the FECH heterozygous mutant mice developed vemurafenib nephrotoxicity within 7 days of treatment, at a time-point when the wild type group demonstrated no obvious renal injury or damage. In all the bar graphs (n=6-8 biologically independent samples), experimental values are presented as mean $\pm$ s.d. The height of error bar = 1 s.d. and $p < 0.05$ was indicated as statistically significant. One-way ANOVA followed by Tukey's multiple-comparison test was carried out and statistical significance is indicated by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 7: CRISPR/Cas9 mediated FECH knockout induces cell death in HK-2 cells. Stable HK-2 cells were generated with either an control or FECH targeted sgRNA. In these stable cells lines, Cas9 expression can be induced by doxycycline treatment. (**A**) In the presence of doxycycline, FECH gene deletion occurs in the cells that express FECH targeted sgRNA. Representative immunoblot depicting successful gene knockout. Blots are representative of three independent experiments. (**B**) Densitometric analysis of FECH protein after 72 hours of doxycycline treatment. (**C**) Control (vector) and FECH sgRNA (FECH^{Dox-KO}) stable cells growing in normal media, were treated with doxycycline and trypan blue based survival assays were performed at 24-72 hours. The results show that FECH knockdown resulted in significant reduction in cell survival. (In all the bar graphs (n=5-6 biologically independent samples) from one out of three independent experiments, all producing similar results, and experimental values are presented as mean $\pm$ s.d. The height of error bar = 1 s.d. and $p < 0.05$ was indicated

as statistically significant. One-way ANOVA followed by Dunnett's test was carried out and statistical significance is indicated by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

References

1. Malyszko J, Tesarova P, Capasso G, Capasso A. The link between kidney disease and cancer: complications and treatment. *The Lancet*. 2020;396(10246):277-287. doi:10.1016/S0140-6736(20)30540-7
2. Bellomo R, Kellum JA, Ronco C. Acute kidney injury. *Lancet*. 2012;380(9843):756-766. doi:10.1016/S0140-6736(11)61454-2
3. Iff S, Craig JC, Turner R, et al. Reduced estimated GFR and cancer mortality. *Am J Kidney Dis*. 2014;63(1):23-30. doi:10.1053/j.ajkd.2013.07.008
4. Schwartz WB, Bennett W, Curelop S, Bartter FC. A syndrome of renal sodium loss and hyponatremia probably resulting from inappropriate secretion of antidiuretic hormone. *Am J Med*. 1957;23(4):529-542. doi:10.1016/0002-9343(57)90224-3
5. Uppal NN, Wanchoo R, Barnett R, Sinha A, Jhaveri KD. Hyponatremia in a Patient With Cancer. *Am J Kidney Dis*. 2020;75(1):A15-A18. doi:10.1053/j.ajkd.2019.09.005
6. Rosner MH, Perazella MA. Acute Kidney Injury in Patients with Cancer. *New England Journal of Medicine*. 2017;376(18):1770-1781. doi:10.1056/NEJMr1613984
7. Launay-Vacher V. Epidemiology of chronic kidney disease in cancer patients: lessons from the IRMA study group. *Semin Nephrol*. 2010;30(6):548-556. doi:10.1016/j.semnephrol.2010.09.003
8. Seethapathy H, Zhao S, Chute DF, et al. The Incidence, Causes, and Risk Factors of Acute Kidney Injury in Patients Receiving Immune Checkpoint Inhibitors. *Clin J Am Soc Nephrol*. 2019;14(12):1692-1700. doi:10.2215/CJN.00990119
9. Jhaveri KD, Sakhiya V, Wanchoo R, Ross D, Fishbane S. Renal effects of novel anticancer targeted therapies: a review of the Food and Drug Administration Adverse Event Reporting System. *Kidney Int*. 2016;90(3):706-707. doi:10.1016/j.kint.2016.06.027

10. Lam AQ, Humphreys BD. Onco-Nephrology: AKI in the Cancer Patient. *Clin J Am Soc Nephrol*. 2012;7(10):1692-1700. doi:10.2215/CJN.03140312
11. Salahudeen AK, Bonventre JV. Onconeurology: The Latest Frontier in the War against Kidney Disease. *JASN*. 2013;24(1):26-30. doi:10.1681/ASN.2012070690
12. Rosner MH, Jhaveri KD, McMahon BA, Perazella MA. Onconeurology: The intersections between the kidney and cancer. *CA Cancer J Clin*. Published online August 27, 2020. doi:10.3322/caac.21636
13. Cambier J-F, Ronco P. Onco-nephrology: glomerular diseases with cancer. *Clin J Am Soc Nephrol*. 2012;7(10):1701-1712. doi:10.2215/CJN.03770412
14. Yaeger R, Corcoran RB. Targeting Alterations in the RAF–MEK Pathway. *Cancer Discov*. 2019;9(3):329-341. doi:10.1158/2159-8290.CD-18-1321
15. Holderfield M, Deuker MM, McCormick F, McMahon M. Targeting RAF kinases for cancer therapy: BRAF-mutated melanoma and beyond. *Nature Reviews Cancer*. 2014;14(7):455-467. doi:10.1038/nrc3760
16. Davies H, Bignell GR, Cox C, et al. Mutations of the BRAF gene in human cancer. *Nature*. 2002;417(6892):949-954. doi:10.1038/nature00766
17. Flaherty KT, Yasothan U, Kirkpatrick P. Vemurafenib. *Nature Reviews Drug Discovery*. 2011;10(11):811-812. doi:10.1038/nrd3579
18. Bollag G, Tsai J, Zhang J, et al. Vemurafenib: the first drug approved for BRAF -mutant cancer. *Nature Reviews Drug Discovery*. 2012;11(11):873-886. doi:10.1038/nrd3847
19. Hauschild A, Grob J-J, Demidov LV, et al. Dabrafenib in BRAF-mutated metastatic melanoma: a multicentre, open-label, phase 3 randomised controlled trial. *Lancet*. 2012;380(9839):358-365. doi:10.1016/S0140-6736(12)60868-X
20. Lacouture ME, Duvic M, Hauschild A, et al. Analysis of Dermatologic Events in Vemurafenib-Treated Patients With Melanoma. *Oncologist*. 2013;18(3):314-322. doi:10.1634/theoncologist.2012-0333
21. Bollag G, Hirth P, Tsai J, et al. Clinical efficacy of a RAF inhibitor needs broad target blockade in BRAF-mutant melanoma. *Nature*. 2010;467(7315):596-599. doi:10.1038/nature09454
22. Jhaveri KD, Sakhiya V, Fishbane S. Nephrotoxicity of the BRAF Inhibitors Vemurafenib and Dabrafenib. *JAMA Oncol*. 2015;1(8):1133-1134. doi:10.1001/jamaoncol.2015.1713
23. Uthuriague C, Thellier S, Ribes D, Rostaing L, Paul C, Meyer N. Vemurafenib significantly decreases glomerular filtration rate. *J Eur Acad Dermatol Venereol*. 2014;28(7):978-979. doi:10.1111/jdv.12322
24. Regnier-Rosencher E, Lazareth H, Gressier L, Avril MF, Thervet E, Dupin N. Acute kidney injury in patients with severe rash on vemurafenib treatment for metastatic melanomas. *Br J Dermatol*. 2013;169(4):934-938. doi:10.1111/bjd.12555

25. Teuma C, Perier-Muzet M, Pelletier S, et al. New insights into renal toxicity of the B-Raf inhibitor, vemurafenib, in patients with metastatic melanoma. *Cancer Chemother Pharmacol.* 2016;78(2):419-426. doi:10.1007/s00280-016-3086-7
26. Hurabielle C, Pillebout E, Stehlé T, et al. Mechanisms Underpinning Increased Plasma Creatinine Levels in Patients Receiving Vemurafenib for Advanced Melanoma. *PLoS One.* 2016;11(3):e0149873. doi:10.1371/journal.pone.0149873
27. Pabla N, Dong Z. Cisplatin nephrotoxicity: mechanisms and renoprotective strategies. *Kidney Int.* 2008;73(9):994-1007. doi:10.1038/sj.ki.5002786
28. Wilhelm S, Carter C, Lynch M, et al. Discovery and development of sorafenib: a multikinase inhibitor for treating cancer. *Nature Reviews Drug Discovery.* 2006;5(10):835-844. doi:10.1038/nrd2130
29. Yang H, Higgins B, Kolinsky K, et al. Antitumor Activity of BRAF Inhibitor Vemurafenib in Preclinical Models of BRAF-Mutant Colorectal Cancer. *Cancer Res.* 2012;72(3):779-789. doi:10.1158/0008-5472.CAN-11-2941
30. Ichimura T, Bonventre JV, Bailly V, et al. Kidney Injury Molecule-1 (KIM-1), a Putative Epithelial Cell Adhesion Molecule Containing a Novel Immunoglobulin Domain, Is Up-regulated in Renal Cells after Injury. *J Biol Chem.* 1998;273(7):4135-4142. doi:10.1074/jbc.273.7.4135
31. Kumar S, Liu J, Pang P, et al. Sox9 Activation Highlights a Cellular Pathway of Renal Repair in the Acutely Injured Mammalian Kidney. *Cell Rep.* 2015;12(8):1325-1338. doi:10.1016/j.celrep.2015.07.034
32. Okusa MD. The inflammatory cascade in acute ischemic renal failure. *Nephron.* 2002;90(2):133-138. doi:10.1159/000049032
33. Bajwa A, Kinsey GR, Okusa MD. Immune mechanisms and novel pharmacological therapies of acute kidney injury. *Curr Drug Targets.* 2009;10(12):1196-1204. doi:10.2174/138945009789753174
34. Ramesh G, Reeves WB. TNF-alpha mediates chemokine and cytokine expression and renal injury in cisplatin nephrotoxicity. *J Clin Invest.* 2002;110(6):835-842. doi:10.1172/JCI15606
35. Rousselle TV, Kuscu C, Kuscu C, et al. FTY720 Regulates Mitochondria Biogenesis in Dendritic Cells to Prevent Kidney Ischemic Reperfusion Injury. *Front Immunol.* 2020;11:1278. doi:10.3389/fimmu.2020.01278
36. Kim JY, Bai Y, Jayne LA, et al. SOX9 promotes stress-responsive transcription of VGF nerve growth factor inducible gene in renal tubular epithelial cells. *J Biol Chem.* 2020;295(48):16328-16341. doi:10.1074/jbc.RA120.015110
37. Chen AP, Ohno M, Giese KP, Kühn R, Chen RL, Silva AJ. Forebrain-specific knockout of B-raf kinase leads to deficits in hippocampal long-term potentiation, learning, and memory. *J Neurosci Res.* 2006;83(1):28-38. doi:10.1002/jnr.20703

38. Iwano M, Plieth D, Danoff TM, Xue C, Okada H, Neilson EG. Evidence that fibroblasts derive from epithelium during tissue fibrosis. *J Clin Invest.* 2002;110(3):341-350. doi:10.1172/JCI15518
39. Gross S, Rahal R, Stransky N, Lengauer C, Hoeflich KP. Targeting cancer with kinase inhibitors. *J Clin Invest.* 2015;125(5):1780-1789. doi:10.1172/JCI76094
40. Davis MI, Hunt JP, Herrgard S, et al. Comprehensive analysis of kinase inhibitor selectivity. *Nat Biotechnol.* 2011;29(11):1046-1051. doi:10.1038/nbt.1990
41. Knight ZA, Lin H, Shokat KM. Targeting the cancer kinome through polypharmacology. *Nat Rev Cancer.* 2010;10(2):130-137. doi:10.1038/nrc2787
42. Li J, Rix U, Fang B, et al. A chemical and phosphoproteomic characterization of dasatinib action in lung cancer. *Nat Chem Biol.* 2010;6(4):291-299. doi:10.1038/nchembio.332
43. Savitski MM, Reinhard FBM, Franken H, et al. Tracking cancer drugs in living cells by thermal profiling of the proteome. *Science.* 2014;346(6205). doi:10.1126/science.1255784
44. Klaeger S, Gohlke B, Perrin J, et al. Chemical Proteomics Reveals Ferrochelatase as a Common Off-target of Kinase Inhibitors. *ACS Chem Biol.* 2016;11(5):1245-1254. doi:10.1021/acscchembio.5b01063
45. Bloomer J, Bruzzone C, Zhu L, Scarlett Y, Magness S, Brenner D. Molecular defects in ferrochelatase in patients with protoporphyria requiring liver transplantation. *J Clin Invest.* 1998;102(1):107-114. doi:10.1172/JCI1347
46. Sprowl JA, Ong SS, Gibson AA, et al. A phosphotyrosine switch regulates organic cation transporters. *Nat Commun.* 2016;7:10880. doi:10.1038/ncomms10880
47. Kim JY, Jayne LA, Bai Y, et al. Ribociclib mitigates cisplatin-associated kidney injury through retinoblastoma-1 dependent mechanisms. *Biochem Pharmacol.* 2020;177:113939. doi:10.1016/j.bcp.2020.113939
48. Goldinger SM, Rinderknecht J, Dummer R, et al. A single-dose mass balance and metabolite-profiling study of vemurafenib in patients with metastatic melanoma. *Pharmacol Res Perspect.* 2015;3(2). doi:10.1002/prp2.113
49. Hatzivassiliou G, Song K, Yen I, et al. RAF inhibitors prime wild-type RAF to activate the MAPK pathway and enhance growth. *Nature.* 2010;464(7287):431-435. doi:10.1038/nature08833
50. Bonventre JV, Yang L. Cellular pathophysiology of ischemic acute kidney injury. *J Clin Invest.* 2011;121(11):4210-4221. doi:10.1172/JCI45161
51. Bhargava P, Schnellmann RG. Mitochondrial energetics in the kidney. *Nat Rev Nephrol.* 2017;13(10):629-646. doi:10.1038/nrneph.2017.107
52. Basile DP. The endothelial cell in ischemic acute kidney injury: implications for acute and chronic function. *Kidney Int.* 2007;72(2):151-156. doi:10.1038/sj.ki.5002312

53. Bullen A, Liu ZZ, Hepokoski M, Li Y, Singh P. Renal Oxygenation and Hemodynamics in Kidney Injury. *Nephron*. 2017;137(4):260-263. doi:10.1159/000477830
54. van den Meiracker AH, Danser AHJ. Mechanisms of Hypertension and Renal Injury During Vascular Endothelial Growth Factor Signaling Inhibition. *Hypertension*. 2016;68(1):17-23. doi:10.1161/HYPERTENSIONAHA.116.07618
55. Sieber J, Wieder N, Clark A, et al. GDC-0879, a BRAFV600E Inhibitor, Protects Kidney Podocytes from Death. *Cell Chem Biol*. 2018;25(2):175-184.e4. doi:10.1016/j.chembiol.2017.11.006
56. Bryer JS, Susztak K. Screening Drugs for Kidney Disease: Targeting the Podocyte. *Cell Chem Biol*. 2018;25(2):126-127. doi:10.1016/j.chembiol.2018.01.018
57. Tracz MJ, Alam J, Nath KA. Physiology and Pathophysiology of Heme: Implications for Kidney Disease. *JASN*. 2007;18(2):414-420. doi:10.1681/ASN.2006080894
58. Walker VJ, Agarwal A. Targeting Iron Homeostasis in Acute Kidney Injury. *Semin Nephrol*. 2016;36(1):62-70. doi:10.1016/j.semnephrol.2016.01.003
59. Scindia PhD Y, Leeds Md J, Swaminathan Md S. Iron Homeostasis in Healthy Kidney and its Role in Acute Kidney Injury. *Semin Nephrol*. 2019;39(1):76-84. doi:10.1016/j.semnephrol.2018.10.006
60. Izzedine H, El-Fekih RK, Perazella MA. The renal effects of ALK inhibitors. *Invest New Drugs*. 2016;34(5):643-649. doi:10.1007/s10637-016-0379-y
61. Kim JY, Bai Y, Jayne LA, et al. A kinome-wide screen identifies a CDKL5-SOX9 regulatory axis in epithelial cell death and kidney injury. *Nat Commun*. 2020;11(1):1924. doi:10.1038/s41467-020-15638-6
62. Pabla N, Dong G, Jiang M, et al. Inhibition of PKC δ reduces cisplatin-induced nephrotoxicity without blocking chemotherapeutic efficacy in mouse models of cancer. *J Clin Invest*. 2011;121(7):2709-2722. doi:10.1172/JCI45586

A **C** **E**

- Control siRNA
- Braf siRNA

B **D** **F**

- Control
- Braf KO

A **B** **C** **D** **E** **F** **G**

A **B** **C** **D** **E** **F** **G**

● Vehicle
● Vemurafenib

A **B** **C** **D** **E**