1 **Neutralization of SARS-CoV-2 VOC 501Y.V2 by human antisera elicited by both** 2 **inactivated BBIBP-CorV and recombinant dimeric RBD ZF2001 vaccines**

3

4 **Authors**

5 Baoying Huang^{1*}, Lianpan Dai^{2*}, Hui Wang³, Zhongyu Hu⁴, Xiaoming Yang³, Wenjie Tan^{1†},
6 George F. Gao^{1,2†}

7

8 **Affiliations**

9 ¹NHC Key Laboratory of Biosafety, National Institute for Viral Disease Control and
10 Prevention, Chinese Center for Disease Control and Prevention, Beijing, 102206, China

11 ²CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of
12 Microbiology, Chinese Academy of Sciences, Beijing, 100101, China.

13 ³Beijing Institute of Biological Products Company Limited, Beijing, China

14 ⁴National Institute for Food and Drug Control, Beijing, 100050, China

15 *These authors contributed equally to this work

16 †Corresponding author. Email: gaof@im.ac.cn (G.F.G.); tanwj@ivdc.chinacdc.cn (W.T.)

17

18

19

20

21

22

23

24 **Abstract**

25 Recently, the emerged and rapidly spreading SARS-CoV-2 variant of concern (VOC)
 26 501Y.V2 with 10 amino acids in spike protein were found to escape host immunity induced
 27 by infection or vaccination. Global concerns have been raised for its potential to affect
 28 vaccine efficacy. Here, we evaluated the neutralization activities of two vaccines developed in
 29 China against 501Y.V2. One is licensed inactivated vaccine BBIBP-CorV and the other one is
 30 recombinant dimeric receptor-binding domain (RBD) vaccine ZF2001. Encouragingly, both
 31 vaccines largely preserved neutralizing titres, with slightly reduction, against 501Y.V2
 32 authentic virus compare to their titres against both original SARS-CoV-2 and the currently
 33 circulating D614G virus. These data indicated that 501Y.V2 variant will not escape the
 34 immunity induced by vaccines targeting whole virus or RBD.

35

36

37

38

39

40

41

42

43

44

45

46 **Maintext**

47 The rollout of vaccines is the hope to control the coronavirus disease 2019 (COVID-19)
 48 pandemic and reboot economy and society(1). However, the recent emergence of severe acute
 49 respiratory syndrome coronavirus-2 (SARS-CoV-2) variants raised global concerns because
 50 of their enhanced transmission and potential viral escape of host immunity elicited by natural
 51 infection or vaccination. Variants containing D614G mutation in spike (S) protein was first
 52 reported from the middle of 2020, which significantly increased the transmission rate and
 53 became dominant in circulating strains ever since(2). Evolved from D614G mutants, recent
 54 circulating isolates from Republic of South Africa (501Y.V2 variant) introduced further
 55 mutations that escape neutralization by COVID-19 convalescent plasma and sera from human
 56 receiving licensed mRNA vaccines expressing SARS-CoV-2 S protein(3-5). It also
 57 dramatically decreased the protective efficacy for trimeric S protein-based vaccine in phase 3
 58 clinical trial in South Africa(6).

59 501Y.V2 variant emerged in more and more countries, and was first isolated in China on
 60 January 6, 2021 from an airline pilot of South African nationality(7). This variant, GDPCC
 61 strain, contains 10 amino acid mutations sites in S protein corresponded to the features of the
 62 variant of concern (VOC) South African 501Y.V2, with 5 (D80A, ΔL242, ΔA243, ΔL244 and
 63 R246I) located at N-terminal domain (NTD), 3 (K417N, E484K and N501Y) in

receptor-binding domain (RBD) , and the other two in CTD2 domain and S1/S2-S2' region (Fig.1).

The effectiveness of current vaccines against this VOC is of high importance to guide the ongoing vaccination program worldwide. To answer this question, we evaluated two representative COVID-19 vaccines developed in China for their neutralizing activities against 501Y.V2 variant. One is BBIBP-CorV, a licensed COVID-19 inactivated vaccine(8, 9). The other one is ZF2001, a protein subunit vaccine targeting S protein RBD currently in phase 3 clinical trials(10, 11). Both vaccines showed good immunogenic in their clinical trials(9, 11). For instance, ZF2001 induced neutralizing GMTs two times greater than that from convalescent samples(11). We chose 12 serum samples for each vaccine from clinical trial participants covering a range of different neutralizing titers (Table 1). The neutralizing activities of these serum samples against live SARS-CoV-2 strains GDPCC (501Y.V2) were measured by microcytopathogenic effect assay. SARS-CoV-2 strains HB02 (WT) (12, 13) and BJ01 (D614G) (14) were tested as the control. Impressively, all 12 serum samples from either ZF2001 or BBIBP-CorV recipients largely preserved neutralization of 501Y.V2 variant, with slightly reduced geometric mean titres (GMTs) compared with their titres against WT or D614G strain (Fig. 2 and Table 1). For ZF2001, the GMTs declined for 1.6-fold from 106.1 (95% CI, 75.0-150.1) to 66.6 (95% CI, 51.0-86.9) (Fig.2A). While for BBIBP-CorV, the decline is also 1.6-fold, GMTs from 110.9(95% CI, 76.7-160.2) to 70.9(95% CI, 50.8-98.8) (Fig.2B). These reductions are significantly less than those reported previously for convalescent plasma (more than 10-folds)(4) or antisera from mRNA vaccine recipients (more

than 6-folds) (3, 5).

These results suggest that the 501Y.V2 variant does not escape the immunity induced by vaccines targeting S protein RBD (ZF2001) or whole virus (BBIBP-CorV). The potential 1.6-fold reduction of neutralizing GMTs should be taken into account for its impact for the clinical efficacy of these vaccines. For both vaccines, antisera neutralize both variant 501Y.V2 and D614G, the one currently circulating globally, without statistical significances. For ZF2001, a slight significance ($P=0.04$) between variant 501Y.V2 and the HB02 may due to the sample selection and size. For the neutralization-reduction discrepancy between our protein-based vaccine and mRNA vaccine needs further investigation in the future.

Acknowledgements

This work was supported by the National Program on Key Research Project of China (2020YFA0907101, 2016YFD0500301), the National Natural Science Foundation of China (NSFC) (82041041, 82061138008), National Mega Projects of China for Major Infectious Diseases (2016ZX10004001-003). Lianpan Dai is supported by Youth Innovation Promotion Association of the CAS (2018113).

101

102

Fig.1. Schematic demonstration of the spike protein of SARS-CoV-2 HB02 (WT), BJ01

(D614G) and GDPCC (501Y.V2). The mutation sites were denoted and marked as cycles,

104

with the point mutation colored in red and the deletion colored in white.

Fig.2. Neutralization titres of 24 antisera from vaccine BBIBP-CorV or ZF2001 recipients against authentic SARS-CoV-2 and its variants, D614G and 501Y.V2. (A-B)

N=12 representative antisera each from ZF2001 (A) and BBIBP-CorV (B) vaccine recipients were tested for their neutralizing activity to authentic SARS-CoV-2 HB02 (WT), BJ01 (D614G) and GDPCC (501Y.V2) via microcytopathogenic effect assay. Shown are the geometric mean with 95% CI. P values were analyzed with One-way ANOVA test.

Table 1: Information for serum samples tested in this study

RBD protein vaccine (ZF2001)				
	50% Neutralization titer			
Serum ID	Clinical trials	HB02 (WT)	BJ01 (D614G)	GDPCC (501Y.V2)
012-6-(5)	128	128	96	64
014-6-(5)	256	256	128	128
024-6-(5)	64	64	48	48
082-6-(5)	96	64	64	32
088-6-(5)	96	48	96	48
102-6-(5)	256	128	128	96
135-6-(5)	384	192	192	96
153-6-(5)	128	128	96	64
154-6-(5)	64	64	64	48
233-6-(5)	256	128	128	96
290-6-(5)	128	64	48	48
308-6-(5)	256	192	128	96
Inactivated vaccine (BBIBP-CorV)				
	50% Neutralization titer			
Serum ID	Clinical trials	HB02 (WT)	BJ01 (D614G)	GDPCC (501Y.V2)
B867-2	192	128	128	96
B850-2	256	192	192	128
B848-2	64	48	64	32
B875-2	96	96	64	48
B841-2	256	192	128	64
B843-2	256	192	128	64
B844-2	192	128	128	96
B856-2	96	64	96	86
B851-2	192	128	96	64
B890-2	64	48	48	32
B845-2	384	256	256	192
B869-2	96	64	96	64

126

127

128 **References**

129

130 1. L. Dai, G. F. Gao, Viral targets for vaccines against COVID-19. *Nature Reviews Immunology*,
131 (2020). doi.org/10.1038/s41577-020-00480-0

132 2. B. Korber et al., Tracking changes in SARS-CoV-2 Spike: Evidence that D614G increases
133 infectivity of the COVID-19 Virus. *Cell* 182, 812-827 e819 (2020).

134 3. K. Wu et al., mRNA-1273 vaccine induces neutralizing antibodies against spike mutants
135 from global SARS-CoV-2 variants. *bioRxiv*, 2021.2001.2025.427948 (2021).

136 4. C. K. Wibmer et al., SARS-CoV-2 501Y.V2 escapes neutralization by South African
137 COVID-19 donor plasma. *bioRxiv*, 2021.2001.2018.427166 (2021).

138 5. P. Wang et al., Increased resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7 to antibody
139 neutralization. *bioRxiv*, 2021.2001.2025.428137 (2021).

140 6.

141 <https://ir.novavax.com/news-releases/news-release-details/novavax-covid-19-vaccine-demonstrates-893-efficacy-uk-phase-3>.
142

143 7. F. Chen et al. A case of new variant COVID-19 first emerging in South Africa detected in
144 airplane pilot — Guangdong Province, China, January 6, 2021. *China CDC Weekly* 3: 28-29
145 (2021).

146 8. H. Wang et al., Development of an inactivated vaccine candidate, BBIBP-CorV, with potent
147 protection against SARS-CoV-2. *Cell* 182, 713-721 e719 (2020).

148 9. S. Xia et al., Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine,
149 BBIBP-CorV: a randomised, double-blind, placebo-controlled, phase 1/2 trial. *Lancet Infect Dis*
150 21, 39-51 (2020).

151 10. L. Dai et al., A universal design of betacoronavirus vaccines against COVID-19, MERS, and
152 SARS. *Cell* 182, 722-733 e711 (2020).

153 11. S. Yang et al., Safety and immunogenicity of a recombinant tandem-repeat dimeric RBD
154 protein vaccine against COVID-19 in adults: pooled analysis of two randomized, double-blind,
155 placebo-controlled, phase 1 and 2 trials. *medRxiv*, 2020.2012.2020.20248602 (2020).

156 12. N. Zhu et al., A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J*
157 *Med* 382, 727-733 (2020).

158 13. W. Tan et al., A novel coronavirus genome identified in a cluster of pneumonia cases —
159 Wuhan, China 2019–2020. *China CDC Weekly* 2, 61-62 (2020).

160 14. W. Tan et al., Reemergent cases of COVID-19 — Xinfadi wholesales Market, Beijing
161 municipality, China, June 11, 2020. *China CDC Weekly* 2, 502-504 (2020).

162

163

164

165

166

167

168

169

170

171

172

173

Supplementary Materials for

174

175

Neutralization of SARS-CoV-2 VOC 501Y.V2 by human antisera

176

elicited by both inactivated BBIBP-CorV and recombinant dimeric

177

RBD ZF2001 vaccines

178

Baoying Huang*, Lianpan Dai*, Hui Wang, Zhongyu Hu, Wenjie Tan†, George F. Gao†

179

180

*These authors contributed equally to this work

181

†Corresponding author. Email: gaof@im.ac.cn (G.F.G.); tanwj@ivdc.chinacdc.cn (W.T.)

182

183

This PDF file includes:

|

184 Materials and Methods

185 Supplemental references

186

187

188

189

190 **Material and methods**

191 **Viruses and titration**

192 The SARS-CoV-2 virus 19nCoV-CDC-Tan-HB02 (short as HB02), 19nCoV-CDC-Tan-BJ01
193 (short as BJ01) and 19nCoV-CDC-Tan-GDPCC (short as GDPCC) were used in our
194 experiments(1-4). Virus titer was determined by a micro-cytopathogenic efficiency (CPE)
195 assay on Vero cells as described previously(5).

196 **Neutralization assay**

197 The serum were inactivated in a 56°C water bath for 30 min, then successively diluted 1:4 to
198 the required concentration by a 2-fold series. An equal volume of challenge virus solution
199 containing 100 CCID₅₀ virus was added. After neutralization in a 37°C incubator for 2 h, a
200 1.0~2.5×10⁵/ml cell suspension was added to the wells and cultured in a CO₂ incubator at 37°C
201 for 4 days. Titers expressed as the reciprocal of the highest dilution protecting 50% cell from
202 virus challenge. A neutralization antibody potency < 1:4 is negative, while that ≥1:4 is
203 positive.

204 **Serum samples from clinical trial participants**

205 For the 12 Serum samples from BBIBP-CorV vaccination(6, 7), vaccine recipients received
206 BBIBP-CorV containing 4 µg total protein on days 0 and 21, blood samples were taken from
207 participants for serology tests 28 days after second vaccination, covering a range of different
208 neutralizing titers, the ClinicalTrials.gov Identifier is NCT04510207.

209 For the 12 Serum samples from ZF2001 vaccination (8), vaccine recipients received ZF2001
210 containing 25 µg vaccine dose on days 0, 30, 60. Blood samples were taken from participants

for serology tests 14 days after third vaccination, covering a range of different neutralizing titers, the ClinicalTrials.gov Identifier is NCT04466085.

Supplemental references

1. N Zhu et al., A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 382,727-733 (2020).
2. W Tan et al., A novel coronavirus genome identified in a cluster of pneumonia cases — Wuhan, China 2019–2020. *China CDC Weekly* 2, 61-62 (2020).
3. W Tan et al., Reemergent cases of COVID-19 — Xinfadi wholesales Market, Beijing municipality, China, June 11, 2020. *China CDC Weekly* 2: 502-504 (2020).
4. F. Chen et al., A case of new variant COVID-19 first emerging in South Africa detected in airplane pilot — Guangdong province, China, January 6, 2021. *China CDC Weekly* 3, 28-29 (2021).
5. MA. Ramakrishnan. Determination of 50% endpoint titer using a simple formula. *World J Virol* 5, 85-86 (2021).
6. H. Wang et al., Development of an inactivated vaccine candidate, BBIBP-CorV, with potent protection against SARS-CoV-2. *Cell* 182, 713-721 e719 (2020).
7. S. Xia et al., Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBIBP-CorV: a randomised, double-blind, placebo-controlled, phase 1/2 trial. *Lancet Infect Dis*, 21, 39-51 (2020).
8. S. Yang et al., Safety and immunogenicity of a recombinant tandem-repeat dimeric RBD protein vaccine against COVID-19 in adults: pooled analysis of two randomized, double-blind, placebo-controlled, phase 1 and 2 trials. *medRxiv*, 2020.2012.2020.20248602 (2020).