Phenotypic and genotypic characterization of antibiotic Resistant Gram Negative Bacteria isolated in Tabuk City, Saudi Arabia

Tarig M.S. Alnour (Ph.D) * ^{a, b, c}; Elmutuz H. Elssaig (Ph.D) ^{a, c}; Eltayib H. Ahmed-Abakur (Ph.D) ^{a, b, c}; Faisel M. Abuduhier (Ph.D) ^{a, b}; Khalid A. S. Alfifi (MSc) ^d; Mohammad S. Abusuliman (MSc) ^e; and Tawfiq Albalawi (MSc) ^f

- a. Medical laboratory technology department, Faculty of Applied Medical Science, University of Tabuk - P.O. Box 741, Tabuk -Postal code 71411, Saudi Arabia
- b. Prince Fahad Research Chair, Department of Medical Laboratory Technology (FAMS), University of Tabuk- P.O. Box 741, Tabuk -Postal code 71411, Saudi Arabia
- c. Faculty of Medical Laboratory Science, Department of Microbiology and Immunology Alzaiem Alazhari University, Khartoum north Postal code 11111, Sudan
- d. Medical laboratory specialist- Supervisor of Microbiology Lab -King Fahd Specialist Hospital, Tabuk, Saudi Arabia khalfifi@moh.gov.sa
- e. Senior specialist -Lab medicine (Microbiology) -King Fahd Specialist Hospital, Tabuk, Saudi Arabia mabusuliman@moh.gov.sa
- f. Senior Specialist in Molecular Diagnostics, Department of laboratory and Blood Bank, King Fahad Specialist Hospital, Tabuk, Saudi Arabia talbalawi150@gmail.com

* Corresponding author:

Dr. Tarig M. S. Alnour

Faculty of Applied Medical Sciences

Department of Medical Laboratory Technology

University of Tabuk

E. mail Address: telnour@ut.edu.sa

Tel: +966535782141

Abstract: 31

Antimicrobial surveillance and identifying the genetic basis of antimicrobial resistance provide 32

important information to optimize patient care. The present study was analytical cross sectional 33

study aimed to determine the prevalence of MDR, XDR, PDR and extended-spectrum β - 34

lactamases genes (SHV, CTX-M and TEM) among Gram-negative bacteria isolated in Tabuk, 35

Saudi Arabia. A total number of 386 non-duplicate Gram-negative isolate were collected. 36

Identification and susceptibility testing were done using automation system (BD Phoenix™). 37

The extracted DNA were subjected to multiplex polymerase chain reaction (PCR). The results 38

showed that only 15 (3.9%) of isolates were fully susceptible, the overall prevalence of XDR, 39

MDR, PDR was 129 (33.4%), 113 (29.3%) and 48(12.4%) respectively. High resistant rate was 40

observed against the antibiotic agents of cephalosporins class 79.3% followed by the agents of 41

penicillins class 69.4%. The most dominant gene was bla SHV which detected in 106/386 42

(27.5%) isolates followed by bla CTX-M 90/386 (23.3%). Bla CTX-M showed significant 43

relation with all used antibiotic except ampicillin/clavulanic acid, aztreonam, cefoxitin, and 44

meropene. The isolates which showed most frequent resistant genes were *Klebsiella pneumoniae* 45

90/124 (72.6%), *A. baumannii* 37/67 (55.2%), and *P.mirabilis* 24/44 (54.5%). These findings 46

underscores the need for optimization of current therapies and prevention of the spread of these 47

organisms. 48

49

50

51

52

53

54

55

56

57

58

59

60

61

Key word: MDR, XDR, PDR, ESBLs, bla SHV, bla TEM, bla CTX-M 62

1. Introduction: 63

Antimicrobial resistance is described as a condition in which the pathogen escape from the stress 64
of the antibiotic exposure (1). The increasing incidence of antimicrobial resistance is the key 65
concern globally and considered main obstacle in the treatment of patients suffering from 66
bacterial infections.(1, 2) It has been estimated that about 1.3 to 2 fold rise in mortality caused 67
by antimicrobial resistant bacteria compared to susceptible infections. (1). 68

A dramatic evolution has occurred in the significance of infections caused by Gram-negative 69
bacteria (GNB) and associated with considerable mortality (1, 2, 3). The efficiency of the current 70
prophylactic and empiric antibiotic treatment is compromised by the emergence of pan drug 71
resistant (PDR), extensively drug resistant (XDR) and multidrug resistant (MDR) Gram-negative 72
bacteria (GNB) (2, 3). The ability of escaping from the antimicrobial effects may be contributed 73
to the nature of these organisms, which are heterogeneous, complex group of plasmids-borne and 74
rapidly evolving enzymes, which are capable of hydrolyzing cephalosporins, penicillins, 75
aztreonam and monobactams (4). 76

The American society of infectious diseases identified six top priority dangerous pathogens 77
producing extended spectrum β -lactamases (ESBLs). Three of these six pathogens are antibiotic 78
resistant Gram-negative bacteria; *Pseudomonas aeruginosa*, *Acinetobacter baumannii* and 79
Enterobacteriaceae (5). ESBLs have been classified into three major groups; bla SHV, bla CTX- 80
M and bla TEM (2), Ting et al., 2013 stated that bla TEM, bla SHV and bla CTX-M genes are 81
super-resistant extended spectrum β -lactamases (6). In 2017, the WHO published list of global 82
priority pathogens, a catalog of twelve species of bacteria grouped according to their antibiotic 83
resistance under three priority tiers; critical, high, and medium. The critical group involved three 84
pathogens that were Gram-negative bacilli; *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, 85
and *Enterobacteriaceae*, (7). The WHO also notified that the level of resistance to antimicrobial 86
drugs used to treat common infections is reaching a crisis point. If world administrations do not 87
control infections in order to slow down the growth of drug resistance, entire populations could 88
be wiped out by superbugs (8, 9). However, the availability of regional information on the 89
resistance rate is fundamental to implementing efficient treatment protocols against infectious 90
pathogens and may help to prevent infections with multidrug resistance pathogens at the local 91
level (10, 11). Therefore, the present study was aimed to determine pattern of antimicrobial 92

resistance and to detect ESBL genes among Gram-negative bacteria isolated in Tabuk city, Saudi Arabia. 93 94

2. Materials and Methods: 95

The present study was analytical cross sectional study, conducted in King Fahad Specialist Hospital and prince Fahad Bin Sultan Research Chair (University of Tabuk), Saudi Arabia. A total number of 386 non-duplicate Gram-negative isolates were collected in order to determine the prevalence of MDR, XDR, PDR and to detect extended-spectrum β -lactamases genes bla SHV, bla CTX-M and bla TEM. 96 97 98 99 100 101

Identification and susceptibility test:

Depending on the origin of the samples, each sample cultured on suitable medium/ media from: MacConkey agar, CLED agar blood agar, Chocolate agar or Brain Heart infusion broth. Then they were incubated aerobically at 37°C for 24 to 48 hours except for the blood culture, which was incubated for 5 to 7 days in broth medium. Growth of corresponding organisms were further sub-cultured for purification purpose. The significant growth was identified to the species level. Identification and susceptibility testing were done using automation system (BD Phoenix™). Identified strains were tested in vitro against several antimicrobial classes including carboxypenicillin (Ticarcillin/Clavulanic acid), Penicillinase resistant penicillin (Ampicillin/Sulbactam and Piperacillin/Tazobactam), Cephalosporins (Ceftazidime and Cefepime), Aztreonam, Carbapenems (Ertapenem, Imipenem and Meropenem), Aminoglycoside (Amikacin, Gentamicin and Tobramycin), Fluoroquinolones (Ciprofloxacin and levofloxacin), Minocycline (Tetracyclin), Glycylcycline (Tigecycline), polymyxin E (Colistin), and sulpha drugs (Trimethoprim/Sulfamethoxazole). Antimicrobial selection for testing depends on types of isolates and site of samples which done automatically by the program. Based on susceptibility test of the above mentioned antibiotic classes the isolates were characterized as MDR, XDR and PDR. 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117

Moreover thirteen agents of five antimicrobial classes which represented the most commonly used antibiotics; carboxypenicillin, penicillinase resistant penicillin, cephalosporins, aztreonam and carbapenems were subjected for further study in order to determine the relation between these antibiotics and ESBL genes, these agents included: ampicillin, ampicillin/clavulanic acid, ticarcillin/clavulanic acid, aztreonam, piperacillin/tazobactam, cefalotin, cefoxitin, ceftazidime, 118 119 120 121 122

ceftriaxone, cefepime, imipenem, meropenem and ertapenem. Quality control and maintenance were achieved according to the manufacturer's guidelines.

The BD Phoenix™ automated identification and susceptibility testing system empowers workflow efficiency using automated nephelometry, which results in a standardized isolate inoculum and a reduction in potential technologist error along with accurate, reliable and rapid detection of known and emerging antimicrobial resistance (12).

Detection of Antimicrobial Resistance Genes

DNA was extracted from whole 386 GNB isolates using boiling technique; few colonies from each isolate were mixed with molecular biology-grade water (Eppendorf, Hamburg, Germany), the mixture were centrifuged at $15,000 \times g$ for 5 min. The supernatant was discharged and the pellet was re-suspended in molecular biology-grade water (Eppendorf, Hamburg, Germany) and subjected to boiling at 100°C in a water bath for 20 min, then cooled and centrifuged at $15,000 \times g$ for 60s before it was stored at -20°C .

Multiplex polymerase chain reaction (PCR) were done to determine the presence of three ESBLs genes encoding bla TEM, bla SHV, bla CTX-M. Each extracted DNA was tested against the three sets of specific primers in a single test (Table 1). Amplification was performed in a final volume of 25 μL containing 5 μL of template DNA, 0.5 μL Taq polymerase, 1.0 μL of each primers and 0.2 μL dNTP mixture (10 mM), and finally the volume was completed to 25 μL by molecular biology-grade to reach volume of 25 μL . The PCR protocol was run according to the following protocol; essential denaturation at 95°C for 5 minutes followed by 40 cycle of denaturation at 95°C for 30s, annealing at 60°C for 30s and extension at 72°C for 1 minutes, the final step was extension at 72°C for 5 minutes. PCR product was run on 2% ethidium bromide agarose gel electrophoresis, and examined with gel imaging system, bands pattern was observed and interpreted according to their size (Table 1).

Analysis: The proportion of resistant for each antibiotic was calculated as the sum of resistant antibiotic relative to the sum of susceptible and resistant. The proportion of resistant class of antimicrobial was represent the mean of resistance of all antimicrobial agents belong to that class. Chi-square tests was performed to determine the relation between ESBLs genes and antibiotic resistant using SPSS version 22. P value <0.05 was considered significant.

The isolates which showed susceptibility to all groups of antimicrobial agents were classified as susceptible, while those showed resistant to one group were classified as mono-resistant. The

isolates resistant to one drugs in two groups were classified as resistant to two antimicrobial group. Multidrug resistant (MDR) denoted when the isolates shown resistance to 3 or more antimicrobial group but susceptible to 2 or more. The US Centers for Disease Control and Prevention (CDC) and European Centre for Disease Prevention and Control (ECDC) have defined bacteria as pan drug resistant (PDR) when they are non-susceptible to all agents in all antimicrobial categories and as extensively drug-resistant (XDR) when they are non-susceptible to at least one agent in all but two or fewer antimicrobial categories (13).

The ethical clearance for this study (UT-86-10-2019) was obtained from research ethics committee, University of Tabuk (Saudi Arabia).

Result:

Table no 2 showed the frequency of isolates along with patterns of occurrence of resistant ESBLs genes; the results revealed that the most common isolates were *Klebsiella pneumoniae* 124(32.1%), followed by *A. baumannii* 67(17.4%), *E.coli* 51(13.2%), *P.aeruginosa* 50(13.0%) and *P.mirabilis* 44(11.2%).

The isolates showed high resistant rate against the antimicrobial agents of cephalosporins group 79.3% followed by the agents of penicillins 69.4% while against the agents of carbapenems they exhibited 32.5 % resistance rate. In term of individual antimicrobial agent; the isolated Gram-negative bacteria revealed a high resistance rate against ampicillin 262(93.2%) followed by aztreonam 87(90.6%) and cefalotin 157(90.2%). Only 15 (3.9%) of isolates were fully susceptible to all used antimicrobials. The overall prevalence of MDR, XDR, PDR was 113 (29.3%), 129 (33.4%) and 48(12.4%) respectively (Table no 3).

Screening for resistance genes showed that, most of Gram-negative isolates harbored the resistance genes 198/386(51.3%) while isolates free from resistant genes were 188 (48.7%) (Table 2). Bla SHV was most dominant gene which detected in 106/386 (27.5%) isolates followed by bla CTX-M and bla TEM which detected in 90/386 (23.3%) and 78/386 (20.2%) isolates, respectively. Single resistant gene was detected in 137/386 (35.5%) isolates, coexistence of two genes were detected in 41/386 (10.6%) isolates while triple genes were present in 20 (5.2%) isolates. Bla CTX-M showed significant relation with all used antibiotic except ampicillin/clavulanic acid, aztreonam and ceftazidime while bla SHV exhibited significant statistic relationship with Piperacillin/ Tazobactam, Cefalotin, ceftazidime, cefepime, imipenem and

meropenem. Bla TEM displayed significant relation only to ampicillin/clavulanic acid, ceftazidime, cefepime and imipenem. Ceftazidime and cefepime agents of cephalosporins class were least effective agents as they showed significant relation to the three resistant genes (Table 4). The isolates which showed the most frequent resistant genes were *K. pneumoniae* 90/124 (72.6%), *A. baumannii* 37/67 (55.2%), *E. coli* 22/51 (43.1%), *P. aeruginosa* 8/42(19.0%) and *P. mirabilis* 24/44 (54.5%) (Table 2).

Discussion:

Bacteria resistant to different classes of antimicrobial agents are a main threat to humanity and high risk, which may return the world towards pre-antimicrobial era (13). While active surveillance systems are set up in many countries in Europe, USA and Asia, little is reported on antimicrobial resistance status among Gram-negative bacteria in the Middle East, Africa and Saudi Arabia (14). The present study determined the prevalence of MDR, XDR, PDR and extended-spectrum β -lactamases genes (TEM, SHV, CTX-M) among Gram-negative bacteria in Saudi Arabia.

The third generation cephalosporin such as ceftazidime and cefoperazone marked by stability to the common beta-lactamases of Gram-negative bacilli and these compounds are highly active against Enterobacteriaceae (15). Although the isolated Gram-negative bacilli in this study showed high resistant rate to the antibiotic agents of cephalosporins class 79.3% followed by the agents of penicillinase resistant penicillins which showed 69.4% resistant, while the agents of carbapenems had least resistant rate; the highest resistance rate was reported against ampicillin (93.2%) followed by aztreonam (90.6%) and cefalotin 157(90.2%). Similar trend of resistance was observed by Ruppé et al 2015 who owing the dramatic increase in the rates of resistance to third-generation cephalosporins to spread of plasmid-borne extended spectrum beta-lactamases (ESBLs) in *Enterobacteriaceae* and to occurrence of sequential chromosomal mutations, which may lead to the overproduction of intrinsic beta-lactamases, hyper-expression of efflux pumps, target modifications and permeability alterations in non-fermenting Gram-negative bacteria (16). The serious finding in our study was emerging of carbapenems resistant. Carbapenems agents considered the most active and potent agents against multidrug-resistant (MDR) (17), this finding is totally contradictory to that reported by Zaman et al., 2015 who determined the susceptibility pattern of Gram-negative bacilli isolated from a Teaching Hospital in Jeddah,

Saudi Arabia and reported (100%) sensitive of enterobacteriaceae to carbapenems (18). However
WHO have been listed carbapenem-resistant *Enterobacteriaceae* among the top tier of the
antibiotic resistant that pose the greatest threat to human health (7).
The overall prevalence of MDR, XDR and PDR in this study was 29.3%, 33.4%, and 12.4%
respectively. Several studies were conducted in Saudi Arabia and showed high resistance rate
among Gram-negative bacteria, most of these studies focused on susceptibility per individual
pathogen (1, 9, 11, 14 , 19, 20) However, Ibrahim 2018, reported higher rate of MDR in
Southwest Saudi Arabia (67.9%) (11). the present study showed high PDR and less XDR
compared to that reported by Mohapatra et al 2018 (13). Increasing antimicrobial resistant in
Saudi Arabia may also be owing to increased cross geographic transmission of drug resistant.
Saudi Arabia is capital of Islamic world and has great number of expatriates, which makes it a
potential center for the import and export of multi resistant strains (21), a recent study by
Leangapichart et al (2016) showed that returned travelers from Hajj had acquired MDR *A. baumannii* and NDM producing *E. coli* during the Hajj event (22).
Our findings showed that more than half (51.3%) of isolated Gram-negative harbored with
resistant genes while isolates free from resistant genes were 48.7%. Only 3.9% of isolates were
fully susceptible to the used antimicrobials. This finding in alignment with Munita and Arias
report (2016) (23) and with Patil et al (2019) results, who showed increase in the resistant rate
among gram-negative pathogen and stated that Gram-negative bacteria are continuously evolving
mechanisms to deactivate clinically important antimicrobial drugs by acquisition of resistance
elements such as bla SHV, bla TEM and bla CTX-M (2). However antimicrobial resistant is an
outcome of multifaceted microbial interactions such as microbial characteristics to gain
resistance genes, selective pressure owing to inappropriate use and widespread of antibiotics,
resistance may arise by the acquisition of de-novo mutation during treatment or by acquisition of
integrative or replicative mobile genetic elements that have evolved over time in microbes in the
natural ecosystem (23).
Our results indicated that bla SHV was most prevalent resistant gene which detected in (27.5%)
of isolates followed by bla CTX-M (23.3%). Similar results indicated bla CTX-M and bla SHV
as the most prevalent genotypes of ESBLs producing gram-negative pathogens were reported in
several countries (2, 15, 24). Asokan et al (2019), reported that the bla CTX-M gene indicating
bacterial evolution due to cover prescription or weak enforcement of existing antibiotics policies

(25). The present study showed that *Klebsiella pneumoniae* (72.6%), *A. baumannii* (55.2%), *P.mirabilis* (54.5%) *E.coli* (43.1%), and *P.aeruginosa* (19.0%), were the highest isolates harbored with of ESBLs genes, these findings were in alignment with Maina et al 2012 (15), Ibrahim 2018 (11) and Asokan et al (2019) (25). However Enterobacteriaceae such as *K. pneumoniae*, *E. coli*, *Proteus spp*, *P. aeruginosa*, were naturally competent and can uptake naked DNA from the environment in suitable conditions (2).

In our study bla CTX-M showed significant association to all used antibiotic agents of cephalosporins, this finding is in alignment with Maina et al (2102), who stated that CTX-M type extended-spectrum β -lactamases (ESBLs) showing resistance to third and fourth-generation cephalosporins and to aztreonam (15).

Conclusion: infections caused by multi-resistant Gram-negative pathogens negatively influence patient outcomes and costs. This study showed that only 3.9% of isolates had susceptibility to all used antibiotics, high resistant rate was observed against the antimicrobial agents of cephalosporins class and penicillinase resistant penicillin. The most dominant gene was bla SHV.

Acknowledgments:

The authors would like to thank deanship of scientific research (DSR) University of Tabuk, Kingdom of Saudi Arabia for sponsoring the current research (project number S-1440-0339). We are very grateful for the Microbiology staff of King Fahad Specialist Hospital for their valuable efforts in sample collection. Thanks are also offered to Prince Fahad Bin Sultan Research Chair and Medical laboratory Technology staff, faculty of medical applied sciences, Tabuk University for their valuable support.

Conflicts of interest:

There are no conflicts of interest

References:

1. Alam, M, Z., Alam, Q., Jiman-Fatani, A, A., shukri,H, A., Haque, A. A surveillance study on the prevalence and antimicrobial resistance pattern among different groups of bacteria isolated from Western province of Saudi Arabia. Biomed Res India. 2017; 28 (2): 898-906.
2. 23Patil S, Chen H , Zhang X , Lian M, Ren P, Wen F. Antimicrobial Resistance and Resistance Determinant Insights into Multi-Drug Resistant Gram-Negative Bacteria Isolates from Paediatric Patients in China. Infect Drug Resist. 2019; 22;12:3625-3634.

	278
3. Paterson, D, L. Impact of Antibiotic Resistance in Gram-Negative Bacilli on Empirical and Definitive Antibiotic Therapy. <i>Clinical Infectious Diseases</i> CID 2008; 47(1): S14-S20	279 280
4. Fernando, M.M.P.S.C., Luke, W.A.N.V., Miththinda, J.K.N.D. <i>et al.</i> Extended spectrum beta lactamase producing organisms causing urinary tract infections in Sri Lanka and their antibiotic susceptibility pattern –A hospital based cross sectional study. <i>BMC Infect Dis</i> 17 , 138 (2017). https://doi.org/10.1186/s12879-017-2250-y	281 282 283 284
5. Lee, J, H., Jeong, S, H. and Lee, S, C. A lack of drugs for antibioticresistant Gram-negative bacteria. <i>Nature Reviews</i> . 2007.	285 286
6. Ting C., Jun A., Shun Z. Detection of the common resistance genes in Gram-negative bacteria using gene chip technology. <i>Indian Journal of Medical Microbiology</i> , (2013) 31(2): 142-147	287 288
7. WHO. Media Centre. News Release. WHO publishes list of bacteria for which new antibiotics are urgently needed. 2017 Available from: https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed (ACCESSED 8.12.2020)	289 290 291 292
8. Perez, F; Adachi, J; Bonomo, R A. Antibiotic-Resistant Gram-Negative Bacterial Infections in Patients with Cancer	293 294
9. Ibrahim, M K; Galal, A M; Al-Turk, I M; Al-Zhrany , K D; Antibiotic resistance in Gram-negative pathogenic bacteria in hospitals' drain in Al-Madina Al-Munnawara. <i>JTUSCI</i> 3: 14-22 (2010)	295 296 297
10. Moolchandani K, Sastry AS, Deepashree R, Sistla S, Harish BN, Mandal J. Antimicrobial Resistance Surveillance among Intensive Care Units of a Tertiary Care Hospital in Southern India. <i>J Clin Diagn Res</i> . 2017 11(2):	298 299 300
11. Ibrahim ME. High antimicrobial resistant rates among Gram-negative pathogens in intensive care units. A retrospective study at a tertiary care hospital in Southwest Saudi Arabia. <i>Saudi Med J</i> . 2018; 39(10):1035-1043. doi:10.15537/smj.2018.10.22944.	301 302 303
12. Evaluation of the BD Phoenix Automated Microbiology System for Identification and Antimicrobial Susceptibility Testing of Enterobacteriaceae	304 305

13. Mohapatra DP, Debata NK, Singh SK. Extensively drug-resistant and pan drug-resistant Gram-negative bacteria in a tertiary-care hospital in Eastern India: A 4-year retrospective study. *Journal of Global Antimicrobial Resistance*, 23 Aug 2018, 15:246-249.
14. 9 Zowawi, H, M. Antimicrobial resistance in Saudi Arabia An urgent call for an immediate action. *Saudi Med J*. 2016; 37 (9):935-940.
15. Maina D, Revathi G, Kariuki S, Ozwara H. Genotypes and cephalosporin susceptibility in extended-spectrum beta-lactamase producing enterobacteriaceae in the community. *J Infect Dev Ctries* 2012; 6(6):470-477.
16. Ruppé, É., Woerther, PL. & Barbier, F. Mechanisms of antimicrobial resistance in Gram-negative bacilli. *Ann. Intensive Care* **5**, 21 (2015). <https://doi.org/10.1186/s13613-015-0061-0>
17. Sheu C, Chang Y, Lin S, Chen Y, and Hsueh P. Infections Caused by Carbapenem-Resistant *Enterobacteriaceae*: An Update on Therapeutic Options. *Front. Microbiol*. 2019 | <https://doi.org/10.3389/fmicb.2019.00080>
18. Zaman R, Aly MM, Helmi N R. Antimicrobial susceptibility pattern of Gram-negative bacilli isolated from a Teaching Hospital in Jeddah, Saudi Arabia. 2015; *AJMR* 9(41), pp. 2145-2158, 14
19. Al Yousef S A. Surveillance of Antibiotic-Resistant Bacteria in King Khalid Hospital, Hafr Al-Batin, Saudi Arabia, During 2013. *Jundishapur J Microbiol*. 2016 September; 9(9):e19552.
20. El-Saed A, Balkhy H H, Alshamrani M M, Aljohani S , Alsaedi A , Al Nasser W , Gammal A E , Almohrij S A , Alyousef Z , Almunif S, Alzahrani M. High contribution and impact of resistant gram negative pathogens causing surgical site infections at a multi-hospital healthcare system in Saudi Arabia, *BMC infectious diseases* 2007–2016. (2020) 20:275
21. Ahmed-Abakur EH, Alnour TM. Detection of multidrug resistant *Mycobacterium tuberculosis* in Tabuk, Saudi Arabia, using genotype MTBDRplus. *Int J Mycobacteriol*. 2019; 8:25-8
22. Leangapichart T, Gautret P, Griffiths K, Belhouchat K, Memish Z, Raoult D, et al. Acquisition of a high diversity of bacteria during Hajj pilgrimage, including *Acinetobacter baumannii* with blaOXA-72, and *Escherichia coli* with blaNDM-5 carbapenemases. *Antimicrob Agents Chemother* 2016. pii: AAC.00669-16.

23. Munita JM, Arias CA. Mechanisms of antibiotic resistance. Microbiol Spectr. 2016;4 (2):1–	335
37. doi:10.1128/microbiolspec.VMBF-0016–2015	336
24. Tian G, Garcia J, Adams-Haduch JM, Evangelista JP, Destura RV, Wang H and Doi Y.	337
CTX-M as the predominant extended-spectrum beta-lactamases among Enterobacteriaceae in	338
Manila, Philippines. J Antimicrob Chemother. 2010; 65: 584-586.	339
25. Asokan G V., Ramadhan T., Ahmed E, Sanad H. WHO Global Priority Pathogens List: A	340
Bibliometric Analysis of MedlinePubMed for Knowledge Mobilization to Infection Prevention	341
and Control Practices in Bahrain.2019; Oman Medical Journal [2019], Vol. 34, No. 3: 184-193.	342
	343
	344
	345
	346
	347
	348
	349
	350
	351
	352
	353
	354
	355
	356
	357
	358

Table 1: the primers sequences of ESBLs genes and corresponding size

Gene	Primer sequences	Size
blaTEM	F-5 -TCGTGTCGCCCTTATTCCCTTTTT-3 R-5-GCGGTTAGCTCCTCC GGTCTC-3	426
blaSHV	F 5-GTGGATGC CGGTGACGAACAGC-3 R 5 -TGGCGCAAAAA GGCAGTCAATCCT-3	212
blaCTX-M	5'TTTGCGATGTGCAGTACCAGTAA3' 5'CGATATCGTTGGTGGTGCCATA3'	619

Isolate	Frequency of isolate	Occurrence of resistant encoding gene				
		none	Patten of occurrence			
			Single gene	Two genes	Three genes	Total
<i>A. baumannii</i>	67/386 (17.4%)	30 (44.8%)	31(46.3%)	4 (6.0%)	2(3.0%)	37/67(55.2%)
<i>C. freundii</i>	3/386 (0.78%)	2 (66.7%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	1/3 (33.3%)
<i>C. youngae</i>	1/386 (0.26%)	1 (100%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
<i>comamonas testosteroni</i>	1/386 (0.26%)	0 (0.0%)	1 (100%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
<i>E. cloacae</i>	7/386 (1.8%)	5 (71.4%)	2 (28.6%)	0 (0.0%)	0 (0.0%)	2/7 (28.6%)
<i>E. coli</i>	51/386 (13.2%)	29 (56.9%)	15 (29.4%)	6(11.8%)	1 (1.2%)	22/51(43.1%)
<i>K. pneumoniae</i>	124/386(32.1%)	34 (27.4%)	45 (36.3%)	29 (23.4%)	16 (12.9%)	90/124(72.6%)
<i>M. morgani</i>	12/386 (3.1%)	9 (75.0%)	3 (25.0%)	0 (0.0%)	0 (0.0%)	3/12 (25.2%)
<i>P. aeruginosa</i>	50/386 (13.0%)	42 (84.0%)	7(14.0%)	1 (2.0%)	0 (0.0%)	8/50 (16.0%)
<i>P. hauseri</i>	2/386 (0.52%)	2 (100%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
<i>P. mirabilis</i>	44v (11.2%)	20 (45.5%)	22 (50.0%)	1 (2.3%)	1 (2.3%)	24/44 (54.5%)
<i>P. stuartii</i>	16/386 (4.1%)	9 (56.3%)	7 (43.8%)	0 (0.0%)	0 (0.0%)	7/16(43.8%)
<i>S. marcescens</i>	8/386 (2.1%)	5 (62.5%)	3 (37.5%)	0 (0.0%)	0 (0.0%)	3/8(37.5%)

Table 2: Frequency of isolates and patterns of occurrence of resistant ESBLs genes

Total	386	188	137	41	20	386 (100%)
-------	-----	-----	-----	----	----	------------

395

396

397

398

399

Table 3: Phenotypic characterization of antibiotic resistance among Gram-negative bacteria

Phenotypic characterization of isolate	Frequency	Percentage
Mono-resistant	28	7.2
Resist to 2 antimicrobial group	53	13.7
MDR	113	29.3
XDR	129	33.4
PDR	48	12.4
Fully Sensitive	15	3.9
Total	386	100%

410

Beta lactam groups	Antimicrobials	Resistance Nr (%)	Genes		
			blaSHV Nr (%)	blaTEM Nr (%)	blaCTX-M Nr (%)
Penicillinase resistant penicillins and carboxypenicillin	Ampicillin	262/282 (93.2)	92/262 (35.1)	60/262 (22.9)	78/262 (29.8) <i>P</i> = 0.005
	Ampicillin/ clavulanic acid	78/141 (55.3)	46/78 (58.9)	21/78 (26.9) <i>P</i> = 0.022	32/78 (41.0)
	Ticarcillin/Clavulanic acid	96/161 (59.6)	24/96 (25)	18/96 (18.8)	21/96 (21.9) <i>P</i> = 0.003
Aztreonam	Aztreonam	87/96 (90.6)	24/87 (27.6)	23/87 (26.4)	11/87 (12.6)
Monobactam / sulbactam	Piperacillin/ Tazobactam	168/249 (67.5)	59/168 (35.1) <i>P</i> = 0.006	41/168 (24.4)	58/168 (34.5) <i>P</i> = 0.000
Cephalosporins	Cefalotin	157/174 (90.2)	61/157 (38.9) <i>P</i> = 0.043	37/157 (23.6)	53/157 (33.8) <i>P</i> = 0.017
	Cefoxitin	88/174 (50.6)	42/88 (47.7)	22/88 (25)	29/88 (33)
	Ceftazidime	325/386 (84.2)	98/325 (30.2) <i>P</i> = 0.016	75/325 (23.1) <i>P</i> = 0.002	87/325 (26.8) <i>P</i> = 0.002
	Ceftriaxone	185/210 (88.1)	64/185 (34.6)	46/185 (24.9)	64/185 (34.6) <i>P</i> = 0.021
	Cefepime	322/386 (83.4)	99/322 (30.7) <i>P</i> = 0.014	75/322 (23.3) <i>P</i> = 0.001	88/322 (27.3) <i>P</i> = 0.001
Carbapenems	Imipenem	201/375 (53.6)	56/201 (27.9) <i>P</i> = 0.003	52/201 (25.9) <i>P</i> = 0.016	51/201 (25.4) <i>P</i> = 0.020
	Meropenem	158/381 (41.5)	55/158 (34.8) <i>P</i> = 0.042	33/158 (20.9)	47/158 (29.7) <i>P</i> = 0.032
	Ertapenem	2/80 (2.5)	0/2 (0)	1/2 (50)	0/2 (0)

411

412

413

414

415

416

417

Table 4: Phenotypic resistance along with frequency of ESBLs genes

418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437