Artificial light at night leads to circadian disruption in a songbird:

integrated evidence from behavioural, genomic and metabolomic data

Davide M. Dominoni^{1,2,*}, Maaïke de Jong^{2,3}, Kees van Oers², Peter O'Shaughnessy¹, Gavin Blackburn⁴, Els Atema², Christa A. Mateman², Pietro B. D'Amelio^{5,6,7}, Lisa Trost⁵, Michelle Bellingham¹, Jessica Clark¹, Marcel E. Visser^{2,8}, Barbara Helm^{1,8}

¹ Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, University Avenue, Glasgow, G12 8QQ UK

² Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, The Netherlands

³ Plant Ecology and Nature Conservation Group, Wageningen University & Research, Wageningen, The Netherlands

⁴ Glasgow Polyomics, Wolfson Wohl Cancer Research Centre, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G61 1BD, UK

⁵ Department of Behavioural Neurobiology, Max Planck Institute for Ornithology, Seewiesen, Germany

⁶ FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch 7701, South Africa

⁷ Centre d'Ecologie Fonctionnelle et Evolutive, University of Montpellier, CNRS, EPHE, IRD, Univ Paul-Valéry Montpellier 3, Montpellier, France

⁸ Groningen Institute of Evolutionary Life Sciences (GELIFES), University of Groningen,
Nijenborgh 7, 9747 AG Groningen, The Netherlands

*Correspondence: Davide M. Dominoni, davide.dominoni@glasgow.ac.uk

Abstract

Globally increasing levels of artificial light at night (ALAN) are associated with shifts in circadian rhythms of behaviour in many wild species. However, it is still unclear whether changes in behavioural timing are underlined by parallel shifts in the molecular clock, and whether such internal shifts may differ between different tissues and physiological pathways, which could highlight circadian disruption. We tackled these questions in a comprehensive study that integrated behavioural, gene expression and metabolomic analyses. We exposed captive male great tits (*Parus major*) to three ALAN intensities or to dark nights, recorded their activity rhythms and obtained mid-day and midnight samples of brain, liver, spleen and blood. ALAN advanced wake-up time, and this shift was paralleled by an advance in the expression of the clock gene *BMAL1* in all tissues, suggesting close links of brain and peripheral clock gene expression with activity rhythms. However, several metabolic and immune genes were desynchronised the shifted *BMAL1* expression, suggesting circadian disruption of behaviour and physiology. This result was reinforced by untargeted metabolomic profiling, which showed that only 9.7% of the 755 analysed metabolites followed the behavioural shift. We suggest circadian as a key mediator of the health impacts of ALAN on wild animals.

Introduction

On our rhythmic planet, organisms have adapted to the change of day and night by evolving circadian rhythms that are highly sensitive to light [1]. The near-ubiquity of circadian rhythms across kingdoms of life suggests major fitness benefits on two grounds. Internally, the circadian system regulates temporal coordination within the body to reduce conflict and overlap between different processes. Externally, the circadian system anticipates environmental fluctuations, enabling organisms to align their behavior and physiology with nature's cycles [1,2], such as the daily alternation of light and darkness. However, globally most humans and wild organisms in their vicinity are now exposed to artificial light at night (ALAN), and thus to a rapidly altered light environment [3,4] that threatens the refined functioning of the circadian system.

In animals, rhythmicity is primarily generated on a molecular level by a transcription-translation feed-back loop (TTFL). This rhythmicity is modulated by multiple interacting systems, including neuronal, endocrine, metabolic and immune pathways [5,6][7]. The orchestration of these processes involves complex interactions between sensory input, central and peripheral clocks, and effector systems [2]. There is increasing evidence that ALAN disrupts these processes, with possible consequences ranging from compromised human health to loss of ecosystem functions [8–10]. In free-living and captive organisms, altered daily and annual activity has been widely reported, and experimental illumination has confirmed causal effects of ALAN [11,12]. Still, it is largely unclear whether the circadian system, its multiple components, and the physiological pathways it coordinates, remain synchronized with activity patterns [13–18]. ALAN has also been shown to induce physiological changes, including in endocrine, immune and metabolic pathways [15,19,20]. These changes could be due to circadian disruption, with possible negative consequences for fitness [9,21]. Addressing these issues requires multi-level

analyses that simultaneously examine effects of ALAN on rhythmic behavior and different physiological pathways [9], but these are currently lacking.

Here we aim to fill this gap by an integrated study of a bird, the great tit (*Parus major*), whose behavioral response to ALAN is well-characterized [11,22–26]. We measured day-night differences in gene transcripts in multiple tissues and in blood metabolites under a realistic range [27,28] of experimental ALAN and in dark controls, and investigated links to behavioral rhythms. The selected genes represented the circadian TTFL (Brain and Muscle ARNT-Like 1, *BMAL1*, alias *ARNTL*; cryptochrome 1, *CRY1*), a clock modulator (*casein kinase 1ε*, *CK1ε*) [29], and endocrine, immune and metabolic pathways putatively affected by circadian disruption (Table S1). Tissues included central pacemaker and memory sites (hypothalamus, where important avian circadian pacemaker components are located [29], and hippocampus; Fig. S1), and metabolic (liver) and immune tissues (spleen). Testes of the same birds were analyzed in a separate study [30]. In contrast to the candidate gene approach, our untargeted metabolomics approach captured both expected and novel effects of ALAN [31]. We aimed to identify whether i) hypothalamic clock gene expression was affected by ALAN, ii) potential temporal shifts in clock gene expression were consistent across tissues, iii) behavioral and clock gene rhythms were aligned, and iv) transcript and metabolite temporal shifts were consistent across physiological pathways. Any inconsistencies in temporal shifts indicate the potential for internal desynchronization, and hence, circadian disruption [9,21].

Great tits are a rewarding study system because their urbanized distribution allows to study ALAN responses also in free-living individuals, because detailed molecular and circadian information is available [32–34], and because like humans, they are diurnal [9,11,22]. We studied 34 male great tits under simulated winter daylength (LD 8.25:15.75 h) in four treatment

groups, ranging from dark night controls to 5 lx (Table S2, S3), and sampled metabolites and transcripts at mid-day (3 h 30 min after lights on; i.e. 3.5 h Zeitgeber time) and midnight (7 h 15 min after lights off; i.e. 15.5 h Zeitgeber time). We chose a study design that enabled detection of rhythmicity and ALAN effects from sampling two time-points 12 h apart [35,36]. The design was enhanced firstly by tracking possible shifts in circadian rhythms by a focal clock gene, *BMAL1*, whose transcription under dark nights in songbirds peaks in the late evening [29]. Secondly, we applied ALAN levels that advance activity of captive great tits by 6 h [22] and thus, if molecular rhythms track behavior, day-night differences at all phase positions are captured.

Our specific predictions are illustrated in Figure 1, which shows expected patterns for *BMAL1*. Under dark nights (Fig. 1A, green curve), during midnight sampling (blue dots) *BMAL1* transcripts will have just passed the peak (maximum), and during mid-day (yellow dots) they will have just passed the trough (minimum). Under our hypothesis, the TTFL matches behavior, and thus, with increasing ALAN (red curves), the *BMAL1* rhythm will also advance. Hence, at midnight *BMAL1* levels will be measured progressively later than the peak, and drop, whereas mid-day levels will be measured closer to the next peak, and hence rise. When combining midnight and mid-day data (Fig. 1B), we thus expected a cross-over of detected *BMAL1* levels. Other rhythmic compounds should show similar patterns, although the point of intersection and precise change of level depends on their phase. In contrast, if the TTFL does not match the behavioral shift by ALAN, compound levels will show as two horizontal lines across ALAN, representing day and night, respectively. Levels of non-rhythmic compounds will fall on a horizontal line, representing both day and night.

Figure 1. Expected clock gene rhythm advance in response to ALAN. Schematic shows ALAN effects on transcript levels of *BMAL1* measured at midnight (blue) and mid-day (yellow). (A) Rhythm of ALAN under dark night shown as green curve; if the gene's rhythm advances (red curves) with increasing ALAN, transcript levels sampled at midnight will drop, whereas those measured at mid-day will rise; horizontal arrows indicate the advance of the *BMAL1* peak. (B) The trends of transcripts with increasing ALAN therefore cross for mid-day vs. midnight sampling.

Results

ALAN advances circadian timing of activity and BMAL1 expression

Daily cycles of activity were strongly affected by the ALAN treatment (GAMM, $p=0.001$, Fig. 2A and Fig. S2; Table S4). In the 5 lx group birds were generally active 6-7 h before lights-on, whereas birds in the other two light treatments (0.5 and 1.5 lx) advanced morning activity to a much lesser extent. This advancement in the onset of morning activity led to 40% of the overall diel activity in the 5 lx group to occur during the night, compared to 11 and 14% in the 0.5 and 1.5 lx groups, and less than 1% in the control dark group. Thus, with increasing ALAN, nocturnal activity also increased (LMM, treatment $p < 0.001$, Fig. 2A and Table S5).

Figure 2. Activity timing is affected by intensity of light at night. The proportion of active 2-min intervals in each treatment group per hour of the day is shown in panel (A) (raw mean $\pm$ SEM, N = 34). Grey background indicates night-time, white background indicates daytime. On the right, we show daily treatment group data (mean $\pm$ SEM), for the timing of (A) evening offset and (B) morning onset of activity (time in min). Activity onset and offset refer to times of lights-on and lights-off, which are shown as the horizontal lines crossing zero in both panels.

Breaking down this average diel profile (Fig. 2A) by time since first exposure to ALAN (i.e., days from start of the experiment to first sampling, days 0 to 18) yields insights into how differences in activity developed, and into circadian mechanisms involved (Fig. 2B-C). Upon exposure to ALAN, the birds' activity onset (Fig. 2C) advanced in all treatment groups. In the groups with intermediate light exposure (0.5 lx, 1.5 lx) the phase-advance occurred instantaneously and to a similar extent (155 and 142 min for the 0.5 and 1.5 lx groups respectively, $P > 0.1$ for this pairwise comparison), but thereafter timing remained stable. The group exposed to 5 lx showed an even larger instantaneous phase advance of an average of almost five hours (mean $\pm$ SEM = 289 ± 21 min), but thereafter continued to gradually phase-advance until reaching a stable phase after 10 days (interaction treatment*day, $p < 0.001$, Fig. 2C, Table S2). The advance until stabilization could equally represent gradual entrainment to an early phase, or temporary free-run of activity, as suggested by periodogram analysis. Indeed, we

found that in the 5 lx group, prior to stabilization, period length deviated from that of all other groups and from 24 h, reaching levels similar to those of free-running conspecifics in an earlier study [37] (mean period length 5 lx group: 23.6 h; LM; Table S6). The individual actograms (Fig. S3) further suggest that the activity rhythm in the 5 lx group may have split into an advancing morning component and a more stably entrained evening component, suggesting internal desynchronization.

Changes in the activity offset were much less pronounced (Fig. 2B). The 5 lx group showed an instantaneous phase-shift, which in contrast to morning activity delayed, rather than advanced, activity compared to the lights-off time. This initial delay was followed by a gradual advance of evening offset, similar to but smaller than that of morning onset. At the end of the experiment birds in the 5 lx group ceased their activity before lights-off, and earlier than other groups (treatment*day, $p < 0.001$, Fig. 2B, Table S5). This advance did not compensate for the earlier onset, as birds in the 5 lx group were more active over the whole 24h than the remaining birds (treatment*day, $p = 0.01$, Table S5).

Hypothalamic BMAL1 expression at night parallels advanced activity onset

We next sought to identify whether the profound shifts in activity patterns were paralleled by corresponding shifts in the pacemaker, measured by expression of *BMAL1* in the hypothalamus. Day-night differences in transcripts of *BMAL1* inverted with increasing ALAN (Fig. S4A), as predicted above (Fig. 1). While *BMAL1* expression was higher at midnight than at mid-day for the control birds, increasing ALAN induced a reversal of this pattern, so that birds in the 5 lx group had much higher expression at mid-day than at midnight (treatment*time, $p < 0.01$, Table S7).

176 To assess whether changes in day-night *BMAL1* gene expression correlated with
177 temporal behavioral shifts, we related *BMAL1* levels to onset of activity of an individual once it
178 had stably shifted in response to the ALAN treatment (Fig. 2B, 2C, after 10 days). Onset was
179 closely predicted by hypothalamic *BMAL1* expression at midnight (Gaussian LM, $p < 0.001$,
180 $R^2 = 0.71$, Fig. 3A). Across ALAN levels, the earliest rising birds had the lowest midnight expression
181 of *BMAL1*. However, the steep linear regression was largely based on differences between ALAN
182 groups in both activity timing (Figs. 2, 3) and *BMAL1* expression (Fig. S4A). Indeed, this
183 relationship was even stronger when we only considered the 0.5, 1.5 and 5 lx group in the
184 analysis (Gaussian LM $p < 0.001$, $R^2 = 0.85$), but the association was not present for the dark
185 control birds (Gaussian LM, $P = 0.87$). Individual midnight *BMAL1* levels were also predictive of
186 mean offset of activity, albeit less strongly so than for onset (Gaussian LM, $p = 0.006$, $R^2 = 0.28$, Fig.
187 3B). Conversely, mid-day *BMAL1* levels did not significantly predict variation in any of the
188 activity traits (Gaussian LMs, $p > 0.1$ and $R^2 < 0.16$ for all measures, Fig. 3C-D).

189

Figure 3. *BMAL1* expression in the hypothalamus predicts the advance of morning activity. mRNA levels of *BMAL1* at midnight correlated with the onset (A) and offset of activity (B), but mid-day levels (C, D) did not. Shown are log-transformed mRNA levels, separated by sampling time (day vs night) and ALAN treatments (blue color gradient). Points represent individual birds (total N = 34), lines and shaded areas represent model fits $\pm$ 95% confidence intervals.

ALAN reverses day-night *BMAL1* expression patterns in multiple tissues

ALAN-induced shifts in *BMAL1*, as detected in the hypothalamus, were remarkably consistent across tissues. Hippocampal *BMAL1* expression profiles resembled those in the hypothalamus (Fig. S5A) and were strongly affected by the interaction of treatment and sampling time ($p < 0.001$, Table S8). Within individuals, mid-day and midnight transcripts in both brain tissues

were closely related (LM, $p < 0.001$, Fig. 4A, Table S9). Also liver *BMAL1* showed similar effects of ALAN on day-night expression profiles (Fig. S6A; time*treatment, $p < 0.001$, Table S10), so that within individuals, hepatic and hypothalamic transcripts also correlated closely (LM, $p < 0.001$, Fig. 4B, Table S9). These findings were consolidated by parallel ALAN effects on *BMAL1* expression in the spleen (Fig. S7A; time*treatment, $p = 0.003$, Table S11), and close individual-level correlation of spleen transcripts with those in hypothalamus (LM, $p = 0.011$, Fig. 4C) and liver (LM, $p = 0.001$, Fig. 4D, Table S9).

Figure 4. ALAN effects on *BMAL1* expression were comparable in different tissues. Correlation of expression patterns of *BMAL1* in different tissues. Shown are log-transformed mRNA levels, separated by sampling time (day vs night) and ALAN treatments (blue color gradient). Points represent individual birds (N = 34). Lines and shaded areas depict model estimated means $\pm$ 95% confidence intervals. Panels show expression levels of hypothalamic *BMAL1* levels in relation to (A) hippocampus, (B) liver and (C) spleen levels, as well as spleen in relation to liver levels (D).

Partial disruption of expression patterns by ALAN in other genes

We next sought to assess whether the same reversal of day-night expression patterns found for *BMAL1* was paralleled in other genes analyzed in the different tissues. We found mixed evidence for this, as in most of the pathways we examined some genes shifted in concert with *BMAL1*, while others did not. This suggests that different pathways were differentially affected by ALAN.

Among clock-related genes, hypothalamic expression levels of *CK1ε*, a clock modulator, was not affected by the light treatment ($p=0.71$). Expression was consistently, although not significantly, higher at mid-day ($p=0.09$, Fig. 5H, Table S7). Similarly, the same gene was not significantly affected by sampling time or treatment in the liver. Expression of hepatic *CK1ε* increased with light intensity, albeit not significantly so ($p=0.078$, Fig. 5P, Table S10), and was not affected by sampling time ($p=0.13$, Table S10). In the liver another circadian gene, *CRY1*, showed no expression trend that aligned with that of *BMAL1* (Fig. 5O). Moreover, *CRY1* was not affected by treatment or sampling time ($P>0.6$ for both variables, Fig. 5O, Table S10).

Among metabolic genes, patterns similar to those in *BMAL1* were evident in *SIRT1*, a gene which is also involved in the modulation of the circadian cycle [38][39] (Table S1). Hypothalamic *SIRT1* showed a clear change of day-night expression with increasing ALAN (Fig. 5E; treatment*time, $p = 0.029$, Table S7), and *SIRT1* mRNA levels were closely related to those of hypothalamic *BMAL1* (LM, $p<0.001$, Table S9). In the liver, the metabolic gene *NRF1* showed a similar response to ALAN as *BMAL1*, with reversed day-night expression in the 5 lx group compared to other groups (treatment*time, $p<0.001$, Fig. 5F, Table S10), and close correlation with *BMAL1* (LM, $p<0.001$). In contrast, another hepatic metabolic gene, *IGF1*, was not significantly affected by light treatment or sampling time (for both, $p>0.11$, Fig. 5Q, Table S10).

In the hippocampus (Table S8), mid-day and midnight levels of the mineralocorticoid receptor, *MR*, decreased significantly with increasing ALAN ($p=0.044$, Fig. 5M). Levels were higher at night than during the day, albeit not significantly so ($p=0.1$). Last, the levels of the glucocorticoid receptor, *GR*, showed no significant relationship with either light treatment or sampling time ($p>0.33$ in both cases, Fig. 5N).

Among immune genes, ALAN affected the hypothalamic mRNA levels of *LY86*, which showed reduced levels with increasing ALAN ($p=0.04$, Fig. 5K, Table S7). Expression of this gene tended to be lower at midnight than mid-day, albeit not significantly so ($p=0.08$). However, the same gene analyzed in the spleen was not affected by either treatment or sampling time ($p>0.7$, Fig. 5L, Table S11). Conversely, another immune gene in the spleen, *TLR4*, showed the same pattern as *BMAL1* (Fig. 5G, time*treatment, $p=0.006$, Table S11).

Last, we also analyzed genes involved in photoperiod seasonal response in the avian brain. *FOXP2*, a gene that in birds is involved in learning, song development and photoperiod-dependent seasonal brain growth, showed no significant trends related to ALAN or sampling time ($p>0.32$ in both cases, Fig. 5J). *DIO2*, a thyroid-axis gene involved in photoperiodic reproductive activation, was also not affected by either ALAN or sampling time ($p>0.45$ for both variables, Fig. 5I).

Figure 5. ALAN effects on gene expression are gene-specific. ALAN does not equally affect all physiological systems. ALAN effects on *BMAL1* (A)-(D) were paralleled by those on three additional genes in the hypothalamus (*SIRT1*), liver (*NRF1*) and spleen (*TLR4*) (E)-(G), but not by other genes analyzed across tissues (H)-(Q). Shown are log-transformed mRNA levels, separated by sampling time (mid-day: yellow; midnight: dark blue). Large symbols $\pm$ SEM connected by lines represent model estimates, whereas small symbols depict raw data points (N = 34 birds).

Metabolomic profiles support only a limited reversal of day-night physiology under ALAN

To explore the different impacts of ALAN on whole-body physiology, we carried out untargeted LC-MS metabolomic analysis and obtained abundance values for 5483 compounds. Out of these, 682 were annotated as known metabolites based on accurate mass and predicted retention time [40] and 73 were identified based on accurate mass measurement and matching retention time to a known standard (within 5%), for a total of 755 metabolites. We ran individual linear mixed models for all these 755 metabolites (correcting for false discovery rate at 5%), and found that 44.1% (333) differed significantly by sampling time, with higher levels at mid-day in 197, and higher levels at midnight in 136 (to see all metabolite tables: <https://doi.org/10.6084/m9.figshare.12927539.v1>). For 29 metabolites we found significant effects of treatment (Table S12). The direction of the treatment effect depended on the metabolite considered. In 11 metabolites, levels decreased with ALAN, while in the remaining 18 metabolites an increase was observed when compared to the dark night control group. Finally, 73 (9.7%) of the 755 metabolites showed significant interaction between treatment and sampling time (Fig. 6 and Table S13; 34 of those also differed by sampling time). As this pattern supported reversal of day-night physiology similar to that shown for activity and *BMAL1* expression, these metabolites were selected for subsequent focal analyses (hereafter named “interactive dataset”).

We dissected variation in the interactive dataset by using two principal component analyses (PCA) on the samples collected at mid-day and midnight (Fig. 6C, D). For mid-day samples, ALAN treatments overlapped considerably (Fig. 6C), although low values of PC1 (26 % of variance explained) aligned with some of the birds in the 1.5 lx and 5 lx treatments. PC1 in the mid-day dataset was heavily loaded with metabolites of Arginine biosynthesis pathway,

including L-Arginine, Homoarginine and L-Glutamate, as well as other important amino acids such as L-Threonine, L-Lysine and L-Tyrosine. Conversely, the midnight samples (Fig. 6D) separated clearly between the 5 lx treatment and the remaining groups. In this midnight PCA, PC1 explained 27% of the variance and was heavily loaded with metabolites of the Glutamate and Arginine pathways, as well as with N-acetyl-L-aspartate. PC2, which explained 21% of variation, was heavily loaded with fatty acids, including Linoleate (to see all factor loading tables: <https://doi.org/10.6084/m9.figshare.12927536.v1>). The contribution of the Arginine pathway was further confirmed by pathway analysis, conducted with Metaboanalyst [41], which indicated “Arginine biosynthesis” as a highly significant pathway in this interactive dataset ($p < 0.001$). “Aminoacyl-tRNA metabolism” ($p < 0.001$), “Histidine metabolism” ($p = 0.005$), and “Alanine, Aspartate and glutamate metabolism” ($p = 0.026$) were also indicated as significant pathways.

We finally investigated whether, just like midnight levels of *BMAL1* expression (Fig. 4), midnight principal components of metabolites correlated with individual activity timing. PC1 strongly predicted the onset of activity via a linear and quadratic relationship ($n = 19$, $p_{\text{linear}} = 0.007$, $p_{\text{quadratic}} = 0.014$, $R^2 = 0.92$, Fig. 6E), but did not explain offset of activity ($p = 0.63$, $R^2 = 0.04$, Fig. 6F). PC2 was related to neither timing trait ($p > 0.2$).

Figure 6. Metabolomics analysis supports ALAN-induced shifts in day-night physiology. The 73 metabolites found to be significantly affected by the interaction of treatment and sampling time (9.6% of all metabolites, interactive dataset) were dissected by means of pathway analysis and principal component analysis. Pathway analysis revealed that the Arginine Biosynthesis pathway was particularly enriched in this dataset. Heatmaps show the top-25 metabolites in the interactive dataset at either mid-day (A) or mid-night (B). Principal component analysis showed considerable overlap between ALAN groups at mid-day (C), whereas ALAN treatment effects were mostly visible at midnight, particularly for the 5 lx group (D). In all PCA plots, points represent individual samples, and ellipses contain 80% of samples in a group. The first PC of the night cluster (E) significantly predicted the onset of activity in the morning (D), but not the offset of activity in the evening (F). In (E) and (F) points represent individual birds (N = 19), and lines and shaded areas represent model fits $\pm$ 95% confidence intervals.

Discussion

Birds advanced the circadian timing of their activity as expected with increasing levels of ALAN, and in parallel the gene expression of our focal clock gene, *BMAL1*, was also advanced in the hypothalamus. Advances in *BMAL1* were consistent across tissues, indicating a shift of the circadian system in tissues implicated in timing, memory, metabolism and immune function. Furthermore, advances in nocturnal *BMAL1* potentially correlated with activity onset at the individual level, consolidating close links between core clock gene expression and behavior. Responses of *BMAL1* expression were paralleled by a minority of other genes. Similarly, only 9.7% of the metabolome followed the same shift observed in *BMAL1*, indicating that most physiological pathways were desynchronized from the circadian system. The emerging picture is that birds shifted their internal clock time under ALAN, but suffered a high degree of internal desynchronization.

On a behavioral level, our findings closely match those of earlier demonstrations of advanced daily activity under ALAN in captivity for several avian species, including the great tit [15,22,24,42]. In the wild, birds also advanced daily activity under ALAN, although to a lesser extent (e.g. [14,26,43]), and often in onset but not offset [25,26,28,44,45]. Previously,

344 behavioral shifts were interpreted as not involving the circadian clock [24]. In an experiment
 345 also on the great tit, Spoelstra and colleagues [24] exposed birds to dark nights and then to
 346 ALAN as in our study. Subsequently, birds were released to constant low-levels of dim light (0.5
 347 lx), where they free-ran. The study found that the birds free-ran from the timing they had shown
 348 under initial dark nights, rather than from their advanced timing under ALAN. Thus, the authors
 349 concluded that the behavioral response to ALAN was due to masking, while the internal clock
 350 remained unchanged [24]. Our molecular data suggest a different conclusion, namely that
 351 within three weeks of ALAN exposure, internal time had phase-advanced in concert with
 352 behavior. These discrepancies are difficult to interpret because inferences of the studies are
 353 based on different criteria (molecular vs. behavioral) and different experimental phases (during
 354 ALAN vs. during ensuing free-run), but it is clear that additional experimental data are needed.

355
 356 **Figure 7. Proportion of shifts in day-night pattern in response to ALAN.** Shown are proportions
 357 of genes (grey) and metabolites (red) whose levels were, or were not, significantly impacted by
 358 the interaction of sampling time and ALAN level.
 359

360 Our transcriptional findings of ALAN-altered rhythmicity gain support from a comparison
 361 of clock gene expression in Tree sparrows (*Passer montanus*) from an illuminated urban and
 362 dark non-urban habitat [46]. Sampled within a day after being brought into captivity, urban
 363 birds showed clear advances in the circadian system, including, as in our birds, in hypothalamic
 364 *BMAL1*. Other experimental studies have also confirmed effects of ALAN on avian rhythms in
 365 brain and other tissues [16,17]. In our study, only some of the investigated regulatory genes
 366 aligned with the ALAN-dependent advances of rhythms in behavior and *BMAL1*. The genes from
 367 metabolic pathways that have close molecular links to the TTFL, *SIRT1* and *NRF1*, mirrored
 368 ALAN-dependent changes in *BMAL1*. However, regulatory genes of immune pathways
 369 responded inconsistently, whereby *TLR4* aligned with *BMAL1* whereas *LY86* did not. The learning
 370 gene, *FOXP2* and the thyroid-activating gene *DIO2* did not mirror the changes in *BMAL1*, nor did
 371 the endocrine genes (*MR*, *GR*, *IGF1*). Conversely, in a complementary study on these same birds,
 372 we observed that ALAN exposure, which also activated the reproductive system, shifted the day-
 373 night expression patterns of corticoid receptors [30].

374 Other experimental studies have confirmed that effects of ALAN on avian rhythms in
 375 brain and other tissues differed between genes and pathways. For example, a study on Zebra
 376 finches (*Taeniopygia guttata*) reported ALAN-induced changes in rhythmic expression of
 377 hypothalamic *CRY1* but not *BMAL1* [16]. This differs from our findings, where advances in
 378 *BMAL1* were not paralleled by *CRY1* [17], and from findings that *BMAL1* and *CRY1*, but not
 379 another TTFL gene, *CLOCK*, advanced in an urban bird [47]. Divergent responses between clock
 380 genes might participate in circadian disruption, and could underlie discrepant behavioral
 381 responses, such as differences between activity onset and offset observed in our study, and in
 382 wild great tits [25,26,44] and other avian species [28,45]. In our study in the 5 lx group, we also

observed splitting of rhythms, which has previously been linked to reproductive activation [48],
a known side-effect of ALAN [13].

Our metabolomic data corroborated our main findings on gene expression. Of the 755
identified metabolites, nearly 50% (333) differed between mid-day and mid-night levels.
However, less than 10 % showed changes in rhythm under ALAN (Fig. 7). These findings confirm
that some, but not all featured pathways aligned with shifts in behavior and *BMAL1*. Our
findings from captive wild birds under ALAN match those from human studies. To identify the
mechanisms by which circadian disruption drives metabolic disorders and other pathologies,
these studies severely disrupted the circadian system by sleep deprivation and shift-work
protocols [31,49,50]. The reported changes in gene expression and metabolite levels were
similar to those of our birds under ALAN, including highly responsive pathways and compounds,
in particular Arginine [50], an amino acid strongly linked to circadian rhythms and innate
immune responses [51]. Glutamate production from arginine is well known [52], and changes in
these two metabolites may be due to changes in energy requirements at the different light
intensities. N-acetyl-aspartate, a metabolite involved in energy production from glutamate [53],
was also observed to follow changes in behavior and *BMAL1*. Both glutamate and arginine have
a variety of biochemical roles [54,55], so further work would be required to determine which of
these functions, if any, are associated to the behavioral and gene expression changes we
observed. While preliminary, this data shows the potential of metabolomic techniques for
furthering this area of research.

Despite our sampling design of only two time-points and low sample sizes, we derived
descriptors of internal time (*BMAL1* expression; metabolomics PC1 of interactive dataset) whose
midnight levels had high predictive power of activity timing. Thereby, we have shown that

internal time can be captured in birds by a single sample of blood or tissue, a frontline ambition of biomedical research [35,36]. Our predictive power was limited to treatment groups and within-ALAN individuals, whereas birds kept under dark nights were highly synchronized to the sudden switch of lights-on.

For wild animals, our study adds to emerging evidence of detrimental effects of ALAN on physiological pathways [9,10,21]. For example, under ALAN molecular markers for sleep deprivation were elevated, hypothalamic expression of genes such as *TLR4* was altered [16], neuronal features in the brain were changed, and cognitive processes and mental health-like states were impaired [16,20,56,57]. Altered hepatic expression of several metabolic genes further suggested negative effects on gluconeogenesis and cholesterol biosynthesis [15]. Consequences of ALAN-induced changes in immune function include increased host competence for infectious disease [58], indicating how effects on individuals may cascade to ecological or epidemiological scales.

Addressing effects of ALAN is therefore urgent [10,59]. Our data contribute to the rising evidence for dose-dependent responses of behavior and physiology [22,30,60], which might allow mitigating against ALAN impacts on wildlife by reducing light intensity [61]. Importantly, we detected substantial effects even at light intensities (0.5 lx) that are typically far exceeded by street illumination, and to which animals are exposed to in the wild [27,28]. These findings transfer to other organisms including plants, insects, and mammals including humans [12,62–65] and call for limits to the ever faster global increase in light pollution [3].

430

431 **Methods**

432 *Data availability*

433 The full details of our methods are presented in the *Supporting* Information document. Raw

434 data, created datasets and R scripts are available via Figshare:

435 [https://figshare.com/projects/Artificial_light_at_night_shifts_the_circadian_system_but_still_l](https://figshare.com/projects/Artificial_light_at_night_shifts_the_circadian_system_but_still_leads_to_physiological_disruption_in_a_wild_bird/88841)
436 [eads_to_physiological_disruption_in_a_wild_bird/88841](https://figshare.com/projects/Artificial_light_at_night_shifts_the_circadian_system_but_still_leads_to_physiological_disruption_in_a_wild_bird/88841).

437

438 *Animals and experimental design*

439 We studied 34 hand-raised, adult male great tits that were kept in individual cages (90 × 50 × 40

440 cm) under simulated natural daylength and ambient temperature of 10 to 14 °C with *ad libitum*

441 access to food and water, as described in [30].

442 The experiment started on February 1st, 2014, when daylength was fixed at 8 h 15 min

443 light and 15 h 45 min darkness. During the day, all birds were exposed to full spectrum daylight

444 by high frequency fluorescent lights emitting ~1000 lx at perch level (Activa 172, Philips,

445 Eindhoven, the Netherlands). During the night, birds were assigned to four treatment groups

446 exposed to nocturnal light intensity of 0 lx (n = 13), 0.5 lx (n = 7), 1.5 lx (n = 7), or 5 lx (n = 7). In

447 composing these groups, we prioritized assigning birds to the dark night group to obtain reliable

448 benchmark data on day-night differences in gene expression. Lights were provided by warm

449 white LED light (Philips, Eindhoven, The Netherlands; for details on the spectral composition of

450 lights, see [22]).

451 On Feb 20th an initial blood sample (~200 µl) was collected from all birds at mid-day for

452 metabolomic profiling. On Feb 22nd birds were randomly assigned to mid-day or midnight groups

for culling to collect tissues for morphological and molecular analyses. The mid-day group was culled on Feb 22nd, whereas culling of the midnight group was divided over two subsequent nights (Feb 22nd: 12 birds; Feb 23rd: 10 birds). Blood was again collected for metabolomic profiling.

All experimental procedures were carried out under license NIOO 13.11 of the Animal Experimentation Committee (DEC) of the Royal Netherlands Academy of Arts and Sciences.

Locomotor activity

Daily activity patterns of each individual bird were measured continuously using micro-switches recorded by a computer, as described in de Jong et al [22]. See Supporting Information for more details.

Gene expression analyses

After culling, organs were extracted, snap-frozen on dry ice, and stored at -80 °C within 10 min of capture.

Brain tissue was cut on a cryostat at -20 °C. We cut sagittal sections throughout the brain (Fig. S1). The hypothalamus and hippocampus were located by the use of the Zebrafish atlas ZEBRA (Oregon Health & Science University, Portland, OR, USA; <http://www.zebrafinchatlas.org>) and isolated from the frozen brain sections either by surgical punches for the hypothalamus (Harris Uni-Core, 3.0 mm), or by scraping the relevant tissue with forceps, for the hippocampus. For the hypothalamus, the edge of the circular punch was positioned adjacent to the midline and ventral edge of the section, just above the optic chiasm, following the procedure of [66]. Hypothalamic and hippocampal tissue was then immediately

added to separate 1.5ml buffer tubes provided by the Qiagen RNeasy micro extraction kit (see below), homogenized and stored at -80 °C until extraction.

Whole spleens were homogenized with a ryboliser and added to 1.5 ml RNeasy micro buffer and stored at -80 °C. For livers, we cut 0.5 g of tissue from each individual liver, homogenized it and added it to 1.5 ml RNeasy micro buffer and stored it at -80 °C. RNA was extracted using the RNeasy micro extraction kit and reverse transcribed it to generate cDNA using a standard kit following the manufacturer's instructions (Superscript III, Invitrogen).

We selected exemplary genes known to be involved in circadian timing, seasonal timing, and in metabolic, immune and endocrine function (Table S1). We analyzed the core clock gene *BMAL1* in all tissues as our primary clock indicator because of the timing of its expression and because of its role as central hub for inter-linking molecular pathways [7]. We also studied a second core clock gene, *CRY1*, in a single tissue, and a clock modulator, *CK1ε*, in two tissues. In the hypothalamus, we also studied two genes involved in seasonal changes (*DIO2*, *FOXP2*), and one metabolic and ageing gene (*SIRT1*). The second metabolic gene, *NRF1*, was studied in the liver. Two immune genes represented different pathways (*LY86*, *TLR4*). Finally, we studied endocrine genes involved in stress signaling in the Hippocampus (*NR3C1* (*alias GR*), *NR3C2* (*alias MR*)) and in tissue homeostasis (*IGF1*), as well as reference genes (for full details see Table S1). Primers were built based on the great tit reference genome build 1.1

(https://www.ncbi.nlm.nih.gov/assembly/GCF_001522545.2) [33] and annotation release 101 (https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Parus_major/101/). Primer design was conducted with Geneious version 10.0.2 [67].

Amplification efficiency of each primer pair was determined through quantitative real-time polymerase chain reaction (RT-qPCR). RT-qPCR was performed on duplicate samples by a 5-

point standard curve. We used reference gene levels to correct for variation in PCR efficiency between samples. Reference gene expression stability was calculated using the application geNorm [68], from which we identified the best pair of reference genes for each tissue. Absolute amounts of cDNA were calculated by conversion of the Ct values ($C \times E^{-Ct}$, with $C=10^{10}$ and $E=2$) [69]. The absolute amounts of the candidate genes were then normalized by division by the geometric mean of the absolute amounts of the reference genes. This step yielded relative mRNA expression levels of the candidate genes. For more details, see the Supporting Information document.

Metabolomics analysis

See Supporting information for initial sample preparation and for additional details. All samples were analyzed on a Thermo Scientific QExactive Orbitrap mass spectrometer running in positive/negative switching mode. Mass spectrometry data were processed using a combination of XCMS 3.2.0 and MZMatch.R 1.0–4 [70]. Unique signals were extracted using the centwave algorithm [71] and matched across biological replicates based on mass to charge ratio and retention time. The final peak set was converted to text for use with IDEOM v18 [72], and filtered on the basis of signal to noise score, minimum intensity and minimum detections, resulting in a final dataset of 755 metabolites.

Statistical analysis

All statistical analyses were conducted in R, version 3.63 [73]. In all models we included treatment as log-transformed light intensity (adding a constant to avoid zero). Details of all statistical analyses can be seen in the Supporting Information document.

To analyze locomotor activity data (i.e. perch-hopping), we first divided the time series of activity into an unstable phase and stable phase (see Supporting information). We used the data in the unstable phase to quantify circadian period length (τ) for each bird, then tested treatment effects using a gaussian linear model (LM). The data in the stable phase were used to test for variation in the proportion of time spent active every hour depending on treatment, using a generalized additive mixed model (GAMM). Finally, we tested for variation in onset time, offset time, nocturnal activity and total daily activity using separate linear mixed models (LMMs).

To examine variation in relative transcript levels, we ran LMs including ALAN treatment, sampling time (two-level factor, day and night), and their interaction as explanatory variables, and mRNA expression levels of the different genes in the different tissues as response variables. Similar models were used to test for relationships in mRNA levels between the same gene in different tissues, or different genes in the same tissue.

To test for variation in the levels of the individual metabolites identified by the LC-MS, we used all data, including the replicated mid-day samples (total $n = 64$). We ran independent LMMs for each metabolite, with metabolite levels as response variable (log transformed and normalized), and treatment, time of day and their interaction as explanatory variables. Moreover, we ran two principal component analyses using only the 73 metabolites found to be significantly affected by the treatment*time interaction in the LMMs described above. The two PCAs were run separately for the individual samples collected at mid-day or midnight. We then used the first two principal components (PC1 and PC2) of the midnight based PCA as explanatory variables in two LMs with onset and offset of activity as response variables, respectively.

References

1. Foster RG, Kreitzmann L. Rhythms of life: The biological clocks that control the daily lives of every living thing. New Haven, CT, USA: Yale University Press; 2004.
2. Helm B, Visser ME, Schwartz W, Kronfeld-Schor N, Gerkema M, Piersma T, et al. Two sides of a coin: Ecological and chronobiological perspectives of timing in the wild. *Philosophical Transactions of the Royal Society B: Biological Sciences*. Royal Society Publishing; 2017. doi:10.1098/rstb.2016.0246
3. Falchi F, Cinzano P, Duriscoe D, Kyba CCM, Elvidge CD, Baugh K, et al. The new world atlas of artificial night sky brightness. *Sci Adv*. 2016;2: e1600377–e1600377. doi:10.1126/sciadv.1600377
4. Davies TW, Smyth T. Why artificial light at night should be a focus for global change research in the 21st century. *Glob Chang Biol*. 2017; early view. doi:10.1111/gcb.13927
5. Borniger JC, Cisse YM, Surbhi, Nelson RJ. Reciprocal Regulation of Circadian Rhythms and Immune Function. *Current Sleep Medicine Reports*. Springer International Publishing; 2017. pp. 93–103. doi:10.1007/s40675-017-0070-7
6. Caratti G, Iqbal M, Hunter L, Kim D, Wang P, Vonslow RM, et al. REVERBa couples the circadian clock to hepatic glucocorticoid action. *J Clin Invest*. 2018;128: 4454–4471. doi:10.1172/JCI96138
7. Ribas-Latre A, Eckel-Mahan K. Interdependence of nutrient metabolism and the circadian clock system: Importance for metabolic health. *Molecular Metabolism*. Elsevier GmbH; 2016. pp. 133–152. doi:10.1016/j.molmet.2015.12.006
8. Dominoni DM, Nelson RJ. Artificial light at night as an environmental pollutant: An integrative approach across taxa, biological functions, and scientific disciplines. *J Exp Zool Part A Ecol Integr Physiol*. 2018;329. doi:10.1002/jez.2241
9. Vetter C. Circadian disruption: What do we actually mean? *European Journal of Neuroscience*. Blackwell Publishing Ltd; 2020. pp. 531–550. doi:10.1111/ejn.14255
10. Falcón J, Torriglia A, Attia D, Viénot F, Gronfier C, Behar-Cohen F, et al. Exposure to Artificial Light

570 at Night and the Consequences for Flora, Fauna, and Ecosystems. *Frontiers in Neuroscience*.
571 Frontiers Media S.A.; 2020. p. 1183. doi:10.3389/fnins.2020.602796

572 11. Ouyang JQ, de Jong M, van Grunsven RHA, Matson KD, Haussmann MF, Meerlo P, et al. Restless
573 roosts – light pollution affects behavior, sleep and physiology in a free-living songbird. *Glob Chang*
574 *Biol*. 2017; 1–8.

575 12. Knop E, Zoller L, Ryser R, Gerpe C, Hörler M, Fontaine C. Artificial light at night as a new threat to
576 pollination. *Nature*. 2017. doi:10.1038/nature23288

577 13. Dominoni D, Quetting M, Partecke J. Artificial light at night advances avian reproductive
578 physiology. *Proc R Soc B Biol Sci*. 2013;280: 20123017. doi:10.1098/rspb.2012.3017

579 14. Dominoni DMM, Helm B, Lehmann M, Dowse HBB, Partecke J. Clocks for the city: circadian
580 differences between forest and city songbirds. *Proc R Soc London B Biol Sci*. 2013;280: 20130593.
581 doi:10.1098/rspb.2013.0593

582 15. Batra T, Malik I, Kumar V. Illuminated night alters behaviour and negatively affects physiology and
583 metabolism in diurnal zebra finches. *Environ Pollut*. 2019;254: 112916.
584 doi:10.1016/J.ENVPOL.2019.07.084

585 16. Batra T, Malik I, Prabhat A, Bhardwaj SK, Kumar V. Sleep in unnatural times: illuminated night
586 negatively affects sleep and associated hypothalamic gene expressions in diurnal zebra finches.
587 *Proc R Soc B Biol Sci*. 2020;287: 20192952. doi:10.1098/rspb.2019.2952

588 17. Yang Y, Liu Q, Wang T, Pan J. Light pollution disrupts molecular clock in avian species: A power-
589 calibrated meta-analysis. *Environ Pollut*. 2020; 114206. doi:10.1016/j.envpol.2020.114206

590 18. Van Der Vinne V, Riede SJ, Gorter JA, Eijer WG, Sellix MT, Menaker M, et al. Cold and hunger
591 induce diurnality in a nocturnal mammal. *Proc Natl Acad Sci U S A*. 2014;111: 15256–15260.
592 doi:10.1073/pnas.1413135111

593 19. Ouyang JQ, Davies S, Dominoni D. Hormonally mediated effects of artificial light at night on
594 behavior and fitness: linking endocrine mechanisms with function. *J Exp Biol*. 2018;221:
595 jeb156893. doi:10.1242/jeb.156893

596 20. Moaraf S, Vistoropsky Y, Pozner T, Heiblum R, Okuliarová M, Zeman M, et al. Artificial light at
597 night affects brain plasticity and melatonin in birds. *Neurosci Lett*. 2020;716: 134639.
598 doi:10.1016/j.neulet.2019.134639

599 21. Moreno CRC, Wright K, Skene DJ, Louzada FM. Phenotypic plasticity of circadian entrainment
600 under a range of light conditions. *Neurobiol Sleep Circadian Rhythm*. 2020;9: 100055.
601 doi:10.1016/j.nbscr.2020.100055

602 22. de Jong M, Jeninga L, Ouyang JQ, van Oers K, Spoelstra K, Visser ME. Dose-dependent responses
603 of avian daily rhythms to artificial light at night. *Physiol Behav*. 2016;155: 172–179.
604 doi:10.1016/j.physbeh.2015.12.012

605 23. Sun J, Raap T, Pinxten R, Eens M. Artificial light at night affects sleep behaviour differently in two
606 closely related songbird species. *Environ Pollut*. 2017;231: 882–889.
607 doi:10.1016/j.envpol.2017.08.098

608 24. Spoelstra K, Verhagen I, Meijer D, Visser ME. Artificial light at night shifts daily activity patterns
609 but not the internal clock in the great tit (*Parus major*). *Proc R Soc B Biol Sci*. 2018;285: 20172751.
610 doi:10.1098/rspb.2017.2751

611 25. Ulgezen ZN, Kämpylä T, Meerlo P, Spoelstra K, Visser ME, Dominoni DM. The preference and costs
612 of sleeping under light at night in forest and urban great tits. *Proc R Soc B Biol Sci*. 2019;286:
613 20190872. doi:10.1098/rspb.2019.0872

614 26. Raap T, Pinxten R, Eens M. Light pollution disrupts sleep in free-living animals. *Sci Rep*. 2015;5:
615 13557. doi:10.1038/srep13557

616 27. Dominoni D, Quetting M, Partecke J. Artificial light at night advances avian reproductive
617 physiology. *Proc R Soc B Biol Sci*. 2013;280. doi:10.1098/rspb.2012.3017

618 28. Dominoni DMD, Carmona-Wagner EOE, Hofmann M, Kranstauber B, Partecke J. Individual-based
619 measurements of light intensity provide new insights into the effects of artificial light at night on
620 daily rhythms of urban-dwelling songbirds. *J Anim Ecol*. 2014;83: 681–692. doi:10.1111/1365-
621 2656.12150

- 622 29. Cassone VM, Paulose JK, Harpole CE, Li Y, Whitfield-Rucker M. Avian circadian organization.
623 Biological Timekeeping: Clocks, Rhythms and Behaviour. Springer (India) Private Ltd.; 2017. pp.
624 241–256. doi:10.1007/978-81-322-3688-7_11
- 625 30. Dominoni D, de Jong M, Bellingham M, O’Shaughnessy P, van Oers K, Robinson J, et al. Dose-
626 response effects of light at night on the reproductive physiology of great tits (*Parus major*):
627 Integrating morphological analyses with candidate gene expression. *J Exp Zool Part A Ecol Integr*
628 *Physiol.* 2018;2018: 1–15. doi:10.1002/jez.2214
- 629 31. Davies SK, Ang JE, Revell VL, Holmes B, Mann a., Robertson FP, et al. Effect of sleep deprivation
630 on the human metabolome. *Proc Natl Acad Sci.* 2014;111: 10761–10766.
631 doi:10.1073/pnas.1402663111
- 632 32. Watson H, Videvall E, Andersson MN, Isaksson C. Transcriptome analysis of a wild bird reveals
633 physiological responses to the urban environment. *Sci Rep.* 2017;7: 1–10. doi:10.1038/srep44180
- 634 33. Laine VN, Gossmann TI, Schachtschneider KM, Garroway CJ, Madsen O, Verhoeven KJ, et al.
635 Evolutionary signals of selection on cognition from the great tit genome and methylome. *Nat*
636 *Commun.* 2016;7: 10474. doi:10.1038/ncomms10474
- 637 34. Capilla-Lasheras P, Dominoni DMDM, Babayan SASA, O’Shaughnessy PJPJ, Mladenova M,
638 Woodford L, et al. Elevated Immune Gene Expression Is Associated with Poor Reproductive
639 Success of Urban Blue Tits. *Front Ecol Evol.* 2017;5: 64. doi:10.3389/fevo.2017.00064
- 640 35. Kasukawa T, Sugimoto M, Hida A, Minami Y, Mori M, Honma S, et al. Human blood metabolite
641 timetable indicates internal body time. *Proc Natl Acad Sci U S A.* 2012;109: 15036–15041.
642 doi:10.1073/pnas.1207768109
- 643 36. Laing EE, Möller-Levet CS, Poh N, Santhi N, Archer SN, Dijk DJ. Blood transcriptome based
644 biomarkers for human circadian phase. *Elife.* 2017;6. doi:10.7554/eLife.20214
- 645 37. Helm B, Visser ME. Heritable circadian period length in a wild bird population. *Proc R Soc B Biol*
646 *Sci.* 2010;277: 3335–3342. doi:10.1098/rspb.2010.0871
- 647 38. Chang H-C, Guarente L. SIRT1 Mediates Central Circadian Control in the SCN by a Mechanism that

648 Decays with Aging. *Cell*. 2013;153: 1448–1460. doi:10.1016/j.cell.2013.05.027

649 39. Trivedi AK, Kumar J, Rani S, Kumar V. Annual life history-dependent gene expression in the
650 hypothalamus and liver of a migratory songbird: Insights into the molecular regulation of seasonal
651 metabolism. *J Biol Rhythms*. 2014;29: 332–345. doi:10.1177/0748730414549766

652 40. Creek DJ, Jankevics A, Breitling R, Watson DG, Barrett MP, Burgess KEV. Toward global
653 metabolomics analysis with hydrophilic interaction liquid chromatography-mass spectrometry:
654 Improved metabolite identification by retention time prediction. *Anal Chem*. 2011;83: 8703–8710.
655 doi:10.1021/ac2021823

656 41. Chong J, Soufan O, Li C, Caraus I, Li S, Bourque G, et al. MetaboAnalyst 4.0: Towards more
657 transparent and integrative metabolomics analysis. *Nucleic Acids Res*. 2018;46: W486–W494.
658 doi:10.1093/nar/gky310

659 42. Dominoni DMD, Goymann W, Helm B, Partecke J. Urban-like night illumination reduces melatonin
660 release in European blackbirds (*Turdus merula*): implications of city life for biological time-keeping
661 of songbirds. *Front Zool*. 2013;10: 60. doi:10.1186/1742-9994-10-60

662 43. Kempenaers B, Borgström P, Loës P, Schlicht E, Valcu M. Artificial night lighting affects dawn song,
663 extra-pair siring success, and lay date in songbirds. *Curr Biol*. 2010;20: 1735–1739.
664 doi:10.1016/j.cub.2010.08.028

665 44. Dominoni DM, Smit JAH, Visser ME, Halfwerk W. Multisensory pollution: Artificial light at night
666 and anthropogenic noise have interactive effects on activity patterns of great tits (*Parus major*).
667 *Environ Pollut*. 2019; 113314. doi:10.1016/j.envpol.2019.113314

668 45. Da Silva A, Samplonius J, Schlicht E, Valcu M, Kempenaers B. Artificial night lighting rather than
669 traffic noise affects the daily timing of dawn and dusk singing in common European songbirds.
670 *Behav Ecol*. 2014;25: 1037–1047. doi:10.1093/beheco/aru103

671 46. Renthlei Z, Trivedi AK. Effect of urban environment on pineal machinery and clock genes
672 expression of tree sparrow (*Passer montanus*). *Environ Pollut*. 2019;255: 113278.
673 doi:10.1016/j.envpol.2019.113278

674 47. Renthlei Z, Borah BK, Gurumayum T, Trivedi AK. Season dependent effects of urban environment
675 on circadian clock of tree sparrow (*Passer montanus*). Photochem Photobiol Sci. 2020 [cited 9
676 Dec 2020]. doi:10.1039/d0pp00257g

677 48. Gwinner F. Testosterone Induces “Splitting” of Circadian Locomotor Activity Rhythms in Birds.
678 Science. 1974. pp. 72–74. doi:10.1126/science.185.4145.72

679 49. Archer SN, Laing EE, Moller-Levet CS, van der Veen DR, Bucca G, Lazar a. S, et al. PNAS Plus: From
680 the Cover: Mistimed sleep disrupts circadian regulation of the human transcriptome. Proc Natl
681 Acad Sci. 2014;111: E682–E691. doi:10.1073/pnas.1316335111

682 50. Skene DJ, Skornyakov E, Chowdhury NR, Gajula RP, Middleton B, Satterfield BC, et al. Separation
683 of circadian- and behavior-driven metabolite rhythms in humans provides a window on peripheral
684 oscillators and metabolism. Proc Natl Acad Sci U S A. 2018;115: 7825–7830.
685 doi:10.1073/pnas.1801183115

686 51. Morris SM. Arginine: Master and commander in innate immune responses. Science Signaling.
687 American Association for the Advancement of Science; 2010. pp. pe27–pe27.
688 doi:10.1126/scisignal.3135pe27

689 52. Morris SM. Enzymes of arginine metabolism. Journal of Nutrition. American Institute of Nutrition;
690 2004. pp. 2743S–2747S. doi:10.1093/jn/134.10.2743s

691 53. Clark JB. N-acetyl aspartate: A marker for neuronal loss or mitochondrial dysfunction.
692 Developmental Neuroscience. Dev Neurosci; 1998. pp. 271–276. doi:10.1159/000017321

693 54. Yelamanchi SD, Jayaram S, Thomas JK, Gundimeda S, Khan AA, Singhal A, et al. A pathway map of
694 glutamate metabolism. J Cell Commun Signal. 2016;10: 69–75. doi:10.1007/s12079-015-0315-5

695 55. Morris SM. Arginine metabolism revisited. J Nutr. 2016;146: 2579S–2586S.
696 doi:10.3945/jn.115.226621

697 56. Taufique SKT, Prabhat A, Kumar V. Illuminated night alters hippocampal gene expressions and
698 induces depressive-like responses in diurnal corvids. Eur J Neurosci. 2018;48: 3005–3018.
699 doi:10.1111/ejn.14157

700 57. Taufique SKT, Prabhat A, Kumar V. Constant light environment suppresses maturation and
701 reduces complexity of new born neuron processes in the hippocampus and caudal nidopallium of
702 a diurnal corvid: Implication for impairment of the learning and cognitive performance. *Neurobiol*
703 *Learn Mem.* 2018;147: 120–127. doi:10.1016/j.nlm.2017.12.001

704 58. Kernbach ME, Newhouse DJ, Miller JM, Hall RJ, Gibbons J, Oberstaller J, et al. Light pollution
705 increases West Nile virus competence of a ubiquitous passerine reservoir species. *Proc R Soc B*
706 *Biol Sci.* 2019;286: 20191051. doi:10.1098/rspb.2019.1051

707 59. Kyba CCM, Kuester T, Sánchez de Miguel A, Baugh K, Jechow A, Hölker F, et al. Artificially lit
708 surface of Earth at night increasing in radiance and extent. *Sci Adv.* 2017;3: e1701528.
709 doi:10.1126/sciadv.1701528

710 60. Bruening A, Hölker F, Franke S, Preuer T, Kloas W. Spotlight on fish: Light pollution affects
711 circadian rhythms of European perch but does not cause stress. *Sci Total Environ.* 2015;511: 516–
712 522. doi:10.1016/j.scitotenv.2014.12.094

713 61. Gaston KJ, Davies TW, Bennie J, Hopkins J. REVIEW: Reducing the ecological consequences of
714 night-time light pollution: options and developments. *J Appl Ecol.* 2012;49: 1256–1266.
715 doi:10.1111/j.1365-2664.2012.02212.x

716 62. Dominoni D, Borniger J, Nelson R. Light at night, clocks and health: from humans to wild
717 organisms. *Biol Lett.* 2016;12: 20160015. doi:10.1098/rsbl.2016.0015

718 63. Bennie J, Davies TW, Cruse D, Gaston KJ. Ecological effects of artificial light at night on wild plants.
719 *J Ecol.* 2016;104: 611–620. doi:10.1111/1365-2745.12551

720 64. van Langevelde F, Braamburg-Annegarn M, Huigens ME, Groendijk R, Poitevin O, van Deijk JR, et
721 al. Declines in moth populations stress the need for conserving dark nights. *Glob Chang Biol.*
722 2018;24: 925–932. doi:10.1111/gcb.14008

723 65. Paksarian D, Rudolph KE, Stapp EK, Dunster GP, He J, Mennitt D, et al. Association of Outdoor
724 Artificial Light at Night with Mental Disorders and Sleep Patterns among US Adolescents. *JAMA*
725 *Psychiatry.* 2020 [cited 27 Aug 2020]. doi:10.1001/jamapsychiatry.2020.1935

726 66. Perfito N, Jeong SY, Silverin B, Calisi RM, Bentley GE, Hau M. Anticipating spring: Wild populations
727 of great tits (*Parus major*) differ in expression of key genes for photoperiodic time measurement.
728 PLoS One. 2012;7. doi:10.1371/journal.pone.0034997

729 67. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: An
730 integrated and extendable desktop software platform for the organization and analysis of
731 sequence data. Bioinformatics. 2012;28: 1647–1649. doi:10.1093/bioinformatics/bts199

732 68. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, et al. Accurate
733 normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal
734 control genes. Genome Biol. 2002;3: research0034.1. doi:10.1186/gb-2002-3-7-research0034

735 69. Dijk F, Kraal-Muller E, Kamphuis W. Ischemia-Induced Changes of AMPA-Type Glutamate Receptor
736 Subunit Expression Pattern in the Rat Retina: A Real-Time Quantitative PCR Study. Investig
737 Ophthalmol Vis Sci. 2004;45: 330–341. doi:10.1167/iops.03-0285

738 70. Scheltema RA, Jankevics A, Jansen RC, Swertz MA, Breitling R. PeakML/mzMatch: A file format,
739 Java library, R library, and tool-chain for mass spectrometry data analysis. Anal Chem. 2011;83:
740 2786–2793. doi:10.1021/ac2000994

741 71. Tautenhahn R, Bottcher C, Neumann S. Highly sensitive feature detection for high resolution
742 LC/MS. BMC Bioinformatics. 2008;9: 1–16. doi:10.1186/1471-2105-9-504

743 72. Creek DJ, Jankevics A, Burgess KEV, Breitling R, Barrett MP. IDEOM: An Excel interface for analysis
744 of LC-MS-based metabolomics data. Bioinformatics. 2012;28: 1048–1049.
745 doi:10.1093/bioinformatics/bts069

746 73. R Development Core Team. R: A language and environment for statistical computing. URL
747 <http://www.R-project.org>. Vienna: R Foundation for Statistical Computing; 2015.
748
749

Acknowledgments: This work was supported by a Wellcome Trust grant to B.H and D.M.D (097821/Z/11/Z), a Marie-Curie Career Integration Grant to B.H. (ECCIG (618578) Wildclocks), a NERC Highlight Topics grant to D.M.D. (NE/S005773/1) and the Dutch Technology Foundation (STW). We thank Kamiel Spoelstra, Takashi Yoshimura and Bill Schwartz for fruitful discussions on the results of this study.

Author Contributions: DMD, MdJ, MEV and BH designed the study. DMD, MdJ, PBD, LT and BH collected the data and samples. DMD, KvO, POS, EA, CAM, MB, JC performed the gene expression assays. GB performed the metabolomics analyses. DMD conducted the statistical analyses. DMD and BH wrote the paper.

Competing Interest Statement: We declare no competing interests.