Touch-triggered bZIP translocation regulates elongation and secondary wall biosynthesis

Joshua H. Coomey^{1,2}, Kirk J.-M. MacKinnon^{1,3}, Pubudu P. Handakumbura^{1,2}, Ian W. McCahill^{1,2}, Gina M. Trabucco^{1,3}, Jessica Mazzola¹, Nicole A. Leblanc¹, Rithany Kheam¹, Miriam Hernandez-Romero^{1,2}, Kerrie Barry⁴, Lifeng Liu⁴, Ji E. Lee⁴, John P. Vogel⁴, Ronan C. O'Malley⁴, James J. Chambers⁵, Samuel P. Hazen^{1,2,3}

¹Biology Department, University of Massachusetts, Amherst, MA 01003, USA,

²Plant Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA,

³Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA,

⁴DOE Joint Genome Institute, Berkeley, CA 94720, USA,

⁵Institute for Applied Life Science, University of Massachusetts, Amherst, MA 01003, USA,

Author for correspondence:

Samuel P. Hazen, hazen@bio.umass.edu

ORCID

Joshua H. Coomey, <https://orcid.org/0000-0001-5866-6671>

Pubudu P. Handakumbura, <https://orcid.org/0000-0002-1708-8315>

Kerrie Barry, <https://orcid.org/0000-0002-8999-6785>

Ji E. Lee, <https://orcid.org/0000-0001-8325-1689>

John P. Vogel, <https://orcid.org/0000-0003-1786-2689>

James J. Chambers, <https://orcid.org/0000-0003-3883-8215>

Samuel P. Hazen, <https://orcid.org/0000-0001-9131-8994>

AUTHOR CONTRIBUTIONS

JHC and SH conceived and designed the study. JHC, PH, MH-R, JJC developed techniques and reagents. JHC, PH, GT, JM, NL, RK, KB, LL, JL, ZV, and RO acquired the data. JHC, KM, and SH analyzed and interpreted the data. JHC, KM, IM, SH drafted the manuscript. SH agrees to serve as the author responsible for contact and ensures communication.

ABSTRACT

Plant growth is mediated by the integration of internal and external cues, perceived by cells and transduced into a developmental program that gives rise to cell division, elongation, and wall thickening. Extra-, inter-, and intra- physical cellular forces contribute to this regulation. Across the plant kingdom thigmomorphogenesis is widely observed to alter plant morphology by reducing stem height and increasing stem diameter. The transcriptome is highly responsive to touch, including components of calcium signalling pathways and cell wall modification. One aspect of this cascade involves gibberellins and bZIP family transcription factors. Here, we present data connecting thigmomorphogenesis with secondary cell wall synthesis by gibberellin (GA) inactivation and bZIP translocation into the nucleus. *Brachypodium distachyon* SECONDARY WALL INTERACTING bZIP (SWIZ) protein translocated into the nucleus in response to mechanical stimulation. This translocation was mitigated by supplement with exogenous bioactive GA, and induced by chemical inhibition of GA synthesis. Misregulation of SWIZ expression resulted in plants hypersensitive to mechanical stimulation, with reduced height and increased secondary wall thickness. Classical touch responsive genes were upregulated in *B. distachyon* following touch, and we observe significant induction of the glycoside hydrolase 17 family, which may be indicative of a grass specific facet of thigmomorphogenesis. SWIZ binding to an E-box variant in exons and introns was associated with immediate activation followed by repression of gene expression. Thus, SWIZ is a transcriptional regulatory component of thigmomorphogenesis, which includes the thickening of secondary cell walls.

INTRODUCTION

Forces both internal and external to a cell influence growth. Turgor pressure in conjunction with anisotropic cell wall dynamics determine direction of growth and thus cell shape. Force perception between neighboring cells plays a critical role in the development and maintenance of tissue form and function, such as the lobed, interlocking, pavement cells on the leaf epidermis, or the developmental hotspots in the apical meristem (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi et al., 2019). Specific inter-cell forces result in dynamic remodelling of the cortical cytoskeleton, with subsequent changes in cellulose microfibril alignment and alterations to other cell wall components such as pectin methyl esterification (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi and Geitmann, 2018; Altartouri et al., 2019; Bidhendi et al., 2019). The classic hallmarks of touch responsive growth, or thigmomorphogenesis, include reduced plant height, increased radial growth in plants with a cambial meristem, increased branching, and delayed flowering time (Jaffe, 1973; Biro et al., 1980; Jaffe et al., 1980; Braam, 2004). These attributes have been leveraged by farmers for hundreds of years. As early as 1680 records show Japanese farmers tread on young wheat and barley seedlings to elicit increased branching, spikes per plant, and grain weight per plant, along with stronger roots (Iida, 2014). This practice, known as mugifumi, continues today with mechanized rollers.

Calcium is a critical cellular signalling molecule, and cytosolic Ca^{2+} fluctuations are induced by mechanical stimuli and implicated in mechanosignaling (Monshausen et al., 2009; Szczegieliński et al., 2012). Mechanical stimulus can significantly remodel gene expression (Braam and Davis, 1990; Braam, 2004; Lee et al., 2005). The so-called *TOUCH* (*TCH*) genes in *Arabidopsis thaliana*, encode calmodulin (*AtTCH1/AtCaM2*), calmodulin-like proteins (*AtTCH2/AtCML24*, *AtTCH3/CML12*), and a xyloglucan endotransglucosylase/hydrolase (*AtTCH4/AtXTH22*) (Braam and Davis, 1990). Touch responsive gene expression patterns often overlap with other stimuli such as dark, cold, and hormone treatment (Polisensky and Braam, 1996; Lee et al., 2005). In addition to calcium binding and signaling, genes related to cell wall modification and a variety of transcription factors and kinases are regulated by mechanical stimulus, as well as genes involved in hormone homeostasis and signalling.

The dynamic control of bioactive cellular gibberellic acid (GA) is linked to thigmomorphogenesis. GA is a growth promoting hormone derived through multiple enzymatic steps from the diterpene pathway, with interconvertible bioactive and inactive species. The bioactive forms of GA interact with protein receptors that regulate transcription factors controlling growth. Mechanical force application results in the inactivation of cellular GA by GA20 oxidases (Lange and Lange, 2015). The addition of exogenous bioactive GA to mechanically stressed plants remedies their height phenotype, and mutations in *Atga20ox* do not show thigmomorphogenic phenotypes under mechanical stress treatment (Lange and Lange, 2015). Recent work studying the translocation dynamics of several bZIP family transcription factors has provided new links for calcium and GA signalling in mechanoperception signalling.

Group I bZIPs are also implicated in mechanosensing. AtVIP1 and related Group I bZIP proteins translocate from the cytoplasm to the nucleus in response to a variety of biotic and abiotic stimuli, including hypo-osmotic conditions (Tsugama et al., 2012; Tsugama et al., 2014; Tsugama et al., 2016). This translocation appears to be dependent on protein phosphorylation, either from MITOGEN ACTIVATED PROTEIN KINASE 3 during pathogen invasion, or via calcium dependent protein kinases (CDPK). AtVIP1 interacts with calmodulins, the calcium binding proteins involved in many calcium signalling events. The Group I bZIP *Nt REPRESSOR OF SHOOT GROWTH* (*NtRSG*) in tobacco plays a role in maintaining GA homeostasis, wherein it translocates from the cytoplasm to the nucleus in response to cellular bioactive GA levels (Igarashi et al., 2001; Ishida et al., 2008; Fukazawa et al., 2010). Under standard conditions, NtRSG is largely localized in the cytoplasm, and to some extent in the nucleus, where it directly activates expression of *NtEnt-kaurene oxidase*, an enzyme at an early step in the GA biosynthesis pathway. After cellular GAs were chemically inhibited, NtRSG translocated to the nucleus and also activated *NtGA20oxidase1* (*NtGA20ox1*), an enzyme further downstream in the GA biosynthesis pathway that is in part responsible for converting GA species to their bioactive form (Fukazawa et al., 2010; Fukazawa et al., 2011). This suggests that nuclear-localized NtRSG acts to promote bioactive GA synthesis in response to low GA conditions. As with AtVIP1, NtRSG translocation relies on CDPK mediated phosphorylation (Ishida et al., 2008; Ito et al.,

2017). Both AtVIP1 and NtRSG associate with 14-3-3 proteins in the cytoplasm while phosphorylated (Ishida et al., 2004; Ito et al., 2014; Ito et al., 2017; Tsugama et al., 2018). Phosphatase activity causes NtRSG and the related AtbZIP29 to dissociate from the 14-3-3 protein and enter the nucleus (Van Leene et al., 2016). *NtRSG* is named for the dwarf phenotype observed when a dominant negative form of the protein is overexpressed. In these dwarf plants, there were reduced levels of bioactive GA and reduced elongation of stem internode cells (Fukazawa et al., 2000).

The development of secondary cell walls is a distinct and critical developmental process in plant growth. Following cellular elongation, some cells, particularly in vascular and structural tissues, increase their mechanical strength by depositing a thick layer of rigid material between the plasma membrane and primary cell wall. These secondary walls are made of crystalline cellulose interwoven with hemicelluloses and phenolic lignin polymers. Although functionally similar, secondary walls in monocotyledonous plants have key differences from those in eudicots, including distinct hemicellulose chemistry and differences in lignin biosynthesis. Grasses also produce mixed-linkage glucans (MLGs), a wall polysaccharide that is rarely found outside the commelinid monocots (Coomey et al., 2020). The synthesis and deposition of secondary wall polymers is tightly controlled spatially and temporally by a well defined transcriptional regulatory network. Evidence in both eudicots and grasses suggest a system of feed-forward loops regulating transcription of wall synthesizing enzymes, with NAC family transcription factors activating wall synthesis genes as well as MYB family and other transcription factors that further promote secondary wall synthesis (McCahill and Hazen, 2019). These networks are very similar between grasses and eudicots, with some components in each that have yet to be described in the other. There are currently no bZIP family members in any secondary wall regulatory model.

Beyond differences at the chemical and regulatory levels, grasses employ a fundamentally different growth mechanism than eudicots. Grasses do not have a cambium layer, and thus no lateral meristem. Stem elongation comes from cellular division and elongation in a discrete series of intercalary meristems, called nodes, with one internode region elongating and pushing up subsequent nodes. Detailed studies of thigmomorphogenesis have been conducted almost exclusively in eudicots. However, recent work in the model cereal grass *Brachypodium distachyon* shows general overlap with conventional dicot thigmomorphogenesis, albeit with some notable differences, such as no change in stem diameter and increased flowering time (Gladala-Kostarz et al., 2020).

Thigmomorphogenesis is a widely observed phenomenon that results in reduced height, increased radial growth, and increased branching. The mechanisms behind this form of growth are not yet fully understood, but involve aspects of hormone regulation, Ca^{2+} signalling, Group I bZIP intracellular translocation, and changes in gene expression. Here we describe the function of a *B. distachyon* bZIP transcription factor, SECONDARY WALL ASSOCIATED bZIP (Bradi1g17700) and its role in touch response and cell wall biosynthesis.

RESULTS

SWIZ is a Group I bZIP transcription factor and candidate cell wall regulator.

To identify genes involved in the regulation of secondary cell wall thickening, transcript abundance was measured in *B. distachyon* leaf, root, and stem tissue (Trabucco et al., 2013). A gene annotated as a bZIP transcription factor, Bradi1g17700, was highly expressed in root and stem relative to leaf (**Fig. 1A**). Bradi1g17700 is also a member of a 112 gene coexpression network (**Supplemental Table S1**) that includes genes highly expressed in the peduncle (Sibout et al., 2017). Phylogenetic analysis of Bradi1g17700, hereinafter referred to as *SECONDARY WALL INTERACTING bZIP* (*SWIZ*), amino acid sequence shows it to be an ortholog of the *A. thaliana* Group I bZIPs (Jakoby et al., 2002; Dröge-Laser et al., 2018), and closely related to *AtbZIP18* and *AtbZIP52* (**Fig. 1B**).

SWIZ translocates into the nucleus in response to mechanical stimulus

Given the translocation dynamics described for other Group I bZIPs, we hypothesized that *SWIZ* protein may similarly translocate within the cell in response to mechanical force. To test this, the roots of transgenic plants overexpressing either *SWIZ* fused to *GFP* (*SWIZ:GFP-OE*) or *GFP* alone (*GFP-OE*) were observed following a mechanical stimulus (**Fig 2A**). In control plants, *GFP* signal was present in both the cytosol and nucleus, which remained constant over the imaging period (**Fig 2B**). *GFP* signal was mostly observed in the cytosol in *SWIZ:GFP-OE* plants, but following mechanical stimulus, nuclear *GFP* signal increased substantially, reaching a peak around 30 min post stimulus and returned to near basal levels by about 60 min (**Fig 2B**).

The dynamics and repeatability of *SWIZ* nuclear translocation were further investigated by sequential stimulus treatments. Touch response to stimulus can saturate at a certain number of treatments (Martin et al., 2010; Leblanc-Fournier et al., 2014; Moulia et al., 2015). To test if *SWIZ* translocation dynamics varied after repeated treatments, we applied mechanical force to *SWIZ:GFP-OE* roots at regular intervals. A second stimulus was given 90 min after the first, and a third at 180 min. Following each mechanical stimulation, *SWIZ* consistently translocated from cytoplasm to nucleus (**Fig 2D**). This suggests that *SWIZ* translocation dynamics are not impacted by repeated stimulus events 90 min apart.

When conducting these translocation assays, root tissue in the field of view was treated with mechanical force and nuclei in that region were tracked and quantified for fluorescence. To determine if the *SWIZ* translocation signal is transmitted outside the specific region of stimulus, two regions of the same *SWIZ:GFP-OE* root separated by 3 cm were simultaneously observed. The stimulated region showed typical *SWIZ:GFP* nuclear signal accumulation and in the region below no translocation was observed (**Fig 2C**). At 120 min, the treatments were reversed, with the lower root region receiving a stimulus while the upper region was unperturbed. The lower region showed *SWIZ:GFP* nuclear translocation while the upper region did not. Thus, the touch stimulus perceived by *SWIZ* is localized to directly perturbed cells.

***B. distachyon* shows conventional thigmomorphogenic phenotypes of reduced height and increased branching.**

Given the touch responsive nature of SWIZ protein translocation and the evidence suggesting a role as a cell wall regulator, we hypothesized that SWIZ may play a role in mediating thigmomorphogenesis. To investigate this, we first established a touch response assay and assessed the thigmomorphogenic response in *B. distachyon*. Wildtype plants were perturbed with a metal bar once every 90 min for two or three weeks (**Supplemental Fig 1**). After the treatment period, all plants were allowed to recover and grow undisturbed until senescence (**Fig 3A**). Two week stressed plants were significantly shorter than controls, and three week stressed plants were shorter still (**Fig 3B**). Despite this difference in height, there was not a significant difference between the groups in terms of aboveground biomass weight (**Fig 3C**). Three week stressed plants had significantly more branches ($p < 0.05$), with a non-significant increase in two week stressed plants (**Fig 3D**).

The effect of repeated mechanical stress on stem biology and interfascicular fiber cell walls was assessed by making transverse sections of the peduncle and the third internode (**Fig 3E**). The mechanical stress assay contacted the growing plant at ~5 cm above the soil, such that the third internode region had already elongated prior to stress treatment, while the developing peduncle was directly contacted by the metal bar during elongation. In this way, the third internode served as an internal control tissue that did not experience direct mechanical perturbation. Sections were stained with phloroglucinol-HCl to detect lignin in secondary cell walls. The degree of staining and stem morphology in the third internode did not appear to be impacted by mechanical perturbation, but the peduncle showed distinct morphological changes in the three week stressed plants (**Fig 3E**). Interfascicular fiber cell walls were more lightly stained, and large, non-lignified pockets containing starch granules were evident around the stem periphery, between the minor bundles (**Supplemental Fig 2**). There was no significant difference in fiber wall thickness between the internode regions or from differing touch treatments (**Fig 3F**).

Thigmomorphogenesis in *SWIZ* genetic reagents

To test the role of SWIZ in regulating thigmomorphogenesis and secondary cell wall development we tested wildtype, *SWIZ-OE*, and *swiz-amiRNA* under two weeks of mechanical perturbation as described above (**Supplemental Fig 3**). In control conditions, there were no differences among genotypes. In mechanically stressed conditions, both *SWIZ-OE* and *swiz-amiRNA* were significantly shorter than wildtype, and *swiz-amiRNA* had significantly more branches (**Fig 4A-C**).

Transverse sections of the stem were made in the third elongated internode and the peduncle and stained with phloroglucinol-HCl (**Fig 4D**). The third internode was not directly contacted during mechanical perturbation, and as expected these sections did not show a difference in staining or fiber wall thickness compared to control conditions. However, both *SWIZ-OE* and *swiz-amiRNA* stained lighter than wildtype, independent of mechanical perturbation. The peduncle was directly

contacted during mechanical perturbation, and exhibited comparable lignin staining in both *SWIZ-OE* and *swiz-miRNA* to wildtype under touch perturbation, while remaining lighter staining than wildtype in control conditions. *SWIZ-OE* touched peduncles showed significantly thicker interfascicular fiber cell walls (**Fig 4E**). Thus, positive or negative modulation of *SWIZ* transcript interferes with lignin accumulation in control conditions, and *SWIZ-OE* wall thickening was more responsive to touch relative to no treatment.

Transcriptional response to touch

To assess the impact of *SWIZ* on gene expression in combination with touch treatment, we measured transcript abundance by sequencing mRNA in the roots of wildtype, *SWIZ-OE*, and *swiz-miRNA*, just prior to touch and at 10, 30, and 60 min following touch treatment (**Fig 5A**). Principal component analysis of transcript abundance shows the greatest amount of variance is attributed to the first principal component, 53%, where samples cluster based on genotype (**Fig 5B**). The second principal component, which accounts for 15% of the variance, distinguished between pre-touch and 10 min following touch, with the last two time points clustering similarly.

The wildtype transcriptome was massively remodelled in response to touch, with 8,902 transcripts differentially expressed ($q < 0.1$) at 10 min post touch, 5,682 transcripts at 30 min, and 7,672 transcripts at 60 min (**Supplemental Tables 2-4**). Touch response studies in *A. thaliana* describe similar changes that include numerous calmodulin (CAM), calmodulin-like (CmL), and xyloglucan endo-transglycosylase/hydrolase (XTH) activity (**Fig 5C**) (Braam and Davis, 1990; Lee et al., 2005). Based on homology with *TOUCH1* (*TCH1*), *TCH2*, and *TCH3*, 79 CAM and CmL genes were identified, the majority of which showed upregulation in wildtype plants following touch treatment (**Supplemental Table 5**). Similarly, homology with *TCH4* identified 37 *B. distachyon* *XTH* and *XTH-like* genes (Eklöf and Brumer, 2010). Of these, 8 genes were upregulated by touch in wildtype plants, including the two with the highest similarity to *TCH4* (**Fig 5D**) (**Supplemental Table 6**). Since xyloglucans are in relatively low abundance in grass cell walls compared to dicots, we also investigated the effect of touch on other families of the glycosyl hydrolase superfamily (Tyler et al., 2010). In particular, GH17 genes were differentially regulated following touch. The GH17 family is broadly described as modifying β -1,3-glucans. We measured transcript abundance for 35 of the 53 annotated GH17 family members, with 8, 11, and 16 members showing differential expression at 10, 30 and 60 min respectively. Fisher's exact test determined significant enrichment of GH17 expression at time 60 ($p = 9.307e-3$) (**Fig 5D, Supplemental Table 7**). Online resources are available to explore the RNA-seq data at <https://hazenlab.shinyapps.io/swiztc/>.

Cell wall genes are downregulated immediately following touch, then induced concurrent with *SWIZ* nuclear translocation and more strongly in *SWIZ-OE*

Genes related to secondary cell wall synthesis were immediately downregulated in wildtype plants following touch, but were subsequently upregulated (**Fig 6**). Xylan, cellulose, MLG, and callose synthesis associated genes were upregulated 30 and 60 min following touch, while lignin

biosynthesis and BAHD acyltransferase genes were upregulated at 60 min. Network analysis of gene expression patterns over time using the iDREM software showed that in *SWIZ-OE* plants, cell wall genes were more rapidly induced (**Supplemental Fig 4A-C**). Prior to touch, nearly all of the cell wall associated genes investigated were significantly downregulated relative to wildtype, then significantly upregulated following touch (**Fig 6**). In *SWIZ-OE*, cell wall related terms were enriched in paths that had stronger upregulation earlier than the same terms in wildtype (**Supplemental Fig 4A-C**). Notably, aspects of cell wall polysaccharide biosynthesis, particularly glucan biosynthesis, were significantly enriched in *SWIZ-OE* path D, while not significantly enriched in wildtype or *swiz-amiRNA* (**Supplemental Table 8**). This is mirrored when looking at specific gene expression, with primary wall associated CESAs, MLG, and callose synthesis genes all significantly upregulated following touch in *SWIZ-OE*. In *swiz-amiRNA*, cell wall related genes were generally not differentially expressed compared to wildtype (**Fig 6**). Thus, cell wall genes were immediately repressed by touch, followed by significant activation. That activation occurs more rapidly in *SWIZ-OE* plants and is associated with the timing of SWIZ protein nuclear translocation.

SWIZ protein binding in the gene body is associated with dynamic changes in gene expression

Next we investigated the direct binding targets of SWIZ protein by performing DNA affinity purification sequencing (DAP-seq) with whole genomic DNA. In total, SWIZ interacted with 2,455 distinct positions in the genome. Those regions were significantly enriched for two sequence motifs, (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G) that are variants of the E-box motif (**Fig 7A**). A plurality of the binding sites were between the translational start and stop sites with 21% of the peaks in introns and 21% in exons (**Fig 7B**). Fewer than 10% of the peaks occurred in UTRs and the promoter (defined as 5 kb upstream of the 5' UTR) and intergenic regions each accounted for 25% of the binding sites. Thus, SWIZ protein preferentially bound to an E-box like motif and often in the gene body (**Fig 7C**).

To investigate the effects of possible SWIZ binding on gene expression, we compared the genes differentially expressed in the *SWIZ-OE* and *swiz-amiRNA* touch response RNA-seq time courses with the promoter and gene body binding targets identified by the DAP-seq assay. Prior to touch, genes with promoter-SWIZ protein interactions were most often downregulated in *SWIZ-OE* and upregulated in *swiz-amiRNA* (**Table 1**). That same trend was observed for gene body targets in untouched plants. However, some SWIZ targets behaved differently following touch. SWIZ promoter binding targets were most often upregulated in *SWIZ-OE* in all three post-touch time points. Thus, SWIZ promoter binding was coincident with repression, but following a touch stimulus, activation was more prominent. However, this difference between untreated and touched was not as pronounced in gene body targets, with more upregulated genes 10 min following touch and more downregulated 30 and 60 min following touch. To further explore these trends we conducted iDREM cluster analysis of the differentially expressed genes that

were also SWIZ protein targets (**Fig 7D**). For all genotypes, the network analysis revealed a pathway to increased gene expression by 60 min following touch. The network pathway trend for both promoter and gene binding targets in wildtype was immediate repression at 10 min followed by increased expression at 30 and 60 min post-stimulus. A unique pathway was observed among gene body binding targets in *SWIZ-OE*; transcript abundance was reduced following touch. Thus, SWIZ binding to a promoter or gene body was more strongly associated with increased expression with the exception of gene body targets in touched *SWIZ-OE* plants, which were more strongly repressed.

Among the genes differentially expressed in response to both touch, *SWIZ-OE* transgene, and bound by SWIZ in the first intron is the mixed glucan synthase *CSLF6* (**Fig 8A**). The transcript abundance of *GA2ox3*, which inactivates bioactive GA, was bound by SWIZ protein in both the gene body and 3'UTR and expression significantly increased 30 min after touch in *SWIZ-OE* (**Fig 8B**). An example of gene body binding repression in Bradi1g30252, the closest ortholog to a wheat transcription factor induced by hypoxia, *TaMYB1* (Lee 2007). This gene was touch induced in wildtype and repressed in *SWIZ-OE* with a binding site in the 5'UTR (**Fig 8C**). The NAC35 transcription factor Bradi2g28020 exhibited a similar expression to *MYB44* and SWIZ bound the promoter region (**Fig 8D**).

SWIZ translocation is sensitive to cellular GA levels, and may act in maintaining GA homeostasis following mechanical stimuli

Mechanical forces have been shown to inactivate cellular GA, and that cellular GA levels dictate cytosolic versus nuclear localizations of *N. benthamiana* Group I bZIPs similar to SWIZ (Fukazawa et al., 2010; Lange and Lange, 2015). To investigate this in relationship to SWIZ, *SWIZ:GFP-OE* roots were treated with bioactive GA or paclobutrazol, an inhibitor of GA synthesis. SWIZ:GFP signal amplitude was reduced in roots treated with bioactive GA, particularly at the 50 and 100 mM level. After 60 min the nuclear signal had returned to near pre-stimulus levels, and a second stimulus was applied (**Fig 9A**). This time all GA treated roots showed a similar reduction in nuclear signal, nearly half of the untreated control. These results suggest that the addition of exogenous GA reduces SWIZ translocation to the nucleus following mechanical force application.

Roots treated with 10 mM paclobutrazol did not show altered translocation dynamics compared to the untreated control, but roots treated with 50 or 100 mM paclobutrazol initially showed greater SWIZ:GFP nuclear localization, even prior to mechanical stimulus (**Fig 9B**). SWIZ:GFP signal in the 100 mM paclobutrazol treated roots was high prior to treatment and remained high even after the other nuclear signals diminished. This suggests that pharmacological reduction of bioactive GA levels triggers SWIZ nuclear translocation even without the application of mechanical force.

Given the role of GA in touch responsive bZIP translocation dynamics, we measured how GA related gene expression changed in response touch. In wildtype following touch, GA synthesis and GA activating genes *KS*, *KAO4*, and *GA20ox4* were downregulated, while the genes encoding GA inactivating enzymes *GA2ox4* and *GA2ox9* were significantly upregulated (**Fig 9C**). In *SWIZ-OE*, *KS* was upregulated prior to touch and 10 min post touch. *GA20ox4*, a gene responsible for synthesizing bioactive GA, was downregulated prior to touch in *SWIZ-OE* plants, then significantly upregulated following touch. At 30 and 60 min *GA2ox4* and *GA2ox9* were upregulated, along with the DELLA protein Rht-1 (**Fig 9C**). Here we show consistent patterns of gene expression associated with the touch inactivation of GA and that SWIZ protein promotes expression of GA synthesis and GA activation genes following mechanical perturbation.

DISCUSSION

Touch stimulus is generally an inhibitor of plant elongation growth, but promotes branching and radial expansion (Jaffe, 1973; Braam, 2004; Chehab et al., 2009). The majority of this work has been done in dicots, where increased radial growth has been associated with greater deposition of secondary wall forming cells, particularly in woody species such as poplar (Biro et al., 1980; Coutand et al., 2009; Börnke and Rocks, 2018; Roignant et al., 2018; Niez et al., 2019). Our understanding of thigmomorphogenesis in grasses is limited, and mostly in the agricultural context of lodging (Shah et al., 2019). While these studies highlight the importance of stem strength and associated cell wall defects, they do relatively little to elucidate the mechanosensitive response. Recent work by Gladala-Kostarz et al (2020) describes grass touch response to both wind and direct mechanical treatment, with an emphasis on cell walls and stem anatomy. Touch treatment significantly increased lignin content, and also altered the relative abundance of wall bound hydroxycinnamates. Wall polysaccharide content, particularly glucose and galactose, also increased. Consistent with increased secondary wall lignification, touched plants had increased wall stiffness and decreased saccharification. Our study of wildtype *B. distachyon* plants with two or three weeks of regular touch revealed a decrease in plant height. We also observed an increase in branching with increasing touch treatment. However, we did not observe a change in total biomass at senescence, perhaps due to compensation by increased branching.

Specific gene expression patterns have become molecular hallmarks of plant touch response, most notably induction of the *TCH* and similar genes. Orthologs of wall modifying XTHs and CAM/CML signalling genes are upregulated in response to mechanical stimulation across species, and as we present here, in *B. distachyon*. Touch elicits major global changes in gene expression, with 2.5% to 10% of the genes assayed in *A. thaliana* and poplar differentially expressed (Lee et al., 2005; Pomiès et al., 2017; Van Moerkercke et al., 2019). In sorghum (*Sorghum bicolor*), leaf sheath imposes mechanical constraints on emerging buds, and removing this dramatically altered gene expression, with 42% of genes differentially expressed over a 9 h period (Liu and Finlayson, 2019). Our observations of short term (<2 h) touch responses within stress related pathways align with observations from poplar and *A. thaliana*, with rapid induction

of *TCH*, biotic, and abiotic stress related genes (Pomiès et al., 2017). Canonical touch responsive genes such as the orthologs of *TCH1-4* and related *XTH* and *CML* genes were upregulated immediately following stimulus. However, we also note the previously unreported touch-induction of GH17 family genes, as well as *CSLF6*, suggesting a role for MLG and β -1,3 glucan modification in grass touch response.

In both poplar and in our analysis of *B. distachyon* touch responses, touch-regulated expression of secondary cell wall related transcripts differed between early and late timepoints. Upon touch, we observed immediate repression of many key cell wall biosynthetic enzymes, followed by significant upregulation one hour after stimulus. In poplar, no significant repression of these transcripts was reported, but terms related to wood or cell wall synthesis were also not enriched among touch-regulated transcripts until 24 or 72 hours after touch (Pomiès et al., 2017). While differences in the method used to quantify gene expression, and in the temporal sampling scheme complicate direct comparisons of plant touch response experiments, this may suggest that delayed induction of secondary cell wall transcripts is a common feature of plant touch responses.

Thigmomorphogenesis, GA physiology, and secondary cell walls

Our understanding of touch responsive gene expression is further informed by the dynamics of SWIZ nuclear translocation. The timing of SWIZ nuclear accumulation following mechanical stimulus was consistent with observations of *A. thaliana* Group I bZIPs (Tsugama et al., 2014; Tsugama et al., 2016). However, by adopting a finer temporal sampling scheme, and quantifying nuclear signal sooner after touch (2 vs. 30 min), our results clarify the rapid and immediate nature of this translocation. Furthermore, we show that the speed and magnitude of SWIZ nuclear accumulation is not diminished over three successive touch treatments. Notably, poplar transcriptional responses to a second touch stimulus were significantly attenuated relative to a single treatment, and in separate experiments, trees subjected to a daily touch treatment over four days no longer showed touch-induced increases in their rates of radial growth (Martin et al., 2010; Pomiès et al., 2017). Thus, if *B. distachyon* is likewise desensitized to repeated touch, the high repeatability of SWIZ translocation might implicate a mechanism downstream of touch perception and bZIP translocation in mediating that shift. Like the touch-responsive transcription factor AtVIP1, SWIZ translocation occurred in the specific area of touch treatment, suggesting a cellular rather than tissue level response. SWIZ also displays a highly similar pattern of GA responsive translocation to the well-studied bZIP NtRSG (Ishida et al., 2004; Fukazawa et al., 2010; Fukazawa et al., 2011). Exogenous bioactive GA dampened SWIZ nuclear translocation, while inhibition of GA synthesis with paclobutrazol induced translocation, even prior to touch. Furthermore, like NtRSG, SWIZ appears to both respond to and regulate GA bioactivity. In wildtype, GA inactivating enzymes GA2ox4 and GA2ox9 were upregulated following touch, consistent with previous reports (Lange and Lange, 2015). In *SWIZ-OE* we also observe touch induced expression of *GA20ox4*, a GA activating enzyme, and as with NtRSG, this response can restore bioactive GA following touch induced degradation (Ishida et al., 2004; Fukazawa et al.,

2010; Fukazawa et al., 2011). Altered GA pathways are a well described mechanism for reduced stem elongation (Lange and Lange, 2006; Nagai et al., 2020), and dwarfing has been described for dominant negative mutants of Group I bZIPs *NtRSG* and *OsRF2A* (Fukazawa et al., 2000; Dai et al., 2003). *SWIZ-OE* and *swiz-miRNA* were both significantly shorter following touch treatment, suggesting that GA misregulation may contribute to this thigmomorphogenic phenotype. Another component of this reduced elongation may be increased secondary cell wall deposition, as this could terminate elongation. Prior to touch, almost the entire monolignol biosynthetic pathway, as well as laccase genes that radicalize monolignols, and BAHD acyltransferases that add hydroxycinnamic acids to wall polymers, are significantly downregulated in *SWIZ-OE* plants. Following touch, most of these genes become significantly upregulated. This activity is consistent with our observations of increased lignin staining in touched *SWIZ-OE* peduncles, and aligns with the compositional data presented by Gladala-Kostarz et al (2020). This activation may be a result of direct binding of SWIZ to modulate expression, or an indirect effect from another transcriptional regulator such as the NAC transcription factor *SWN5*, capable of activating the full developmental program for secondary cell wall synthesis (Valdivia et al., 2013), which is significantly upregulated in *SWIZ-OE* following touch.

While the E-box (CANNTG) was initially described as a bHLH binding motif, many bZIP groups in *A. thaliana* have been shown to bind similar sequences. The two closest *A. thaliana* orthologs of SWIZ, bZIP18 and 52, both bind (A/C)CAG(G/C)(T/C), while VIP1 binds (A/C)CAGCT(G/A) (O'Malley et al., 2016). Using the same technique, we identified the SWIZ binding motifs as (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G). Both of these sequences are similar to those bound by *A. thaliana* orthologs, although the presence of an ambiguous nucleotide in the core of one variant is not previously reported for other Group I bZIPs. Among genes bound by SWIZ *in vitro*, we observed both activation and repression following touch, suggesting a complex regulatory function for SWIZ. Furthermore, SWIZ direct binding sites were found in both promoter regions and in the body of genes and most often in genes activated by touch. SWIZ gene body binding targets tended to be repressed in *SWIZ-OE* plants following touch, without either being clearly associated with up or down regulation. AtVIP1 and AtbZIP29 have both been described as activators, particularly of genes related to biotic and abiotic stress response, cell cycle control, vascular development, and cell wall biogenesis (Yin et al., 1997; Ringli and Keller, 1998; Pitzschke et al., 2009; Van Leene et al., 2016). AtbZIP18 is suggested to have repressive roles (Tsumaga et al., 2012; Gibalová et al., 2017).

bZIPs are known to homo- and heterodimerize through their leucine zipper domains (Schütze et al., 2008), and in *A. thaliana* the combinatorial interactions of different bZIP groups have been fairly well described (Deppmann et al., 2004; Ehlert et al., 2006; Grigoryan and Keating, 2006; Weltmeier et al., 2006; Schütze et al., 2008). These interactions can have a synergistic effect on transcriptional activity, and can result in unique binding interactions (Schütze et al., 2008; Van Leene et al., 2016). It is conceivable that this heterodimerization may explain some aspects of

our results, such as why so many of the cell wall genes that are upregulated in *SWIZ-OE* after touch are significantly downregulated prior to touch. One hypothesis is that SWIZ, as with many other bZIPs, can regulate targets as a heterodimer. The abundance of SWIZ protein in the cytoplasm in *SWIZ-OE* plants prior to touch may act as a sink, sequestering interacting partners that would otherwise be regulating targets independent of a touch stimulus. Similarly, redundant or compensatory function among similar bZIPs may explain why the *swiz-amiRNA* plants did not show significant and opposite phenotypes from the *SWIZ-OE*. Further experimentation is needed to elucidate this bZIP regulatory network in response to touch and other stimuli.

Conclusions

Proteins orthologous to SWIZ in other species have been implicated in cell wall development and remodelling, but not cell wall thickening. SWIZ appears to control stem height and cell wall thickening in response to mechanical stimuli, which regulates SWIZ translocation into the nucleus by acting on the cellular bioactive GA pool. Touch has been shown to decrease cellular bioactive GA levels in *A. thaliana*, and Group I bZIPs are known to translocate to the nucleus in response to touch and to low bioactive GA. Direct mechanical stimulation and chemical inhibition of GA biosynthesis both caused SWIZ to translocate from the cytoplasm into the nucleus. This translocation can be reduced by supplementing with exogenous bioactive GA. GA biosynthesis and signalling genes were upregulated in *SWIZ-OE* lines following touch, indicating that SWIZ may act in managing GA homeostasis following mechanical perturbation. Altering SWIZ function may impact this GA homeostasis and result in reduced elongation following touch. Touch significantly remodeled the *B. distachyon* transcriptome, with notable changes in wall polysaccharide biosynthetic gene expression not previously reported. Enhanced SWIZ function through overexpression amplified the expression of cell wall related genes and synthesis of interfascicular fiber walls in touched tissues. Greater secondary wall synthesis in response to touch may also explain reduced stem elongation in *SWIZ-OE*. Together, the evidence presented here connects mechanosensitive bZIP translocation with dynamic changes in cellular GA levels that have been observed separately in other systems and shows a link between touch response and secondary cell wall transcriptional regulation (**Fig 10**).

MATERIALS AND METHODS

Phylogenetic analysis

Protein sequences described for *A. thaliana*, *B. distachyon*, and *O. sativa* (Liu and Chu, 2015) as Group II bZIPs were selected and searched against the most recent genome annotations in Phytozome v12.1 (<https://phytozome.jgi.doe.gov>). The *Nicotiana tabacum* homologs NtRSGa and NtRSGb were also added to the analysis. Protein sequences were aligned using the MAFFT service for multiple sequence alignment with the iterative refinement method L-INS-I (Kato et al., 2019). The alignment was used to construct the phylogenetic tree using a maximum likelihood analysis with a bootstrap resampling value of 1000 in W-IQ-TREE (Trifinopoulos et

al., 2016). All of the proteins included in the phylogenetic analysis are described in
Supplemental File 1.

Plant transformation

Overexpression and artificial microRNA transgenes and transgenic plants were constructed as previously described (Handakumbura et al., 2013). The full length coding sequence of Bradi1g17700 was amplified from cDNA and cloned into the pOL001 ubigate ori1 binary expression vector (Handakumbura et al., 2018) to make the *SWIZ-OE* transgene. The coding sequence lacking the stop codon was amplified and fused in frame with the coding sequence of *Aequorea victoria* enhanced GFP in the pOL001 vector to generate *SWIZ:GFP-OE*. Artificial microRNA constructs were generated as previously described (Handakumbura et al., 2013) to include a targeting sequence specific to the Bradi1g17700 transcript, CAGCCTGAACCTTCCCAGTCC. This sequence was identified using the WMD3 webtool (Schwab et al., 2006; Warthmann et al., 2008). The modified amiRNA sequence was cloned into pOL001 for overexpression.

Translocation assay

Seeds were surface sterilized and grown vertically on 1X MS media, pH 5.7, without sucrose for 6d at 28°C in the dark. After 6 d, seedlings were moved to treatment plates containing 1X MS media, pH 5.7, plus varying concentrations (0, 10, 50, or 100mM) of GA4, paclobutrazol. Roots were positioned such that they were completely submerged in media and relatively flat along the imaging plane. After a 6h incubation, plants were stimulated and imaged.

All observations were made on a Nikon A1R scanning confocal microscope using a Plan Apo 10x 0.5NA objective. Root areas to be observed were located by eye using transmitted light and then confirmed under confocal conditions using a 488 nm laser and green filtered PMT detector. The X, Y, and Z coordinate stage locations of each region were programmed into the Nikon NIS Elements 5.2 Advanced Research V6 software package for automated imaging. After all target regions were programmed, 30 min of imaging began pre-treatment, with images captured every 2 min.

To elicit the touch response, the observed root region was gently probed for 5 sec with a blunt metal probe while observing through the eyepiece. Images were captured for 60-90 min post treatment. For experiments with multiple stimulus events, the timelapse sequence was paused and roots were probed as described for the relevant stimulus events.

Analysis of GFP signal was done using the Nikon NIS Elements Advanced Research V5 software package. For each time series, the frame showing maximal nuclear signal was selected and used as a reference point. Using the General Analysis tool, a channel for GFP signal was established and thresholded for intensity and particle size to identify the nuclear regions. These regions were added to the timelapse image series as a binary layer and then converted to static regions of interest for quantification. The GFP intensity under each nuclear region of interest was calculated for the course of the timelapse and the average signal from each nucleus was plotted for intensity over time.

Plant growth

Accession Bd21-3 was used for all experiments. Seeds were stratified on wet paper towel wrapped in foil to exclude light for 10 days at 4°C before being planted in Promix BX potting mix in SC10 Ray Leach Cone-tainers (Stuewe & Sons Inc). Plants were grown in a Percival PGC-15 growth chamber with day/night conditions of 20h light at 24°C and 4h dark at 18°C, respectively.

Thigmomatic construction and operation

The Thigmomatic is a basic robotic device that sweeps plants with a metal bar at regular intervals to elicit a touch response. The device was constructed from aluminum V-Slot linear rail (Openbuilders Partstore, Monroeville, NJ) and bracket joints for the upright supports (20x20 mm), cross bars (20x20 mm), and tracks (20x40 mm). Two gantry carts ride along the 20x40 mm V-Slot linear rails, connected by a 6.35 mm diameter metal rod bolted to the carts. Their movement is powered by a belt driven linear actuator system using a NEMA 17 stepper motor with a 12V 18W AC/DC power supply. The stepper motor provides fine spatial control over the gantry cart position with bi-directional motion. Motor function is controlled by a Raspberry Pi 3B microcomputer equipped with a stepper motor HAT (Adafruit Industries, New York). The Thigmomatic was programmed to cover a specified distance in one direction once every 90 min.

Transverse stem sections and histology

The main stem of senesced plants was taken and the internode of interest removed and embedded in 8% agarose. A Leica VT1000 Vibratome was used to make 55 µm thick transverse sections. Histochemical staining was carried out using phloroglucinol-HCl as previously described (Matos et al., 2013). Images were obtained at 4, 10, and 20X using a Nikon Eclipse E200MV R light microscope and PixelINK 3 MP camera. Transverse sections imaged at 20x were used for cell wall thickness measurements. Interfascicular fiber cells separated by one cell layer from the mestome cells on the phloem side of major vascular bundles were targeted for measurement. Using ImageJ, lines were drawn across two walls of adjoining cells. The resulting line length was divided by two to give one cell wall width. Approximately 15 measurements were made for each plant.

RNA extraction and quantification and analysis

Seedlings were grown for 6 d on vertical 1X MS agar plates. Touch treatment was performed as described above using a metal probe along the entire length of the root. Untouched samples were collected immediately before touch treatment began, and touched samples were collected after the designated time. Three roots were pooled per biological replicate, and RNA was extracted using the Qiagen RNeasy Plant Mini Kit with on-column DNA digestion with RNase-free DNase I (Qiagen). Strand-specific libraries were prepared using the Illumina TruSeq kit. Libraries were sequenced using Illumina technology. Sequences were processed as previously described (MacKinnon et al., 2020) and then checked for differential expression using the likelihood ratio test. Specific contrasts (i.e., wildtype vs *SWIZ-OE*) were identified and compared using the Wald test. Statistical enrichment of gene families was assessed using Fisher's exact test. Raw read data were deposited in the European Nucleotide Archive for public access (Accession no.: XYZ).

DNA affinity purification sequencing

DNA affinity purification was carried out as previously described (Handakumbura et al., 2018). In brief, transcription factor coding sequences were HALO tagged and mixed with Bd21 genomic DNA for *in vitro* binding. Protein-DNA was crosslinked, fragmented, immunoprecipitated using the HALO antibody, barcoded, and sequenced. Reads were mapped back to the Bd21 genome using HiSat2 (Kim et al., 2015) to identify binding target loci. Peak calling and motif analysis was done using HOMER v4.10 (Hypergeometric Optimization of Motif EnRichment) suite (Heinz et al., 2010). Motif enrichment was calculated against the hypergeometric distribution; the significance threshold was set to $p < 0.05$. The nearest annotated gene to a bound peak was used for GO analysis.

Gene Ontology analysis

Phytozome was used to find orthologs for all *B. distachyon* v3.1 genes as the reciprocal best match to *A. thaliana* TAIRv10 protein sequences. Arabidopsis gene identifiers were submitted to g:Profiler (Raudvere et al., 2019) for KEGG and Wiki pathway enrichment analysis.

iDREM network analysis

Probabilistic graphical models that predict diverging gene expression paths and the points at which regulators may influence those paths were generated using iDREM (Ding et al., 2018). Briefly, this software applies an input-output hidden Markov model to time course gene expression data overlaid with static regulatory information, in this case SWIZ protein-DNA interactions identified from DAP-seq. GO analysis, described above, was also applied to the gene sets in each path identified by iDREM.

SUPPLEMENTAL MATERIALS

Supplemental Table S1. Cluster 56 from network analysis of gene expression atlas (Sibout et al. 2017) includes Bradi1g17700.

Supplemental Table S2. Differentially expressed genes in wildtype Bd21-3 root tissue 10 min after touch treatment.

Supplemental Table S3. Differentially expressed genes in wildtype Bd21-3 root tissue 30 min after touch treatment.

Supplemental Table S4. Differentially expressed genes in wildtype Bd21-3 root tissue 60 min after touch treatment.

Supplemental Table S5. Calmodulin and calmodulin-like touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S6. XTH and XTH-like touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S7. Glycoside hydrolase touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S8. GO analysis of genes in iDREM identified pathways derived from differential gene expression following touch in wildtype, swiz-amiRNA, and SWIZ-OE root tissue.

Supplemental Figure 1. Thigmomatic, an automated system to administer touch stimulus. (A) Overview of Thigmomatic inside a Percival PGC-15 growth chamber showing linear rail based frame (1), gantry carts (2), NEMA17 stepper motor (3), 12V 18W AC/DC power supply (4), and Raspberry Pi 3b microcomputer (5). (B) Thigmomatic making contact with a *Brachypodium distachyon* plant.

Supplemental Figure 2. Regions in the periphery of the stem without secondary cell walls containing starch granules. Lugol's iodine stained starch granules of peduncle transverse sections of *Brachypodium distachyon* treated with three weeks of mechanical stress. Arrow indicates an example starch granule. Scale bar = 100 μ m.

Supplemental Figure 3. Description of SWIZ transgenic reagents. (A) Diagram of SWIZ overexpression transgenes. The *ZmUbi* promoter was used to drive expression of the SWIZ coding sequence, either alone or fused in frame with *eGFP*. (B) Artificial microRNA construct for targeted SWIZ transcript knockdown. The maize ubiquitin promoter was used to drive expression of an amiRNA that has homology to the first SWIZ exon and forms a hairpin after transcription. The 21-mer nucleotide sequence used for target specificity is listed, with red

postions showing hairpin mismatches that are predicted to interact with the DICER complex. Relative level of *SWIZ* transcript measured by RT-qPCR in whole stem tissue was collected 1 d after inflorescence emergence in *SWIZ:GFP-OE* (C) and *swiz-amiRNA* (D) lines. LB, left border; *ZmUbi* prom, maize ubiquitin promoter; Hyg, hygromycin phosphotransferase gene; NOS, nopaline synthase terminator; RB, right border. ns: $p > 0.05$, : *: $p < 0.05$.

Supplemental Figure 4. Network analysis of differential gene expression paths and enriched GO terms. (A) Path determinations from iDREM time course analysis. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch. (B) Heatmap of selected GO term enrichment of genes from iDREM path analysis. Color indicates qvalue score, with a cutoff of 0.1. Terms without a colored block are not significantly enriched in that path.

Supplemental File 1. Gene names and amino acid sequences used in the *SWIZ* bZIP Group I phylogenetic analysis.

ACCESSION NUMBERS

AtbZIP18 (At2g40620), AtbZIP29 (At4g38900), AtbZIP52 (At1g06850), AtTCH1/AtCaM2 (At2g41110), AtTCH2/AtCML24 (At5g37770), AtTCH3/CML12 (At2g41100), AtTCH4/AtXTH22 (At5g57560), CAD1 (Bradi3g17920), CESA4 (Bradi4g28350), COMT6 (Bradi3g16530), CSLF6 (Bradi3g16307), GA2ox3 (Bradi2g50280), MYB44 (Bradi1g30252), NAC35 (Bradi2g28020), NtRSGa (Niben101Scf01150g00005), NtRSGb (Niben101Scf02191g00014), *SWIZ* (Bradi1g17700), VIP1 (At1g43700).

ACKNOWLEDGEMENTS

This work was supported by the Office of Science, Biological and Environmental Research, Department of Energy (DE-FG02-08ER64700DE and DE-SC0006641) and The National Science Foundation Division of Integrative Organismal Systems (NSF IOS-1558072) to S.P.H.. The work conducted by the US DOE Joint Genome Institute is supported by the Office of Science of the US Department of Energy under Contract no. DE-AC02-05CH11231. The microscopy data was gathered in the Light Microscopy Facility and Nikon Center of Excellence at the Institute for Applied Life Sciences, UMass Amherst with support from the Massachusetts Life Sciences Center.

Table 1. The count of differentially expressed genes relative to untouched wildtype plants that were SWIZ DAP-seq targets in either the promoter or gene body regions.

Genotype	Time after touch	Promoter		Gene body	
		Up	Down	Up	Down
	(min)	percent (count)			
<i>SWIZ-OE</i>	0	42.0 (103)	58.0 (142)	48.0 (286)	52.0 (310)
	10	57.0 (73)	43.0 (55)	53.0 (167)	47.0 (148)
	30	59.4 (79)	40.6 (54)	41.2 (136)	58.8 (194)
	60	64.2 (77)	35.8 (43)	45.7 (128)	54.3 (152)
<i>swiz-amiRNA</i>	0	67.1 (47)	32.9 (23)	60.0 (78)	40.0 (52)
	10	56.1 (37)	43.9 (29)	55.9 (76)	44.1 (60)
	30	65.0 (13)	35.0 (7)	29.4 (5)	70.6 (12)
	60	53.8 (7)	46.2 (6)	27.8 (5)	72.2 (13)

FIGURE LEGENDS

Figure 1. SWIZ is a Group I bZIP highly expressed in maturing stem and root. (A) *SWIZ* transcript abundance in *Brachypodium distachyon* leaf, root, and stem tissue measured by microarray. Mean +/- standard deviation of three biological replicates. (B) Phylogeny analysis of amino acid sequences from *B. distachyon* (green), *Oryza sativa* (blue), *Arabidopsis thaliana* (red), *Nicotiana tabacum* (orange) shows *SWIZ* (black dot). The phylogeny was reconstructed using maximum likelihood analysis with a bootstrap resampling value of 1000. The numbers labeled on the branches are the posterior probability to support each clade. The protein sequences, full locus IDs, and plant species with genome source are provided in Supplemental File 1.

Figure 2. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to stimulus and 30 min post stimulus. Roots were observed immediately following mechanical perturbation. (B) Quantification of nuclear signal. n=14-20 nuclei. (C) *SWIZ* translocation occurred in the local area of the stimulus. At 30 min, stimulus was applied to an upper region of the root, while at 120 min it was applied to a lower region approximately 3 cm below. n=109,

184 nuclei respectively for upper and lower regions. (D) *SWIZ:GFP-OE* roots were imaged by confocal microscopy with stimulus applied in the field of view at 0, 90, and 180 min. $n=126$ nuclei. (B-D) Images were taken every 2 min. Nuclear GFP signal was quantified in selected nuclei at each time point. The average nuclear GFP signal is represented by the line with error bars indicating standard error of the mean.

Figure 3. *Brachypodium distachyon* displays classic thigmomorphogenic phenotypes. One-week-old wildtype plants were treated with touch stimulus every 90 min. (A) Left to right, plants that experienced no stress, two weeks stress, three weeks stress and were then allowed to recover, imaged one week after the end of treatment (B) height, (C) aboveground non-grain biomass weight, and (D) branch number were measured at senescence. ns: $p > 0.05$, *: $p < 0.05$, $n = 5$ plants per treatment. (E) Transverse sections of the peduncle or third internode were taken from control, 2 week stressed, and 3 week stressed plants and stained with phloroglucinol-HCl. (F) Quantification of interfascicular fiber wall thickness. Scale bar = 100 μm . $n = 3$ plants per treatment.

Figure 4. *SWIZ* transgenes enhance thigmomorphogenic response. Wildtype, *SWIZ-OE*, and *swiz-amiRNA* plants were placed under control conditions or received two weeks of mechanical stress every 90 min. After senescence, plants were phenotyped for (A) height, (B) branch number, and (C) aboveground biomass weight. ns: $p > 0.05$, *: $p < 0.05$. $n = 5$ plants per treatment. (D) The main stem was collected and transverse sections made of the peduncle and third elongated internode. Sections were stained with phloroglucinol-HCl. Scale bar = 100 μm . (E) Quantification of interfascicular fiber wall thickness. $n = 4$ to 6 plants per genotype, per treatment.

Figure 5. Transcriptome analysis of touch response in *Brachypodium distachyon* roots. (A) Root tissue was sampled just prior to touch ($t=0$), and at 10, 30, and 60 min following touch treatment in wildtype, *SWIZ-OE*, and *swiz-amiRNA*. (B) Principal component analysis of variance in gene expression across samples shows the greatest difference corresponding to genotype, with the second greatest corresponding to time after touch. (C) Canonical, as well as novel touch responsive genes are upregulated in *B. distachyon* following touch. Closest orthologs of the *Arabidopsis thaliana* *TOUCH* genes encoding calmodulin-like (CML) and xyloglucan endo-transglycosylase/hydrolases (XTH) are upregulated following touch, as are previously unreported members of the glycosyl hydrolase 17 (GH17) family. Full list of gene expression for *TOUCH* and glycoside hydrolases defined by Tyler et al. (2010) in Supplemental Tables 5,6, and 7.

Figure 6. Gene expression analysis of cell wall related genes. Log fold-change of gene expression in wildtype, *SWIZ-OE*, and *swiz-amiRNA* relative to wildtype expression at time 0, pre-touch. Bar color indicates class of cell wall gene. Error bars indicate standard deviation of

three biological replicates. Asterisks represent significantly differentially expressed genes, q -value < 0.1 for Wald test of each time and genotype compared to wild type at time 0.

Figure 7. DNA affinity purification sequencing to determine SWIZ binding sites. (A) Top two most statistically enriched sequence motifs in SWIZ binding sites. (B) Distribution of binding sites across genomic features, relative to primary transcripts of the *Brachypodium distachyon* annotation v 3.1. (C) Relative distribution of binding sites centered on the transcriptional start site (TSS, blue dashed line), transcriptional termination site (TTS, red dashed line) represents the average length of all annotated transcripts, approximately 4.5 kb away from the TSS. (D) Path determinations from iDREM time course analysis of differentially expressed genes that also have DAP-seq binding sites. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch.

Figure. 8 SWIZ binding targets differentially expressed in response to touch and SWIZ. Gene expression over time of selected genes with SWIZ binding sites. Line graphs are the average transcript abundance of three biological replicates for each time point. Binding site determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on the gene model.

Figure 9. SWIZ translocation is dependent on bioactive GA levels and impacts GA related gene expression. (A) SWIZ translocation to the nucleus is dampened by the addition of exogenous GA4. (B) SWIZ translocation to the nucleus is enhanced by the addition of paclobutrazol (PBZ), a GA synthesis inhibitor. The average nuclear GFP signal is represented by the trend line, with error bars indicating standard deviation of the mean. $n = 85-125$ (A) and $n = 64-142$ (B) nuclei per treatment. Gene expression analysis of GA related genes. (C) Log fold change of gene expression in wildtype samples, relative to their expression at time 0, pre touch. (D) Log fold change of gene expression in SWIZ OE and swiz amiRNA samples, relative to their expression in wildtype at the same time points. Bar color indicates relevant function in GA synthesis and signalling.

Figure 10. SWIZ regulates cell wall synthesis and GA homeostasis in response to mechanical stimulus. Prior to touch, SWIZ is localized to the cytosol and partially in the nucleus. The bioactive GA pool is depleted in response to touch, triggering SWIZ to translocate to the nucleus where it positively regulates cell wall synthesis and GA activating enzymes.

REFERENCES

Altartouri B, Bidhendi AJ, Tani T, Suzuki J, Conrad C, Chebli Y, Liu N, Karunakaran C, Scarcelli G, Geitmann A (2019) Pectin chemistry and cellulose crystallinity govern

765 pavement cell morphogenesis in a multi-step mechanism. *Plant Physiol* **181**: 127–141

766 **Bidhendi AJ, Altartouri B, Gosselin FP, Geitmann A** (2019) Mechanical stress initiates and
767 sustains the morphogenesis of wavy leaf epidermal cells. *Cell Rep* **28**: 1237–1250.e6

768 **Bidhendi AJ, Geitmann A** (2018) Finite element modeling of shape changes in plant cells.
769 *Plant Physiol* **176**: 41–56

770 **Biro RL, Hunt ER, Erner Y, Jaffe MJ** (1980) Thigmomorphogenesis: changes in cell division
771 and elongation in the internodes of mechanically-perturbed or ethrel-treated bean plants.
772 *Ann Bot* **45**: 655–664

773 **Börnke F, Rocks T** (2018) Thigmomorphogenesis – Control of plant growth by mechanical
774 stimulation. *Sci Hort* **234**: 344–353

775 **Braam J** (2004) In touch: plant responses to mechanical stimuli. *New Phytol* **165**: 373–389

776 **Braam J, Davis RW** (1990) Rain-, wind-, and touch-induced expression of calmodulin and
777 calmodulin-related genes in *Arabidopsis*. *Cell* **60**: 357–364

778 **Chehab EW, Eich E, Braam J** (2009) Thigmomorphogenesis: a complex plant response to
779 mechano-stimulation. *J Exp Bot* **60**: 43–56

780 **Coomey JH, Sibout R, Hazen SP** (2020) Grass secondary cell walls, *Brachypodium distachyon*
781 as a model for discovery. *New Phytol* **227**: 1709–1724

782 **Coutand C, Martin L, Leblanc-Fournier N, Decourteix M, Julien J-L, Moulia B** (2009)
783 Strain mechanosensing quantitatively controls diameter growth and PtaZFP2 gene
784 expression in poplar. *Plant Physiol* **151**: 223–232

785 **Dai S, Petrucci S, Ordiz MI, Zhang Z, Chen S, Beachy RN** (2003) Functional analysis of
786 RF2a, a rice transcription factor. *J Biol Chem* **278**: 36396–36402

787 **Deppmann CD, Acharya A, Rishi V, Wobbes B, Smeekens S, Taparowsky EJ, Vinson C**
788 (2004) Dimerization specificity of all 67 B-ZIP motifs in *Arabidopsis thaliana*: a
789 comparison to *Homo sapiens* B-ZIP motifs. *Nucleic Acids Res* **32**: 3435–3445

790 **Ding J, Hagood JS, Ambalavanan N, Kaminski N, Bar-Joseph Z** (2018) iDREM: Interactive
791 visualization of dynamic regulatory networks. *PLoS Comput Biol* **14**: e1006019

792 **Dröge-Laser W, Snoek BL, Snel B, Weiste C** (2018) The *Arabidopsis* bZIP transcription factor
793 family-an update. *Curr Opin Plant Biol* **45**: 36–49

794 **Ehlert A, Weltmeier F, Wang X, Mayer CS, Smeekens S, Vicente-Carbajosa J, Dröge-**

795 **Laser W** (2006) Two-hybrid protein-protein interaction analysis in Arabidopsis protoplasts:
796 establishment of a heterodimerization map of group C and group S bZIP transcription
797 factors. *Plant J* **46**: 890–900

798 **Eklöf JM, Brumer H** (2010) The XTH gene family: an update on enzyme structure, function,
799 and phylogeny in xyloglucan remodeling. *Plant Physiol* **153**: 456–466

800 **Fukazawa J, Nakata M, Ito T, Matsushita A, Yamaguchi S, Takahashi Y** (2011) bZIP
801 transcription factor RSG controls the feedback regulation of NtGA20ox1 via intracellular
802 localization and epigenetic mechanism. *Plant Signal Behav* **6**: 26–28

803 **Fukazawa J, Nakata M, Ito T, Yamaguchi S, Takahashi Y** (2010) The transcription factor
804 RSG regulates negative feedback of NtGA20ox1 encoding GA 20-oxidase. *Plant J* **62**:
805 1035–1045

806 **Fukazawa J, Sakai T, Ishida S, Yamaguchi I, Kamiya Y, Takahashi Y** (2000) Repression of
807 shoot growth, a bZIP transcriptional activator, regulates cell elongation by controlling the
808 level of gibberellins. *Plant Cell* **12**: 901–915

809 **Gibalová A, Steinbachová L, Hafidh S, Bláhová V, Gadiou Z, Michailidis C, Müller K,**
810 **Pleskot R, Dupřáková N, Honys D** (2017) Characterization of pollen-expressed bZIP
811 protein interactions and the role of ATbZIP18 in the male gametophyte. *Plant Reprod* **30**:
812 1–17

813 **Gladala-Kostarz A, Doonan JH, Bosch M** (2020) Mechanical stimulation in *Brachypodium*
814 *distachyon*: implications for fitness, productivity and cell wall properties. *Plant Cell*
815 *Environ* **43**: 1314–1330

816 **Grigoryan G, Keating AE** (2006) Structure-based prediction of bZIP partnering specificity. *J*
817 *Mol Biol* **355**: 1125–1142

818 **Hamant O, Heisler MG, Jönsson H, Krupinski P, Uyttewaal M, Bokov P, Corson F, Sahlin**
819 **P, Boudaoud A, Meyerowitz EM, et al** (2008) Developmental patterning by mechanical
820 signals in Arabidopsis. *Science* **322**: 1650–1655

821 **Handakumbura P, Matos D, Osmont K, Harrington M, Heo K, Kafle K, Kim S, Baskin T,**
822 **Hazen S** (2013) Perturbation of *Brachypodium distachyon* *CELLULOSE SYNTHASE A4* or
823 *7* results in abnormal cell walls. *BMC Plant Biol* **13**: 131

824 **Handakumbura PP, Brow K, Whitney IP, Zhao K, Sanguinet KA, Lee SJ, Olins J,**
825 **Romero-Gamboa SP, Harrington MJ, Bascom CJ, et al** (2018) *SECONDARY WALL*
826 *ASSOCIATED MYB1* is a positive regulator of secondary cell wall thickening in
827 *Brachypodium distachyon* and is not found in the Brassicaceae. *Plant J* **96**: 485–699

828 **Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H,**
829 **Glass CK** (2010) Simple combinations of lineage-determining transcription factors prime
830 *cis*-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**: 576–
831 589

832 **Igarashi D, Ishida S, Fukazawa J, Takahashi Y** (2001) 14-3-3 proteins regulate intracellular
833 localization of the bZIP transcriptional activator RSG. *Plant Cell* **13**: 2483–2497

834 **Iida H** (2014) Mugifumi, a beneficial farm work of adding mechanical stress by treading to
835 wheat and barley seedlings. *Front Plant Sci* **5**: 453

836 **Ishida S, Fukazawa J, Yuasa T, Takahashi Y** (2004) Involvement of 14-3-3 signaling protein
837 binding in the functional regulation of the transcriptional activator REPRESSION OF
838 SHOOT GROWTH by gibberellins. *Plant Cell* **16**: 2641–2651

839 **Ishida S, Yuasa T, Nakata M, Takahashi Y** (2008) A tobacco calcium-dependent protein
840 kinase, CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in
841 response to gibberellins. *Plant Cell* **20**: 3273–3288

842 **Ito T, Ishida S, Oe S, Fukazawa J, Takahashi Y** (2017) Autophosphorylation affects substrate-
843 binding affinity of tobacco Ca²⁺-dependent protein kinase1. *Plant Physiol* **174**: 2457–2468

844 **Ito T, Nakata M, Fukazawa J, Ishida S, Takahashi Y** (2014) Scaffold function of Ca²⁺-
845 dependent protein kinase: tobacco Ca²⁺-DEPENDENT PROTEIN KINASE1 transfers 14-
846 3-3 to the substrate REPRESSION OF SHOOT GROWTH after phosphorylation. *Plant*
847 *Physiol* **165**: 1737–1750

848 **Jaffe MJ** (1973) Thigmomorphogenesis: The response of plant growth and development to
849 mechanical stimulation : With special reference to *Bryonia dioica*. *Planta* **114**: 143–157

850 **Jaffe MJ, Biro R, Bridle K** (1980) Thigmomorphogenesis: Calibration of the parameters of the
851 sensory function in beans. *Physiol Plant* **49**: 410–416

852 **Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T,**
853 **Parcy F, bZIP Research Group** (2002) bZIP transcription factors in Arabidopsis. *Trends*
854 *Plant Sci* **7**: 106–111

855 **Katoh K, Rozewicki J, Yamada KD** (2019) MAFFT online service: multiple sequence
856 alignment, interactive sequence choice and visualization. *Brief Bioinform* **20**: 1160–1166

857 **Kim D, Langmead B, Salzberg SL** (2015) HISAT: a fast spliced aligner with low memory
858 requirements. *Nat Methods* **12**: 357

- 859 **Lange MJ, Lange T** (2006) Gibberellin biosynthesis and the regulation of plant development.
860 *Plant Biol* **8**: 281–290
- 861 **Lange MJP, Lange T** (2015) Touch-induced changes in Arabidopsis morphology dependent on
862 gibberellin breakdown. *Nat Plants* **1**: 14025
- 863 **Leblanc-Fournier N, Martin L, Lenne C, Decourteix M** (2014) To respond or not to respond,
864 the recurring question in plant mechanosensitivity. *Front Plant Sci* **5**: 401
- 865 **Lee D, Polisensky DH, Braam J** (2005) Genome-wide identification of touch- and darkness-
866 regulated Arabidopsis genes: a focus on calmodulin-like and XTH genes. *New Phytol* **165**:
867 429–444
- 868 **Liu R, Finlayson SA** (2019) Sorghum tiller bud growth is repressed by contact with the
869 overlying leaf. *Plant Cell Environ* **42**: 2120–2132
- 870 **Liu X, Chu Z** (2015) Genome-wide evolutionary characterization and analysis of bZIP
871 transcription factors and their expression profiles in response to multiple abiotic stresses in
872 *Brachypodium distachyon*. *BMC Genomics* **16**: 227
- 873 **MacKinnon KJ-M, Cole BJ, Yu C, Coomey JH, Hartwick NT, Remigereau M-S, Duffy T,**
874 **Michael TP, Kay SA, Hazen SP** (2020) Changes in ambient temperature are the prevailing
875 cue in determining *Brachypodium distachyon* diurnal gene regulation. *New Phytol* **227**:
876 1649–1667
- 877 **Martin L, Leblanc-Fournier N, Julien J-L, Moulia B, Coutand C** (2010) Acclimation
878 kinetics of physiological and molecular responses of plants to multiple mechanical loadings.
879 *J Exp Bot* **61**: 2403–2412
- 880 **Matos DA, Whitney IP, Harrington MJ, Hazen SP** (2013) Cell walls and the developmental
881 anatomy of the *Brachypodium distachyon* stem internode. *PLoS One* **8**: e80640
- 882 **McCahill IW, Hazen SP** (2019) Regulation of cell wall thickening by a medley of mechanisms.
883 *Trends Plant Sci* **24**: 853–866
- 884 **Monshausen GB, Bibikova TN, Weisenseel MH, Gilroy S** (2009) Ca²⁺ regulates reactive
885 oxygen species production and pH during mechanosensing in Arabidopsis roots. *Plant Cell*
886 **21**: 2341–2356
- 887 **Moulia B, Coutand C, Julien J-L** (2015) Mechanosensitive control of plant growth: bearing the
888 load, sensing, transducing, and responding. *Front Plant Sci* **6**: 52
- 889 **Nagai K, Mori Y, Ishikawa S, Furuta T, Gamuyao R, Niimi Y, Hobo T, Fukuda M, Kojima**

890 **M, Takebayashi Y, et al** (2020) Antagonistic regulation of the gibberellic acid response
891 during stem growth in rice. *Nature* **584**: 109–114

892 **Niez B, Dlouha J, Moulia B, Badel E** (2019) Water-stressed or not, the mechanical acclimation
893 is a priority requirement for trees. *Trees* **33**: 279–291

894 **O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti**
895 **A, Ecker JR** (2016) Cistrome and epicistrome features shape the regulatory DNA
896 landscape. *Cell* **165**: 1280–1292

897 **Pitzschke A, Djamei A, Teige M, Hirt H** (2009) VIP1 response elements mediate mitogen-
898 activated protein kinase 3-induced stress gene expression. *Proc Natl Acad Sci U S A* **106**:
899 18414–18419

900 **Polisensky DH, Braam J** (1996) Cold-shock regulation of the Arabidopsis TCH genes and the
901 effects of modulating intracellular calcium levels. *Plant Physiol* **111**: 1271–1279

902 **Pomiès L, Decourteix M, Franchel J, Moulia B, Leblanc-Fournier N** (2017) Poplar stem
903 transcriptome is massively remodelled in response to single or repeated mechanical stimuli.
904 *BMC Genomics* **18**: 300

905 **Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J** (2019) g:Profiler: a
906 web server for functional enrichment analysis and conversions of gene lists. *Nucleic Acids*
907 *Res* **47**: W191–W198

908 **Ringli C, Keller B** (1998) Specific interaction of the tomato bZIP transcription factor VSF-1
909 with a non-palindromic DNA sequence that controls vascular gene expression. *Plant Mol*
910 *Biol* **37**: 977–988

911 **Roignant J, Badel É, Leblanc-Fournier N, Brunel-Michac N, Ruelle J, Moulia B,**
912 **Decourteix M** (2018) Feeling stretched or compressed? The multiple mechanosensitive
913 responses of wood formation to bending. *Ann Bot* **121**: 1151–1161

914 **Schütze K, Harter K, Chaban C** (2008) Post-translational regulation of plant bZIP factors.
915 *Trends Plant Sci* **13**: 247–255

916 **Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D** (2006) Highly specific gene
917 silencing by artificial microRNAs in Arabidopsis. *Plant Cell* **18**: 1121–1133

918 **Shah L, Yahya M, Shah SMA, Nadeem M, Ali A, Ali A, Wang J, Riaz MW, Rehman S, Wu**
919 **W, et al** (2019) Improving Lodging Resistance: Using Wheat and Rice as Classical
920 Examples. *Int J Mol Sci*. doi: 10.3390/ijms20174211

921 **Sibout R, Proost S, Hansen BO, Vaid N, Giorgi FM, Ho-Yue-Kuang S, Legée F, Cézart L,**
922 **Bouchabké-Coussa O, Soulhat C, et al** (2017) Expression atlas and comparative
923 coexpression network analyses reveal important genes involved in the formation of lignified
924 cell wall in *Brachypodium distachyon*. *New Phytol* **215**: 1009–1025

925 **Szczegielniak J, Borkiewicz L, Szurmak B, Lewandowska-Gnatowska E, Statkiewicz M,**
926 **Klimecka M, Cieśła J, Muszyńska G** (2012) Maize calcium-dependent protein kinase
927 (ZmCPK11): local and systemic response to wounding, regulation by touch and components
928 of jasmonate signaling. *Physiol Plant* **146**: 1–14

929 **Trabucco GM, Matos DA, Lee SJ, Saathoff A, Priest H, Mockler TC, Sarath G, Hazen SP**
930 (2013) Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid *O*-
931 methyltransferase in *Brachypodium distachyon*. *BMC Biotechnol* **13**: 61

932 **Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ** (2016) W-IQ-TREE: a fast online
933 phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* **44**: W232–W235

934 **Tsugama D, Liu S, Fujino K, Takano T** (2018) Calcium signalling regulates the functions of
935 the bZIP protein VIP1 in touch responses in *Arabidopsis thaliana*. *Ann Bot* **122**: 1219–1229

936 **Tsugama D, Liu S, Takano T** (2012) A bZIP protein, VIP1, is a regulator of osmosensory
937 signaling in *Arabidopsis*. *Plant Physiol* **159**: 144–155

938 **Tsugama D, Liu S, Takano T** (2016) The bZIP protein VIP1 Is involved in touch responses in
939 *Arabidopsis* roots. *Plant Physiol* **171**: 1355–1365

940 **Tsugama D, Liu S, Takano T** (2014) Analysis of functions of VIP1 and its close homologs in
941 osmosensory responses of *Arabidopsis thaliana*. *PLoS One* **9**: e103930

942 **Tyler L, Bragg JN, Wu J, Yang X, Tuskan GA, Vogel JP** (2010) Annotation and comparative
943 analysis of the glycoside hydrolase genes in *Brachypodium distachyon*. *BMC Genomics* **11**:
944 600

945 **Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle A,**
946 **Ludynia M, Traas J, Boudaoud A, et al** (2012) Mechanical stress acts via katanin to
947 amplify differences in growth rate between adjacent cells in *Arabidopsis*. *Cell* **149**: 439–451

948 **Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J** (2013)
949 Regulation of secondary wall synthesis and cell death by NAC transcription factors in the
950 monocot *Brachypodium distachyon*. *J Exp Bot* **64**: 1333–1343

951 **Van Leene J, Blomme J, Kulkarni SR, Cannoot B, De Winne N, Eeckhout D, Persiau G,**
952 **Van De Slijke E, Vercruysse L, Vanden Bossche R, et al** (2016) Functional

953 characterization of the Arabidopsis transcription factor bZIP29 reveals its role in leaf and
954 root development. *J Exp Bot* **67**: 5825–5840

955 **Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey**
956 **MG, Lama S, Singh KB, Ljung K, et al** (2019) A MYC2/MYC3/MYC4-dependent
957 transcription factor network regulates water spray-responsive gene expression and
958 jasmonate levels. *Proc Natl Acad Sci U S A* **116**: 23345–23356

959 **Warthmann N, Chen H, Ossowski S, Weigel D, Hervre P** (2008) Highly specific gene
960 silencing by artificial miRNAs in rice. *PLoS One* **3**: e1829

961 **Weltmeier F, Ehlert A, Mayer CS, Dietrich K, Wang X, Schütze K, Alonso R, Harter K,**
962 **Vicente-Carbajosa J, Dröge-Laser W** (2006) Combinatorial control of Arabidopsis
963 proline dehydrogenase transcription by specific heterodimerisation of bZIP transcription
964 factors. *EMBO J* **25**: 3133–3143

965 **Yin Y, Zhu Q, Dai S, Lamb C, Beachy RN** (1997) RF2a, a bZIP transcriptional activator of the
966 phloem-specific rice tungro bacilliform virus promoter, functions in vascular development.
967 *EMBO J* **16**: 5247–5259

968

Figure 1. SWIZ is a Group I bZIP highly expressed in maturing stem and root. (A) SWIZ transcript abundance in *Brachypodium distachyon* leaf, root, and stem tissue measured by microarray. Mean $\pm$ standard deviation of three biological replicates. (B) Phylogeny analysis of amino acid sequences from *B. distachyon* (green), *Oryza sativa* (blue), *Arabidopsis thaliana* (red), *Nicotiana tabacum* (orange) shows SWIZ (black dot). The phylogeny was reconstructed using maximum likelihood analysis with a bootstrap resampling value of 1000. The numbers labeled on the branches are the posterior probability to support each node. The protein sequences, full locus IDs, and plant species with genome source are provided in Supplemental File 1.

Figure 2. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to stimulus and 30 min post stimulus. Roots were observed immediately following mechanical perturbation. (B) Quantification of nuclear signal. $n=14-20$ nuclei. (C) *SWIZ* translocation occurred in the local area of the stimulus. At 30 min, stimulus was applied to an upper region of the root, while at 120 min it was applied to a lower region approximately 3 cm below. $n=109, 184$ nuclei respectively for upper and lower regions. (D) *SWIZ:GFP-OE* roots were imaged by confocal microscopy with stimulus applied in the field of view at 0, 90, and 180 min. $n=126$ nuclei. (B-D) Images were taken every 2 min. Nuclear GFP signal was quantified in selected nuclei at each time point. The average nuclear GFP signal is represented by the line with error bars indicating standard error of the mean.

Figure 3. *Brachypodium distachyon* displays classic thigmomorphogenic phenotypes. One-week-old wildtype plants were treated with touch stimulus every 90 min. (A) Left to right, plants that experienced no stress, two weeks stress, three weeks stress and were then allowed to recover, imaged one week after the end of treatment (B) height, (C) aboveground non-grain biomass weight, and (D) branch number were measured at senescence. ns: $p > 0.05$, *: $p < 0.05$, $n = 5$ plants per treatment. (E) Transverse sections of the peduncle or third internode were taken from control, 2 week stressed, and 3 week stressed plants and stained with phloroglucinol-HCl. (F) Quantification of interfascicular fiber wall thickness. Scale bar = 100 μm . $n = 3$ plants per treatment.

Figure 4. SWIZ transgenes enhance thigmomorphogenic response. Wildtype, SWIZ-OE, and swiz-amiRNA plants were placed under control conditions or received two weeks of mechanical stress every 90 min. After senescence, plants were phenotyped for (A) height, (B) branch number, and (C) aboveground biomass weight. ns: $p > 0.05$, *: $p < 0.05$. $n = 5$ plants per treatment. (D) The main stem was collected and transverse sections made of the peduncle and third elongated internode. Sections were stained with phloroglucinol-HCl. Scale bar = 100 μ m. (E) Quantification of inter fascicular fiber wall thickness. $n = 4$ to 6 plants per genotype, per treatment.

Figure 6. Gene expression analysis of cell wall related genes. Log fold-change of gene expression measured by RNA-seq in wildtype, *SWZ-OE*, and *swiz-amiRNA* relative to wildtype expression at time 0, pre-touch. Bar color indicates class of cell wall gene. Error bars indicate standard deviation of three biological replicates. Asterisks represent significantly differentially expressed genes, q-value < 0.1 for Wald test of each time and genotype compared to wild type at time 0.

Figure 7. DNA affinity purification sequencing to determine SWIZ binding sites. (A) Top two most statistically enriched sequence motifs in SWIZ binding sites. (B) Distribution of binding sites across genomic features, relative to primary transcripts of the *Brachypodium distachyon* annotation v 3.1. (C) Relative distribution of binding sites centered on the transcriptional start site (TSS, blue dashed line), transcriptional termination site (TTS, red dashed line) represents the average length of all annotated transcripts, approximately 4.5 kb away from the TSS. (D) Path determinations from iDREM time course analysis of differentially expressed genes that also have DAP-seq binding sites. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch.

Figure. 8 SWIZ binding targets differentially expressed in response to touch and SWIZ. Gene expression over time of selected genes with SWIZ binding sites. Line graphs are the average transcript abundance of three biological replicates for each time point. Binding site determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on the gene model.

Figure 10. SWIZ regulates cell wall synthesis and GA homeostasis in response to mechanical stimulus. Prior to touch, SWIZ is localized to the cytosol and partially in the nucleus. The bioactive GA pool is depleted in response to touch, triggering SWIZ to translocate to the nucleus where it positively regulates cell wall synthesis and GA activating enzymes.

Parsed Citations

- Altartouri B, Bidhendi AJ, Tani T, Suzuki J, Conrad C, Chebli Y, Liu N, Karunakaran C, Scarcelli G, Geitmann A (2019) Pectin chemistry and cellulose crystallinity govern pavement cell morphogenesis in a multi-step mechanism. *Plant Physiol* 181: 127–141**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bidhendi AJ, Altartouri B, Gosselin FP, Geitmann A (2019) Mechanical stress initiates and sustains the morphogenesis of wavy leaf epidermal cells. *Cell Rep* 28: 1237–1250.e6**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bidhendi AJ, Geitmann A (2018) Finite element modeling of shape changes in plant cells. *Plant Physiol* 176: 41–56**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Biro RL, Hunt ER, Erner Y, Jaffe MJ (1980) Thigmomorphogenesis: changes in cell division and elongation in the internodes of mechanically-perturbed or ethrel-treated bean plants. *Ann Bot* 45: 655–664**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Börnke F, Rockscht T (2018) Thigmomorphogenesis – Control of plant growth by mechanical stimulation. *Sci Hortic* 234: 344–353**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Braam J (2004) In touch: plant responses to mechanical stimuli. *New Phytol* 165: 373–389**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Braam J, Davis RW (1990) Rain-, wind-, and touch-induced expression of calmodulin and calmodulin-related genes in *Arabidopsis*. *Cell* 60: 357–364**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Chehab EW, Eich E, Braam J (2009) Thigmomorphogenesis: a complex plant response to mechano-stimulation. *J Exp Bot* 60: 43–56**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Coomes JH, Sibout R, Hazen SP (2020) Grass secondary cell walls, *Brachypodium distachyon* as a model for discovery. *New Phytol* 227: 1709–1724**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Coutand C, Martin L, Leblanc-Fournier N, Decourteix M, Julien J-L, Moulia B (2009) Strain mechanosensing quantitatively controls diameter growth and *PtaZFP2* gene expression in poplar. *Plant Physiol* 151: 223–232**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Dai S, Petrucci S, Ordiz MI, Zhang Z, Chen S, Beachy RN (2003) Functional analysis of RF2a, a rice transcription factor. *J Biol Chem* 278: 36396–36402**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Deppmann CD, Acharya A, Rishi V, Wobbes B, Smeeckens S, Taparowsky EJ, Vinson C (2004) Dimerization specificity of all 67 B-ZIP motifs in *Arabidopsis thaliana*: a comparison to *Homo sapiens* B-ZIP motifs. *Nucleic Acids Res* 32: 3435–3445**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Ding J, Hagood JS, Ambalavanan N, Kaminski N, Bar-Joseph Z (2018) iDREM: Interactive visualization of dynamic regulatory networks. *PLoS Comput Biol* 14: e1006019**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Dröge-Laser W, Snoek BL, Snel B, Weiste C (2018) The *Arabidopsis* bZIP transcription factor family—an update. *Curr Opin Plant Biol* 45: 36–49**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Ehlert A, Weltmeier F, Wang X, Mayer CS, Smeeckens S, Vicente-Carbajosa J, Dröge-Laser W (2006) Two-hybrid protein-protein interaction analysis in *Arabidopsis* protoplasts: establishment of a heterodimerization map of group C and group S bZIP transcription factors. *Plant J* 46: 890–900**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Eklöf JM, Brumer H (2010) The XTH gene family: an update on enzyme structure, function, and phylogeny in xyloglucan remodeling. *Plant Physiol* 153: 456–466**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fukazawa J, Nakata M, Ito T, Matsushita A, Yamaguchi S, Takahashi Y (2011) bZIP transcription factor RSG controls the feedback regulation of *NtGA20ox1* via intracellular localization and epigenetic mechanism. *Plant Signal Behav* 6: 26–28**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fukazawa J, Nakata M, Ito T, Yamaguchi S, Takahashi Y (2010) The transcription factor RSG regulates negative feedback of *NtGA20ox1* encoding GA20-oxidase. *Plant J* 62: 1035–1045**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fukazawa J, Sakai T, Ishida S, Yamaguchi I, Kamiya Y, Takahashi Y (2000) Repression of shoot growth, a bZIP transcriptional activator, regulates cell elongation by controlling the level of gibberellins. *Plant Cell* 12: 901–915**

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gibalová A, Steinbachová L, Hafidh S, Bláhová V, Gadiou Z, Michailidis C, Müller K, Pleskot R, Dupřáková N, Honys D (2017) Characterization of pollen-expressed bZIP protein interactions and the role of AtbZIP18 in the male gametophyte. Plant Reprod 30: 1–17

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gladala-Kostarz A, Doonan JH, Bosch M (2020) Mechanical stimulation in Brachypodium distachyon: implications for fitness, productivity and cell wall properties. Plant Cell Environ 43: 1314–1330

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grigoryan G, Keating AE (2006) Structure-based prediction of bZIP partnering specificity. J Mol Biol 355: 1125–1142

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hamant O, Heisler MG, Jönsson H, Krupinski P, Uyttewaal M, Bokov P, Corson F, Sahlin P, Boudaoud A, Meyerowitz EM, et al (2008) Developmental patterning by mechanical signals in Arabidopsis. Science 322: 1650–1655

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Handakumbura P, Matos D, Osmont K, Harrington M, Heo K, Kafle K, Kim S, Baskin T, Hazen S (2013) Perturbation of Brachypodium distachyon CELLULOSE SYNTHASE A4 or 7 results in abnormal cell walls. BMC Plant Biol 13: 131

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Handakumbura PP, Brow K, Whitney IP, Zhao K, Sanguinet KA, Lee SJ, Olins J, Romero-Gamboa SP, Harrington MJ, Bascom CJ, et al (2018) SECONDARY WALL ASSOCIATED MYB1 is a positive regulator of secondary cell wall thickening in Brachypodium distachyon and is not found in the Brassicaceae. Plant J 96: 485–699

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. Mol Cell 38: 576–589

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Igarashi D, Ishida S, Fukazawa J, Takahashi Y (2001) 14-3-3 proteins regulate intracellular localization of the bZIP transcriptional activator RSG. Plant Cell 13: 2483–2497

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Iida H (2014) Mugifumi, a beneficial farm work of adding mechanical stress by treading to wheat and barley seedlings. Front Plant Sci 5: 453

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ishida S, Fukazawa J, Yuasa T, Takahashi Y (2004) Involvement of 14-3-3 signaling protein binding in the functional regulation of the transcriptional activator REPRESSION OF SHOOT GROWTH by gibberellins. Plant Cell 16: 2641–2651

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ishida S, Yuasa T, Nakata M, Takahashi Y (2008) A tobacco calcium-dependent protein kinase, CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in response to gibberellins. Plant Cell 20: 3273–3288

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ito T, Ishida S, Oe S, Fukazawa J, Takahashi Y (2017) Autophosphorylation affects substrate-binding affinity of tobacco Ca²⁺-dependent protein kinase1. Plant Physiol 174: 2457–2468

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ito T, Nakata M, Fukazawa J, Ishida S, Takahashi Y (2014) Scaffold function of Ca²⁺-dependent protein kinase: tobacco Ca²⁺-DEPENDENT PROTEIN KINASE1 transfers 14-3-3 to the substrate REPRESSION OF SHOOT GROWTH after phosphorylation. Plant Physiol 165: 1737–1750

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jaffe MJ (1973) Thigmomorphogenesis: The response of plant growth and development to mechanical stimulation : With special reference to Bryonia dioica. Planta 114: 143–157

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jaffe MJ, Biro R, Bridle K (1980) Thigmomorphogenesis: Calibration of the parameters of the sensory function in beans. Physiol Plant 49: 410–416

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T, Parcy F, bZIP Research Group (2002) bZIP transcription factors in Arabidopsis. Trends Plant Sci 7: 106–111

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. Brief Bioinform 20: 1160–1166

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods 12: 357

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lange MJ, Lange T (2006) Gibberellin biosynthesis and the regulation of plant development. Plant Biol 8: 281–290

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lange MJ, Lange T (2015) Touch-induced changes in Arabidopsis morphology dependent on gibberellin breakdown. Nat Plants 1: 14025

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Leblanc-Fournier N, Martin L, Lenne C, Decourteix M (2014) To respond or not to respond, the recurring question in plant mechanosensitivity. Front Plant Sci 5: 401

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee D, Polisensky DH, Braam J (2005) Genome-wide identification of touch- and darkness-regulated Arabidopsis genes: a focus on calmodulin-like and XTH genes. New Phytol 165: 429–444

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu R, Finlayson SA (2019) Sorghum tiller bud growth is repressed by contact with the overlying leaf. Plant Cell Environ 42: 2120–2132

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu X, Chu Z (2015) Genome-wide evolutionary characterization and analysis of bZIP transcription factors and their expression profiles in response to multiple abiotic stresses in Brachypodium distachyon. BMC Genomics 16: 227

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

MacKinnon KJ-M, Cole BJ, Yu C, Coomey JH, Hartwick NT, Remigereau M-S, Duffy T, Michael TP, Kay SA, Hazen SP (2020) Changes in ambient temperature are the prevailing cue in determining Brachypodium distachyon diurnal gene regulation. New Phytol 227: 1649–1667

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Martin L, Leblanc-Fournier N, Julien J-L, Moulia B, Coutand C (2010) Acclimation kinetics of physiological and molecular responses of plants to multiple mechanical loadings. J Exp Bot 61: 2403–2412

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Matos DA, Whitney IP, Harrington MJ, Hazen SP (2013) Cell walls and the developmental anatomy of the Brachypodium distachyon stem internode. PLoS One 8: e80640

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McCahill IW, Hazen SP (2019) Regulation of cell wall thickening by a medley of mechanisms. Trends Plant Sci 24: 853–866

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Monshausen GB, Bibikova TN, Weisenseel MH, Gilroy S (2009) Ca²⁺ regulates reactive oxygen species production and pH during mechanosensing in Arabidopsis roots. Plant Cell 21: 2341–2356

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Moulia B, Coutand C, Julien J-L (2015) Mechanosensitive control of plant growth: bearing the load, sensing, transducing, and responding. Front Plant Sci 6: 52

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nagai K, Mori Y, Ishikawa S, Furuta T, Gamuyao R, Niimi Y, Hobo T, Fukuda M, Kojima M, Takebayashi Y, et al (2020) Antagonistic regulation of the gibberellic acid response during stem growth in rice. Nature 584: 109–114

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Niez B, Dlouha J, Moulia B, Badel E (2019) Water-stressed or not, the mechanical acclimation is a priority requirement for trees. Trees 33: 279–291

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR (2016) Cistrome and epicistrome features shape the regulatory DNA landscape. Cell 165: 1280–1292

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pitzschke A, Djamei A, Teige M, Hirt H (2009) VP1 response elements mediate mitogen-activated protein kinase 3-induced stress gene expression. Proc Natl Acad Sci U S A 106: 18414–18419

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Polisensky DH, Braam J (1996) Cold-shock regulation of the Arabidopsis TCH genes and the effects of modulating intracellular calcium levels. Plant Physiol 111: 1271–1279

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pomiès L, Decourteix M, Franchel J, Moulia B, Leblanc-Fournier N (2017) Poplar stem transcriptome is massively remodelled in response to single or repeated mechanical stimuli. BMC Genomics 18: 300

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web server for functional enrichment analysis and conversions of gene lists. Nucleic Acids Res 47: W191–W198

- Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Ringli C, Keller B (1998) Specific interaction of the tomato bZIP transcription factor VSF-1 with a non-palindromic DNA sequence that controls vascular gene expression. Plant Mol Biol 37: 977–988**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Roignant J, Badel É, Leblanc-Fournier N, Brunel-Michac N, Ruelle J, Moulia B, Decourteix M (2018) Feeling stretched or compressed? The multiple mechanosensitive responses of wood formation to bending. Ann Bot 121: 1151–1161**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Schütze K, Harter K, Chaban C (2008) Post-translational regulation of plant bZIP factors. Trends Plant Sci 13: 247–255**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D (2006) Highly specific gene silencing by artificial microRNAs in Arabidopsis. Plant Cell 18: 1121–1133**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Shah L, Yahya M, Shah SMA, Nadeem M, Ali A, Ali A, Wang J, Riaz MW, Rehman S, Wu W, et al (2019) Improving Lodging Resistance: Using Wheat and Rice as Classical Examples. Int J Mol Sci. doi: 10.3390/ijms20174211**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Sibout R, Proost S, Hansen BO, Vaid N, Giorgi FM, Ho-Yue-Kuang S, Legée F, Cézart L, Bouchabké-Coussa O, Soulhat C, et al (2017) Expression atlas and comparative coexpression network analyses reveal important genes involved in the formation of lignified cell wall in Brachypodium distachyon. New Phytol 215: 1009–1025**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Szczegieliński J, Borkiewicz L, Szurmak B, Lewandowska-Gnatowska E, Statkiewicz M, Klimecka M, Cieśła J, Muszyńska G (2012) Maize calcium-dependent protein kinase (ZmCPK11): local and systemic response to wounding, regulation by touch and components of jasmonate signaling. Physiol Plant 146: 1–14**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Trabucco GM, Matos DA, Lee SJ, Saathoff A, Priest H, Mockler TC, Sarath G, Hazen SP (2013) Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid O-methyltransferase in Brachypodium distachyon. BMC Biotechnol 13: 61**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ (2016) W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res 44: W232–W235**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Tsugama D, Liu S, Fujino K, Takano T (2018) Calcium signalling regulates the functions of the bZIP protein VIP1 in touch responses in Arabidopsis thaliana. Ann Bot 122: 1219–1229**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Tsugama D, Liu S, Takano T (2012) A bZIP protein, VIP1, is a regulator of osmosensory signaling in Arabidopsis. Plant Physiol 159: 144–155**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Tsugama D, Liu S, Takano T (2016) The bZIP protein VIP1 is involved in touch responses in Arabidopsis roots. Plant Physiol 171: 1355–1365**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Tsugama D, Liu S, Takano T (2014) Analysis of functions of VIP1 and its close homologs in osmosensory responses of Arabidopsis thaliana. PLoS One 9: e103930**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Tyler L, Bragg JN, Wu J, Yang X, Tuskan GA, Vogel JP (2010) Annotation and comparative analysis of the glycoside hydrolase genes in Brachypodium distachyon. BMC Genomics 11: 600**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle A, Ludynia M, Traas J, Boudaoud A, et al (2012) Mechanical stress acts via katanin to amplify differences in growth rate between adjacent cells in Arabidopsis. Cell 149: 439–451**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J (2013) Regulation of secondary wall synthesis and cell death by NAC transcription factors in the monocot Brachypodium distachyon. J Exp Bot 64: 1333–1343**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Van Leene J, Blomme J, Kulkarni SR, Cannoot B, De Winne N, Eeckhout D, Persiau G, Van De Slijke E, Vercruysse L, Vanden Bossche R, et al (2016) Functional characterization of the Arabidopsis transcription factor bZIP29 reveals its role in leaf and root development. J Exp Bot 67: 5825–5840**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey MG, Lama S, Singh KB, Ljung K, et al (2019) A MYC2/MYC3/MYC4-dependent transcription factor network regulates water spray-responsive gene expression and jasmonate levels.**

Proc Natl Acad Sci U S A 116: 23345–23356

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Warthmann N, Chen H, Ossowski S, Weigel D, Hervre P (2008) Highly specific gene silencing by artificial miRNAs in rice. PLoS One 3: e1829

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Weltmeier F, Ehler A, Mayer CS, Dietrich K, Wang X, Schütze K, Alonso R, Harter K, Vicente-Carbajosa J, Dröge-Laser W (2006) Combinatorial control of Arabidopsis proline dehydrogenase transcription by specific heterodimerisation of bZIP transcription factors. EMBO J 25: 3133–3143

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yin Y, Zhu Q, Dai S, Lamb C, Beachy RN (1997) RF2a, a bZIP transcriptional activator of the phloem-specific rice tungro bacilliform virus promoter, functions in vascular development. EMBO J 16: 5247–5259

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)