Title: Deficits in Behavioral and Neuronal Pattern Separation in Temporal Lobe Epilepsy

Abbreviated Title: Pattern Separation and Mnemonic Discrimination in Epilepsy

Authors and Affiliations: A.D. Madar^{1,2}, J.A. Pfammatter¹, J. Bordenave³, E.I. Plumley³, S. Ravi¹, M. Cowie¹, E.P. Wallace^{1,3}, B.P. Hermann³, R.K. Maganti³, M.V. Jones¹

¹Department of Neuroscience, University of Wisconsin-Madison, 53703

²Department of Neurobiology, Grossman Institute for Neuroscience, Quantitative Biology and Human Behavior, University of Chicago, 60637

³Department of Neurology, University of Wisconsin-Madison, School of Medicine and Public Health

Corresponding authors: madar@uchicago.edu, mathewjones@wisc.edu

Number of pages: 59

Number of figures: 9

Number of words: Abstract (158), Introduction (650), Discussion (1499)

Conflicts of interest statement: The authors declare no conflict of interest.

Acknowledgements: We thank Lin Lin who helped start behavioral tests in mice in the Jones lab and Drs. C.E.L Stark and M.A. Yassa for allowing us to use software and materials they developed. This work was supported by the US National Institutes of Health (MVJ: RO1 NS075366), the University of Wisconsin Institute for Clinical and Translational Research (MVJ.; NIH/NCATS UL1TR000427), the US Department of Defense (RKM: PR161864) and Lily's Fund for Epilepsy Research (ADM.; 2015 fellow).

Abstract:

In temporal lobe epilepsy, the ability of the dentate gyrus to limit excitatory cortical input to the hippocampus breaks down, leading to seizures. The dentate gyrus is also thought to help discriminate between similar memories by performing pattern separation, but whether epilepsy leads to a breakdown in this neural computation, and thus to mnemonic discrimination impairments, remains unknown. Here we show that temporal lobe epilepsy is characterized by behavioral deficits in mnemonic discrimination tasks, in both humans and mice (C57Bl6 males, systemic low-dose kainate model). Using a recently developed assay in brain slices of the same epileptic mice, we reveal a decreased ability of the dentate gyrus to perform certain forms of pattern separation. This is due to a subset of granule cells with abnormal bursting that can develop independently of early EEG abnormalities. Overall, our results linking physiology, computation and cognition in the same mice advance our understanding of episodic memory mechanisms and their dysfunction in epilepsy.

Keywords:

Patch-clamp, low-dose systemic kainate model, kainic acid, interictal spikes, novelty recognition, firing rate, spiking patterns, similarity metrics, input-output, entorhinal cortex, perforant path, EPSC, excitation/inhibition ratio, dentate gate, Hebb-Marr theory

Introduction:

Temporal lobe epilepsy (TLE) is characterized by recurring focal seizures originating in or near the hippocampus (Toyoda et al., 2013), microcircuit pathologies in brain regions including the hippocampus (Alexander et al., 2016) and memory-related cognitive deficits (Helmstaedter et al., 2003; Zhao et al., 2014). Although the hippocampus is a nexus for episodic memory that is critically affected during TLE, the relationship between TLE and hippocampus-dependent memory is insufficiently understood.

Episodic memory formation is thought to involve storage of neural representations in area CA3 of the hippocampus via Hebbian plasticity at recurrent excitatory synapses of coactivated cells (Rolls, 2010). In this view, a partial cue reactivates a subset of the CA3 ensemble that in turn recruits the other neurons of the original pattern resulting in recall of the original event. However, recurrent excitation is problematic because it theoretically a) predisposes the network to over-excitation that could trigger seizures (Traub and Wong, 1982; Le Duigou et al., 2014) and b) limits the number of patterns that can be stored without overlap (Rolls, 2010). Overlapping memory representations would in turn lead to interference during recall and thus cognitive confusion. To solve these problems, it was proposed that the dentate gyrus (DG) of the hippocampus acts as a) a gate and b) a pattern separator, so that similar cortical representations are transformed into sparse and dissimilar patterns before reaching CA3 (O'Reilly and McClelland, 1994; Hsu, 2007; Dengler and Coulter, 2016).

The function of DG as a gate for cortical activity to prevent seizure generation fails in TLE (Hsu, 2007; Krook-Magnuson et al., 2015; Dengler and Coulter, 2016; Lu et al., 2016). Granule cells (GCs), the output neurons of DG, lose their usual sparseness due to network reorganization (Artinian et al., 2011; Dengler and Coulter, 2016; Dengler et al., 2017) making cortical excitation

easier to propagate to CA3 (Behr et al., 1998; Patrylo et al., 1999; Ouedraogo et al., 2016). Even in healthy animals, repeated excitation of GCs induces seizures (Krook-Magnuson et al., 2015) and prolonged stimulation of DG causes TLE (Sloviter, 1983).

An alternate view of the DG function, developed in parallel to the concept of dentate gate, is that it performs pattern separation: the transformation of similar cortical patterns into dissimilar hippocampal representations. This process theoretically supports mnemonic discrimination, the ability to distinguish between similar memories. Indeed, DG lesions impair mnemonic discrimination both in rodents (Treves et al., 2008; Kesner and Rolls, 2015; Kesner et al., 2016) and humans (Yassa et al., 2011; Baker et al., 2016; Bennett and Stark, 2016; Dillon et al., 2017). Moreover, computational models (Chavlis and Poirazi, 2017) and recent experiments (Berron et al., 2016; Knierim and Neunuebel, 2016; Madar et al., 2019a, 2019b; Woods et al., 2020) suggest that the DG circuitry supports multiple forms of pattern separation (Santoro, 2013). Whether such computations underlie mnemonic discrimination remains unclear.

Unsurprisingly, TLE negatively impacts hippocampal-dependent memory in humans (Coras et al., 2014) and rodents (Grötcke et al., 2008; Chauvière et al., 2009; Müller et al., 2009; Inostroza et al., 2013; Lenck-Santini and Scott, 2015). However, the effect of TLE specifically on DG computations and DG-dependent cognition is understudied. It was only recently reported that patients with TLE are impaired on mnemonic discrimination tasks (Reyes et al., 2018; Poch et al., 2020) and that TLE causes deficits in DG-dependent object location memory in mice (Bui et al., 2018). A computational model has also suggested that hippocampal pathologies in TLE would degrade DG pattern separation (Yim et al., 2015) but the hypotheses that 1) TLE causes a breakdown in DG neural pattern separation and 2) that such a failure causes mnemonic discrimination impairments remain experimentally untested.

Here we investigated, in humans and mice, whether TLE is characterized by deficits in mnemonic discrimination and then recorded, in brain slices from the same mice, the spiking patterns of single GCs in response to parametrically varied afferent stimulation in order to gauge neuronal pattern separation.

Materials and Methods:

Human Behavior. A mnemonic similarity task (*Stark et al., 2019*), also known as a behavioral pattern separation (BPS) task (*Stark et al., 2013*), was administered to 15 patients in the University of Wisconsin-Madison Epilepsy Monitoring Unit. Under an approved Institutional Review Board protocol and after obtaining informed consent, the task was administered in conjunction with a standard neuropsychiatric evaluation and during electroencephalographic (EEG) recording, both of which are part of standard practice for diagnosing the patients' seizures. Patients were 18-65 years old, male and female. Only patients with a preliminary diagnosis of TLE were included in our analysis. As controls, 20 subjects without epilepsy, recruited to match the patients' age and sex distributions (family members of the patients when possible, or other volunteers), were also tested. Consent documents, medical records and primary data are on file in a secure location within the Dept. of Neurology. Data were deidentified prior to analysis. Subjects received no compensation for participation.

The visual, object recognition-based BPS task is described in Yassa et al. (2011) and has been further validated by demonstrating mnemonic discrimination deficits during normal aging and multiple neurologic and psychiatric disorders (*Stark et al., 2019*). It was implemented on a laptop by trained neuropsychiatric postdoctoral fellows (J.B. and E.I.P.) using software distributed freely by the Stark lab (<http://faculty.sites.uci.edu/starklab/mnemonic-similarity-task-mst/>).

Briefly, participants viewed a series of pictures of everyday objects: During Phase 1, 128 different images were presented and the subject was asked to classify each as "indoor" or "outdoor", simply to engage the subject's attention. During Phase 2, a new series of 192 images was presented: 1/3 repeated from Phase 1 (*Repeated*), 1/3 new but similar to images from Phase 1 (*Lure*), and 1/3 new and completely different from Phase 1 (*Novel*). Participants were asked to classify each image of the Phase 2 set as "Old", "Similar", or "New". BPS was evaluated with a *discrimination index* computed as the difference between $p(\text{"Similar"}|\text{Lure})$ and $p(\text{"Similar"}|\text{Novel})$, as in past research (Yassa et al., 2011; Stark et al., 2019).

Mouse Experiments. Male mice (C57BL6J, 5-6 weeks old) were received from Envigo (formerly Harlan, Madison, WI), housed in groups and allowed to acclimate to their new home environment for one or two weeks. Animals then underwent epilepsy induction with kainic acid (KA) injections (see below) (J.A.P.). A control group of mice was injected with saline and another received no injection: both were pooled together and considered as the control group because our analyses did not reveal any difference. 7-9 weeks after injection, mice started 4 weeks of behavioral testing on the BPS task. After completion of behavioral testing, animals (~18-20 weeks old) were transferred to a different building for EEG implantation. Each animal was recorded continuously for three days before being transferred back to the original building and sacrificed for slice electrophysiology (~20-24 weeks old). After each building-to-building transfer, mice were allowed a period of acclimation ranging from two days to two weeks. Behavioral testing (J.A.P, S.R. and M.C.), EEG recordings (E.P.W.) and slice electrophysiology (A.D.M) were performed by different researchers, all of whom were blind to KA or Control treatments.

Epilepsy induction. During the induction process, when not being handled, mice were individually housed in enclosed ~150 cm³ acrylic cubicles with opaque sides and clear front portals

with holes to allow air exchange, food and bedding. Animals were then randomly assigned to KA or Control treatment, ear punched for identification and weighed. Animals were induced for epileptogenesis using the repeated low-dose kainate method (*Hellier et al., 1998; Sharma et al., 2018*): a series of intraperitoneal injections based on the following schedule. Animals first received 10 mg/kg of kainic acid (Tocris Bioscience, UK) mixed in 1x PBS (8 ml per 10 mg) prepared from PBS tablets (Dot Scientific, Michigan) with deionized distilled water and filter sterilized (Millipore, Burlington, MA). Control animals were injected with an equivalent volume of saline made of 1x PBS. Animals continued to receive injections at 5mg/kg every 20 minutes until status epilepticus (SE) occurred (same number of saline injections in control mice). Some animals received alternating 5mg/kg and 2.5 mg/kg injections after the initial 10 mg/kg dose, but we found this schedule took more time than 5 mg/kg injections and our survival rate (>90% across all cohorts) was no different between the two schedules. Animals were considered to be in SE when displaying persistent behavioral seizures of level 4-5 on the Racine Scale (Racine, 1972), less than 5 minutes apart, for a minimum of 30 minutes. Animals received 4-9 injections depending on tolerance to kainic acid and injection schedule. After the injection schedule, animals were given fresh apple slices, monitored until SE ceased (assessed from normal posturing, within 1-1.5 hours), and returned to group housing. They were monitored, weighed and animals weighing less than their preinjection levels were given 0.4 ml of 1x PBS via intraperitoneal injection on a daily basis until their weight exceeded preinjection level. No animal injected with saline ever experienced status epilepticus, lost weight, or required recovery injections of 1x PBS. Mice remained in the vivarium for 7-9 weeks after injections, allowing time for the development of epilepsy in KA mice – the latent period in the KA model of TLE has previously been evaluated as 10-30 days before the first spontaneous electrographic seizure, and $\sim 11 \pm 5.4$ weeks before the first spontaneous

convulsive seizure (*Lévesque and Avoli, 2013*). During this time some KA mice became aggressive and many of the aggressor animals were removed from group housing and caged individually.

Mouse Behavior. At 14-16 weeks of age, mice started behavioral testing on a variant of a novelty recognition-based BPS task that is well established in rodents to report mnemonic discrimination abilities (*van Hagen et al., 2015*). Our protocol lasted 4 weeks, including 1 week of habituation and 3 weeks of trials. During the first week of habituation, each mouse was gently handled by the experimenter for five minutes every day. On the second week mice were split into three groups (A, B, and C). Each group went through a three-day schedule where animals were habituated on the first and second days and performed the object location task on the third day. Groups were staggered by one day each, such that group A was scheduled Monday-Wednesday, B Tuesday-Thursday and C Wednesday-Friday. Animals were housed following a 12:12 light-dark cycle and always handled or tested between 1 pm and 4pm (light period).

The arena used for the BPS task was a 63.5 x 63.5 x 15 cm open topped Plexiglas box wrapped on the outside and bottom with black felt cloth. On each side of the arena were different images serving as proximal cues for orientation. The room also had numerous distinct distal cues (e.g. the video camera placed over the center of the arena, shelving, etc.), that were kept unchanged. During behavioral testing, the room was lit with a single bank of overhead fluorescent lamps.

For the BPS task, animals underwent three phases of exploration in the arena, each lasting 3 min and separated by 1.5 min of single housing in a solitary chamber. In Phase 1 (habituation) the arena was empty, in Phase 2 (sampling) the arena contained two identical objects in the center, and in Phase 3 (testing) one object had been randomly selected and displaced by a distance of 7, 14 or 21 cm. The arena and objects were cleaned with 70% ethanol before and after each phase.

During Phases 2 and 3, the experimenter recorded the amount of time that the mouse spent interacting with each object with two silent stopwatches. The mouse was considered to be interacting with an object if it was within 1 cm of it and oriented towards, sniffing, scratching, or on top of the object. These three phases were repeated three times with a different distance each week, with the order of distances randomized. Objects used for the task were ~6 cm tall block style plastic colored figures or metal cylinders, with a footprint of ~2 cm in diameter.

All phases of the experiment were recorded with a 1080p webcam positioned above the arena. During all phases, the experimenter sat in the corner of the room out of the line of sight of the mouse. The experimenter was consistent in appearance and smell (e.g., same experimenter for a given cohort, wearing same white lab coat and gloves of the same color each day; use of scented soap, etc. was minimized).

BPS Analysis. The discrimination ratio for each trial was computed as $(T_{\text{moved}} - T_{\text{unmoved}}) / T_{\text{total}}$ where T_{moved} is the time spent exploring the moved object, T_{unmoved} is the time spent exploring the unmoved object and T_{total} is the sum of T_{moved} and T_{unmoved} . The exploration times used to calculate the reported discrimination ratio were manually collected by experimenters during the BPS task. We (M.V.J. and J.A.P.) also developed a home-written program using video recordings for motion-tracking and trajectory analysis of each mouse during behavioral testing. Discrimination ratios computed from automated tracking data were well correlated to the ratios from manually recorded times, and led to similar results ($R^2 = 0.68$, $T(117) = 16.07$, $P < 0.001$). Automated tracking data were also used to assess a) distance traveled as an estimate of motility and exploration and b) tendency to stay close to walls (i.e., thigmotaxis) as an estimate of anxiety. Thigmotaxis (*Simon et al., 1994*) was evaluated as the proportion of time spent within 6.35 mm of the wall (10% of the arena width).

Mouse electroencephalography (EEG). EEG electrode implantation was performed at ~18-20 weeks of age for all animals, following a previously established protocol (*Wallace et al., 2015*). Briefly, mice were anesthetized with isoflurane and stainless steel screw electrodes were implanted in the skull (bregma +1.5 mm and 1 mm right, bregma -3 mm and 1 mm left, and lamda -1 mm at midline). Two stainless steel braided wires were placed in the nuchal muscles for electromyography (EMG) recording. After a 72 hour recovery, we transferred mice into individual tethered EEG acquisition chambers and allowed a >12 hour acclimation period. We acquired EEG and EMG signals continuously for 3 days. Recordings were digitized with an XLTek amplifier (XLTEK, USA) sampled at 1024 Hz. Ad libitum access to food and water was ensured. Note that the resulting EEG dataset was previously reported and analyzed in Pfammatter et al. (2018).

Interictal spike (IIS) analysis. In humans and most animal models of acquired TLE, overt seizures are relatively rare (often much less than once per day) (*Lévesque et al., 2016*). In contrast, nonconvulsive and subclinical epileptiform events such as interictal spikes (IISs) can be very frequent, up to hundreds per day. Therefore, in order to assess epileptiform activity in KA animals, we used a modification of our previously published principal components (PC)-based method to quantify IISs (*Pfammatter et al., 2018*) and calculated an "Hourly IIS index" for each animal. All detected high-amplitude EEG events from 9 Ctrl and 15 KA animals were projected into the space spanned by their first three PCs. The main modification here is that, instead of using a Gaussian Mixture Model (GMM) to assign events to 'clusters', we simply gridded the PC space into 'voxels' and computed relevant quantities within each voxel exactly as we computed those same quantities within GMM 'clusters' previously: 1) The probability that events within a voxel are characteristic of an epileptogenic treatment (i.e., KA) was computed as the voxel-wise proportion of events coming from KA mice, 2) Only the voxels above chance level were considered specific to

epileptogenic treatment and probabilities were scaled accordingly, 3) The Hourly IIS Index of a given animal was defined as the average frequency of detected events weighted by the scaled probabilities of each event's voxel. This new voxel-based method has the advantages to make no assumptions about the structure of the data in PC space or about the number of clusters to fit. Instead, the main free parameter is now the voxel volume: we selected a size of 10 cubic PC units following the same optimization procedure described in Pfammatter et al. (2018) in order to avoid overfitting. Extensive exploration (not shown) revealed that the final results are similar to the GMM method and are not importantly affected by moderate changes in voxel volume.

Slice electrophysiology. Mice used for electrophysiology were p115-p182 at the time of experiment (mean $\pm$ SEM: p141 $\pm$ 4 days; no difference in age between treatments: KA n = 13, Ctrl n = 6, U-test: P = 0.9, Z = 0.2, rank sum = 246). Age was also not correlated with any summary statistics presented in this study.

Adult mice were euthanized by transcardial perfusion with oxygenated PBS under isoflurane anesthesia, before decapitation and brain extraction. 400 μ m horizontal slices of the ventral and intermediate hippocampus were prepared as detailed in Madar et al. (2019a). After slicing in a sucrose-based cutting solution (Yi et al., 2015), slices were transferred to an incubation chamber filled with 50% cutting solution and 50% artificial cerebrospinal fluid (aCSF) at 37°C for 30 minutes, then room temperature. Patch-clamp recordings were done in a chamber submerged with aCSF containing (in mM) 125 NaCl, 25 NaHCO₃, 2.5 KCl, 1.25 NaH₂PO₄, 2 CaCl₂, 1 MgCl₂, and 25 D-Glucose, flowing at 5 ml/min and saturated with a gas mixture of 95% O₂ and 5% CO₂. Stimulation was applied through a double-barreled "theta" pipette filled with aCSF. Patch pipettes were filled with an intracellular solution of the following composition (in mM): 135 K-gluconate, 5 KCl, 0.1 EGTA, 10 HEPES, 20 Na-Phosphocreatine, 2 Mg₂-ATP, 0.3 Na-GTP, 0.25

CaCl₂ adjusted to pH 7.3 with KOH and 310 mOsm with H₂O, leading to a 2-5 MΩ pipette resistance in aCSF. Whole-cell patch-clamp recordings of single DG GCs in response to electric stimulation of the outer molecular layer (perforant path) were performed as detailed in Madar et al. (2019a), and the stimulation protocols used to test neural pattern separation were the same as in Madar et al. (2019b). Briefly, input sets used for stimulation were composed of five (type 1) or ten (type 2 and 3) spiketrains (two seconds long), delivered sequentially (separated by five seconds of pause) and repeated ten or five times, respectively, in order to yield fifty output spiketrains. The stimulation pipette was placed >100μm lateral to the recorded GC to avoid direct stimulation of GC dendrites, with the baseline membrane potential held at -70 mV for current and voltage-clamp recordings.

Intrinsic electrophysiological properties of recorded GCs were the following (mean ± SEM for Ctrl / KA): resting membrane potential $V_{rest} = -78.0 \pm 1.9 / -81.4 \pm 1.2$ mV; membrane resistance $R_m = 139 \pm 17 / 178 \pm 16$ MΩ; membrane capacitance $C_m = 18 \pm 1.2 / 17 \pm 0.7$ pF. There were no significant differences between control and KA mice (U-tests: P = 0.1, 0.2, 0.7; Z = 1.6, -1.2, 0.4; rank sums = 294, 198.5, 252.5 respectively).

Neural pattern separation analysis. Similarity between spiketrains was assessed as in Madar et al. (2019a, b), using three metrics based on dividing spiketrains into time bins of a specific duration τ_w : the Pearson's correlation coefficient (R), the normalized dot product (NDP) and the scaling factor (SF). NDP and SF consider spiketrains as vectors of spike count per bin and measure the angle (NDP is the cosine) and the ratio of norms (SF) between the two vectors. R considers vectors of deviation from mean spike count, which loses the original angle between the vectors of raw spike counts and leads to considering common periods of silence as correlated, unlike NDP. Both R and NDP are sensitive to binwise synchrony, whereas SF is more sensitive to

differences in firing rate and burstiness (see Madar et al. 2019b for a detailed discussion on the different neural codes assumed by each metric). To further test the role of various spiketrain features in mediating pattern separation, we also considered neural codes purely focused on either 1) the average firing rate (FR), 2) the compactness (1 - proportion of time bins with at least one spike) and 3) the occupancy (average number of spikes in bins with at least one spike). These were computed as in Madar et al. (2019b) and the degree of pattern separation (or convergence, for negative values) was evaluated as the difference between the dispersion of the output spiketrains minus the dispersion of the input spiketrains. The dispersion was computed as the mean absolute value of pairwise differences for each spiketrain feature (FR, Compactness or Occupancy) over all spiketrains of a set, excluding self-comparisons and, in the case of the output sets, all comparisons between output spiketrains resulting from the same input spiketrain. When using a similarity metric S (R, NDP or SF), pattern separation was assessed following the same logic. First, the similarity between each pair of spiketrains of a given set was computed. S_{input} was the average for all pairs of an input set (excluding self-comparisons), and S_{output} was the average of all pairs in output sets excluding comparisons between output spiketrains coming from repetitions of the same input train. This yielded a single output similarity value for a given output set, and the degree of pattern separation was thus $S_{\text{input}} - S_{\text{output}}$. To gain a finer view, we also performed a pairwise analysis (as in Madar et al. 2019b) where each of the pairwise S_{input} of an input set were distinguished (10 pairs for input sets of type 1, 45 pairs for input sets of type 2 and 3). The pairwise S_{output} (pw S_{output}) was thus the average similarity across all pairs of output spiketrains resulting from a given pair of input spiketrains. Pattern separation was computed as $\text{pw } S_{\text{input}} - \text{pw } S_{\text{output}}$.

Experimental design and statistical analysis. Sample sizes correspond to the standards of each field (human behavior, mouse behavior, slice electrophysiology) such that large effects could

be detected. Data analysis was performed using MATLAB (Mathworks, Natick, MA, USA). The Lilliefors test was used to verify the normality of data distributions (*Lilliefors, 1967*). Parametric or nonparametric statistical tests were appropriately used to assess significance (p-value < 0.05). Throughout the Results section, KW ANOVA corresponds to the nonparametric Kruskal-Wallis analysis of variance, U-test corresponds to the Wilcoxon rank-sum test equivalent to the Mann-Whitney U-test, and KS test corresponds to the two-sample Kolmogorov-Smirnov test (sidedness is specified in the legends).

To analyze performance of mice on the BPS task we performed a two-way ANOVA with the Matlab function *anova* based on a linear mixed-effects model built using *fitlme*, with the distance of the moved object as a continuous fixed effect, animal treatment (Ctrl vs. KA) as a categorical fixed effect, and animal identity nested within object distances as a random effect to account for repeated measurements. The same procedure was used to test for differences in distances travelled and thigmotaxis.

In the pattern separation analysis, to determine whether (S_{input} , S_{output}) distributions were significantly different, we performed an analysis of the covariance (ANCOVA) using separate parabolic or linear regression models, implemented in MATLAB with a custom-written code following the method described in (Motulsky and Ransnas, 1987), as in Madar et al. (2019a, b)

For the non-linear regression between pBurst and pattern separation, we used the Matlab function *fitnlm* with a three-parameter model function $f(x) = a/(x + b) + c$.

Code accessibility. Raw data for all figures are available at:

<https://www.ebi.ac.uk/biostudies/studies/SBSST353>. Code to read and analyze the data is available at: <https://github.com/jessePfammatter/DeficitsBehavNeuronalPatternSep> and <https://github.com/antoinemadar/PatSepSpikeTrains>

Results:

Behavioral pattern separation deficits in TLE. Mnemonic discrimination is often called behavioral pattern separation (BPS) because it is hypothesized to be the behavioral outcome of neural pattern separation. It is the ability to discriminate between similar experiences (event, environment, object, etc.) that occurred at different times (*Santoro, 2013*).

We first investigated the effect of TLE on mnemonic discrimination in humans. We used the *Mnemonic Similarity Task* developed by the Stark lab, an established DG-dependent BPS task (Baker et al., 2016; Stark et al., 2019) where participants must distinguish between similar images presented at different times (**Figure 1A, Materials and Methods – Human behavior**). Patients previously diagnosed with TLE showed a severe deficit (~50%) compared to nonepileptic subjects in their ability to correctly identify objects as being similar but not identical (**Figure 1**). These results confirmed our hypothesis that TLE impairs DG-dependent mnemonic discrimination, thus warranting further study of the impact of TLE on DG computations that might explain this cognitive deficit.

To investigate the effect of TLE at the behavioral, computational and cellular levels, we turned to the common post-kainate animal model of acquired TLE (KA, **Methods – Mouse experiments**). An automated detection algorithm of interictal spikes (IISs) allowed us to quantify epileptiform activity in each animal even in the absence of seizures (**Methods – IIS analysis** and see Pfammatter et al. 2018). Although we recorded seizures in only two KA animals (Pfammatter et al. 2018), most KA mice developed IISs (**Figure 2**), demonstrating epileptogenesis. The detected IISs belonged to two distinct groups (**Figure 2D**), both having waveform features typical

of epilepsy (**Figure 2E-F**) (White et al., 2010; Chauvière et al., 2012; Huneau et al., 2013; Pfammatter et al., 2018).

Prior to EEG recordings, mice were subjected to an object-location novelty-recognition BPS task (**Methods – Mouse Behavior**) where their ability to discriminate between a moved object versus an identical unmoved object was measured for different object displacement distances (**Figure 3A**). Our results demonstrate that, on average, KA mice have lower object-location mnemonic discrimination than control mice over all object distances (**Figure 3B-C**). Control analyses revealed that KA mice have no locomotor deficits (assessed by the total distance travelled) and do not display higher anxiety levels (assessed by the time spent near walls) during the sampling and testing phases (**Figure 3D-E**). Moreover, there was no correlation between individual mnemonic discrimination ratios and total distance moved during the testing phase ($R^2 = 28\%$, $T(118) = 1.86$, $P = 0.066$) or time spent near walls ($R^2 = 2\%$, $T(118) = 0.50$, $P = 0.62$). These results suggest that differences between KA and control mice in the BPS task were not due to differences in motor ability, exploration or anxiety.

TLE is thus accompanied by mnemonic discrimination deficits in both humans and mice.

DG computational impairments in TLE. To test our hypothesis that DG pathologies in TLE lead to a breakdown of the neural pattern separation function of the DG, which in turn would translate into the cognitive impairments we observed, we measured pattern separation in hippocampal slices from the same KA and control mice used above in behavioral and EEG experiments. Although the slice preparation isolates the DG from its natural inputs, it provides the advantage that applied inputs and recorded outputs can be known with certainty, a necessary condition for quantifying circuit-level pattern separation (Santoro, 2013; Madar et al. 2019a). Briefly, the assay has three steps (**Figure 4A-B** and see Madar et al. 2019a, b): 1) ensembles of

stimulus patterns (simulating afferent input spiketrains) are generated, with known degrees of similarity to each other. These spiketrains are then fed into the DG by stimulating the lateral perforant path. 2) The response of a single GC is recorded in whole-cell current-clamp. 3) The similarity between the output spiketrains is compared to the similarity between the input spiketrains, revealing the degree of separation or convergence (**Figure 4C** and see **Material and Methods – Neural pattern separation analysis**).

We tested pattern separation levels in response to three types of input sets. Input sets of type 1 were constituted of Poisson spiketrains with a 10 Hz mean firing rate designed to have a prespecified average similarity as measured by the Pearson's correlation coefficient (R) (**Figure 4B**). For an input set with spiketrains correlated by 77% between each other (timescale: $\tau_w = 20$ ms), the output spiketrains of GCs from epileptic mice had a higher average correlation (R_{output}) than GCs from control mice (**Figure 4C**). This demonstrates a decrease in pattern separation in the DG of epileptic mice. Exploring a wider range of input correlations shows that the pattern separation function of the DG of KA mice is generally impaired compared to the normal pattern separation function (**Figure 4D**). Furthermore, a decrease in pattern separation was observed at multiple timescales (millisecond to second) and using multiple similarity metrics that assume different neural codes (**Figure 4E**, see **Material and Methods – Neural pattern separation analysis**). The impairment is most noticeable using the NDP metric, showing that DG output patterns are less orthogonalized in KA animals at all timescales, whereas differences in pattern separation via scaling (SF metric) are small but significant at large timescales.

An input set of type 2, made of Poisson spiketrains similar in terms of R but with varying firing rates (**Figure 5A1**), and an input set of type 3, with 10.5 Hz uncorrelated trains ($R = 0$) with varying burstiness (**Figure 5B1**), were designed to explore a wide range of input similarities as

measured by SF and to characterize DG computations on inputs with a variety of statistical structures (Madar et al., 2019b). As we have shown before, the normal DG exhibits low pattern separation via scaling for highly similar inputs, and significant pattern convergence for dissimilar inputs (Madar et al., 2019b). In KA mice, GCs show a decrease in both pattern convergence and separation via scaling: the DG computation function is steeper, closer to the identity line, meaning that the DG of KA mice is weaker at transforming the similarity of its inputs in general (**Figure 5A1 and B1**). When similarity is measured with R or NDP (binwise synchrony code) instead of SF (neural code focused on FR and bursting), these experiments confirmed that pattern separation is decreased in TLE (**Figure 5A2**) and that this deficit is strongest when inputs are highly similar (i.e., when pattern separation is theoretically most important) and weaker when inputs are already dissimilar (**Figure 5B2**).

Pathological spiking patterns in a subset of GCs from epileptic mice. What features of GC output spiketrains are changed in TLE to explain the difference in DG computations between KA and control mice? On visual inspection, we noticed that many KA neurons occasionally fired short bursts of action potentials after a single input pulse (**Figure 6A**), which is unusual for control GCs (Madar et al., 2019a). To determine whether there was a difference in the spiking patterns of KA and control GCs, we measured three spiketrain features: 1) the average firing rate (FR) of a GC across a full recording set (fifty spiketrains), 2) the probability of spiking after a single input pulse (SP) and 3) the probability of bursting (p(Burst), i.e., more than one spike) after a single input pulse. Our results show that on average GCs from KA mice fire more faithfully after an input (**Figure 6B**) and also tend to fire in bursts (**Figure 6C-D**), which together lead to a higher FR (**Figure 6B, E**).

Closer inspection suggests that the distributions of SP, p(Burst) and FR are different between KA and controls, with larger variance and upper tails for GCs from KA mice (**Figure 6B-E**). Although we did not formally test for discrete clusters, visual inspection of Figure 6E shows that some GCs from KA animals have higher SP, p(Burst) and FR than any control GC. Thus, it appears that, following an epileptogenic insult, only a subset of GCs displayed pathological characteristics amidst a background of seemingly normal GCs.

What is the origin of the pathological spiking patterns? We first controlled that stimulation intensities were comparable between treatment groups and were not causing the abnormal spiking patterns (**Figure 7A**). We next asked whether the abnormal firing came from changes in the synaptic drive. To answer this question, we measured the synaptic input-output relation (Ewell and Jones, 2010) by recording GCs first under current-clamp then under voltage-clamp, in response to the same pattern separation stimulus protocol (**Figure 7B1-2**). Our data show that the average excitatory drive per neuron was not different between KA and control groups, and that it does not predict burstiness (**Figure 7B3**). A finer grained analysis on the individual currents suggests that although the excitatory drive and the spiking output are correlated, high amplitude EPSCs are neither sufficient nor necessary to elicit a burst (**Figure 7B4**). Bursting does not seem to result from other changes in excitatory input-output coupling either (**Figure 7B5**), which suggest that inhibition might be involved. GCs are indeed subjected to strong tonic, feedforward and feedback inhibition that controls the sparseness of their activity (Coulter and Carlson, 2007; Ewell and Jones, 2010; Pardi et al., 2015; Lee et al., 2016) and is altered in TLE (Alexander et al., 2016; Dengler and Coulter, 2016). We did not record evoked IPSCs directly but, in normal mice, partial block of inhibition elevates GC firing rates and causes bursts (Madar et al., 2019b) similar to what we observed here in KA mice. Although indirect, our results thus converge to support the

hypothesis that pathological spiking arises from an excitation/inhibition imbalance in a subset of GCs, likely driven by a decrease in inhibition.

How do high FR and bursts relate to the DG computational impairment measured in KA mice? We first tested the hypothesis that higher GC firing rates and burstiness yield less pattern separation as measured with R (**Figure 8A**). In GCs of young mice, FR and pattern separation are related, but loosely (Madar et al., 2019a), whereas here in adult mice there is a strong linear relationship, such that GCs with pathologically high FR exhibit pathologically low pattern separation. Similarly, abnormal bursting corresponds to abnormally low levels of pattern separation. This analysis relates cell-wise spiking features (FR or pBurst averaged across all spike trains of a recorded neuron) to a form of pattern separation that is not theoretically concerned with such features (Madar et al., 2019b).

Other forms of pattern separation, more directly related to FR or bursting, can in theory also be performed: for example by increasing spiketrain-to-spiketrain *variability* in FR or burstiness of the output neuron, even if the average quantities were identical (Madar et al., 2019b). Because pathological GCs only occasionally fired bursts, we asked whether TLE could affect such forms of pattern separation. To test this, we measured the spiketrain-to-spiketrain variability in FR as well as in two complementary measures of burstiness (occupancy and compactness, see **Materials and Methods – Neural pattern separation analysis**). For input sets of type 1 (10 Hz Poisson), GC output variability was slightly lower in terms of burstiness or FR. This weak type of pattern convergence disappeared in GCs from KA mice (**Figure 8B**). For input sets of type 2 and 3, no significant difference between KA and Ctrl was detected (**Figure 8C-D**). Overall, Figure 8 suggests that TLE reduces DG neural pattern separation mostly by raising average levels of FR and burstiness, rather than through spiketrain-to-spiketrain variations of those features.

Electrographic, behavioral and computational pathologies in individual mice. We have

shown that, on average, KA mice develop EEG abnormalities (**Figure 2**), suffer mnemonic discrimination impairments (**Figure 3**) and have a bursting subpopulation of GCs with pattern separation deficits (**Figures 4-8**). Because we often performed all of the aforementioned experiments in the same mice, we next asked how those different epilepsy-related pathologies are linked at the individual level (**Figure 9**). In a simple framework where behavioral impairments are caused by computational deficits that are due to reorganization of DG network function that also leads to an increase in epileptiform EEG events, one could expect a simple relationship between all the variables we measured. Interestingly, our data suggest a more complex view. For example, animals with few interictal spikes may still harbor some pathological neurons (**Figure 9A**) and be highly impaired at the BPS task (**Figure 9B**). This shows that both DG network pathologies and mnemonic impairments can occur independent of, or before, early EEG abnormalities. Conversely, both normal and pathological GCs were recorded in mice with more advanced EEG pathology, which confirms that those two subpopulations can coexist in the same KA animal (**Figure 9A**). Mice with clearly epileptiform EEG activity can also sometimes perform normally on the BPS task (**Figure 9B**). As expected from theory we found a correlation between pattern separation in single GCs and mnemonic discrimination, but only for the behavioral test with medium interference levels (**Figure 9C**). When considering behavioral performance over all interference levels, the variability in individual behavior and in cell sampling prevents us to definitively conclude, but we note that KA animals with the largest overall cognitive impairments all had at least one pathological recorded GC (**Figure 9C**). Finally, in cases where EEG, BPS and patch-clamp data were all obtained from each animal, the combination of these measurements yielded a very obvious separation between normal and KA subjects (**Figure 9D**). Note that considering an

EEG index based on individual clusters of IISs identified in Figure 2D did not change any of the conclusions stated above. Overall, our results suggest that epilepsy-related pathologies do not all develop in concert and that computational, behavioral and electrographic measures provide complementary informative dimensions that, together, better assess the epileptic state than any single dimension or pair of dimensions.

Discussion

Cognitive and computational deficits in TLE. We provide the first experimental evidence that TLE is characterized by both mnemonic discrimination impairments and neuronal pattern separation deficits. Our data in human patients (Figure 1) is consistent with recent studies showing that TLE leads to spatial or object mnemonic discrimination impairments (Reyes et al., 2018; Poch et al., 2020). Reyes and colleagues employed a spatial memory task whereas Poch and colleagues used an object-recognition task akin to ours. However, the specific task that we used is known to be DG-dependent in humans (Baker et al., 2016; Stark et al., 2019), allowing us to conclude that impairments in TLE patients are likely due to DG alterations. Our behavioral experiment in mice (Figure 3) also complements a previous study (Bui et al., 2018) by testing short-term (minutes) rather than long-term (24h) object-location memory and by testing multiple, parameterized levels of mnemonic interference, as recommended for rigorous testing of mnemonic discrimination (Hunsaker and Kesner, 2013; Liu et al., 2016). Although not demonstrated, the DG-dependency of our task in mice is likely because: 1) variants of this task are DG-dependent (Bui et al., 2018; Pofahl et al., 2020), 2) in rodents, DG supports discrimination for short to long term memory, as long as the task is not too easy or difficult (Gilbert et al., 2001; Engin et al., 2015; Pofahl et al., 2020), 3) we tested several degrees of difficulty and found a graded response and an overall

deficit in TLE. Our results and the literature together suggest that mnemonic discrimination deficits are shared by humans and mice with TLE, occur across different memory domains and modalities, and point to DG malfunctions.

The impact of TLE-related DG pathologies on neural pattern separation had so far been investigated only in a computational model (Yim et al., 2015). This model suggested that mossy fiber sprouting and increased synaptic transmission from the perforant path to GCs can theoretically lead to a breakdown of pattern separation. We experimentally confirmed that TLE leads to a deficit in neural pattern separation (Figure 4). However, the deficit we observed is not as large, possibly because the model 1) considered synaptic transmission as deterministic rather than probabilistic and plastic, 2) assessed a different form of pattern separation (patterns were defined as population ensembles of very short spiketrains, thus focusing on a population code whereas we investigated temporal codes) and 3) considered severe degrees of epilepsy.

The role of sparsity in neuronal pattern separation and DG gating. We found that GCs from epileptic mice exhibit pathological spiking consistent with a breach of the DG gatekeeper function (Figure 6) that leads to decreased neuronal pattern separation (Figure 8A). As noted by others, the gating and pattern separation functions of DG may be related (Dengler and Coulter, 2016) because the filtering of incoming cortical activity into a sparser output can be a mechanism underlying pattern separation based on population codes (O'Reilly and McClelland, 1994; Dieni et al., 2016; Cayco-Gajic and Silver, 2019; Braganza et al., 2020). Our results suggest that maintaining a sparse output also supports pattern separation of spiketrains in the time domain, at the level of single GCs.

Rare investigations of the spiking output of single GCs in epileptic tissue have reported that single stimulation of the perforant path sometimes leads GCs to fire bursts of spikes (Lynch

et al., 2000; Kobayashi and Buckmaster, 2003). We expand on this by showing that bursts also emerge in response to complex, naturalistic stimulation patterns (**Figures 6 and 7**). It is important to note that occasional bursting is observed in normal GCs in vivo (Pernía-Andrade and Jonas, 2014), but this is very unusual in slices (Mongiat et al., 2009; Ewell and Jones, 2010; Zhang et al., 2012; Dieni et al., 2016; Madar et al., 2019a). In our experiments, bursting GCs therefore demonstrate a breach of the dentate gate. Because bursts are known to be more efficient at driving spiking in CA3 pyramidal neurons (Henze et al., 2002), a single GC firing bursts will have a higher probability of making downstream CA3 pyramidal cells fire. This could have deleterious effects on memory encoding and promote seizures, as more active CA3 neurons would 1) increase the chance of overlap between memory representations (Madar et al., 2019a, 2019b; Lee and Lee, 2020) and 2) overexcite a recurrent excitatory circuit, a mechanism of seizure generation (Traub and Wong, 1982; Santhakumar et al., 2005; de la Prida et al., 2006). Thus, our findings are consistent with the idea that pathological firing or bursting of even a subset of GCs could simultaneously contribute to deficits in mnemonic discrimination and epileptiform activity in TLE.

Our study also provides new insight on the mechanisms underlying GCs increased ability to burst by testing the influence of the excitatory drive (**Figure 7B**). The main source of excitation to GCs, the perforant path, is sometimes assumed to be augmented in TLE (Yim et al., 2015) given evidence of increased release probability at the lateral perforant path synapse (Scimemi et al., 2006). Our results do not directly contradict this, but we did not find a difference in the average synaptic current evoked in GCs by Poisson input spiketrains applied to the perforant path (**Figure 7B3**). More work is required to carefully test potential TLE-related changes in the coupling between perforant path and GCs, as well as EPSC kinetics and short-term dynamics (Madar et al., 2019a), but it is clear that the size of excitatory drive alone is not a good predictor of bursting

propensity in GCs (**Figure 7B3-4**). We also previously showed that partial blockade of inhibition can produce GC bursting and pattern separation deficits (Madar et al., 2019b). Thus, in accordance with past research (Kobayashi and Buckmaster, 2003), we suggest that the strong decrease in synaptic inhibition characteristic of TLE (Dengler and Coulter, 2016; Dengler et al., 2017) is a likely driver of pathological bursting.

Perhaps the most intriguing insight we bring on the failure of the DG as a gate and pattern separator in TLE is that this failure occurs only in a subset of GCs, upon a background of apparently normal GCs in the same epileptic network. This finding invites many more questions: In the epileptic DG, is there a spectrum of spiking patterns among GCs going from normal to pathological, or are there two discrete types? What is the ratio of normal to pathological GCs and does it evolve during epileptogenesis? Do GCs become pathological due to changes in intrinsic properties or networking? Our results suggest deficits in inhibition, but additional mechanisms could be in play (Kelly and Beck, 2017). Anatomical or biophysical differences between GCs could be related to different birthdates of adult-born GCs relative to an epileptogenic brain insult, resulting in only a subset of GCs integrating abnormally into the DG network (Kron et al., 2010). Alternatively, pathological GCs could develop from a more susceptible and active subclass of mature GCs (Erwin et al., 2020). In any case, the discovery that the GC population in the epileptic DG is not homogeneous in terms of spiking patterns is a crucial step towards developing new treatments that would specifically target pathological neurons to avoid deleterious side-effects.

The link between mnemonic discrimination, pattern separation and TLE. For thirty years the DG has been hypothesized to support mnemonic discrimination by performing neuronal pattern separation (Treves et al., 2008), but despite important efforts (McHugh et al., 2007; Marrone et al., 2011; Ramirez et al., 2013; Yokoyama and Matsuo, 2016; Woods et al.,

2020), no experiment has yet directly connected DG input-output computations with their role in memory. Here, we confirmed our previous work showing that the DG itself performs temporal pattern separation (Madar et al., 2019a, b) and we showed that a deficit in pattern separation is associated with a deficit in mnemonic discrimination, on average (Figure 3, 4) and possibly at the individual level (Figure 9C). This is correlative evidence, linking non-simultaneous experiments, we thus cannot exclude that other DG or non-DG processes (Bernier et al., 2017; Guise and Shapiro, 2017; Hainmueller and Bartos, 2020) are impaired in KA mice and partially lead to the observed cognitive deficits. A full demonstration of the causal link between DG pattern separation and mnemonic discrimination will require measuring and manipulating DG computations in vivo during episodic memory tasks, but our study is a preliminary step toward understanding how DG circuit-level computations relate to a high-level cognitive function and how these processes fail in TLE. A mechanistic understanding of this relation may lead to new early diagnosis tools (e.g. cognitive tests like in Figure 1) as well as therapies alleviating memory disorders in TLE. For example, our finding that a subset of GCs with pattern separation deficits can develop without EEG hallmarks of epilepsy (Figure 9A) suggests a mechanism for memory impairments that often occur early during epileptogenesis (Jones et al., 2007; Chauvière et al., 2009; Witt and Helmstaedter, 2015). Similarly, the coexistence of pathological and normal neurons could explain memory impairments in patients without hippocampal sclerosis (Rayner et al., 2019).

601 (C) The same data from B, presented as cumulative frequency distributions. A non-parametric
602 one-way KW ANOVA on the indices grouped by treatment confirms a highly significant deficit
603 in visual pattern separation memory for human subjects with TLE ($P = 0.0343$, $\chi^2(34) = 4.4809$).
604 (D) TLE and Ctrl groups were properly age matched (unpaired two-sided T-test: $P = 0.9411$, $T(31)$
605 $= -0.0745$).
606 (E) There were slightly more women than men in the control group, but the gender difference was
607 not significant between the two treatments (one-way KW ANOVA on gender values, 1 or 2,
608 grouped by treatment: $P = 0.0663$, $\chi^2(32) = 3.3724$).

Figure 2. Kainate-injected mice have epileptiform electrographic events with a range of event frequency

(A) The Hourly Interictal Spikes (IIS) index, a proxy for epilepsy severity based on EEG recordings (Pfammatter et al., 2018), was calculated for 9 Ctrl (black) and 15 KA (red) animals. KA animals have a significantly higher Hourly IIS index than Ctrl animals (U-test: $P = 0.009$, $Z = 2.207$, rank sum = 225).

(B) To compute the index, high-amplitude events (200 ms around a spike) were identified from left frontal EEG and projected on the first three principal components (PC) of the ensemble of events from all mice. Each dot here is an EEG event, red when from a KA animal and black when from a Ctrl. Notice the 'fingers' of red KA dots extending from the central cluster of mixed Ctrl and KA data points.

(C) Using a voxel gridding of the PC space (voxel size: 10 cubic units), we calculated the probability $P(KA)$ for each event to be specific to KA treatment. The hourly IIS index of a mouse is the average number of detected events/h, weighted by their probability of being specific to the KA treatment.

(D) Events with $P(KA) > 0.80$. Two visually identified groups (red ellipses) were separated along the PC1 axis (group a < -10 and group b > -10).

(E) Each row shows example events from groups identified in D and from the remaining events in C. EEG amplitude was normalized and reported in terms of standard deviation (SD).

(F) Average waveforms from group a (top), b (middle), and events with $P(KA) < 0.80$. Despite morphological differences, group a and b both show a wave following the detected spike, which is a known marker of epilepsy in TLE (Chauvière et al., 2012; Huneau et al., 2013).

Figure 3. Kainate-injected mice have deficits in mnemonic discrimination

(A) Object location BPS task arena setup. The experiment was run in three phases of three minutes. In Phase 1, a mouse was allowed to explore the empty square arena. In Phase 2, the same mouse was allowed to explore the same arena but with two identical objects (position indicated by grey circles). In Phase 3, one randomly chosen object was moved either 7, 14 or 21 cm from its original position and the mouse was allowed to explore again. Between phases, the mouse was placed in a solitary chamber for 1.5 minutes. Each mouse performed the task once at each distance with different objects, in randomized order, with a week in between trials.

(B) Performance at the BPS task is evaluated with a discrimination ratio computed for each animal at each distance. Healthy mice generally prefer to explore objects with novel characteristics (such as change in relative position) and are thus expected to spend more time with the moved object if they notice it has changed location. Discrimination is thus quantified as the difference between the time exploring the moved object and the time exploring the unmoved object. A two-way ANOVA with repeated measures shows that mice discriminate better when objects are moved farther apart, and that Ctrl mice ($n = 29$) discriminate better than KA mice ($n = 27$) (Distance Object Moved: $P = 0.002$, $F(1,118) = 9.84$; Treatment: $P = 0.027$, $F(1,118) = 5.02$).

(C) Subset of the data in B plotted in the space of discrimination at each distance, allowing a clearer picture of individual performance. Each animal is represented by a single dot in this 3D space (42 animals with records at the three distances: 24 KA and 18 Ctrl). For each data point, the Mahalanobis distance (De Maesschalck et al., 2000) from the Ctrl centroid is represented by the dot size, and is also displayed in the inset (mean $\pm$ SEM). Mahalanobis distances are larger for KA than Ctrl animals, confirming that the KA population displays mnemonic discrimination deficits (two-sided T-test: $P = 0.003$, $T(40) = -3.15$).

(D,E) Control analysis to determine whether there were motor impairments or anxiety/curiosity differences between KA and Ctrl mice, across phase 1 (habituation), 2 and 3 of the BPS task. In each panel, individual mice are represented by 1-3 data points, as in B, depending on how many conditions (Distance Object Moved) they were tested on. Mean +/- SEM. **(D)** Distance traveled was used as a proxy for motor ability. During Phase 1 and Phase 2, control and KA mice moved similar distances ($P = 0.507$, $F(1, 120) = 0.443$ and $P = 0.070$, $F(1, 122) = 3.35$). During Phase 3 KA mice moved farther than Ctrl mice ($P = 0.023$, $F(1, 124) = 5.34$). **(E)** Percent of time near wall was used as a proxy for anxiety. During Phase 1, KA animals spent more time near the maze walls compared to Ctrl ($P = 0.03$, $F(1, 120) = 4.82$). During Phases 2 and 3, Ctrl and KA animals spent a similar amount of time near walls ($P = 0.68$, $F(1, 122) = 0.173$ and $P = 0.35$, $F(1, 124) = 0.881$). F-tests correspond to one-way ANOVAs with repeated measures, with Treatment as a categorical fixed effect.

Figure 4. Epileptic mice have neural pattern separation deficits

(A) Histology of the DG in a horizontal slice (Cresyl violet/Nissl staining; scale bar: 250 μ m), overlaid with a schematic of the experimental setup: a theta pipette in the ML is used to focally stimulate the outer molecular layer (input) while a responding GC is recorded via whole-cell patch-clamp (output). GCL: granule cell layer, H: hilus, ML: molecular layer. Solid lines represent dendrites and dashed lines axons.

(B) Current-clamp recordings of the membrane potential of two different GCs (Ctrl and KA) in response to the same set of input trains. An input set is constituted of five different trains of electrical pulses following a Poisson distribution with an average rate of 10 Hz. The Pearson's correlation coefficient (R) between two input trains is computed with a binning window (τ_w) of 20 ms (*Left*). R_{input} is the average of the ten pairwise coefficients, diagonal excluded. After converting the GC recordings to vectors of binned spike counts, the pairwise coefficients and their average (R_{output}) can be computed the same way (Madar et al, 2019b). Note that the input set was repeated 10 times to yield output sets of fifty sweeps (only one repeat is shown).

(C) The R_{output} of GCs from kainate-injected animals is higher, on average, than in GCs from controls. Data points correspond to single output sets (KA: 27 recordings from 24 GCs; Ctrl: 12 recordings from 11 GCs), all being responses to the same input set as in B. Black crosses with error bars are means $\pm$ SEM. The asterisk signals significance: U-test comparing the medians: $P = 0.011$, $Z = -2.541$, rank sum = 156; two-sided two-sample KS test comparing the distribution shapes: $P = 0.0037$, $D = 0.583$. The dashed line corresponds to R_{input} (0.77): any value below the line implies effective pattern separation. Thus, GCs from kainate animals exhibit less pattern separation than controls.

(D) Pattern separation was investigated over a wide range of input correlations by using four different input sets of five 10 Hz Poisson trains (at $\tau_w = 20$ ms: $R_{\text{input}} = 0.24, 0.45, 0.77, 0.95$). Crosses and error bars correspond to mean $\pm$ SEM (pw R_{input} , pw R_{output}) across multiple recordings (KA: 6-24 GCs per input set; Ctrl: 4-11 GCs). Data points below the identity line (dashed) correspond to pattern separation. The distributions between KA and Ctrl are significantly different (ANCOVA with separate parabolic models fitting the data points and not the means: $P < 0.0001$, $F(3,714) = 15.485$. Solid curves are the parabolic models used for the ANCOVA.

699 (E) Levels of pattern separation measured using different similarity metrics (S: R, NDP and SF,
700 see Methods) and timescales. Cumulative frequency distributions of the distance of ($p_w S_{input}$, p_w
701 S_{output}) data points to the identity line in pattern separation graphs like in E. Positive values of the
702 x-axis correspond to pattern separation, and negative values to pattern convergence. Insets show
703 medians. For R and NDP, distributions are significantly shifted to the left, showing that GCs from
704 KA exhibit less decorrelation and less orthogonalization (ANCOVA as in D, but using separate
705 linear models: $P < 0.01$ for $\tau_w = 5$ to 1000 ms. For R/NDP and $\tau_w = 10, 20, 100, 250$ ms, $P =$
706 $<0.0001/<0.0001$, $<0.0001/<0.0001$, $0.001/<0.0001$, $0.011/0.0001$, $F(2, 716) = 15.2/18.2,$
707 $20.3/23.4, 6.87/15.97, 4.5/9.4$). For SF, scaling levels are weakly but significantly shifted to less
708 separation at large τ_w (SF: $P > 0.2$ except at 250 ms and 500 ms, with $P = 0.026, 0.046$, $F(2, 716)$
709 $= 3.65, 3.09$ respectively).

Figure 5. Multiple DG computations are affected by epilepsy

Our standard input sets (type 1, Figure 4) consisted of 10 Hz Poisson trains. Two other input sets (type 2 and 3) were designed to explore single GCs responses to inputs with diverse structures and statistics, both with a wide range of pairwise similarity as measured by SF (in contrast to input sets of type 1).

(A1) Left: Input set type 2 was constituted of spiketrains following a Poisson distribution, each with a different firing rate (FR), but with an average R_{input} constrained around 0.75 ($\tau_w = 10$ ms). *Middle and Right:* Pattern separation graphs showing the pairwise output spiketrain similarity as a function of the pairwise input similarity, as measured by SF with two different timescales, averaged across multiple GCs (KA: 16, Ctrl: 5). Distributions are significantly different, suggesting in particular that epilepsy causes a decrease in pattern convergence (via scaling) for low input similarities (ANCOVA with separate linear models, for $\tau_w = 20$ and 250 ms respectively : $P = 0.0016$ and < 0.0001 , $F(2, 941) = 6.5$ and 10.4). **(A2)** Levels of pattern separation for type 2 inputs, measured using different similarity metrics and timescales as in Figure 4. It confirms that epilepsy decreases the separation of similar Poisson input spiketrains as measured by R and NDP, consistent with Figure 4, and shows that differences in terms of SF, although small, are significant (ANCOVA as in A1: R: $P < 0.0001$ for $\tau_w = 20$ ms up to 1000 ms, $P = 0.02$ and 0.18 for 5 and 10 ms; NDP and SF: $P < 0.025$ for τ_w up to 1000 ms. Detailed statistics for $\tau_w = 5, 10, 20, 50, 100, 250, 500$ and 1000 ms respectively: R, $P = 0.0243, 0.1820, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, F(2, 941) = 3.7, 1.7, 10.2, 15.1, 24.1, 36.0, 50.2, 27.0$; NDP, $P = 0.0022, 0.0240, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, F(2, 941) = 6.2, 3.7, 13.8, 17.4, 21.4, 26.0, 21.0, 16.8$; SF, $P = 0.0002, 0.0012, 0.0016, 0.0240, 0.0073, <0.0001, <0.0001, 0.0002, F(2, 941) = 8.7, 6.7, 6.5, 3.7, 4.9, 10.9, 10.4, 8.7$).

734 **(B1)** *Left*: Input set type 3 was constituted of spiketrains with 21 spikes (FR = 10.5Hz) that were
735 distributed among bins to produce trains with varying burstiness (Madar et al., 2019b). R is close
736 to 0 for all pairs. *Middle and Right*: same analysis as in A1 (KA: 8 GCs, Ctrl: 3 GCs). Distributions
737 are significantly different, suggesting again that epilepsy causes a decrease in pattern convergence
738 via scaling, for low input similarities (ANCOVA for $\tau_w = 20$ and 100 ms: $P < 0.0001$ and $F(2, 491)$
739 = 10.3 and 13.8 respectively). **(B2)** Same analysis as in A2. The directions of impairments are the
740 same as in A2, showing that epilepsy decreases pattern separation in terms of R and NDP but
741 slightly improves it in terms of SF. The shift via scaling is weak but significant at all timescales,
742 and larger at the longest τ_w . ANCOVA: R, $P < 0.05$ for $\tau_w = 5, 10$ and 500 ms; NDP and SF, $P <$
743 0.007 for τ_w up to 500 ms. Detailed statistics for $\tau_w = 5, 10, 20, 50, 100, 250, 500$ and 1000 ms
744 respectively: R, $P = 0.0451, 0.0004, 0.3727, 0.7357, 0.9340, 0.4271, 0.0012, 0.2889, F(2, 491) =$
745 3.1, 8.1, 1.0, 0.3, 0.1, 0.8, 6.8, 1.2; NDP, $P = <0.0001, <0.0001, 0.0003, 0.0012, <0.0001, 0.0069,$
746 $<0.0001, 0.4154, F(2, 491) = 11.1, 24.3, 8.4, 6.8, 11.2, 5.0, 14.6, 0.9, SF, P = <0.0001, <0.0001,$
747 $<0.0001, <0.0001, <0.0001, 0.0065, 0.0002, 0.0003, F(2, 941) = 42.1, 20.9, 10.3, 13.6, 13.8, 5.1,$
748 8.5, 8.1.

Figure 6. In epileptic mice, a subpopulation of GCs shows pathological spiking patterns

(A) Example of current-clamp recordings in GCs from a kainate-injected (KA) vs a control animal in response to the same Poisson input train, illustrating that some GCs from KA spiked more, and had a tendency to fire short bursts (2-4 spikes riding a single EPSP, see inset).

(B-E) Some GCs from KA exhibited larger firing rates due to a higher probability of spiking at least once after an input spike (spiking probability), and sometimes a higher probability of spiking more than once between two input spikes ($p(\text{Burst})$) than GCs from controls. The firing rate, spiking probability and $p(\text{Burst})$ of a neuron were all computed as the average over the fifty sweeps of an output set from a pattern separation experiment (input set type 1, see Figure 4B).

(B) Left: data points correspond to the same recordings as in Figure 4C (KA: 27; Ctrl: 12). Black dash and error bars are mean $\pm$ SEM. A U-test was used to compare the medians, showing that on average FR and SP are higher in KA (FR: $P = 0.0415$, $Z = -2.04$, rank sum = 172.5; SP: $P =$

0.0613, $Z = -1.87$, rank sum = 168.5). *Right*: Frequency distributions of the same data. A one-sided two-sample KS test shows that the KA distribution has a larger tail in both cases (FR and SP: $P = 0.0465$, $D = 0.4074$), indicating that a subset of KA GCs are pathological.

(C) Same as in B for $p(\text{Burst})$. Note that the left graph has a $\log_{10}$ scale showing all data points, whereas the middle graph has a linear scale zoomed in (i.e. not showing the 5 largest KA values). The graph on the right has the same scale as the middle graph. Together, they suggest that in KA, there might be a healthy population of GCs coexisting with a different population of pathologically bursty GCs (U-test: $P = 0.0153$, $Z = -2.16$, rank sum = 168.5; KS test: $P = 0.0259$, $D = 0.4444$).

(D) Some bursts were detected by our algorithm in GCs from Ctrl mice, but the vast majority of those were due to the temporal summation of two EPSPs resulting from input spikes occurring close in time. In contrast, a large number of bursts in GCs from KA mice come after an isolated input spike. *Top*: Time interval between the two input spikes (inter-spike-interval, ISI) preceding a given burst. Data points correspond to all detected bursts from the recordings in C. U-test: $P = 0.0003$, $Z = -3.58$, rank sum = 5906.2. *Bottom*: Cumulative frequency distributions of the same data. A one-sided two-sample KS test demonstrates that the distributions are different ($P < 0.0001$, $D = 0.2699$).

(E) Same data as in B and C: the elbow in the distribution of SP, $p(\text{Burst})$ and FR values visually defines two clusters of neurons. One cluster contains GCs from both KA and Ctrl mice (with GCs from both groups spanning a large range of SP values but low $p(\text{Burst})$) and can be considered "normal". In contrast, the other cluster, with highest SP combined with high $p(\text{Burst})$, contains only GCs from KA animals and can thus be considered "pathological".

Figure 7. Pathological bursting is not explained by aberrant excitation

(A) Bursting and high firing rates were not an artefact of stimulation differences: The current intensity of the electric stimulation did not significantly differ between KA and Ctrl (same recordings as in Figure 4C and 6B-E, U-test: $P = 0.2991$, $Z = 1.04$, rank sum = 274.5) and stimulus intensity was a poor predictor of pathological firing patterns (as indicated by the low R^2 values of linear regressions. For FR/SP/pBurst: $R^2 = 12.75\%/11.69\%/4.35\%$, $F(2,37) = 5.4/4.9/1.7$, $P = 0.0257/0.0332/0.2027$).

(B) For a subset of neurons in A (KA: 12; Ctrl: 6), we followed the current-clamp recording with a voltage-clamp recording ($V_{\text{hold}} = -70\text{mV}$) in response to the same pattern separation protocol in

793 order to assess the excitatory synaptic drive and its relationship with spiking behavior. The
794 stimulus electrode location and current intensity was unchanged. **(B1)** Example of current-clamp
795 (top) and voltage-clamp (bottom) recording in the same GC from a Ctrl animal. Only the first 500
796 ms of the ten responses to repetitions of the first input train shown in Figure 4B are displayed. *Top*:
797 action potentials are truncated at 0 mV. *Bottom*: stimulation artefacts and occasional unclamped
798 spikes were blanked. **(B2)** The ten EPSCs (grey) associated to first pulse of input trains in B1 and
799 the mean EPSC (black). We assessed the excitatory drive as the maximum inward current in the
800 interval between each input pulse minus the current baseline (i.e., mode of the current over the full
801 sweep). P(Spike) was defined as the probability of spiking during this interval across the ten
802 sweeps of the current-clamp recording. **(B3)** *Top*: The mean EPSC density for each neuron was
803 not significantly different between KA and Ctrl (U-test: $P = 0.5532$, rank sum = 50; two-sided
804 two-sample KS test: $P = 0.1935$, $D = 0.4167$), suggesting that pathological spiking cannot be
805 explained by larger EPSCs in KA mice. *Bottom*: Indeed, no clear relationship exists between
806 p(Burst) and the mean current density, as some GCs with a great synaptic drive do not exhibit
807 much bursting, and vice versa. **(B4)** Consistently, the peak current density associated to each inter-
808 input-interval in a recording is not well correlated to the corresponding number of spikes in the
809 current-clamp trace, for both Ctrl (top) and KA (bottom). For example, excitatory currents of the
810 same magnitude can be associated to 0, 1, 2, 3 or 4 spikes. EPSC amplitude is thus not sufficient
811 to explain the spiking output, which suggests that other factors, like reduced inhibition, are likely
812 implicated in causing pathological bursts. **(B5)** *Top*: The average EPSC density between two input
813 spikes is plotted against the corresponding probability of spiking p(Spike), as in Ewell and Jones
814 (2010). The resulting input-output (IO) distribution is fitted with a sigmoid (two GCs
815 representative of KA and Ctrl are shown). *Bottom*: In a subset of GCs, mostly from KA, The IO

816 distribution appears disorganized and sometimes difficult to fit. A proxy of this disorganization is
 817 the standard deviation (SD) of the current density, averaged across each level of $p(\text{Spike})$
 818 excluding $p(\text{Spike}) = 0$ and 1. The average SD for a given GC was plotted against the propensity
 819 of that neuron to fire bursts. There is no clear relationship between these two quantities, suggesting
 820 that: 1) decoupling between the excitatory drive and spiking probability is not directly related to
 821 bursting and 2) a subset of GCs from KA with apparently healthy spiking patterns could already
 822 exhibit a more subtle but pathological IO disorganization.

Figure 8. Pathological spiking patterns explain pattern separation differences in epilepsy

(A) Neurons with high average firing rates and burstiness exhibit lower pattern separation (computed as $R_{\text{input}} - R_{\text{output}}$). Same data as in Figure 4C and 6B-C. There is a strong linear relationship between firing rates and decorrelation (Grey line, $R^2 = 63.68\%$, $F(2,37) = 64.9$, $P < 0.0001$). The relationship between $p(\text{Burst})$ and decorrelation is better described by an inverse function with a horizontal asymptote (see Methods – Software and statistics): $R^2 = 53.1\%$, R^2 adjusted for 3 parameters = 50.5% , $F(3,36) = 20.4$, $P < 0.0001$.

(B-D) Mean $\pm$ SEM pattern separation levels across recordings when considering burstiness codes (Compactness or Occupancy) or a rate code (spiking frequency across a 2s sweep) instead of the binwise synchrony code assumed by R. Compactness is the proportion of time bins with at least one spike, whereas occupancy is the average number of spikes in a bin. Pattern separation corresponds to more dispersion in compactness, occupancy or firing rate in the output spiketrains than the input spiketrains. Negative values mean there is pattern convergence. U-tests comparing medians of GCs from KA and Ctrl groups were performed (p-values in panel).

(B) In response to input sets of type 1 (10 Hz Poisson trains, same recordings as in A and Figure 4), GCs exhibit low levels of convergence, if any, in terms of compactness, occupancy and rate codes, and these levels are slightly but significantly shifted to less convergence in GCs from KA for $\tau_w = 10$ ms. This means that the relationships between firing rate or burstiness and pattern decorrelation observed in A is not due to variations of the firing rate or burstiness across sweeps. Detailed statistics at 10, 20, 50, 100, 250 ms, Compactness: $P = 0.0482, .1892, .1628, .2925, .7584$, $Z = -1.97, -1.97, -1.31, -1.39, 0.31$, rank sum = 745, 745, 801, 794, 939 ; Occupancy: $P = 0.0177, 0.2586, 0.3624, 0.1425, 0.0211$, $Z = -2.37, -1.13, -0.91, -1.47, -2.31$, rank sum = 712.5, 816.5, 835, 788, 717; FR: $P = 0.0553$, $Z = -1.92$, rank sum = 750.

847 (C) In response to an input set of type 2 (Poisson trains with different firing rates, same recordings
848 as in Figure 5A), GCs from KA and Ctrl do not show significant differences ($P > 0.2$ for all
849 timescales and codes). Detailed statistics at 10, 20, 50, 100, 250 ms, Compactness: $P = 0.7726$,
850 $0.7726, 0.8365, 0.9014, 0.2312$, $Z = -0.29, -0.29, -0.21, 0.12, 1.20$, rank sum = 51, 51, 52, 57, 70;
851 Occupancy: $P = 1, 1, 0.9671, 0.5915, 0.2006$, $Z = 0, 0, 0.04, -0.54, -1.28$, rank sum = 55, 55, 56,
852 48, 39; FR: $P = 0.9014$, $Z = -0.12$, rank sum = 53.

853 (D) In response to an input set of type 3 (10 Hz trains with varying compactness and occupancy,
854 same recordings as in Figure 5B), GCs exhibit convergence in terms of compactness and
855 occupancy and separation in terms of rate codes. Only computations in terms of compactness and
856 firing rate are mildly shifted in GCs from KA for $\tau_w = 10$ ms. Detailed statistics at 10, 20, 50, 100,
857 250 ms, Compactness: $P = 0.0848$, rank sum = 9; Occupancy: $P = 0.8, 0.7758, 0.9212, 0.3758, 1$,
858 rank sum = 19.5, 20, 17, 23, 18; FR: $P = 0.0848$, rank sum = 9.

Figure 9. EEG, behavioral and computational measures provide complementary insights about epileptic pathology in individual animals.

(A-D) Scatter plots relating the electrographic measure of IISs developed in Figure 2 (black axis labels), measures from patch-clamp recordings of individual GCs from the same animals (purple axis labels) and behavioral discrimination measures described in Figure 3 (green axis labels). Not all three types of records are available for every animal, which leads to differences in the samples shown in each panel.

(A) Data points correspond to individual GCs from animals in which both EEG and patch-clamp data were recorded. Each animal has a unique average hourly IIS index value. Only a subset of GCs are outside of the Ctrl range (i.e. pathological). For example, in the mouse with the highest IIS index (~ 12), only one of the three recorded GCs exhibited a pathological FR (~ 8 Hz, left panel)

and all had a similar R_{output} slightly above the Ctrl range (~ 0.5 , right panel). In contrast, some GCs with abnormally high FR ($\sim 6-12$ Hz) and R_{output} ($\sim 0.5-0.6$) came from animals with low IIS index ($\sim 0-2$).

(B) Data points correspond to individual animals in which both EEG and behavior were measured. Discrimination Impairment is the Mahalanobis distance computed in Figure 3C. In KA animals, low IIS index can correspond to either normal or abnormal mnemonic discrimination, and despite having a high IIS index, some animals have normal discrimination.

(C) Data points correspond to individual GCs from animals in which both behavior and patch-clamp data were recorded. *Left*: Significant linear correlation (grey line) between behavioral discrimination after a 14 cm object displacement (Figure 3B) and pattern separation in DG ($R^2 = 0.0355$, $F(1,21) = 11.02$, $P = 0.0034$). No obvious relationship for the other conditions (7 cm and 21 cm). *Right*: Performance on all displacement distances are taken in account. The relationship is unclear, but all KA mice with large impairments had at least one pathological GC.

(D) Data points correspond to individual GCs from animals in which EEG, behavior and patch-clamp data were all collected. Stems show which GCs were recorded from each animal. The lower right corner would correspond to the highest pathology in electrographic, behavioral and computational categories. Despite any heterogeneity or apparent ambiguity when only one or two of these measures are considered (above), when all three measures are available for each animal, the KA group is very clearly separable from the Ctrl group.

References

- Alexander A, Maroso M, Soltesz I (2016) Organization and control of epileptic circuits in temporal lobe epilepsy. *Prog Brain Res* 226:127–154.
- Artinian J, Peret A, Marti G, Epsztein J, Crépel V (2011) Synaptic kainate receptors in interplay with INaP shift the sparse firing of dentate granule cells to a sustained rhythmic mode in temporal lobe epilepsy. *J Neurosci Off J Soc Neurosci* 31:10811–10818.
- Baker S, Vieweg P, Gao F, Gilboa A, Wolbers T, Black SE, Rosenbaum RS (2016) The Human Dentate Gyrus Plays a Necessary Role in Discriminating New Memories. *Curr Biol CB* 26:2629–2634.
- Behr J, Lyson KJ, Mody I (1998) Enhanced propagation of epileptiform activity through the kindled dentate gyrus. *J Neurophysiol* 79:1726–1732.
- Bennett IJ, Stark CEL (2016) Mnemonic discrimination relates to perforant path integrity: An ultra-high resolution diffusion tensor imaging study. *Neurobiol Learn Mem* 129:107–112.
- Bernier BE, Lacagnina AF, Ayoub A, Shue F, Zemelman BV, Krasne FB, Drew MR (2017) Dentate Gyrus Contributes to Retrieval as well as Encoding: Evidence from Context Fear Conditioning, Recall, and Extinction. *J Neurosci Off J Soc Neurosci* 37:6359–6371.
- Berron D, Schütze H, Maass A, Cardenas-Blanco A, Kuijf HJ, Kumaran D, Düzel E (2016) Strong Evidence for Pattern Separation in Human Dentate Gyrus. *J Neurosci Off J Soc Neurosci* 36:7569–7579.
- Bielefeld P, van Vliet EA, Gorter JA, Lucassen PJ, Fitzsimons CP (2014) Different subsets of newborn granule cells: a possible role in epileptogenesis? *Eur J Neurosci* 39:1–11.
- Braganza O, Mueller-Komorowska D, Kelly T, Beck H (2020) Quantitative properties of a feedback circuit predict frequency-dependent pattern separation. *eLife* 9.
- Bui AD, Nguyen TM, Limouse C, Kim HK, Szabo GG, Felong S, Maroso M, Soltesz I (2018) Dentate gyrus mossy cells control spontaneous convulsive seizures and spatial memory. *Science* 359:787–790.

915 Cayco-Gajic NA, Silver RA (2019) Re-evaluating Circuit Mechanisms Underlying Pattern Separation.
916 Neuron 101:584–602.

917 Chauvière L, Doublet T, Ghestem A, Siyoucef SS, Wendling F, Huys R, Jirsa V, Bartolomei F, Bernard C
918 (2012) Changes in interictal spike features precede the onset of temporal lobe epilepsy. Ann Neurol
919 71:805–814.

920 Chauvière L, Raftafi N, Thinus-Blanc C, Bartolomei F, Esclapez M, Bernard C (2009) Early deficits in spatial
921 memory and theta rhythm in experimental temporal lobe epilepsy. J Neurosci Off J Soc Neurosci
922 29:5402–5410.

923 Chavlis S, Poirazi P (2017) Pattern separation in the hippocampus through the eyes of computational
924 modeling. Synap N Y N 71.

925 Coras R, Pauli E, Li J, Schwarz M, Rössler K, Buchfelder M, Hamer H, Stefan H, Blumcke I (2014)
926 Differential influence of hippocampal subfields to memory formation: insights from patients with
927 temporal lobe epilepsy. Brain J Neurol 137:1945–1957.

928 Coulter DA, Carlson GC (2007) Functional regulation of the dentate gyrus by GABA-mediated inhibition.
929 Prog Brain Res 163:235–243.

930 de la Prida LM, Huberfeld G, Cohen I, Miles R (2006) Threshold behavior in the initiation of hippocampal
931 population bursts. Neuron 49:131–142.

932 De Maesschalck R, Jouan-Rimbaud D, Massart DL (2000) The Mahalanobis distance. Chemom Intell Lab
933 Syst 50:1–18.

934 Dengler CG, Coulter DA (2016) Normal and epilepsy-associated pathologic function of the dentate gyrus.
935 Prog Brain Res 226:155–178.

936 Dengler CG, Yue C, Takano H, Coulter DA (2017) Massively augmented hippocampal dentate granule cell
937 activation accompanies epilepsy development. Sci Rep 7:42090.

938 Dieni CV, Panichi R, Aimone JB, Kuo CT, Wadiche JI, Overstreet-Wadiche L (2016) Low excitatory
939 innervation balances high intrinsic excitability of immature dentate neurons. *Nat Commun* 7:11313.

940 Dillon SE, Tsivos D, Knight M, McCann B, Pennington C, Shiel AI, Conway ME, Newson MA, Kauppinen
941 RA, Coulthard EJ (2017) The impact of ageing reveals distinct roles for human dentate gyrus and CA3
942 in pattern separation and object recognition memory. *Sci Rep* 7:14069.

943 Engin E, Zarnowska ED, Benke D, Tsvetkov E, Sigal M, Keist R, Bolshakov VY, Pearce RA, Rudolph U (2015)
944 Tonic Inhibitory Control of Dentate Gyrus Granule Cells by $\alpha 5$ -Containing GABAA Receptors Reduces
945 Memory Interference. *J Neurosci Off J Soc Neurosci* 35:13698–13712.

946 Erwin SR, Sun W, Copeland M, Lindo S, Spruston N, Cembrowski MS (2020) A Sparse, Spatially Biased
947 Subtype of Mature Granule Cell Dominates Recruitment in Hippocampal-Associated Behaviors. *Cell*
948 *Rep* 31:107551.

949 Ewell LA, Jones MV (2010) Frequency-tuned distribution of inhibition in the dentate gyrus. *J Neurosci Off*
950 *J Soc Neurosci* 30:12597–12607.

951 Gilbert PE, Kesner RP, Lee I (2001) Dissociating hippocampal subregions: double dissociation between
952 dentate gyrus and CA1. *Hippocampus* 11:626–636.

953 Gröticke I, Hoffmann K, Löscher W (2008) Behavioral alterations in a mouse model of temporal lobe
954 epilepsy induced by intrahippocampal injection of kainate. *Exp Neurol* 213:71–83.

955 Guise KG, Shapiro ML (2017) Medial Prefrontal Cortex Reduces Memory Interference by Modifying
956 Hippocampal Encoding. *Neuron* 94:183-192.e8.

957 Hainmueller T, Bartos M (2020) Dentate gyrus circuits for encoding, retrieval and discrimination of
958 episodic memories. *Nat Rev Neurosci* 21:153–168.

959 Hellier JL, Patrylo PR, Buckmaster PS, Dudek FE (1998) Recurrent spontaneous motor seizures after
960 repeated low-dose systemic treatment with kainate: assessment of a rat model of temporal lobe
961 epilepsy. *Epilepsy Res* 31:73–84.

962 Helmstaedter C, Kurthen M, Lux S, Reuber M, Elger CE (2003) Chronic epilepsy and cognition: a
963 longitudinal study in temporal lobe epilepsy. *Ann Neurol* 54:425–432.

964 Henze DA, Wittner L, Buzsáki G (2002) Single granule cells reliably discharge targets in the hippocampal
965 CA3 network in vivo. *Nat Neurosci* 5:790–795.

966 Hsu D (2007) The dentate gyrus as a filter or gate: a look back and a look ahead. *Prog Brain Res* 163:601–
967 613.

968 Huneau C, Benquet P, Dieuset G, Biraben A, Martin B, Wendling F (2013) Shape features of epileptic
969 spikes are a marker of epileptogenesis in mice. *Epilepsia* 54:2219–2227.

970 Hunsaker MR, Kesner RP (2013) The operation of pattern separation and pattern completion processes
971 associated with different attributes or domains of memory. *Neurosci Biobehav Rev* 37:36–58.

972 Inostroza M, Brotons-Mas JR, Laurent F, Cid E, de la Prida LM (2013) Specific impairment of “what-
973 where-when” episodic-like memory in experimental models of temporal lobe epilepsy. *J Neurosci*
974 Off J Soc Neurosci 33:17749–17762.

975 Jones JE, Watson R, Sheth R, Caplan R, Koehn M, Seidenberg M, Hermann B (2007) Psychiatric
976 comorbidity in children with new onset epilepsy. *Dev Med Child Neurol* 49:493–497.

977 Kelly T, Beck H (2017) Functional properties of granule cells with hilar basal dendrites in the epileptic
978 dentate gyrus. *Epilepsia* 58:160–171.

979 Kesner RP, Kirk RA, Yu Z, Polansky C, Musso ND (2016) Dentate gyrus supports slope recognition
980 memory, shades of grey-context pattern separation and recognition memory, and CA3 supports
981 pattern completion for object memory. *Neurobiol Learn Mem* 129:29–37.

982 Kesner RP, Rolls ET (2015) A computational theory of hippocampal function, and tests of the theory:
983 new developments. *Neurosci Biobehav Rev* 48:92–147.

984 Knierim JJ, Neunuebel JP (2016) Tracking the flow of hippocampal computation: Pattern separation,
985 pattern completion, and attractor dynamics. *Neurobiol Learn Mem* 129:38–49.

986 Kobayashi M, Buckmaster PS (2003) Reduced inhibition of dentate granule cells in a model of temporal
987 lobe epilepsy. *J Neurosci Off J Soc Neurosci* 23:2440–2452.

988 Kron MM, Zhang H, Parent JM (2010) The developmental stage of dentate granule cells dictates their
989 contribution to seizure-induced plasticity. *J Neurosci Off J Soc Neurosci* 30:2051–2059.

990 Krook-Magnuson E, Armstrong C, Bui A, Lew S, Oijala M, Soltesz I (2015) In vivo evaluation of the
991 dentate gate theory in epilepsy. *J Physiol* 593:2379–2388.

992 Le Duigou C, Simonnet J, Teleńczuk MT, Fricker D, Miles R (2014) Recurrent synapses and circuits in the
993 CA3 region of the hippocampus: an associative network. *Front Cell Neurosci* 7:262.

994 Lee C-H, Lee I (2020) Impairment of Pattern Separation of Ambiguous Scenes by Single Units in the CA3
995 in the Absence of the Dentate Gyrus. *J Neurosci Off J Soc Neurosci* 40:3576–3590.

996 Lee C-T, Kao M-H, Hou W-H, Wei Y-T, Chen C-L, Lien C-C (2016) Causal Evidence for the Role of Specific
997 GABAergic Interneuron Types in Entorhinal Recruitment of Dentate Granule Cells. *Sci Rep* 6:36885.

998 Lenck-Santini P-P, Scott RC (2015) Mechanisms Responsible for Cognitive Impairment in Epilepsy. *Cold*
999 *Spring Harb Perspect Med* 5.

1000 Lévesque M, Avoli M (2013) The kainic acid model of temporal lobe epilepsy. *Neurosci Biobehav Rev*
1001 37:2887–2899.

1002 Lévesque M, Avoli M, Bernard C (2016) Animal models of temporal lobe epilepsy following systemic
1003 chemoconvulsant administration. *J Neurosci Methods* 260:45–52.

1004 Lilliefors HW (1967) On the Kolmogorov-Smirnov Test for Normality with Mean and Variance Unknown. *J*
1005 *Am Stat Assoc* 62:399–402.

1006 Liu KY, Gould RL, Coulson MC, Ward EV, Howard RJ (2016) Tests of pattern separation and pattern
1007 completion in humans-A systematic review. *Hippocampus* 26:705–717.

1008 Lu Y, Zhong C, Wang L, Wei P, He W, Huang K, Zhang Y, Zhan Y, Feng G, Wang L (2016) Optogenetic
1009 dissection of ictal propagation in the hippocampal-entorhinal cortex structures. Nat Commun
1010 7:10962.

1011 Lynch M, Sayin U, Golarai G, Sutula T (2000) NMDA receptor-dependent plasticity of granule cell spiking
1012 in the dentate gyrus of normal and epileptic rats. J Neurophysiol 84:2868–2879.

1013 Madar AD, Ewell LA, Jones MV (2019a) Pattern separation of spiketrains in hippocampal neurons. Sci
1014 Rep 9:5282.

1015 Madar AD, Ewell LA, Jones MV (2019b) Temporal pattern separation in hippocampal neurons through
1016 multiplexed neural codes. PLoS Comput Biol 15:e1006932.

1017 Marrone DF, Adams AA, Satvat E (2011) Increased pattern separation in the aged fascia dentata.
1018 Neurobiol Aging 32:2317.e23-32.

1019 McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK, Lowell BB, Fanselow MS, Wilson
1020 MA, Tonegawa S (2007) Dentate gyrus NMDA receptors mediate rapid pattern separation in the
1021 hippocampal network. Science 317:94–99.

1022 Mongiat LA, Espósito MS, Lombardi G, Schinder AF (2009) Reliable activation of immature neurons in the
1023 adult hippocampus. PloS One 4:e5320.

1024 Motulsky HJ, Ransnas LA (1987) Fitting curves to data using nonlinear regression: a practical and
1025 nonmathematical review. FASEB J Off Publ Fed Am Soc Exp Biol 1:365–374.

1026 Müller CJ, Grötcke I, Bankstahl M, Löscher W (2009) Behavioral and cognitive alterations, spontaneous
1027 seizures, and neuropathology developing after a pilocarpine-induced status epilepticus in C57BL/6
1028 mice. Exp Neurol 219:284–297.

1029 O'Reilly RC, McClelland JL (1994) Hippocampal conjunctive encoding, storage, and recall: avoiding a
1030 trade-off. Hippocampus 4:661–682.

1031 Ouedraogo DW, Lenck-Santini P-P, Marti G, Robbe D, Crépel V, Epsztein J (2016) Abnormal UP/DOWN
1032 Membrane Potential Dynamics Coupled with the Neocortical Slow Oscillation in Dentate Granule
1033 Cells during the Latent Phase of Temporal Lobe Epilepsy. *eNeuro* 3.

1034 Pardi MB, Ogando MB, Schinder AF, Marin-Burgin A (2015) Differential inhibition onto developing and
1035 mature granule cells generates high-frequency filters with variable gain. *eLife* 4:e08764.

1036 Patrylo PR, Schweitzer JS, Dudek FE (1999) Abnormal responses to perforant path stimulation in the
1037 dentate gyrus of slices from rats with kainate-induced epilepsy and mossy fiber reorganization.
1038 *Epilepsy Res* 36:31–42.

1039 Pernía-Andrade AJ, Jonas P (2014) Theta-gamma-modulated synaptic currents in hippocampal granule
1040 cells in vivo define a mechanism for network oscillations. *Neuron* 81:140–152.

1041 Pfammatter JA, Bergstrom RA, Wallace EP, Maganti RK, Jones MV (2018) A predictive epilepsy index
1042 based on probabilistic classification of interictal spike waveforms. *PLOS ONE* 13:e0207158.

1043 Poch C, Toledano R, García-Morales I, Prieto A, García-Barragán N, Aledo-Serrano Á, Gil-Nagel A, Campo
1044 P (2020) Mnemonic discrimination in patients with unilateral mesial temporal lobe epilepsy relates
1045 to similarity and number of events stored in memory. *Neurobiol Learn Mem* 169:107177.

1046 Pofahl M, Nikbakht N, Haubrich AN, Nguyen T, Masala N, Braganza O, Macke JH, Ewell LA, Golcuk K, Beck
1047 H (2020) Dentate gyrus population activity during immobility drives formation of precise memories.
1048 *bioRxiv*:2020.03.05.978320.

1049 Ramirez S, Liu X, Lin P-A, Suh J, Pignatelli M, Redondo RL, Ryan TJ, Tonegawa S (2013) Creating a false
1050 memory in the hippocampus. *Science* 341:387–391.

1051 Rayner G, Tailby C, Jackson G, Wilson S (2019) Looking beyond lesions for causes of neuropsychological
1052 impairment in epilepsy. *Neurology* 92:e680–e689.

1053 Reyes A, Holden HM, Chang Y-HA, Uttarwar VS, Sheppard DP, DeFord NE, DeJesus SY, Kansal L, Gilbert
1054 PE, McDonald CR (2018) Impaired spatial pattern separation performance in temporal lobe epilepsy

1055 is associated with visuospatial memory deficits and hippocampal volume loss. *Neuropsychologia*
1056 111:209–215.

1057 Rolls ET (2010) A computational theory of episodic memory formation in the hippocampus. *Behav Brain*
1058 *Res* 215:180–196.

1059 Santhakumar V, Aradi I, Soltesz I (2005) Role of mossy fiber sprouting and mossy cell loss in
1060 hyperexcitability: a network model of the dentate gyrus incorporating cell types and axonal
1061 topography. *J Neurophysiol* 93:437–453.

1062 Santoro A (2013) Reassessing pattern separation in the dentate gyrus. *Front Behav Neurosci* 7:96.

1063 Scimemi A, Schorge S, Kullmann DM, Walker MC (2006) Epileptogenesis is associated with enhanced
1064 glutamatergic transmission in the perforant path. *J Neurophysiol* 95:1213–1220.

1065 Sharma S, Puttachary S, Thippeswamy A, Kanthasamy AG, Thippeswamy T (2018) Status Epilepticus:
1066 Behavioral and Electroencephalography Seizure Correlates in Kainate Experimental Models. *Front*
1067 *Neurol* 9:7.

1068 Simon P, Dupuis R, Costentin J (1994) Thigmotaxis as an index of anxiety in mice. Influence of
1069 dopaminergic transmissions. *Behav Brain Res* 61:59–64.

1070 Sloviter RS (1983) “Epileptic” brain damage in rats induced by sustained electrical stimulation of the
1071 perforant path. I. Acute electrophysiological and light microscopic studies. *Brain Res Bull* 10:675–
1072 697.

1073 Stark SM, Kirwan CB, Stark CEL (2019) Mnemonic Similarity Task: A Tool for Assessing Hippocampal
1074 Integrity. *Trends Cogn Sci* 23:938–951.

1075 Stark SM, Yassa MA, Lacy JW, Stark CEL (2013) A task to assess behavioral pattern separation (BPS) in
1076 humans: Data from healthy aging and mild cognitive impairment. *Neuropsychologia* 51:2442–2449.

1077 Toyoda I, Bower MR, Leyva F, Buckmaster PS (2013) Early activation of ventral hippocampus and
1078 subiculum during spontaneous seizures in a rat model of temporal lobe epilepsy. *J Neurosci Off J Soc*
1079 *Neurosci* 33:11100–11115.

1080 Traub RD, Wong RK (1982) Cellular mechanism of neuronal synchronization in epilepsy. *Science*
1081 216:745–747.

1082 Treves A, Tashiro A, Witter MP, Moser EI (2008) What is the mammalian dentate gyrus good for?
1083 *Neuroscience* 154:1155–1172.

1084 van Hagen BTJ, van Goethem NP, Lagatta DC, Prickaerts J (2015) The object pattern separation (OPS)
1085 task: a behavioral paradigm derived from the object recognition task. *Behav Brain Res* 285:44–52.

1086 Wallace E, Kim DY, Kim K-M, Chen S, Blair Braden B, Williams J, Jasso K, Garcia A, Rho JM, Bimonte-
1087 Nelson H, Maganti R (2015) Differential effects of duration of sleep fragmentation on spatial
1088 learning and synaptic plasticity in pubertal mice. *Brain Res* 1615:116–128.

1089 White A, Williams PA, Hellier JL, Clark S, Dudek FE, Staley KJ (2010) EEG spike activity precedes epilepsy
1090 after kainate-induced status epilepticus. *Epilepsia* 51:371–383.

1091 Witt J-A, Helmstaedter C (2015) Cognition in the early stages of adult epilepsy. *Seizure* 26:65–68.

1092 Woods NI, Stefanini F, Apodaca-Montano DL, Tan IMC, Biane JS, Kheirbek MA (2020) The Dentate Gyrus
1093 Classifies Cortical Representations of Learned Stimuli. *Neuron*.

1094 Yassa MA, Mattfeld AT, Stark SM, Stark CEL (2011) Age-related memory deficits linked to circuit-specific
1095 disruptions in the hippocampus. *Proc Natl Acad Sci U S A* 108:8873–8878.

1096 Yi F, Catudio-Garrett E, Gábel R, Wilhelm M, Erdelyi F, Szabo G, Deisseroth K, Lawrence J (2015)
1097 Hippocampal “cholinergic interneurons” visualized with the choline acetyltransferase promoter:
1098 anatomical distribution, intrinsic membrane properties, neurochemical characteristics, and capacity
1099 for cholinergic modulation. *Front Synaptic Neurosci* 7:4.

1100 Yim MY, Hanuschkin A, Wolfart J (2015) Intrinsic rescaling of granule cells restores pattern separation
1101 ability of a dentate gyrus network model during epileptic hyperexcitability. *Hippocampus* 25:297–
1102 308.

1103 Yokoyama M, Matsuo N (2016) Loss of Ensemble Segregation in Dentate Gyrus, but not in
1104 Somatosensory Cortex, during Contextual Fear Memory Generalization. *Front Behav Neurosci*
1105 10:218.

1106 Zhang W, Huguenard JR, Buckmaster PS (2012) Increased excitatory synaptic input to granule cells from
1107 hilar and CA3 regions in a rat model of temporal lobe epilepsy. *J Neurosci Off J Soc Neurosci*
1108 32:1183–1196.

1109 Zhao F, Kang H, You L, Rastogi P, Venkatesh D, Chandra M (2014) Neuropsychological deficits in
1110 temporal lobe epilepsy: A comprehensive review. *Ann Indian Acad Neurol* 17:374–382.

1111