Male genital lobe morphology affects the chance to copulate in *Drosophila pachea*

running title: *D. pachea* mate competition analysis

Bénédicte M. Lefèvre¹, Diane Catté¹, Virginie Courtier-Orgogozo¹, Michael Lang¹

¹Team “Evolution and Genetics”, Institut Jacques Monod, CNRS, UMR7592, Université de Paris,
15 rue Hélène Brion, 75013 Paris

*corresponding author: michael.lang@ijm.fr

running title: *D. pachea* mate competition analysis

8 Abstract

9 **Introduction:** Male genitalia are thought to ensure transfer of sperm through direct physical contact
 10 with female during copulation. However, little attention has been given to their pre-copulatory role
 11 with respect to sexual selection and sexual conflict. Males of the fruitfly *Drosophila pachea* have a
 12 pair of asymmetric external genital lobes, which are primary sexual structures and stabilize the
 13 copulatory complex of female and male genitalia. We wondered if genital lobes in *D. pachea* may
 14 have a role before or at the onset of copulation, before genitalia contacts are made.

15 **Results:** We tested this hypothesis with a *D. pachea* stock where males have variable lobe lengths.
 16 In 92 mate competition trials with a single female and two males, females preferentially engaged
 17 into a first copulation with males that had a longer left lobe and that displayed increased courtship
 18 vigor. In 53 additional trials with both males having partially amputated left lobes of different
 19 lengths, we observed a weaker and non-significant effect of left lobe length on copulation success.
 20 Courtship durations significantly increased with female age and when two males courted the female
 21 simultaneously, compared to trials with only one courting male. In addition, lobe length did not
 22 affect sperm transfer once copulation was established.

23 **Conclusion:** Left lobe length affects the chance of a male to engage into copulation. The
 24 morphology of this primary sexual trait may affect reproductive success by mediating courtship
 25 signals or by facilitating the establishment of genital contacts at the onset of copulation.

26 keywords: *Drosophila pachea*, primary sexual trait, mating behaviour, mate-competition
 27 experiments, genitalia

28 **Introduction**

29 Males and females exhibit different reproduction strategies due to higher energy costs of
30 larger female gametes compared to smaller and more abundant male gametes [1]. This implies
31 male-male intrasexual mate-competition for siring the limited female gametes. In turn, females may
32 optimize reproduction by choosing males that confer survival and fecundity benefits to her and to
33 the offspring. This dynamics has been formalized into genetic models [2, 3] involving female
34 preferences for particular male characters and male mate competition, and can contribute to the
35 rapid evolution of female preferences and male sexual attributes. There is also evidence for male
36 mate-choice and female intra-sexual selection [4–6], showing that traditional sex-roles can change
37 between species and in individuals under diverse conditions. In general, sexual selection implies a
38 sexual conflict whenever male and female reproductive fitness strategies differ [7].

39 Across animals with internal fertilization, genitalia are usually the most rapidly evolving
40 organs [8]. Several hypotheses propose that the evolution of genitalia is based on sexual selection
41 and a consequence of sexual conflict between males and females at different levels of reproduction,
42 including competition of sperm from different males inside the female (sperm competition) [9],
43 female controlled storage and usage of sperm to fertilize eggs (cryptic female choice) [8, 10, 11], or
44 sexually antagonistic co-evolution of aggressive and defensive morphologies and/or behaviors
45 between male and female to control fertilization decisions [12, 13]. Male courtship is well known as
46 a behaviour to attract females before copulation begins, but has also been reported to occur during
47 or even after copulation [14, 15]. It is thought to be a widespread and key aspect of copulation in
48 the cryptic female choice scenario, where males stimulate the female to utilize his own sperm [8,
49 10]. A variety of insect species were reported where males had evolved elaborated male genital
50 structures that are apparently used for courtship throughout copulation to stimulate the female
51 through tapping or other physical stimuli [14, 15].

Traditionally, sexual traits are categorized as primary or secondary. Genital structures are considered “primary” when they are directly used for the transfer of gametes during copulation or when they contribute to the complexing of female and male copulatory organs [8, 16]. Other traits that differ between sexes and that are linked to reproduction are considered “secondary” sexual traits. For primary sexual traits, sexual selection is traditionally thought to act during or after copulation [8, 15], while secondary sexual traits can be involved in pre-copulatory mate-choice via visual, auditory or chemical long range signalling [17–25].

Primary genitalia could also be possibly involved in pre-copulatory mate choice. Supposing that a particular male genital trait enhances or favours female fecundity, it might become preferred by the female during pre-copulatory courtship or at the moment of genitalia coupling [26–28]. This preference could rely on direct benefits to reduce energy investment or predation risk caused by unsuccessful copulation attempts, or on indirect male genetic quality to be inherited into the offspring generation. Several examples of primary genitalia used in pre-copulatory courtship signalling are known in vertebrates. For example, male and female genital display, swelling, or coloration changes are associated with sexual activity [29–31] and penile display was also observed in lizards [32]. Human male penis size was found to influence male attractiveness to women [33]. In diverse mammalian species, females have also evolved external and visible structures that resemble male genitals, a phenomenon called andromimicry [31, 34, 35]. These structures are likely used in visual signaling, so it is possible that the male analogous parts also do. Furthermore, females of some live-bearing fish species were reported to prefer to mate with males with large gonopodia [36, 37]. These cases involve visual stimuli and large genital organs. Only a few studies examined the roles of primary genitalia in invertebrates during courtship or at the onset of copulation before phallus intromission is achieved [26–28, 38, 39]. Longer male external genitalia of the water strider *Aquarius remigis* were associated with higher mating success [28]. In addition, male genital spine morphology of *Drosophila bipectinata* and *Drosophila ananassae* were reported to evolve in

response to direct competition among males for securing mates in a so called ‘scramble competition’ scenario [26, 27]. In general, the role of primary sexual traits in establishing copulation or their role during courtship needs further investigation.

The fruitfly species *Drosophila pachea* is a promising model to study the evolution of primary sexual traits, especially with respect to the evolution of left-right asymmetry. Male *D. pachea* have an asymmetric aedeagus (intromission organ) and a pair of asymmetric external lobes with the left lobe being approximately 1.5 times longer than the right lobe [40–42] (Figure 1). The genital lobes have likely evolved during the past 3-6 Ma and are not found in closely related species [43]. *D. pachea* couples also mate in a right-sided copulation position where the male rests on top of the female abdomen with the antero-posterior midline shifted about 6-8 degrees to the right side of the female midline [41, 42, 44]. Male and female genitalia form an asymmetric complex during copulation and the asymmetric lobes stabilize this genital complex [44]. Furthermore, *D. pachea* is among the *Drosophila* species that produce the longest (giant) sperm [45, 46], and their ejaculates contain in average just about 40 sperm cells [47]. Thus, a particular right-sided mating position could potentially be associated with optimal transfer of giant sperm during copulation [42].

The aim of this study was to test if the asymmetric genital lobes of *D. pachea* would have an effect on pre-copulatory mate-choice, in addition to their role in stabilizing the complex of male and female genitalia during copulation. Previously, we found that males originating from one of our *D. pachea* laboratory stocks possess short and rather symmetric lobes [41] (Figure 1), while others have the typical size asymmetry. The variability in left genital lobe length that we observed within this fly stock enabled us to test if lobe length might affect pre-copulatory courtship or mate competition. We selected *D. pachea* males with short left lobes and produced a stock with an increased variance of left lobe length. Next, we tested sibling males of this “selection stock” in mate-competition assays for their success to engage first into copulation with a single female. We also tested whether copulation success in our assay would be affected by male courtship vigour.

Then, we surgically shortened the length of the left lobe in males that had developed long left lobes, to further test whether left lobe length affects copulation success. Finally, we assessed whether lobe length affects ejaculate allocation into female storage organs.

Results

Genital lobe lengths differ between *D. patchea* stocks

In our laboratory stock 15090-1698.01 (Additional datafile 1: Table S1), *D. patchea* males display a characteristic left-right size asymmetry of genital lobes with the left lobe being consistently larger than the right lobe (Figure 1a,c, Figure 2a, Additional datafile 1: Table S2, Additional datafile 2). Similarly, in stock 15090-1698.02, most males reveal larger left lobes, but a few individuals are observed with particularly small lobes that are rather symmetric in length (Figure 1b-d, Figure 2b). We selected for males with short lobe lengths and crossed them with sibling females for 50 generations (Methods). The resulting stock was named “selection stock”. We observed an increased variance of left lobe length compared to the source stock (Levene’s test: selection stock / 15090-1698.02: $DF = 1/147$, $F = 11.506$, $P = 0.0008913$) (Figure 2c, Additional datafile 2). In contrast, the right lobe length differed only marginally among the two stocks (Additional datafile 1: Table S2) and the variances of right lobe lengths were not significantly different among them (Levene’s test, $DF = 1/147$, $F = 0.5164$, $P = 0.4735$).

Male genital left lobe length, male courtship and failed copulation attempts have an effect on copulation success

We performed a mate competition assay (Methods) with two males and a single female to test if differences in genital lobe length and courtship behavior may influence the chance of a male to copulate first with a single female. This first engagement into copulation will be referred to as “copulation success”. We filmed courtship and copulation behavior (Additional datafile 1: Figure

S1) of two males of the selection stock and a single female of stock 15090-1698.01 or 15090-1698.02. We annotated 111 trials where both males simultaneously courted the female (Methods) (51 and 60 trials with females of stocks 15090-1698.01 or 15090-1698.02, respectively). Then, we dissected the genitalia of both males to determine lobe lengths. We examined 54 additional trials with two courting males from the selection stock that had partially amputated left lobes and females of stock 15090-1698.01. This was done to test if other characters might co-vary with left lobe length and could potentially also have an effect on copulation success. We shortened the left lobes of both males in order to be able to neglect possible effects of the caused wound. The relative courtship activity or vigor of the two males in each trial was compared by quantifying periods that each male spent touching the female ovipositor or the ground next to it with the proboscis (mouth-parts), hereafter referred to as “licking behavior” (Methods). Licking was chosen over other behaviors because it occurs throughout courtship of *D. pachea* (Additional datafile 1: Figure S2) and was relatively easy to spot. We counted the sum duration of these periods (sum licking duration) and the number of licking periods (number of licking sequences). A copulation was considered to take place, once a male had mounted the female abdomen and achieved to settle into an invariant copulation position (Methods).

In the majority of trials (130/165), the first mounting attempt of a male resulted in a stable copulation. We did not observe any male behavior that could indicate an attempt of the non-copulating male to directly usurp the female as observed for *D. ananassae* or *D. bipectinata* [26, 27]. However, in 35 trials, a total of 51 failed mounting attempts were observed (Methods). These were not proportionally higher among trials with lobe amputated males (9 failed mounting attempts in 7 / 54 trials 13%) compared to unmodified males (42 failed mounting attempts in 28 / 111 trials 25%). Thus, the partial lobe amputation probably did not decrease the ability of a male to mount a female. Only 15/51 failed mounting attempts lasted 8 seconds or longer. Among those, the non-mounted male continued to court the female in 10/15 attempts. However, this was also observed in

131/165 successful mounting attempts and these proportions are not significantly different (χ^2 -test, $\chi^2 : 1.4852$, $df = 1$, $P = 0.2230$). Altogether, mounting failure was not found to be significantly associated with the courtship activity of the other male.

Different variables were tested to have an effect on male copulation success by using a Bradley Terry Model [48] (Methods). In trials with unmodified males, copulation success was positively associated with left lobe length and the sum licking duration of the males during courtship (Table 1). The effect of sum licking duration is illustrated when comparing the ratio of sum licking duration and total courtship duration between the copulating and the non-copulating male (Figure 3a). This ratio was higher in the copulating male compared to the non-copulating male. In particular, we identified a positive interaction of failed mounting attempts and female age, indicating that the negative effect of failed mounting attempts is pronounced in young females but not in older females. However, we observed failed mounting attempts in only 35 trials (Additional datafile 1: Figure S3) and the model estimates were sensitive to outlier observations (Additional datafile 1: Figure S10). Therefore, we relied on a simpler but more robust model that only takes sum licking duration and left lobe length into account (Table 1). More observations are required to evaluate the interaction of failed mounting attempts and female age.

In trials with males that underwent lobe surgery, left lobe length was not significantly associated with copulation success (Table 2) or with male courtship behaviors. Thus, we cannot exclude the possibility that left lobe length co-varies with yet another unknown selected trait that reveals an advantage for the male to copulate. Since the copulating male still showed an increased sum licking duration compared to the non-copulating male (Figure 3), the non-significant model fit may thus reflect insufficient statistical power due to a smaller number of observations compared to trials with unmodified males.

Courtship duration is increased when two males court a female simultaneously

Regardless of the stock for the female used, courtship durations appeared to be longer when both males courted the female simultaneously compared to trials where only one male courted (Mann-Whitney-Wilcoxon test, $W = 1906.5$, $N = 48/165$, $P = 4.7 \times 10^{-8}$) (Figure 3b). We also observed that courtship duration was strikingly increased in females that were older than 11 days (after emerging from the pupa) compared to younger females (Additional datafile 1: Figure S4) (Mann-Whitney test, $W = 660$, females younger than 12 days / females 12 days or older = 33 / 132, $P = 6.333 \times 10^{-10}$). Thus, older females may be less likely to copulate and may also discriminate less between males based on failed mounting attempts (see above).

Lobe length is not associated with the amount of ejaculate in female sperm storage organs

To test whether lobe length influences the amount of ejaculate deposited into the female after copulation, we dissected a random subset of females (54 of stock 15090-1698.01 and 24 of stock 15090-1698.02) in trials with non-modified lobes that revealed a copulation. Neither right lobe length, left lobe length, sum licking duration or female age (days after emerging from the pupa) was significantly associated with ejaculate content in the spermathecae (Table 3). However, male adult age and copulation duration had a negative effect. Given that *D. pachea* males become fertile about 13 days after emerging from the pupa [49], our results indicate an optimal period of male ejaculate transfer at the beginning of their reproductive period. Copulation duration had a negative effect on sperm transfer (Table 3), indicating that failed sperm transfer may result in prolonged attempts and longer copulation duration.

193 Discussion

194 Left lobe length affects the chance of a male to copulate

195 We observed that most females copulated first with the male that had the longer left lobe
 196 when two males courted simultaneously in our trials. These results suggest that a long left lobe
 197 might not only stabilize an asymmetric genital complex during copulation [44], but also increases
 198 the chance of a male to engage into copulation. With our data, we cannot exclude that the sensing of
 199 the lobe length by the female occurs just before intromission, while male and female genitalia are
 200 contacting each other or if it is used even during courtship before genital contacts establish. Other
 201 factors, unrelated to lobe length, can also bias copulation. For example, females might prefer to
 202 copulate with males of larger body size or that display particular courtship behaviors. In our study,
 203 we found no significant effect of tibia length on copulation success, which was used as an
 204 approximation for overall body size. However, the mutation(s) associated with lobe length variation
 205 in this stock are unknown. It is possible that one or several alleles associated with lobe length still
 206 segregate in this stock, even after 50 generations of inbreeding. If such segregating mutations have
 207 pleiotropic effects on other male characters (morphological, physiological or behavioural), these
 208 traits are expected to co-vary together with lobe length and the effect of lobe length that we detected
 209 might actually be due to the other factors. In trials with artificially shortened left lobes, males were
 210 taken from the selection stock but had the expected wild-type lobe asymmetry. Left lobe length
 211 difference was then artificially introduced in those males, so that it would not be expected to co-
 212 vary with the relative phenotypic expression of the supposed underlying mutation. In such trials, left
 213 lobe length after artificial length reduction was not associated with copulation success so that we
 214 cannot rule out the possibility that left lobe length in our initial experiment with unmodified males
 215 affects copulation success indirectly via a pleiotropic mutation affecting both, lobe length and other
 216 traits. It is possible that we failed to detect a significant effect because of a reduced variation in left
 217 lobe length among the lobe-modified males, compared to the initial experiment with non-ablated

males. In addition, we could not measure left lobe length on living males before lobe amputation. Therefore, we could not test whether the lobe length "before" length reduction influences male mating success. Furthermore, lobe amputation did not only cause a length reduction of the left lobe but also removed the bristles located at the distal tip. Laser ablation experiments have shown that these bristles contribute to the stabilization of the mating complex during copulation [44]. The bristles may also be important in pre-copulatory events, but we could not measure this since they were removed from both males in our trials. Future mate-choice experiments could be carried out with an unmodified male and a male where only those bristles are ablated [44].

Male intra-sexual competition is potentially reflected by male courtship intensity

Previous analyses in *D. ananassae* and *D. bipectinata* relate male genital spines to pre-copulatory abilities to achieve a copulation due to increased male intrasexual competitiveness in so called "scramble competition" or vigorous copulation attempts with females that are optionally already copulating with another male [26, 27]. In *D. pachea*, this behavior was not observed and most mounting events of a male resulted in a stable copulation. Once a copulation started, the non-copulating male was never observed to separate the couple. Instead, it continued to display courtship behaviors such as licking, wing vibration or touching the female side in the majority of cases. In addition, male mounting was only observed upon female wing opening and oviscapt protrusion. This indicates that mounting in *D. pachea* is a complex behavior involving both sex partners. One possibility might be that upon mounting the chance of a *D. pachea* male to engage into copulation thus may largely depend on male-female communication rather than on male-male competition.

Total courtship durations were increased in trials where both males courted the female simultaneously compared to trials with only one male courting. This indicates that females potentially require more time to accept one of the males for copulation in cases where two males court simultaneously. Alternatively, longer courtship durations might indeed rely on male-male

competition, causing mutual disturbances in courtship display. We detected a strong association of copulation success with the sum licking duration, which we used as an approximation for overall courtship activity or vigor of each male. This is achieved by maintaining its position close to the female ovipositor in competition with the other male, suggesting that licking behavior in part reports male-intrasexual conflict.

Potential female benefits gained from choosing males with long left lobes

Female mate choice on male sexual traits was suggested to rely at least in part on the way a sexual trait is displayed or moved [50]. This implies that a certain quantity but also accuracy of locomotor activity would matter in courtship and might better reflect overall male quality and “truth in advertising” than a morphological character or ornament alone. Our data suggest that male settling upon mounting the female might be crucial for a successful copulation since we identified a negative effect of failed mounting attempts on copulation success and a positive interaction of failed mounting attempts with female age, possibly indicating that young females are more discriminative compared to older females with respect to male mounting performance. However, this effect was only based on 35 trials (Additional datafile 1, Figure S3) and model estimates were sensitive to outliers. Future work must further evaluate this trend. Female mate choice is hypothesized to be based on direct benefits, affecting fecundity and survival of the female, but also on indirect benefits that relate to genetic quality, fecundity and survival of the progeny [50, 51]. Left lobe length and male mounting performance might therefore influence female mate-choice through enhancing the quality of male courtship signals and by enhancing the coupling efficiency of male and female genitalia upon mounting. It remains to be resolved what these courtship signals could be and how they might be presented to the female. We did not observe direct contact of the female with male genitalia prior to the male mounting attempts at copulation start. A possible pre-copulatory signal mediated by genital lobes could therefore be visual or vibratory. During *Drosophila* courtship, males perform abdomen shakes (“quivers”) that generate substrate-borne vibratory signals [52].

Perhaps the quiver frequency or amplitude could be affected by the length of the left lobe, thus producing a modified vibration signal. Alternatively, asymmetric lobes might be associated with lateralized courtship. For example, the beetle species *Sitophilus oryzae* and *Tribolium confusum* were observed to perform left-biased copulation attempts, which led to higher mating success over right-biased males [53]. However, in our study the male always mounted the female from the rear, along the female midline. We also noted that the licking side upon mounting was most commonly the rear of the female abdomen, but changed frequently in response to female abdomen movements and due to the presence of the other male. Therefore, we did not identify any lateralized male pre-copulatory behavior. This is in contrast with the lateralized posture of *D. pachea* males later during copulation, with the male being shifted about 6-8° to the right side of the female abdomen [41, 42, 44].

Alternatively, left lobe length might potentially associate with direct copulation benefits for male and female by ensuring the efficiency of ejaculate transfer during copulation. Males of *D. pachea* produce giant sperm and previous estimates revealed that a *D. pachea* male transmits only about 44 ± 6 sperm cells per copulation [47], while the maximum female sperm storage capacity in the spermathecae was estimated to be much higher and to be about 304 sperm cells [47]. Wild-caught *D. pachea* females were found to contain sperm from at least 3-4 males based on spermathecae filling levels [47]. As asymmetric lobes stabilize the genital complex [44], they might have a positive effect on the amount of sperm transferred per copulation. However, we have not observed any effect of lobe length on ejaculate presence in the spermathecae after copulation. We note that we did not test for the presence of seminal fluid in the female outside of the spermathecae. Also, our quantification method was not precise and sample size was limited. Accurate direct sperm counts requires radio-labeling methods [47], which was not applicable in our experimental setup. In future approaches, improved sperm counts should be applied to test for a correlation between lobe length, female-male copulation complex stability and sperm quantity in female spermathecae.

Based on our data, we note that left lobe length affects the chance of a male to establish genital contacts that lead to a firm complex of female and male genitalia, congruent with previous observations in other insects [26–28, 54]. In particular, the male genital spines in *D. ananassae* and *D. bipectinata* were observed to play a role in pre-copulatory events but had no measurable influence later on with respect to sperm transfer or fecundity [26, 27, 54].

Interestingly, copulation duration negatively affected ejaculate presence in the female. This observation suggests that the couple separates once successful sperm transfer is achieved. However, not all copulations reveal sperm transfer. In occasions of failure the copulation duration may therefore be prolonged to further attempt to mate. Copulation duration varies largely among *Drosophila* species from fractions of a minute to up to about 2 h in *D. acanthoptera* [42, 49]. Copulation duration in *D. pachea* is thus comparatively long with respect to other species of the genus, pointing to presumably complex copulatory interactions affecting sperm transfer. Future studies must analyze the internal processes of sperm transfer during *D. pachea* copulation to better understand this process and its possible limitations.

Indirect paternal effects on the growth and survival of offspring could potentially play a role in *D. pachea* female choice for males with long left lobes. It has been demonstrated that mate choice based on particular male sexual characters correlates with offspring survival in diverse species [18, 20–23, 55]. However, our experiment did not allow us to test for these effects because females were sacrificed after the mating experiment and no progeny was obtained. We assessed ejaculate filling levels instead of the amount of female progeny because *D. pachea* females rarely lay eggs after a single insemination and about four copulations are necessary for a female to start oviposition [56]. To test for indirect effects, future studies should ideally use progeny males from wild caught *D. pachea* females and test if subtle lobe length variation in those males would correlate with offspring survival in single couple crosses. A similar approach was used by Hoikkala

et al. (1998) [55], who found that *Drosophila montana* courtship song frequency of wild caught males correlated with the survival rate of the male's progeny.

Age of reproductive activities appear to be non-overlapping in sibling males and females

The amount of ejaculate present in the female spermathecae after a single copulation was negatively affected by the age of the copulating male. Similarly, it was previously found that the number of progeny in single couple crosses (presumably involving multiple copulations) was also dependent on male adult age in *D. pachea* [57]. Males in *D. pachea* need about 10-14 days at 25° C after emergence to become sexually mature [49, 56, 57]. This is related to adult testis growth and production of giant sperm [47]. It was shown that the relatively long time to reach male sexual maturity impacts the proportion of sexually active *D. pachea* adults (operational sex ratio), which is female-biased [49]. Our findings suggest that males potentially have a maximum fertility period at the beginning of their sexual maturity at approximately 13-16 days after emerging from the pupa. The detected reduction of transferred ejaculate in males older than 16 days implies an additional male reproductive limitation.

It was hypothesized that the delay in male sexual maturity might potentially lead to decreased sibling mating in *D. pachea* [56]. In *D. melanogaster*, sibling matings yield fewer progeny compared to crosses with unrelated individuals [57]. In our study, the female was not a sibling of the two males. Courtship durations were shortest in trials with “young” virgin females, between 6 and 11 days. Given that females reach sexual maturity at about 4 days after emerging from the pupa [49, 57], our observation suggests that young females engage more quickly into copulation compared to older females. Overall, female and male maximum reproductive activities appear to be non-overlapping among siblings of opposite sex and similar age. Such an effect can be relevant in the wild if a cactus rot is colonized by a single previously fertilized female.

Conclusion

Longer male left genital lobes potentially enhance the probability of *D. pachea* males to engage into copulation, which is expected to confer a reproductive advantage over males with shorter left lobes. The chance of a male to copulate first with a female was also affected by the relative male courtship activity and potentially by mounting attempt failures, especially for females at the beginning of their reproductive activity. The evolution of left lobe length might not only relate to its previously known function as a stabilizing device during copulation, but also through pre-copulatory mate choice by increasing the chances to copulate and to increase the coupling efficiency of genitalia upon mounting. Lobes do not appear to impact sperm transfer in later copulatory events. In addition, reproductive ages are non-overlapping between the two sexes potentially decreasing the amount of sibling mating in this species.

Methods

Fly stock establishment and maintenance

Two *D. pachea* stocks 15090-1698.01 and 15090-1698.02 were retrieved from the San Diego Drosophila Species Stock Center (now The National Drosophila Species Stock Center, College of Agriculture and Life Science, Cornell University, USA) (Additional datafile 1: Table S1). Flies were maintained in 25 x 95 mm plastic vials containing 10 mL of standard Drosophila medium (60 g/L brewer's yeast, 66.6 g/L cornmeal, 8.6 g/L agar, 5 g/L methyl-4-hydroxybenzoate and 2.5% v/v ethanol) and a ~ 10 x 50 mm piece of bench protection sheet (Bench guard). As *D. pachea* requires 7-dehydrocholesterol for proper development [58–60], we mixed the medium of each vial with 40 µL of 5 mg/mL 7-dehydrocholesterol, dissolved in ethanol (standard *D. pachea* food). Flies were kept at 25°C inside incubators (Velp) with a self-made light installation for a 12 h light: 12h dark photo-periodic cycle combined with a 30-min linear illumination change between

light (1080 lumen) and dark (0 lumen). We used males of stock 15090-1698.02 to generate a new stock with increased proportions of males with short genital lobes. For this, we chose 3 males with apparently symmetric (aberrant) genital lobes and crossed them with 3-4 sibling virgin females. We repeated the selection with the progeny for a total of 36 generations. Then, we discarded males with clearly visible asymmetric (wild-type) lobes from the progeny for another 14 generations to derive the final stock (selection stock).

Virgin fly selection

Virgin flies at 0-1 d after emerging from the pupa were CO₂ anaesthetised on a CO₂-pad (INJECT+MATIC Sleeper) under a stereo-microscope Stemi 2000 (Zeiss), separated according to sex and maintained in groups of 20-30 individuals. Males and females were raised until reaching sexual maturity, about 14 days for males and 4 days for females at 25° C [49]. This allowed us to use virgin individuals in each mating experiment. Males were anaesthetised on the CO₂ pad (see above), sorted according to lobe morphology (asymmetric and symmetric lobes) and isolated into single vials at least two days before each mating experiment took place.

Mate-competition assay

We used a single virgin female of wild-type stock 15090-1698.01 or 15090-1698.02 and two sibling males of the selection stock that visually differed in lobe length when inspected with a binocular microscope (Additional datafile 1: Figure S5). This selection was done to increase the average pairwise difference of lobe lengths between the two males. Specimens were introduced into a white, cylindrical mating cell (Additional datafile 1: Figure S1) with a diameter of 20 mm, a depth of 4 mm and a transparent 1 mm Plexiglas top-cover. Optionally, mating cells were concave with a diameter of 20 mm and a depth of 4 mm at the center (Additional datafile 3: trials 1-16). Flies were transferred without CO₂ anaesthesia using a fly aspirator: a 7 mm diameter silicone tube closed at

the tip with cotton and a 1000 μ L wide bore (3 mm) micro-tip. Movie recordings were started as soon as the chamber was immediately put under the camera (see below).

All mating trials were carried out inside a temperature and humidity controlled climate chamber at 25° C $\pm$ 0.1° C and 80% $\pm$ 5% or 60% $\pm$ 5% (trials 1-15) humidity. We used digital cameras MIRAZOOM MZ902 (OWL) or DigiMicro Profi (DNT) to record copulation and courtship behaviour. The MIRAZOOM MZ902 (OWL) camera was mounted on a modified microscope stand (191348, Conrad) equipped with a 8-cm LED white light illumination ring (EB-AE-COB-Cover, YM E-Bright) and a platform to hold four individual cylindrical mating cells (Additional datafile 1: Figure S1). Trial 16 (Additional datafile 3) was filmed with the OWL camera and a concave shaped mating cell that was put on a flat plastic cover on top of the microscope stand. For trials 1-15, we used concave shaped mating cells that were put onto the stand of the DigiMicro Profi (DNT) camera. Data was acquired with the programs Cheese (version 3.18.1) (<https://wiki.gnome.org/Apps/Cheese>) or GUVVIEW (version 0.9.9) GTK UVC (trials 1-15) on an ubuntu linux operating system in webm or mkv format. Up to four trials were recorded in a single movie, which was split after recording and converted into mp4 format with Avidemux 2.6 (<http://www.avidemux.org/>) to obtain a single movie per experiment.

We waited for a copulation to take place between one of the males and the female (see below). The mating cell was then shortly recovered from the climate chamber in order to remove the non-copulating male with an aspirator and to transfer it into a 2-mL reaction tube filled with 70% ethanol. This was performed after the copulating male appeared to have settled into a stable mating posture on the female abdomen, which was controlled by live-imaging. The copulating male and the female were isolated from the mating cell after copulation had ended and were also isolated into single 2-mL reaction tubes filled with 70% ethanol. Optionally, the female was kept alive for 12-24 h in a vial containing 5-10 mL grape juice agar (24 gr/L agar, 26 gr/L sucrose, 120 mg/L Tegosept, 20% ν /_v grape juice, and 1.5% ν /_v ethanol). The presence of eggs on the plate was

411 systematically checked but never observed. Females were finally sacrificed to prepare
412 spermathecae.

413 **Left lobe surgery**

414 Epandrial lobe surgery was done on 5-6 day-old *D. patchea* adult males of the selection stock
415 following Rhebergen et al. (2016). Males were anaesthetised on a CO₂ pad (see above) and then
416 further immobilized with a small copper wire, which was slightly pressed onto the male abdomen.
417 The left epandrial lobe was shortened to various lengths with micro dissection scissors (SuperFine
418 Vannas Scissors, World Precision Instruments). Flies were let to recover for at least 7 days on
419 standard *D. patchea* food at 25° C in groups. No mortality was observed in males that underwent
420 lobe surgery, similar to what was observed by Rhebergen et al. (2016). Partially amputated lobes
421 ranged in length from lacking at least their most distal tip to more reduced stumps whose length
422 approximated the corresponding right lobe.

423 **Adult dissection and imaging**

424 Adults were dissected in water with forceps (Forceps Dumont #5, Fine Science Tool) inside
425 a transparent 25 mm round dissection dish under a stereo-microscope (Zeiss Stemi 2000). Male
426 genitalia were isolated by piercing the abdomen with the forceps between the genital arch and the
427 A6 abdominal segment and thereby separating the genitalia from the abdomen. We also dissected
428 the left anterior leg of each dissected male. Female spermathecae were recovered after opening the
429 ventral abdomen with forceps and removal of the gut and ovaries. The spermathecae were isolated
430 and separated according to left and right sides and immediately examined using a microscope (see
431 below). All dissected tissues were stored in 200 µL storage solution (glycerol:acetate:ethanol, 1:1:3)
432 at 4° C.

433 For imaging, male genitalia were transferred into a dissection dish filled with pure glycerol
434 and examined with a VHX2000 (Keyence) microscope equipped with a 100-1000x VH-Z100W

(Keyence) zoom objective at 300-400 fold magnification. Genitalia were oriented to be visible in posterior view. The left and right lateral spines, as well as the dorsal edge of the genital arch were aligned to be visible in the same focal plane. In some preparations, the genital arch broke and lobes were aligned without adjusting the position of the dorsal genital arch. The experiment was discarded from analysis in cases where lobes could not be aligned. Male lobe lengths were measured on acquired images as the distance between the base of each lateral spine and the tip of each lobe (Figure 1c,d) using ImageJ version 1.50d (<https://imagej.nih.gov/ij>). Male legs were put on a flat glycerol surface with the inner side of the tibia facing to the camera. Legs were imaged at 200 fold magnification with the VHX 2000 microscope (Keyence) as described above.

We prepared the female paired sperm storage organs (spermathecae) at 12 h - 24 h after copulation (54 females from stock 15090-1698.01 and 24 females from stock 15090-1698.02, additional file 2) and examined the presence of ejaculate. *D. pachea* has giant sperm [47], which makes direct sperm counts difficult. Therefore, we determined an apparent average spermathecae filling level per female based on visual inspection, similar to Jefferson (1977) [57]. Female spermathecae were arranged on the bottom of the transparent plastic dissection dish, filled with water. Images were acquired using transmission light in lateral view with the Keyence VHX2000 microscope at 400-500 fold magnification. Ejaculate was directly visible inside the transparent spermathecae. Ejaculate filling levels were annotated for each spermatheca separately to match three different categories: 0, 1/6 or 1/3 of its total volume. Then, the average filling level for both spermathecae was calculated for each female.

Annotation of courtship and copulation behaviour

Videos were analysed with the OpenShot Video Editor software, version 1.4.3 (<https://www.openshot.org>) to annotate the relative timing of courtship and copulation in our experiments (Additional datafile 1: Figure S2, Additional datafile 3). Data was manually entered into spreadsheets. We annotated the beginning of male courtship as the start of at least three

consecutive male courtship behaviours according to Spieth (1952) [61], such as male touching the female abdomen with the forelegs, wing vibration, male following the female, and male licking behavior (see above). We estimated licking behavior throughout courtship of each male in trials where both males courted the female simultaneously. Licking occurs abundantly throughout courtship of *D. pachea* (Additional datafile 1, Figure S3) and was therefore chosen to assess the relative courtship contribution of the two males. We calculated the sum duration of all licking events per male and trial and the number of licking sequences. However, this behavior could not be quantified in 7 trials because the males changed positions very fast so that they could not be unambiguously distinguished (Additional datafiles 3,4). In all analysed trials, licking behaviour was observed throughout *D. pachea* courtship from shortly after courtship start to the start of copulation (Additional datafile 1: Figure S2). Both males reached at least once the licking position in all trials (Additional datafile 1: Table S3). This indicates that the female had physical contact with the proboscis of both males before copulation started (Additional datafile 1: Figure S2).

The beginning of copulation was defined as the moment when the male mounted the female abdomen. However, we only scored a copulation start if the male also settled into an invariant copulation posture. Upon mounting, the male moves its abdomen tip forth and back along the female oviscape. This behavior lasts for up to about 3 minutes and during this time the male grasps the female wings and abdomen with its legs. This behavior was previously described as settling period and ends with a characteristic right-sided, invariant copulation posture [41, 42, 44]. Any copulation that ended within the first minutes after mounting without settling was considered to be a “failed mounting attempt” (Additional datafile 4). The end of copulation was considered as the moment when male and female genitalia were separated and the male had completely descended with the forelegs from the female abdomen.

Copulation duration (Additional datafile 1: Table S3) was comparable to previous analyses of *D. pachea* copulation durations [41, 44, 47, 57]. We observed that copulation ended upon isolation of the non-copulating male in 3 trials with females of stock 15090-1698.01 and in 4 trials with females of stock 15090-1698.02, probably indicating premature copulation end due to specimen handling. However, the estimates were included into the dataset and had little impact on average copulation duration when comparing mean and median values.

Mate-competition analysis

Our aim was to compare at least 50 mate-competition trials, where both males courted the female simultaneously and where copulation was observed. In total, we carried out 98 and 89 trials with females of stocks 15090-1698.01 and 15090-1698.02, respectively (Additional datafile 1: Table S2). We performed another 62 trials with males that underwent surgical manipulation of the left lobe and females of stock 15090-1698.01 (Additional datafile 1: Table S4). In total, we removed 36 trials from the analysis because either copulation was not observed until 1 h after recording started (21 trials), flies escaped, got injured or died inside the mating cell (4 trials) or the dissections of male genitalia failed (11 trials, Additional datafile 1: Table S4). We thus examined 152 trials with unmodified males and 61 trials with males that had undergone genital surgery, (see below, Additional file 1: Table S4). Among those, we observed that both males courted the female simultaneously in 111 and 54 trials, respectively. Statistical assessment with the Bradley Terry Model (see below) further required a complete dataset so that we excluded 20 additional trials with missing values, 19 trials with unmodified males (left lobe length: 2 trials, right lobe length: 2 trials, licking behavior and/or failed mounting attempts: 7 trials, tibia length: 8 trials) and one trial with lobe-modified males (no information on tibia length). In total, we considered 92 trials with unmodified males and 53 trials with lobe-modified males.

Data analysis was performed in R version 3.6 [62]. Male copulation success (see above) was evaluated with a Bradley Terry Model [48] using the R package BradleyTerry2 [63]. This model is

suitable for logit fits to pairwise comparison data since it takes into account the special non-random (pairwise) structure of the dataset. We incorporated data (Additional datafile 3) into a list with three objects (Pacdata, available on DRYAD): 1) “contest” with the pairings of males, a trial ID and a factor indicating trials with males that had modified left genital lobes, 2) trial-specific variables “conditions” containing female age, female stock, courtship duration and a factor indicating which male initiated courtship, 3) male-specific predictor variables “male_predictors” with left and right genital lobe lengths, male tibia length, sum licking duration, number of licking sequences and failed mounting attempts. We developed the statistical model (Additional datafile 1: Tables S5-11) by evaluating the effect of different variables on copulation success and by comparing pairs of nested models, using Chi-square tests with the generic anova() function [63]. Outlier observations were considered when the standardized model residuals were greater than 2. The effect of corresponding trials on model fits was evaluated by comparison to additional model estimates without these outlier trials. We estimated correlations of male specific predictor variables and among numeric trial specific variables with the corrplot R package (Additional datafile 1: Figure S6). We found a significant correlation of the male predictor variables ‘sum licking duration’ and ‘number of licking sequences’. In addition, the trial specific variables ‘female age’ and ‘courtship duration’ were significantly correlated (Additional datafile 1: Figure S4). Correlated parameters were not incorporated into the same model for assessing the best fitting model (Additional datafile 1: Tables S5-S12). We tested trial specific variables in interaction with all male-specific variables, but only found an interaction of female age with failed mounting attempts in the dataset with unmodified males (Additional datafile 1: Figure S4, Table S8, Table 1). We also tested for interactions between male specific predictor variables but did not find any. The best fitting model for the datasets was: copulation success ~ sum licking duration + left_lobe length.

Spermathecae filling levels were evaluated with a linear model using the lm() function: spermatheca filling level ~ left lobe length + right lobe length + sum licking duration + Number of

licking sequences + copulation duration + female age + male age, and significance of each variable was evaluated with an F-test using anova().

List of abbreviations

Ma	million years ago
DF	degree of freedom
N	number of observations
P	probability

Declarations

Ethics approval and consent to participate

“Not applicable” --- This research focuses on invertebrate insects. There are no ethical considerations mentioned for these species according to EU Directive 86/609-STE123.

Consent for publication

“Not applicable”

Availability of data and material

The data sets supporting the results of this article are available in the DRYAD repository, https://datadryad.org/stash/share/jzbHRT070QV55ejT_BxIcopEoG-wVDjRI_HpQK0ergY

Competing interests

The authors declare no competing financial interests

Funding

This study was supported by the CNRS and by a grant of the European Research Council under the European Community’s Seventh Framework Program (FP7/2007-2013 Grant Agreement

no. 337579) given to VCO. BL was supported by a pre-doctoral fellowship Sorbonne Paris Cité of the Université Paris 7 Denis Diderot and by a fellowship "Initiatives d'excellence" (Idex ANR-18-IDEX-0001) of the Labex *Who Am I?* project (ANR-11-LABX-0071). The funding body had no influence on the design of the study, on collection, analysis, and interpretation of data, and in writing the manuscript.

Authors' contributions

ML and BL designed the experiments. BL and DC recorded fly copulation and performed light microscopy analysis of *Drosophila* male genitalia. BL and ML analysed the data. ML, VCO and BL wrote the manuscript. All authors have read and approved the manuscript.

Acknowledgements

We thank Therese A. Markow for discussions. We are very grateful to Alexandre Peluffo for help with the statistical analysis, Claus Lang for technical support and Juliette Royer for dissections of *D. pachea* male genitalia.

References

1. Bateman AJ. Intra-sexual selection in *Drosophila*. *Heredity* (Edinb). 1948;2:349–68.
2. Fisher RA. *The Genetical Theory of Natural Selection*. Oxford: Clarendon Press. 272 p. 1930.
3. Lande R. Models of speciation by sexual selection on polygenic traits. *Proc Natl Acad Sci*. 1981;78:3721–5.
4. Bonduriansky R. The evolution of male mate choice in insects: a synthesis of ideas and evidence. *Biol Rev*. 2001;76:305–39.
5. Leonard J, Córdoba-Aguilar A. *The evolution of primary sexual characters in animals*. Oxford University Press; 2010.
6. Edward DA, Chapman T. The evolution and significance of male mate choice. *Trends Ecol Evol*. 2011;26:647–54.
7. Kokko H, Jennions MD. The Relationship between Sexual Selection and Sexual Conflict. *Cold Spring Harb Perspect Biol*. 2014;6.

- 580 8. Eberhard WG. Sexual selection and animal genitalia. Harvard University Press; 1985.
- 581 9. Birkhead TR, Pizzari T. Postcopulatory sexual selection. *Nat Rev Genet.* 2002;3.
- 582 10. Thornhill R. Cryptic Female Choice and Its Implications in the Scorpionfly *Harpobittacus*
583 *nigriceps*. *Am Nat.* 1983;122:765–88.
- 584 11. Eberhard WG. Evolution of genitalia: theories, evidence, and new directions. *Genetica.*
585 2010;138:5–18.
- 586 12. Arnqvist G. Comparative evidence for the evolution of genitalia by sexual selection. *Nature.*
587 1998;393:784–6.
- 588 13. Chapman T, Arnqvist G, Bangham J, Rowe L. Sexual conflict. *Trends Ecol Evol.* 2003;18:41–7.
- 589 14. Eberhard WG. Copulatory Courtship and Cryptic Female Choice in Insects. *Biol Rev.*
590 1991;66:1–31.
- 591 15. Eberhard WG. Evidence for widespread courtship during copulation in 131 species of insects
592 and spiders, and implications for cryptic female choice. *Evolution (N Y).* 1994;48:711–33.
- 593 16. Ghiselin MT. The distinction between primary and secondary sexual characters. In: Leonard JL,
594 Córdoba-Aguilar A, editors. *The Evolution of Primary Sexual Characters in Animals*. New York:
595 Oxford University Press; 2010. p. 9–14.
- 596 17. Andersson M. Female choice selects for extreme tail length in a widowbird. *Nature.*
597 1982;299:818–20.
- 598 18. von Schantz T, Göransson G, Andersson G, Fröberg I, Grahn M, Helgée A, et al. Female choice
599 selects for a viability-based male trait in pheasants. *Nature.* 1989;337:166–9.
- 600 19. Hill GE. Plumage coloration is a sexually selected indicator of male quality. *Nature.*
601 1991;350:337–9.
- 602 20. Norris K. Heritable variation in a plumage indicator of viability in male great tits *Parus major*.
603 *Nature.* 1993;362:537–9.
- 604 21. Petrie M. Improved growth and survival of offspring of peacocks with more elaborate trains.
605 *Nature.* 1994;371:598–9.
- 606 22. Hasselquist D, Bensch S, von Schantz T. Correlation between male song repertoire, extra-pair
607 paternity and offspring survival in the great reed warbler. *Nature.* 1996;381:229–32.
- 608 23. Welch AM, Semlitsch RD, Gerhardt HC. Call Duration as an Indicator of Genetic Quality in
609 Male Gray Tree Frogs. *Science (80-).* 1998;280:1928 LP – 1930.
- 610 24. Birch M. Pre-courtship use of abdominal brushes by the nocturnal moth, *Phlogophora*
611 *meticulosa* (L.) (Lepidoptera: Noctuidae). *Anim Behav.* 1970;18:310–6.
- 612 25. Sreng L. Cockroach mating behaviors, sex pheromones, and abdominal glands (Dictyoptera:
613 Blaberidae). *J Insect Behav.* 1993;6:715–35.

- 614 26. Polak M, Rashed A. Microscale laser surgery reveals adaptive function of male intromittent
615 genitalia. *Proc R Soc B*. 2010;277.
- 616 27. Grieshop K, Polak M. The precopulatory function of male genital spines in *Drosophila*
617 *ananassae* [Doleschall] (Diptera: Drosophilidae) revealed by laser surgery. *Evolution* (N Y).
618 2012;66.
- 619 28. Bertin A, Fairbairn DJ. One tool, many uses: precopulatory sexual selection on genital
620 morphology in *Aquarius remigis*. *J Evol Biol*. 2005;18:949–61.
- 621 29. Ploog DW, MacLean PD. Display of penile erection in squirrel monkey (*Saimiri sciureus*).
622 *Anim Behav*. 1963;11:32–9.
- 623 30. Wickler W. Ursprung und biologische Deutung des Genitalprasentierens männlicher Primaten. *Z*
624 *Tierpsychol*. 1966;23:422–37.
- 625 31. Miller EH. Genitalic traits of mammals. In: Leonard J, Córdoba-Aguilar A, editors. *The*
626 *evolution of primary sexual characters in animals*. New York: Oxford University Press; 2010. p.
627 471–93.
- 628 32. Bohme W. The Tucano Indians of Colombia and the Iguanid Lizard *Plica plica*: Ethnological,
629 Herpetological and Ethological Implications. *Biotropica*. 1983;15:148–50.
- 630 33. Mautz BS, Wong BBM, Peters RA, Jennions MD. Penis size interacts with body shape and
631 height to influence male attractiveness. *Proc Natl Acad Sci*. 2013;110:6925–30.
- 632 34. Estes RD. The significance of horns and other male secondary sexual characters in female
633 bovids. *Appl Anim Behav Sci*. 1991;29:403–51.
- 634 35. Muller MN, Wrangham R. Sexual Mimicry in Hyenas. *Q Rev Biol*. 2002;77:3–16.
- 635 36. Kahn AT, Mautz B, Jennions MD. Females prefer to associate with males with longer
636 intromittent organs in mosquitofish. *Biol Lett*. 2010;6:55–8.
- 637 37. Langerhans RB, Layman CA, DeWitt TJ. Male genital size reflects a tradeoff between attracting
638 mates and avoiding predators in two live-bearing fish species. *Proc Natl Acad Sci*. 2005;102:7618–
639 23.
- 640 38. Moreno-García M, Cordero C. On the function of male genital claspers in *Stenomacra*
641 *marginella* (Heteroptera: Largidae). *J Ethol*. 2008;26:255–60.
- 642 39. Michiels NK. Mating Conflicts and Sperm Competition in Simultaneous. In: Birkhead TR,
643 Møller AP, editors. *Sperm competition and sexual selection*. New York: Academic Press; 1998. p.
644 219–254.
- 645 40. Pitnick S, Heed W. New Species of Cactus-Breeding *Drosophila* (Diptera: Drosophilidae) in the
646 Nannoptera Species Group. *Ann Entomol Soc Am*. 1994;87:307–10.
- 647 41. Lang M, Orgogozo V. Distinct copulation positions in *Drosophila patcha* males with symmetric
648 or asymmetric: External genitalia. *Contrib to Zool*. 2012;81:87–94.

- 649 42. Acurio A, Rhebergen FT, Paulus SE, Courtier-Orgogozo V, Lang M. Repeated evolution of
650 asymmetric genitalia and right-sided mating behavior in the *Drosophila* nanoptera species group.
651 BMC Evol Biol. 2019;19:109.
- 652 43. Lang M, Polihronakis Richmond M, Acurio AE, Markow TA, Orgogozo V. Radiation of the
653 *Drosophila* nanoptera species group in Mexico. J Evol Biol. 2014;27:575–84.
- 654 44. Rhebergen FT, Courtier-Orgogozo V, Dumont J, Schilthuisen M, Lang M. *Drosophila pachea*
655 asymmetric lobes are part of a grasping device and stabilize one-sided mating. BMC Evol Biol.
656 2016;16:176.
- 657 45. Pitnick S, Markow TA, Spicer GS. Delayed male maturity is a cost of producing large sperm in
658 *Drosophila*. Proc Natl Acad Sci. 1995;92:10614–8.
- 659 46. Pitnick S, Markow T, Spicer GS. Evolution of Multiple Kinds of Female Sperm-Storage Organs
660 in *Drosophila*. Evolution (N Y). 1999;53:1804–22.
- 661 47. Pitnick S, Markow TA. Male gametic strategies: sperm size, testes size, and the allocation of
662 ejaculate among successive mates by the sperm-limited fly *Drosophila pachea* and its relatives. Am
663 Nat. 1994;143:785–819.
- 664 48. Bradley RA, Terry ME. Rank Analysis of Incomplete Block Designs: I. The Method of Paired
665 Comparisons. Biometrika. 1952;39 3/4:324–45.
- 666 49. Pitnick S. Operational sex ratios and sperm limitation in populations of *Drosophila pachea*.
667 Behav Ecol Sociobiol. 1993;33:383–91.
- 668 50. Byers J, Hebets E, Podos J. Female mate choice based upon male motor performance. Anim
669 Behav. 2010;79:771–8.
- 670 51. Kirkpatrick M, Ryan MJ. The evolution of mating preferences and the paradox of the lek.
671 Nature. 1991;350:33–8.
- 672 52. Fabre CCG, Hedwig B, Conduit G, Lawrence PA, Goodwin SF, Casal J. Substrate-Borne
673 Vibratory Communication during Courtship in *Drosophila melanogaster*. Curr Biol. 2012;22:2180–
674 5.
- 675 53. Benelli G, Romano D, Stefanini C, Kavallieratos NG, Athanassiou CG, Canale A. Asymmetry
676 of mating behaviour affects copulation success in two stored-product beetles. J Pest Sci (2004).
677 2017;90:547–56.
- 678 54. Grieshop K, Polak M. Evaluating the post-copulatory sexual selection hypothesis for genital
679 evolution reveals evidence for pleiotropic harm exerted by the male genital spines of *Drosophila*
680 *ananassae*. J Evol Biol. 2014;27.
- 681 55. Hoikkala A, Aspi J, Suvanto L. Male courtship song frequency as an indicator of male genetic
682 quality in an insect species, *Drosophila montana*. Proc R Soc London Ser B Biol Sci.
683 1998;265:503–8.

- 684 56. Pitnick SS. Sexual selection and sperm production in *Drosophila*. Thesis (Ph. D.)--Arizona State
685 University, 1992.
- 686 57. Jefferson M. Breeding biology of *Drosophila pachea* and its relatives. 1977.
- 687 58. Heed WB, Kircher HW. Unique Sterol in the Ecology and Nutrition of *Drosophila pachea*.
688 Science (80-). 1965;149:758–61.
- 689 59. Lang M, Murat S, Clark AG, Gouppil G, Blais C, Matzkin LM, et al. Mutations in the
690 *neverland* gene turned *Drosophila pachea* into an obligate specialist species. Science (80-).
691 2012;337:1658–61.
- 692 60. Warren JT, Wismar J, Subrahmanyam B, Gilbert LI. *Woc* (*without children*) gene control of
693 ecdysone biosynthesis in *Drosophila melanogaster*. Mol Cell Endocrinol. 2001;181:1–14.
- 694 61. Spieth HT. Mating behavior within the genus *Drosophila* (Diptera). Bull Am Museum Nat Hist.
695 1952;99.
- 696 62. R Core Team. R: A language and environment for statistical computing. R Foundation for
697 Statistical Computing, Vienna, Austria. 2013. 2014.
- 698 63. Turner H, Firth D. Bradley-Terry Models in R: The BradleyTerry2 Package. J Stat Software,
699 Artic. 2012;48:1–21.

700 **Supporting Information**

701 **Additional datafile 1:** Supplementary Figures and Tables, **Figure S1:** Camera device for movie
702 recording, **Figure S2:** Courtship and copulation duration in competition mating experiments,
703 **Figure S3:** Failed copulation attempts compared to female age, **Figure S4:** Courtship duration
704 increases with female age, **Figure S5:** Pairwise lobe length differences of males used in the
705 competition mating experiment, **Figure S6:** Pearson correlations of male specific predictor
706 variables, **Table S1:** *Drosophila pachea* resources, **Table S2:** Genital lobe lengths in *D. pachea*
707 stocks, **Table S3:** Courtship duration, male licking behavior and copulation duration in trials where
708 both males courted the female, **Table S4:** Competition mating trials used for courtship analysis,
709 **Tables S5-S12:** Bradley Terry Model Fit.

710 **Additional datafile 2:** Raw measurements of lobe lengths in different *D. pachea* stocks

711 **Additional datafile 3:** Measurements of mating behaviour, genital lobe lengths and spermathecae
712 filling levels of individuals used in the competition mating experiments.

713 **Additional datafile 4:** Dataset for male licking behavior and male mounting.

714 Tables

715 **Table 1: Bradley–Terry (BT) model examining the effects of left lobe length, sum licking**
716 **duration and failed mounting attempts on copulation success**, in 92 trials with unmodified males.
717 Model: copulation success ~ sum licking duration + left lobe length, null deviance: 127.539 on 92 degrees
718 of freedom, residual deviance: 98.416 on 90 degrees of freedom. Influential trials 87 and 98 were detected to
719 have only marginal effects on model estimates.

variable	estimate	std. error	z value	P (> z)
sum licking duration	0.714251	0.206293	3.462	0.000536
left lobe	0.007511	0.003254	2.308	0.020981

720 **Table 2: Bradley–Terry (BT) model examining the effects of left lobe length, sum licking**
721 **duration, and number of licking sequences on copulation success**, in 53 trials with males that had
722 surgically modified left genital lobes. Model: copulation success ~ sum licking duration + left lobe length,
723 null deviance: 73.474 on 53 degrees of freedom, residual deviance: 68.163 on 51 degrees of freedom

variable	estimate	std. error	z value	P (> z)
sum licking duration	0.241981	0.166135	1.457	0.145
left lobe	0.007070	0.004615	1.532	0.126

724 **Table 3: Male age and copulation duration affect spermathecae filling levels.** Linear model
725 (spermathecae filling level ~ sum licking duration + left lobe length + right lobe length + copulation
726 duration + female age + male age), 32 observations were deleted due to missingness: sum licking
727 duration 30 values, left lobe length 1 value, right lobe length 1 value, multiple R²: 0.4975, adjusted
728 R²: 0.4201, F-statistic: 6.434 on 6 and 39 DF, P-value: 8.956e-05

variable	F value	P
sum licking duration	0.8658	0.357838
left lobe length	0.0447	0.833711
right lobe length	2.4460	0.125901
copulation duration	11.3116	0.001737
female age	0.3300	0.568960
male age	23.6079	1.954e-05

729

Figure Legends

Figure 1: *Drosophila patchea* male genital lobes. (a) Male of stock 15090-1698.02 in ventral view and with asymmetric lobes. (b) Male of the selection stock with apparently symmetric lobes. The scale bars correspond to 200 μ m. (c,d) Posterior view of a dissected male terminalia, (c) stock 15090-1698.02 with asymmetric lobes (d) selection stock. The scale bars correspond to 100 μ m, dots indicate lobe length measurement points, red arrows point to the lobe tips and black arrows indicate the lateral spines.

Figure 2: Genital lobe lengths differ between *D. patchea* stocks. Lengths of the left and right epandrial lobe lobes are presented for (a) stock 15090-1698.01, (b) stock 15090-1698.02, and (c) the selection stock. Sibling males of those used in the mating experiment are shown in panel (c). Each point represents one male. The variance of left lobe length is increased in the selection stock compared to the source stock 15090-1698.02 (Levene's test: selection stock / 15090-1698.02: DF = 1/147, F = 11.506, P = 0.0009), while the variance of the right lobe length were not significantly different (Levene's test, DF = 1/147, F = 0.5164, P = 0.4735).

Figure 3: Courtship durations and relative courtship activity. (a) The ratio of sum licking duration and total courtship duration in each trial shows that the copulating male (cop, grey box-plots) had a higher average sum licking duration compared to the non-copulating male (non-cop, white box-plots). Each point represents one male. (b) The total courtship duration was shorter in trials with only one male displaying courtship signs (one, white box-plots) compared to trials with both males courting simultaneously (both, grey box-plots) (Mann-Whitney-Wilcoxon test, female stock 15090-1698.01, non modified males, W = 385, N = 25/51, P = 0.0053 ; females of stock 15090-1698.02, W = 219, N = 16/60, P = 0.0009 ; females stock 15090-1698.01, modified males, W = 62.5, N = 7/54, P = 0.0044). Horizontal bars indicate pairwise comparisons with P-values ** < 0.01 or *** < 0.001. Each point represents one mating experiment.

a **b** **c**

sum licking duration / total courtship duration