Extracellular DNA (eDNA) enables early detection of the phenotypic switch of *Pseudomonas* sp. during biofilm development

Fatemeh Bajoul Kakahi¹, Dingrong Kang¹, Daniel C. Volke², Nicolas T. Wirth², Pablo I. Nickel², Frank Delvigne^{1*}

¹Terra Research and Teaching Centre, Microbial Processes and Interactions (MiPI), Gembloux Agro-Bio Tech, University of Liege, Gembloux, Belgium.

²The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, 2800 Kongens Lyngby, Denmark.

*Corresponding author: F.Delvigne@uliege.be

Abstract

Microbial populations undergo phenotypic switching as a response to environmental perturbations. For instance, some bacteria switch from a planktonic lifestyle to biofilm, resulting in altered physiological properties such as increased robustness depending on the conditions. However, the precise detection of phenotypic switching events during the bacterial life cycle is still a technical challenge. Propidium iodide (PI) is one of the most frequently used fluorescence indicators for assessing cell viability based on membrane permeability, yet PI-stained cells sometimes display a red-but-not-dead phenotype. In *Escherichia coli*, this phenomenon is connected to modulation of porins in the outer membrane (OM) to adapt OM permeability according to nutrient availability. In this study, we explored PI staining to assess phenotypic changes in *Pseudomonas* sp. during biofilm development. We show that this switch is linked to excretion of extracellular DNA (eDNA), rather than modification of OM permeability. Confocal laser scanning microscopy (CLSM) enabled direct visualization of red fluorescent clusters outside intact membranes of viable cells, suggesting that PI binds eDNA. Besides, the occurrence of PI-positive sub-populations was correlated with biofilm formation in the model bacterium *Pseudomonas putida* KT2440. Engineered derivatives thereof with altered biofilm-forming capabilities exposed a whole

continuum of phenotypic states involving planktonic cells and aggregates, and were identified according to the dynamic change of PI-positive cells with flow cytometry analysis. Our results demonstrate that PI is a fast, convenient and versatile staining for eDNA to rapidly monitor the phenotypic switching of *Pseudomonas* sp. during transitions in the bacterial lifestyle.

Graphical abstract

Running title: Tracking eDNA in *Pseudomonas* phenotypic switch

Keywords: Propidium iodide (PI), Flow cytometry, Extracellular DNA (eDNA), Biofilm, Aggregation, Sub-population, Phenotypic switching

Introduction

In natural ecosystems, many bacteria are able to switch between two modes of growth: on the one hand, a unicellular lifestyle, in which the cells are swimming freely, also called planktonic, and on the other hand, a multicellular lifestyle, in which the cells are sessile and form biofilms (Berlanga and Guerrero, 2016). Phenotypic switching between these two states is an important phenomenon involved in evolutionary development of bacteria and contributing to single-cell, as well as whole population fitness (Sousa *et al.*, 2011). While bacteria convert to a sessile state, the expression of several genes is altered causing a phenotype distinguishable from planktonic growing cells (Sousa *et al.*, 2011). Many microbial processes are dependent on phenotypic switching (e.g., persistence, utilization of alternative carbon source, competence...), leading to the formation of a heterogeneous, diverse and dynamic population that can overcome stresses and colonize new environments (Schreiber *et al.*, 2016)(Ackermann, 2015)(Eldar and Elowitz, 2010)(Delvigne *et al.*, 2014). Biofilm formation is indeed an efficient microbial strategy for survival. Upon switching from planktonic to biofilm mode of growth, bacteria go through several complex physiological, metabolic, and phenotypic diversifications (Drenkard, 2003). During biofilm growth, bacteria adopt a biofilm-specific phenotype, fundamentally different from that expressed in the corresponding planktonic cells. This switch to the biofilm-specific phenotype can trigger mechanisms responsible for antimicrobial resistance, enhanced virulence, and persistence (Mah and O'Toole, 2001). Our understanding of phenotypic switching in the biofilm field is still limited. However, it is critical to understand how phenotypic switching leads to the formation of biofilms, the understanding of which has important consequences from both a fundamental (i.e., microbial physiology) and applied (i.e., bioprocesses) point of view. *Pseudomonas putida* KT2440 is a non-pathogenic soil bacterium endowed with the ability to adapt to a large variety of physicochemical and nutritional niches (Benedetti *et al.*, 2016)(Nikel *et al.*, 2014), and able to form biofilms depending on the environmental conditions (Volke and Nikel, 2018). Biofilm formation is under control of a complex gene regulatory network (GRN) that orchestrates the phenotypic switching mechanism (Huang *et al.*, 2019)(Thomas *et al.*, 2014). Due to the complexity of the GRN driving

biofilm formation in *P. putida*, it is difficult to rely on a robust, single-cell proxy to characterize the phenotypic switching involved in the process.

Propidium iodide (PI) is widely used for bacterial viability staining; it can only cross compromised bacterial membranes and is therefore considered an indicator of membrane integrity (Shi *et al.*, 2007). However, recent studies demonstrated that PI might provide false-dead signals entering viable cells in some bacteria. Rosenberg *et al.* reported that PI staining is not indicative of membrane integrity in some instances, but rather binds extracellular DNA (eDNA), which is one of the components of bacterial extracellular polymeric substances (EPS) (Rosenberg *et al.*, 2019).

In most cases, the biofilm matrix represents around 90% of the total biofilm biomass and is mainly composed of polysaccharides, lipids, proteins, and eDNA (Fulaz *et al.*, 2019). The contribution of eDNA for the biofilm's three-dimensional structure varies significantly among species (Beitelshees *et al.*, 2018). Over the past few years, a great deal of attention was given to eDNA as one of the essential components of *Pseudomonas* sp.'s biofilm. In *Pseudomonas aeruginosa*, eDNA was found to represent a main component of biofilms. eDNA has also been reported to significantly affect the structural integrity of the biofilm matrix and protect bacterial cells in biofilms from physical stress, antibiotics, and detergents (Fulaz *et al.*, 2019). Moreover, eDNA can be used as a source of nutrients for living cells and spread genetic material within the biofilm as well as planktonic populations (Chimileski *et al.*, 2014)(Brown *et al.*, 2015). In fact, eDNA plays a critical role in several steps of biofilm formation, including early bacterial attachment, microcolony formation and aggregation, and in determining the overall biofilm architecture (Qin *et al.*, 2007)(Das *et al.*, 2010)(Whitchurch *et al.*, 2002). Recent reports suggest that the release of eDNA triggers bacterial surface adhesion by changing electrostatic and hydrophobic interactions between the bacterium and the surface and promotes early bacterial attachment and aggregation (Das *et al.*, 2010)(Das *et al.*, 2011)(Liu *et al.*, 2008). Another study suggests that differences in the concentration of eDNA between Gram-negative (e.g. *Pseudomonas aeruginosa* PA14, *Aeromonas hydrophila*, and *Escherichia coli*) and Gram-positive species (e.g. *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Enterococcus faecalis*) explains the variations in aggregation and biofilm formation behavior (Das *et al.*, 2014). Gram-negative bacterial strains displayed significant drops in aggregation when eDNA was removed through DNase I treatment (Das *et al.*, 2014).

In this work, the unique characteristics of eDNA was targeted in order to monitor biofilm switching dynamics at a single cell level. In particular, by using PI staining, we validated the importance of eDNA through different population states (i.e., planktonic phase, aggregation, and biofilm) and its effect on phenotypic diversification. We demonstrate that early biofilm development can be detected in planktonic cultures according to the degree of aggregation. Moreover, we demonstrate that PI staining is not associated with modifications in the composition of the OM but depends on the release of eDNA. The present study provides information on biofilm formation in *P. putida* KT2440 with both a biofilm-defective mutant ($\Delta lapA$) and biofilm over-producing derivative (overexpressing a mutated, hyperactive diguanylate cyclase), as well as in a natural isolate i.e., *Pseudomonas composti*. By applying a simple staining procedure relying on the use of PI, we were able to track phenotypic heterogeneity during the switch from planktonic to sessile lifestyle. Furthermore, we demonstrated that this property can be applied to identify and track biofilm-related phenotypic switching in a continuous culture system.

Results and discussion

eDNA binds PI and allows for the detection of subpopulations involved in phenotypic switching related to biofilm formation

In order to investigate the link between a PI-positive subpopulation and eDNA, we performed a systematic analysis based on flow cytometry (FC) and confocal laser scanning microscopy (CLSM). Firstly, we conducted a 24-h batch phase mini-bioreactor cultivation of *P. putida*. Biofilm samples were collected at the end of the fermentation and treated with DNase I to remove eDNA; the untreated sample was used as control. The samples were stained with PI and analyzed by FC. In the FC analysis, the microbial population was divided into two subpopulations, according to PI uptake (PI-positive and PI negative). The PI-negative subpopulation had fluorescence values of around 10^3 arbitrary units. These cells were considered to be unstained and thus having an intact, not permeable outer membrane. The bacterial subpopulation, which was partially stained and with

fluorescence values around 10^5 arbitrary units was classified as PI-positive. Flow cytometry analysis showed a significant difference in the PI-positive sub-population before and after treatment with DNase. The R1 region identified on the FC profiles corresponds to non-stained cells that were used as a negative control (Figure 1A). The majority of microbial cells were located in the R1 region before treated by the DNase enzyme, implied PI can bind with most of them (62.6%) (Fig. 1B). In contrast, PI-positive sub-populations in biofilm decrease remarkably to 32% after treating with DNase, which could indicate that cells lose the capacity to bind PI after removing eDNA (Figure 1C).

To further verify the effect of eDNA on PI-positive sub-population, we performed confocal laser microscopy imaging on planktonic, aggregation, and biofilm samples with double staining (e.g., PI and SYTO9, TOTO-1 and SYTO60). Double-stained planktonic cells with PI and SYTO9 showed they were generally co-localized. We noticed green fluorescing cores under red-stained shells of individual sections and confirmed PI staining not being indicative of membrane integrity but rather staining of eDNA, which is present outside of intact membrane (Figure 2A and 2B), except for the most strongly red cells that lacked green signal and were apparently true dead signals. We visualized the percent of eDNA on aggregation and biofilm samples by combining SYTO60 and TOTO-1 using CLSM to produce high-quality images of eDNA. SYTO60 stains bacterial cell red, whereas TOTO-1 stains eDNA green (Figure 2C and 2D). We observed a high amount of eDNA in both samples tested. Moreover, to further prove the role of eDNA in PI-staining, we visualized DNase treated biofilm samples with CLSM. The results show that PI signals in a non-treated sample with DNase enzyme are intensively high compare to a sample that has been treated with DNase (Figure 2E and 2F).

Next, we investigated a possible correlation between eDNA and iron concentration and PI-positive subpopulation in *Pseudomonas* sp. Indeed, iron has been shown to play a critical role in altering the eDNA production and biofilm formation in diverse bacteria (Yang *et al.*, 2007)(Allesen-Holm *et al.*, 2006)(Banin *et al.*, 2006)(Binnenkade *et al.*, 2014)(Oh *et al.*, 2018).

We have designed a matrix of iron concentrations gradient in a 96-well plate format. The extent of biofilm formation and quantity of eDNA has been assessed by crystal violet (CV) staining and fluorescent dye-based quantification, respectively. We found that biofilm formation of

Pseudomonas decreased with increasing iron concentrations (Figure 3A), which is in accordance with the previous study (Yang *et al.*, 2007). Moreover, the quantity of eDNA was assessed by using PicoGreen. The fluorescence intensity was measured immediately after adding the stain. Results demonstrated that the quantity of eDNA increased by decreasing iron concentration (Figure 3B).

In order to verify that the PI staining results are the direct consequence of biofilm formation, we genetically altered *P. putida* to either decrease or increase biofilm formation. To this end, we knocked out *lapA*, coding for the initial attachment protein. This strain, in the following called $\Delta lapA$, is unable to attach to abiotic surfaces. Therefore, biofilm formation is severely reduced (Boyd *et al.*, 2014). In order to increase biofilm formation, we transformed *P. putida* with the plasmid pS638::DGC-244, creating strain DGC. The introduced hyperactive di-guanylate cyclase increases the concentration of the second messenger cyclic-di-GMP (Christen *et al.*, 2006). The concentration of this second messenger is a major regulator for biofilm formation i.e., high c-di-GMP concentrations favor an adhesive lifestyle, while a low c-di-GMP concentration leads to a planktonic lifestyle (Gjermansen *et al.*, 2006). As the basal expression of the di-guanylate cyclase led to a substantial increase in biofilm development, we refrained from supplementing the inducer to minimize the perturbation of the cells.

In addition, the percentage of PI positive-subpopulation in the presence of high (100 μ M) and low (5 μ M) iron (Fe^{3+}) concentration was investigated by FC in all four *Pseudomonas* strains. We have noticed in all tested strains, except *P. putida* $\Delta lapA$, PI-positive subpopulation and biofilm formation increased as the iron concentration decreased (Figure 3C). Given the association of eDNA with the biofilm matrix, we assumed a significant difference in the percentage of PI-positive subpopulation in the presence of high and low iron concentration could be due to a change in the quantity of eDNA present in the sample. For *P. putida* $\Delta lapA$, as a non-biofilm former strain, we did not notice any change in PI- subpopulation or biofilm formation ability.

Considering the above-mentioned outcome, we evaluated the link between the biofilm-forming capacity and PI-positive sub-population by monitoring dynamic changes of PI-positive subpopulation during biofilm development. Results show a significant difference in PI-positive subpopulation among all tested strains. We observed high PI-positive percentage in DGC and *P.*

composti by comparison with *P. putida* wild-type and the $\Delta lapA$ derivative (Figure 4A). In addition, the PI-positive fraction in all cases was increased upon mid-exponential growth phase and decreased at the end of the batch phase (Figure 4A), which could be due to the release of eDNA and in the subsequent increase of aggregation at the beginning of the batch phase and decrease the amount of small aggregate in planktonic phase by switching to biofilm phase at the end of a phase (Figure S2). Meanwhile, we have evaluated the biofilm formation ability of all four strains using a crystal violet microtiter plate assay (Figure 4B). We have shown that DGC, and *P. composti* are strong biofilm former and produce accordingly more eDNA than the other strains tested. Hereafter, due to the strong biofilm formation capability of DGC strain, planktonic cells, aggregates, and biofilm samples of this strain have been collected during the batch phase, and corresponding PI-positive fraction of samples have been compared. FC analysis indicates a higher PI-positive percentage (41.8% and 30.8%) in biofilm and aggregation samples, respectively, by comparison with the planktonic phase (7.5%) (Figure 4C).

OM permeability and porin composition are not implied in phenotypic switching related to biofilm formation

Previous experiments have pointed out that PI binding could be related to a change in outer membrane (OM) composition (Sassi *et al.*, 2019). Indeed, it has been shown that particular porin deletion in *E. coli* can lead to increased OM permeability according to PI staining (Brognaux *et al.*, 2014) (Delvigne *et al.*, 2011)(Brognaux *et al.*, 2013). A systematic analysis of the effects of the group B porins on the outer membrane permeability of *P. putida* was then performed. Our aim was to investigate whether PI staining could also be related to modifying the porin composition in *Pseudomonas* OM. To this end, we constructed the single and multiple deletion mutants for all genes encoding OprB porins (OprB-I, OprB-II, and OprB-III) and measured the corresponding PI-positive subpopulation fraction through various time point starting from the late stationary phase. The proportion of PI-stained cells in the early stationary phase was low (0.2%-2%, data not shown). We have noticed that only 1-5% subpopulation of cells was stained with PI across all time points, and that they exhibited the red fluorescence in FC in all of the tested mutants (Figure 5). This result

indicates that single, double, and even triple knockouts of the *oprB* genes have no significant effect on the PI-positive subpopulation and that PI staining is not related to increased OM permeability for *P. putida*.

Cell auto-aggregation is an important step involved in phenotypic switching leading to biofilm formation

To investigate the impact of pre-formed aggregates in the planktonic phase on early biofilm formation, we have monitored aggregation development during the batch phase cultivation for strains under study. The auto-aggregation properties of four strains *P. putida*, *P. putida* $\Delta lapA$, *P. putida* DGC, and *P. composti* were measured over 24 h based on their sedimentation characteristics (Figure S3). Our results identified three auto-aggregation types, which have defined as follows. Strongly auto-aggregation (Agg⁺) strain (DGC) showed significantly ($P < 0.01$) higher auto-aggregation percentages ($\geq 90\%$) and cells clumped together immediately, forming a precipitate and resulting in a clear solution; non-auto-aggregation (Agg⁻) strain ($\Delta lapA$), which was unable to auto-aggregate (auto-aggregation percentage $< 21\%$) and produced consistent turbidity; and moderate Agg $\times$ strains (*P. putida* and *P. composti*) showing auto-aggregation percentages between 20% to 70%, their suspension showing both a precipitate and constant turbidity (Table S1). The result indicates that the biofilm-forming capability of strains under this study is highly related to auto-aggregation ability. The strong auto-aggregation strain DGC produces significantly higher biofilm compare to other tested strains.

On the other hand, in the same strains, aggregation in the planktonic phase was much more noticeable than for the two additional strains, which probably results in higher eDNA in the planktonic phase. In addition, to study the dynamic of aggregation formation during the batch phase, samples have been taken every 2 h and analyzed by FC. We have gated aggregation from the planktonic phase based on FSC and SSC signals. Flow cytometry analysis indicated the presence of aggregates phenotypic particles with a wide forward scatter distribution, indicative of variable particle sizes. These were distinguishable from single cells by their significantly bigger side- and forward- scatter values (Figure 6A). The degree of aggregation was measured for *P.*

putida, *P. putida* $\Delta lapA$, and *P. composti* during the batch phase (Figure 6B). However, in the case of the DGC strain, the size of the aggregates was too high to be detected by FC. The aggregation percentage was significantly higher in the natural isolate *P. composti* by comparison with *P. putida* and *P. putida* $\Delta lapA$, which were also producing less biofilm. In *P. putida* and *P. composti*, we have noticed an increase of aggregation at the beginning of exponential growth, followed by a decrease upon entry into stationary phase together with the appearance of biofilm on the wall of the bioreactor. However, this phenomenon is hardly observed for *P. putida* Δlap , in accordance with a very low cell aggregation.

Monitoring the phenotypic switching dynamics of *Pseudomonas* based on online flow cytometry allow for investigating biofilm lifecycle at a high temporal resolution

As indicated in the previous sections, PI has been identified as a useful biomarker to follow phenotypic switching from the planktonic phase to sessile life of *P. putida*. Accordingly, automated FC was then used for tracking this process during continuous culture with feast-to-famine cycles. This type of nutrient cycle is known to stimulate c-di-GMP synthesis and alter biofilm formation (Kalia *et al.*, 2013)(Monds *et al.*, 2007)(Gjermansen *et al.*, 2010)(Gjermansen *et al.*, 2005). The result pointed out that most of the cells are in a planktonic state during the batch phase (first 5 h), and few cells are in aggregation (< 7%). Expectedly, the percentage of the corresponding PI-positive subpopulation was low (< 5%), suggesting reduced eDNA release at this stage.

Besides these observations, a significant increase of the PI-positive subpopulation was observed after several feast to famine cycles. A gradual formation of a dense biofilm layer on the bioreactor wall was observed accordingly (Figure 7A and 7C). Intriguingly, the FC results show that the percentage of aggregation in the planktonic phase is fluctuating i.e., it rises during the first 20 h of continuous culture and then decreases. This non-monotonic behavior can be attributed to the accumulation of cells in the biofilm matrix during the first phase, making them impossible to detect by online FC. During the second stage, cells are released from the biofilm and can again be detected by FC (Figure 7B). The biofilm lifecycle can then be quantified at a very high time resolution based on this experimental set-up.

Discussion

Studies about phenotypic diversification of microbial populations attract a lot of attention because of the possible new functionalities exhibiting diversified populations. A switch between planktonic and sessile growth is believed to result in a phenotypic change in bacteria. Several bacterial species benefit from phenotypic switching, which is used to generate population diversity, increase bacterial fitness, and adapt to fluctuating environments (Sousa *et al.*, 2011).

Biofilm development in *P. putida* undergoes various structural and metabolic changes that are tightly controlled through a complex GRN, including phenotypic switch. The current state of knowledge in biofilm development in *P. putida* has been limited to the systematic use of confocal microscopy on *P. putida* biofilms grown in flow cells or various strategies to classify and characterize genes involved in each stage of biofilm development (Fazli *et al.*, 2014)(Klausen *et al.*, 2006). Despite the growing understanding of the biofilm cycle's structural and mechanistic details, little is known of the phenotypic switch during biofilm development in *P. putida*. As a result of the complexity of the GRN governing biofilm formation in *P. putida*, it is relatively challenging to depend on a robust single cell proxy to describe the phenotypic switching that takes place in this process. In the present work, we have identified PI as a useful biomarker to follow the cells switching from the planktonic phase to sessile life of *P. putida*. Our result from FC and CSLM indicates PI can bind to the eDNA present in a particular growth phase. The presence of eDNA, which is actively released through delegated cells, has gained significant interest (Okshevsky *et al.*, 2015). Recent studies showed PI could significantly overestimate dead cell counts in the presence of eDNA (Rosenberg *et al.*, 2019). Gião and Keevil noticed that co-staining of *Listeria monocytogenes* biofilms in tap water with PI and SYTO9 gives a false red signal, which is not an indication of dead cells. Still, this phenomenon it is due to the presence of eDNA (Gião and Keevil, 2014). Gallo also noted a similar picture staining the biofilm cells of *Salmonella typhimurium* strain carrying the reporter $P_{csgBA}::gfp$ with PI. They confirmed green cells were surrounded by a red corona, demonstrating that eDNA was localized around live cells (Gallo *et al.*, 2015). CLSM results look similar to what had been previously shown by Vilain and colleagues for *Bacillus cereus*

biofilm on glass wool. They indicated that nucleic acid was linked with the cell exterior, where PI had free access (Vilain *et al.*, 2009).

We have shown that modifying the amount of eDNA by considering different iron concentrations directly affects PI-positive sub-population. Iron has been demonstrated to be essential in the modulation of eDNA production, for instance, by triggering prophage induction in *Shewanella oneidensis* (Binnenkade *et al.*, 2014), mediating cell lysis during biofilm formation in *Streptococcus pneumoniae* (Trappetti *et al.*, 2011), or, in the case of *P. aeruginosa*, regulating eDNA production in response to both the *pqs* quorum-sensing system and changes in the external iron concentration (Allesen-Holm *et al.*, 2006)(Yang *et al.*, 2007). Our findings are in line with studies that showed how high concentrations of iron suppress *P. aeruginosa* biofilm formation in both microtitre plate cultivations and flow-chamber systems (Musk *et al.*, 2005). Simultaneously, low iron concentrations were reported to be necessary for *P. aeruginosa* biofilm (microcolony) formation in an artificial sputum medium (Sriramulu *et al.*, 2005). In *Campylobacter jejuni*, iron supplementation increased the accumulation of total reactive oxygen species (ROS) and the production of eDNA and extracellular polysaccharides, which leads to stimulating biofilms formation (Oh *et al.*, 2018).

Additionally, we observed that PI-positive sub-populations increase by manipulating the biofilm formation ability of *Pseudomonas* sp. In light of eDNA interfering with viability staining, we assume that the higher presence of eDNA leads to higher PI-positive fractions in strains producing more biofilm. We have also shown that the percentage of PI-positive fraction is more elevated in biofilm and aggregation samples as compared to planktonic cells. Since the quantity of eDNA is higher in the aggregation and biofilm sample, the observed difference could arise due to the binding between eDNA and PI.

In traditional bacterial biofilm development models, biofilm is generally assumed to originate from individual cells seeded on a surface. However, many biofilm-forming bacteria tend to aggregate in the planktonic phase; therefore, there is a chance that many natural and infective biofilms may be rising entirely or partially from pre-formed cell aggregates. Yet how these aggregates influence biofilm initiation and development is not clear. Given the affinity of many bacteria to aggregate and the common observation of aggregates in various environmental situations(Burmølle *et al.*,

2010)(Monier and Lindow, 2003)(Stoodley *et al.*, 2002) , it seems likely that natural biofilms are often initiated from pre-formed aggregates. *P. aeruginosa*, upon growth in liquid culture, forms large aggregates containing massively packed viable cells and eDNA(Schleheck *et al.*, 2009). For marine biofilms, the increased adhesiveness and surface conditioning of planktonic multicellular aggregates have been shown to accelerate bacterial attachment to a surface in early biofilm initiation (Bar-Zeev *et al.*, 2012) . Similarly, for *P. aeruginosa*, an increased affinity toward aggregation, probably an alternative for greater stickiness, has also been associated with increased biofilm formation (Häußler *et al.*, 2003)(Déziel *et al.*, 2001). We have shown that it is possible to detect early biofilm formation in the planktonic phase by determining the degree of aggregation by FC. Our result shows auto-aggregation ability of *Pseudomonas* sp. was highly correlated to their biofilm formation ability.

Previous observations made in *E. coli* by using mutants of the KEIO library have been confirmed that porin suppression increases the occurrence of a partially PI-stained phenotype (Sassi *et al.*, 2019). Although the mechanisms behind this increase in OM permeability are unknown, modifications in outer membrane porins composition have been hypothesized since porin removal has been shown to up and down-regulate other porins expression (Yang *et al.*, 2011). Recently, it has been demonstrated that *E. coli* is capable of modifying the porin composition to adapt its OM permeability under nutrient limitation. Thus, *E. coli* differentiates into two subpopulations displaying individual phenotypic features (non-permeabilized and permeabilized cells) during glucose limitation (Sassi *et al.*, 2019). In the case of *P. putida*, porins organization and regulation have been far less characterized. When compared with *E. coli*, the porins of strain KT2440 are organized in different gene and functional clusters, including membrane stabilization, transport of specific substrates, cell structure determination, and pore formation (Hancock and Brinkman, 2002). Among them, OprB has been more thoroughly explored and shows high homology with OprB from *P. aeruginosa*, which has been suggested to be involved in glucose uptake (Raneri *et al.*, 2018). However, our results have shown that PI staining in *Pseudomonas* sp. is not associated with the modification of OM composition but is more likely depending on the release of eDNA.

It is known that nutrient conditions affect bacterial populations' transition between planktonic state or free-floating aggregates and surface-attached biofilms (Schleheck *et al.*, 2009)(McDougald *et*

al., 2012)(Paananen *et al.*, 2003)(Gjermansen *et al.*, 2010). In this work, we have used automated FC for tracking the dynamics of the PI-positive subpopulation of *P. putida* upon exposure to successive feast-to-famine cycles. Such cycling conditions promoted phenotypic switching between planktonic cells, aggregates, and biofilm cells. The dynamic data acquired in this work allows for tracking phenotypic switching based on PI staining, taking to account switching to a stress condition effect the release of eDNA and biofilm formation (Ibáñez de Aldecoa *et al.*, 2017)(Lysis, 2018). Increasing the PI-positive sub-population also increased biofilm cells appearance on the wall of bioreactor. FC results show online profiling of aggregates during cultivation time, which helps us to monitor the phenotypic switch better. Taken together, the data gathered in this work point out that the combination of online FC, continuous cultivation system and appropriate single-cell proxy provides the user with the overall snapshot of phenotypic change during the biofilm formation process. Further investigation would be required in order to fully understand the mechanisms behind this phenotypic switch, for which our present method can be further expanded and used to this end.

Conclusion

In the present study, PI was shown to be able bind with the viable cells from different phenotypes of *Pseudomonas* sp. We have demonstrated that the ubiquity of eDNA made it an ideal target for monitoring phenotypic switching of *Pseudomonas* in the biofilm development process by a simple staining procedure with PI. Hence, this study opens novel avenues to possible applications of PI as a single-cell proxy allowing us to capture the "key" subpopulation of cells involved in biofilm formation. In this sense, this work paves the way to further understand the phenotypic diversification mechanism involved in biofilm formation. This body of knowledge can be exploited to control phenotypic switch during transition between planktonic, aggregates, and biofilm states and subsequently control biofilm formation, e.g., in bioprocesses relying on 'catalytic' biofilms.

Methods

Strains and medium composition

Bacterial strains used in this study are listed in Table 1. All strains were maintained in 25% (v/v) glycerol at -80°C in working seed vials (2 mL). Prior to experiments, one colony of each bacterium was used to inoculate 10 mL of lysogeny broth (LB) medium (10 g L^{-1} NaCl, 5 g L^{-1} yeast extract, and 12 g L^{-1} tryptone) and grown for 6 h with shaking at 30°C . Precultures and cultures of all bacteria were done in M9 minimal medium (33.7 mM Na_2HPO_4 , 22.0 mM KH_2PO_4 , 8.55 mM NaCl, 9.35 mM NH_4Cl , 1 mM MgSO_4 , and 0.3 mM CaCl_2), complemented with a trace elements (13.4 mM EDTA, 3.1 mM $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, 0.62 mM ZnCl_2 , 76 μM $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$, 42 μM $\text{CoCl}_2 \cdot 2\text{H}_2\text{O}$, 162 μM H_3BO_3 , and 8.1 μM $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$), 1 μg L^{-1} biotin and 1 μg L^{-1} thiamin) and supplemented with glucose (5 g L^{-1}) as the main carbon source (pH = 7.2). For iron-dependent experiments, a concentrated FeCl_3 solution was added to the media to give the final ferric iron concentrations described in the text. For strain DGC, the media was supplemented with gentamycin at a final concentration of 10 μg mL^{-1} .

Plasmid construction and genetic manipulations

Plasmid pS638::DGC was constructed by amplifying the hyperactive diguanylate cyclase mutant A0244 from *Caulobacter crescentus* (Christen *et al.*, 2006), with the primer pair P1 and P2. The resulting amplicon and vector pSEVA638 were digested with *Bam*HI and *Sac*I (FD0054 and FD1133, Thermo Fisher Scientific). The fragments were purified (NucleoSpin Gel and PCR Clean-up Columns, Macherey Nagel) and ligated with T4 DNA ligase (EL0014, Thermo Fisher Scientific). *E. coli* DH5 α was transformed with the ligation mixture and the cell suspension was plated on a gentamycin-selective plate. Subsequently, the constructed pS638::DGC (where *DGCA*⁰²⁴⁴ was placed under transcriptional control of the inducible *XylS/Pm* expression system) plasmid was isolated from a single colony and its correctness was confirmed by sequencing. All the plasmids are listed in Table 2.

The plasmids for deletion of *lapA* and the porins *oprB-I*, *oprB-II* and *oprB-III* were constructed according to (Wirth *et al.*, 2020b). In short, ~500 bp homology arms (HA) flanking the gene coding sequences were amplified from chromosomal DNA of *P. putida* KT2440 using the primer pairs

P3/P4 (*lapA*_HA1), P5/P6 (*lapA*_HA1), P7/P8 (*oprB-I*_HA1), P9/P10 (*oprB-I*_HA2), P11/P12 (*oprB-II*_HA1), P13/P14 (*oprB-II*_HA2), P15/P16 (*oprB-III*_HA1) and P17/P18 (*oprB-III*_HA2), and assembled into the suicide vector pSNW2 employing the USER cloning method (Cavaleiro *et al.*, 2015). The purified plasmids were introduced into stationary *P. putida* KT2440 cells by electroporation and selected on LB agar medium supplemented with kanamycin (50 $\mu\text{g mL}^{-1}$). The corresponding pSNW2 derivative, now fully integrated into the bacterial chromosome, was resolved by transforming the cells with the auxiliary plasmid pQURE6 and selection on gentamicin (10 $\mu\text{g mL}^{-1}$) and 3-methylbenzoate (1 mM) (Volke *et al.*, 2020). Resolved strains (GFP-negative and kanamycin-sensitive) were tested for the desired genotype by colony PCR (OneTaq® 2X Master Mix with Standard Buffer, New England Biolabs).

Mini-bioreactor cultivations

A bioreactor system from 2MAG© block, equipped with eight positions for fermentation vessels of 16 mL total volume and a working volume of 12-mL bioreactors, was used in these experiments. The bioreactor block is equipped with a magnetic inductive drive with two independent heat exchangers integrated into the bioreactor block, one for temperature control for the reaction mixture and the second to control the headspace's cooling of the bioreactors to prevent evaporation. The system is also equipped with fluorometric sensor spots for pH and dissolved oxygen (DO) positioned at the bottom of each reactor (MiniReaktor HTBD, Presens, Regensburg, Germany). We performed 24-h batch phase fermentations in order to test quickly and screen on PI-positive dynamic of strains under study, up to 6 replicates simultaneously. Samples have been taken every 1-2 h and immediately analyzed by FC.

Sample preparation

Adherent cells were harvested by scraping from the wall of the bioreactor vessel with a cell scraper and resuspended in PBS. In the case of aggregation, 5 mL culture was aliquoted into a 15-mL Falcon tube, and aggregates were allowed to sediment for 10 min. Then, aggregates were carefully picked up from the bottom of the tube by using sterile cut pipette tips, and transferred into 1.5-mL Eppendorf tubes. A quick spin at 1,400 rpm was applied for 15 s, and the supernatant was removed.

Both aggregates and biofilm samples were resuspended in sterile PBS and vortexed for 30 s. Before FC analysis, samples were gently sonicated (2× for 12 s, with 25% amplitude) to disperse clumps into single cells without causing cell lysis. A planktonic sample has been filtered through a sterile syringe filter (5 µm) to eliminate aggregates before FC analysis.

Bacteria staining protocols

The four fluorescent stains used in this study were: (i) PI (P4170, Sigma-Aldrich); a stock solution was prepared at 1 mg mL⁻¹ in sterile Milli-Q water and used at a final concentration of 1.5 µM in sterilized PBS; (ii) TOTO™-1 iodide (T3600, Invitrogen™ Thermo Fisher Scientific); a stock solution was prepared in DMSO (at 1 mM) and used at a final concentration of 2 µM in sterilized PBS; (iii) SYTO™9 (S34854, Invitrogen™ Thermo Fisher Scientific); a stock solution was prepared in DMSO at 5 mM and used at a final concentration of 5 µM in sterilized PBS; and (iv) SYTO™60 (S11342, Invitrogen™ Thermo Fisher Scientific); a stock solution was prepared in DMSO at 5 mM and used at a final concentration of 10 µM in sterilized PBS. All bacterial samples were stained right before FC analysis by adding 1 µL of the PI stock solution to 1 mL of a cell suspension in PBS (1×10⁷ cells mL⁻¹). The stained samples were incubated for 10 min in the dark at room temperature and analyzed by FC (live-dead gating was done based on heat-killed bacteria at 80°C for 1 h).

Flow cytometry

FC analysis of PI-stained bacteria was carried out using BD Accuri™ C6 device (BD Biosciences). Cell counts, red fluorescence (FL3), and Forward scatter (FSC) values were determined with FC. The software settings were as following: Fluidics, medium; Threshold, 20,000 on FSC-H; Run with limits, 10,000 events. Gating of dead and alive signal populations was executed on Propidium iodide (FL3-A; 670 nm LP) scatter plot. FC analysis of PI staining bacteria was carried out on planktonic, aggregation, and biofilm samples. Cells were diluted to an appropriate density with filtered 1x phosphate-buffered saline (1× PBS; 8 g L⁻¹ NaCl, 0.2 g L⁻¹ KCl, 1.4 g L⁻¹ Na₂HPO₄, and 0.2 g L⁻¹ KH₂PO₄, pH 7.2) stained with PI and analyzed by flow cytometry. Online flow cytometry platform (Accuri C6 BD Biosciences San Jose CA USA) has been used in combination with a

pulsed continuous cultivation system. The platform has been described in detail in a previous study (Sassi *et al.*, 2019). Briefly, sample processing following multiple steps, including sample acquisition and online staining, dilution threshold, and online FC analysis. Flow cytometry data were exported as an FCS file and processed by a custom Python script version (0.5.0).

DNase enzyme Treatment

Biofilm cells were prepared according to the protocol mentioned above. Then, the number of cells in the sample was adjusted to (1×10^7 cells mL^{-1}), and samples were resuspended in 500 μL of $1 \times$ DNase I buffer (10 mM Tris-HCl, pH = 7.5, 2.5 mM MgCl_2 , and 0.1 mM CaCl_2) with or without DNase I (final concentration 160 U mL^{-1} , Roche, cat. # 04716728001) and were incubated at 37°C for 3 h. After incubation, samples were pelleted by centrifugation at 8500 rpm for 10 min, resuspended in PBS, stained by PI, and analyzed by FCM.

Confocal laser scanning microscopy (CLSM)

All samples were analyzed by confocal laser-scanning microscope (LSM) LSM880 Airyscan super-resolution system (Carl Zeiss, Oberkochen, Germany). The images were taken with the Plan-Apochromat $63 \times / 1.4$ Oil objective. We used an excitation wavelength of 488 nm and emission at 500-550 nm for green fluorescence, and an excitation wavelength of 561 nm and emission at 580-615 nm for red fluorescence. Images were acquired continuously at a pixel resolution of 0.04 μm (regular Airyscan mode) in XY and 1- μm interval in Z step-size using the piezo drive. Before CLSM analysis, staining was performed with one or more fluorescent dyes in the following combinations: PI, SYTO9+PI, and TOTO1+SYTO60. An aliquot of 10-15 μL of 1:1 stain mixture solution has been added to a sample of planktonic, aggregate, and biofilm and incubated under exclusion of light (10 min for PI, SYTO9+PI, and 20 min for TOTO1+SYTO60).

Auto-aggregation assay

Auto-aggregation tests were adapted from the protocol published by Kos *et al.* (Kos *et al.*, 2003). For the auto-aggregation assay, the strains studied were first grown in M9 minimal medium for 18 h. After centrifugation (10,000 rpm, 10 min), pellets of cells were resuspended, washed twice with

1 $1 \times \text{PBS}$, and finally resuspended to $1 \times 10^8 \text{ CFU mL}^{-1}$ in the same medium. Then, suspensions were
2 vortexed and incubated at room temperature for 15 h. Samples were harvested during different
3 times along the incubation experiment (0, 1, 5, 9, and 15 h). Each time, a 1-mL aliquot from the
4 top of the suspensions was carefully removed and its absorbance (A) was read at 600 nm in a
5 spectrophotometer. Auto-aggregation was calculated using the following formula:

$$\text{Auto-aggregation (\%)} = [(A_0 - A_t)/A_0] \times 100$$

6
7
8
9 where A_0 indicates the absorbance at time 0 h and A_t indicates the absorbance every hour, up to 15
10 h.

11 **Quantification of biofilm growth and eDNA in biofilms in 96-well plates**

12 Two-day biofilm formation of *Pseudomonas* strains was determined by crystal violet staining using
13 a 96-well plate lid with pegs extending into each well (Nunc-TSP lid, Invitrogen™ Thermo Fisher
14 Scientific). Briefly, precultures were grown overnight at 30°C to an $\text{OD}_{600 \text{ nm}}$ of 1.0. The cell
15 suspensions were then adjusted to an $\text{OD}_{600 \text{ nm}}$ of 0.1 in M9 medium. A total of 160 μL cell
16 suspension were added to each well. Fresh medium was used as negative control. The plates were
17 sealed with parafilm and incubated with shaking at 180 rpm at 30°C. The biofilm biomass was
18 quantified with a crystal violet staining assay modified from previously reported CV assays (Ren
19 *et al.*, 2014). CV quantification was performed on the pegs of the Nunc-TSP lid culture system.
20 Briefly, after 48 h of cultivation, the peg lids were taken out and washed three times using PBS.
21 Subsequently, the peg lids were placed in plates with 180 μL of an aqueous 1% (w/v) CV solution.
22 Then, the lids were washed with PBS three times after staining for 20 min. Subsequently, the peg
23 lids with crystal violet stain were placed into a new microtiter plate with 200 μL of 33% (w/v)
24 glacial acetic acid in each well for 15 min. The optical density at 590 nm of each sample was
25 measured by a microplate reader (Tecan SPARK, Männedorf, Switzerland). The quantity of eDNA
26 associated with *Pseudomonas* biofilm was examined in 96-well black plates (Cell culture
27 microplate, $\mu\text{CLEAR}^{\text{®}}$) in triplicates parallel with the biofilm quantification. The 2-day biofilm
28 culture was rinsed three times with sterile distilled water. eDNA was quantified by QuantiFluor
29

dsDNA dye (QuantiFluor dsDNA System, Promega, Madison, WI, USA) according to manufacturer's protocol. Briefly, eDNA in each sample was mixed with 200 μ L of freshly prepared QuantiFluor dsDNA dye in TE buffer. Wells with QuantiFluor dsDNA dye were incubated for 5 min before measuring the fluorescence intensity (excitation wavelength = 504 nm and emission wavelength = 531 nm) using a Tecan microplate reader. Lambda DNA (Invitrogen™ Molecular Probes) was used to generate a calibration curve for each run.

Pulsed continuous cultivation with online flow cytometry

P. putida KT2440 was cultivated in a stirred bioreactor (Biostat B-Twin, Sartorius) in continuous mode with glucose feed pulses (0.5 g L⁻¹ per pulse) at a frequency of one pulse per hour. Data acquisition was performed every 12 min *via* a coupled, online FC analysis. A 1-L volume of M9 minimal medium (containing 5 g L⁻¹ glucose) was inoculated by overnight preculture (OD₆₀₀ = 0.3) in the 2-L capacity bioreactor. The pH and temperature were maintained at 7.2 and 30°C, respectively. Stirring was set at 800 rpm and aeration flow rate to 1 L min⁻¹ (1 vvm). The drop in DO marked the depletion of oxygen and the start of continuous cultivation mode. M9 minimal medium was fed without carbon source into the bioreactor at a dilution rate $D = 0.1$ h⁻¹. After 1 h, a pulse of glucose was injected in order to increase the concentration as fast as possible (within 30 s) to a global glucose concentration of 0.5 g L⁻¹. The microbial population was tracked by an online flow cytometry device as previously described (Nguyen *et al.*, 2021)(Sassi *et al.*, 2019). Briefly, a 0.8-mL sample was taken from the bioreactor each 12 min, automatically diluted and stained with PI, and then automatically injected into an Accuri C6 FC for detection. Before processing, the independent .fcs files obtained through online FC were compacted in a dataframe (.pkl file extension) based on the Pandas package (<https://pandas.pydata.org/>) from Python. This file was used for generating the plots with the evolution of PI-positive and aggregates subpopulations, as well as for generating individual dotplots with forward scatter (FSC) as x -axis and PI fluorescence as y -axis. These dotplots were further assembled into a single .avi movie file based on the ImageJ software (Rueden *et al.*, 2017) shown as Supplemental material (Movies S1). Raw .fcs data were deposited on FlowRepository and can be accessed by following the link:

<https://flowrepository.org/id/RvFrYohjOYhgmxWvC6PUFdME2OZX5BGPNYvVLJkpEa1NHzwVN5sllGgBYFgsRyHp>

References

- Ackermann, M. (2015) A functional perspective on phenotypic heterogeneity in microorganisms. *Nat Rev Microbiol* **13**: 497–508.
- Allesen-Holm, M., Barken, K.B., Yang, L., Klausen, M., Webb, J.S., Kjelleberg, S., et al. (2006) A characterization of DNA release in *Pseudomonas aeruginosa* cultures and biofilms. *Mol Microbiol* **59**: 1114–1128.
- Bagdasarian, M., Lurz, R., Rückert, B., Franklin, F.C.H., Bagdasarian, M.M., Frey, J., and Timmis, K.N. (1981) Specific-purpose plasmid cloning vectors II. Broad host range, high copy number, RSF 1010-derived vectors, and a host-vector system for gene cloning in *Pseudomonas*. *Gene* **16**: 237–247.
- Banin, E., Brady, K.M., and Greenberg, E.P. (2006) Chelator-induced dispersal and killing of *Pseudomonas aeruginosa* cells in a biofilm. *Appl Environ Microbiol* **72**: 2064–2069.
- Bar-Zeev, E., Berman-Frank, I., Girshevitz, O., and Berman, T. (2012) Revised paradigm of aquatic biofilm formation facilitated by microgel transparent exopolymer particles. *Proc Natl Acad Sci U S A* **109**: 9119–9124.
- Beitelshes, M., Hill, A., Jones, C.H., and Pfeifer, B.A. (2018) Phenotypic variation during biofilm formation: Implications for anti-biofilm therapeutic design. *Materials (Basel)* **11**: 1–18.
- Benedetti, I., de Lorenzo, V., and Nikel, P.I. (2016) Genetic programming of catalytic *Pseudomonas putida* biofilms for boosting biodegradation of haloalkanes. *Metab Eng* **33**: 109–118.
- Berlangua, M. and Guerrero, R. (2016) Living together in biofilms: The microbial cell factory and its biotechnological implications. *Microb Cell Fact* **15**: 1–11.
- Binnenkade, L., Teichmann, L., and Thormanna, K.M. (2014) Iron triggers λ So prophage induction and release of extracellular DNA in *Shewanella oneidensis* MR-1 biofilms. *Appl Environ Microbiol* **80**: 5304–5316.
- Boyd, C.D., Jarrod Smith, T., El-Kirat-Chatel, S., Newell, P.D., Dufrêne, Y.F., and O'Toole, G.A. (2014) Structural features of the *Pseudomonas fluorescens* biofilm adhesin LapA required for LapG-dependent cleavage, biofilm formation, and cell surface localization. *J Bacteriol* **196**: 2775–2788.
- Brognaux, A., Bugge, J., Schwartz, F.H., Thonart, P., Telek, S., and Delvigne, F. (2013) Real-time monitoring of cell viability and cell density on the basis of a three dimensional optical reflectance method (3D-ORM): Investigation of the effect of sub-lethal and lethal injuries. *J Ind Microbiol Biotechnol* **40**: 679–686.
- Brognaux, A., Francis, F., Twizere, J.C., Thonart, P., and Delvigne, F. (2014) Scale-down effect on the extracellular proteome of *Escherichia coli*: Correlation with membrane permeability and modulation according to substrate heterogeneities. *Bioprocess Biosyst Eng* **37**: 1469–1485.
- Brown, H.L., Reuter, M., Hanman, K., Betts, R.P., and Van Vliet, A.H.M. (2015) Prevention of biofilm formation and removal of existing biofilms by extracellular dnases of *Campylobacter jejuni*. *PLoS One* **10**: 1–21.
- Burmølle, M., Thomsen, T.R., Fazli, M., Dige, I., Christensen, L., Homøe, P., et al. (2010) Biofilms in chronic infections - A matter of opportunity - Monospecies biofilms in multispecies infections. *FEMS Immunol Med Microbiol* **59**: 324–336.
- Cavaleiro, A.M., Kim, S.H., Seppälä, S., Nielsen, M.T., and Nørholm, M.H.H. (2015) Accurate DNA Assembly and Genome Engineering with Optimized Uracil Excision Cloning. *ACS*

- 1 *Synth Biol* **4**: 1042–1046.
- 2 Chimileski, S., Franklin, M.J., and Papke, R.T. (2014) Biofilms formed by the archaeon
- 3 *Haloferax volcanii* exhibit cellular differentiation and social motility, and facilitate
- 4 horizontal gene transfer. *BMC Biol* **12**..
- 5 Christen, B., Christen, M., Paul, R., Schmid, F., Folcher, M., Jenoe, P., et al. (2006) Allosteric
- 6 Control of Cyclic di-GMP Signaling. *J Biol Chem* **281**: 32015–32024.
- 7 Das, T., Krom, B.P., Van Der Mei, H.C., Busscher, H.J., and Sharma, P.K. (2011) DNA-
- 8 mediated bacterial aggregation is dictated by acid-base interactions. *Soft Matter* **7**: 2927–
- 9 2935.
- 10 Das, T., Sehar, S., Koop, L., Wong, Y.K., Ahmed, S., Siddiqui, K.S., and Manefield, M. (2014)
- 11 Influence of calcium in extracellular DNA mediated bacterial aggregation and biofilm
- 12 formation. *PLoS One* **9**: 1–11.
- 13 Das, T., Sharma, P.K., Busscher, H.J., Van Der Mei, H.C., and Krom, B.P. (2010) Role of
- 14 extracellular DNA in initial bacterial adhesion and surface aggregation. *Appl Environ*
- 15 *Microbiol* **76**: 3405–3408.
- 16 Delvigne, F., Brognaux, A., Francis, F., Twizere, J.C., Gorret, N., Sorensen, S.J., and Thonart, P.
- 17 (2011) Green fluorescent protein (GFP) leakage from microbial biosensors provides useful
- 18 information for the evaluation of the scale-down effect. *Biotechnol J* **6**: 968–978.
- 19 Delvigne, F., Zune, Q., Lara, A.R., Al-Soud, W., and Sørensen, S.J. (2014) Metabolic variability
- 20 in bioprocessing: Implications of microbial phenotypic heterogeneity. *Trends Biotechnol* **32**:
- 21 608–616.
- 22 Déziel, E., Comeau, Y., and Villemur, R. (2001) Initiation of biofilm formation by *Pseudomonas*
- 23 *aeruginosa* 57RP correlates with emergence of hyperpiliated and highly adherent phenotypic
- 24 variants deficient in swimming, swarming, and twitching motilities. *J Bacteriol* **183**: 1195–
- 25 1204.
- 26 Drenkard, E. (2003) Antimicrobial resistance of *Pseudomonas aeruginosa* biofilms. *Microbes*
- 27 *Infect* **5**: 1213–1219.
- 28 Eldar, A. and Elowitz, M.B. (2010) Functional roles for noise in genetic circuits. *Nature* **467**:
- 29 167–173.
- 30 Fazli, M., Almlblad, H., Rybtke, M.L., Givskov, M., Eberl, L., and Tolker-Nielsen, T. (2014)
- 31 Regulation of biofilm formation in *Pseudomonas* and *Burkholderia* species. *Environ*
- 32 *Microbiol* **16**: 1961–1981.
- 33 Fulaz, S., Vitale, S., Quinn, L., and Casey, E. (2019) Nanoparticle–Biofilm Interactions: The
- 34 Role of the EPS Matrix. *Trends Microbiol* **27**: 915–926.
- 35 Gallo, P.M., Rapsinski, G.J., Wilson, R.P., Oppong, G.O., Sriram, U., Goulian, M., et al. (2015)
- 36 Amyloid-DNA Composites of Bacterial Biofilms Stimulate Autoimmunity. *Immunity* **42**:
- 37 1171–1184.
- 38 Gião, M.S. and Keevil, C.W. (2014) *Listeria monocytogenes* can form biofilms in tap water and
- 39 enter into the viable but non-cultivable state. *Microb Ecol* **67**: 603–611.
- 40 Gjermansen, M., Nilsson, M., Yang, L., and Tolker-Nielsen, T. (2010) Characterization of
- 41 starvation-induced dispersion in *Pseudomonas putida* biofilms: genetic elements and
- 42 molecular mechanisms. *Mol Microbiol* **75**: 815–826.
- 43 Gjermansen, M., Ragas, P., Sternberg, C., Molin, S., and Tolker-Nielsen, T. (2005)
- 44 Characterization of starvation-induced dispersion in *Pseudomonas putida* biofilms. *Environ*
- 45 *Microbiol* **7**: 894–904.
- 46 Gjermansen, M., Ragas, P., and Tolker-Nielsen, T. (2006) Proteins with GGDEF and EAL
- 47 domains regulate *Pseudomonas putida* biofilm formation and dispersal. *FEMS Microbiol*
- 48 *Lett* **265**: 215–224.
- 49 Hancock, R.E.W. and Brinkman, F.S.L. (2002) Function of *Pseudomonas* porins in uptake and
- 50 efflux. *Annu Rev Microbiol* **56**: 17–38.
- 51 Häußler, S., Ziegler, I., Löttel, A., Götz, F. V., Rohde, M., Wehmhöner, D., et al. (2003) Highly
- 52 adherent small-colony variants of *Pseudomonas aeruginosa* in cystic fibrosis lung infection.
- 53 *J Med Microbiol* **52**: 295–301.
- 54 Huang, M.Y., Woolford, C.A., May, G., Joel Mcmanus, C., and Mitchell, A.P. (2019) Circuit

- diversification in a biofilm regulatory network. *PLoS Pathog* **15**: 1–26.
- Ibáñez de Aldecoa, A.L., Zafra, O., and González-Pastor, J.E. (2017) Mechanisms and regulation of extracellular DNA release and its biological roles in microbial communities. *Front Microbiol* **8**: 1–19.
- Kalia, D., Merey, G., Nakayama, S., Zheng, Y., Zhou, J., Luo, Y., et al. (2013) Nucleotide, c-di-GMP, c-di-AMP, cGMP, cAMP, (p)ppGpp signaling in bacteria and implications in pathogenesis. *Chem Soc Rev* **42**: 305–341.
- Klausen, M., Gjermansen, M., Kreft, J.U., and Tolker-Nielsen, T. (2006) Dynamics of development and dispersal in sessile microbial communities: Examples from *Pseudomonas aeruginosa* and *Pseudomonas putida* model biofilms. *FEMS Microbiol Lett* **261**: 1–11.
- Kos, B., Šušković, J., Vuković, S., Šimpraga, M., Frece, J., and Matošić, S. (2003) Adhesion and aggregation ability of probiotic strain *Lactobacillus acidophilus* M92. *J Appl Microbiol* **94**: 981–987.
- Liu, H.H., Yang, Y.R., Shen, X.C., Zhang, Z.L., Shen, P., and Xie, Z.X. (2008) Role of DNA in bacterial aggregation. *Curr Microbiol* **57**: 139–144.
- Lysis, E.S.B. (2018) crossm. **84**: 1–18.
- Mah, T.F.C. and O'Toole, G.A. (2001) Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol* **9**: 34–39.
- Martínez-García, E., Aparicio, T., Goñi-Moreno, A., Fraile, S., and De Lorenzo, V. (2015) SEVA 2.0: An update of the Standard European Vector Architecture for de-/re-construction of bacterial functionalities. *Nucleic Acids Res* **43**: D1183–D1189.
- McDougald, D., Rice, S.A., Barraud, N., Steinberg, P.D., and Kjelleberg, S. (2012) Should we stay or should we go: Mechanisms and ecological consequences for biofilm dispersal. *Nat Rev Microbiol* **10**: 39–50.
- Monds, R.D., Newell, P.D., Gross, R.H., and O'Toole, G.A. (2007) Phosphate-dependent modulation of c-di-GMP levels regulates *Pseudomonas fluorescens* Pf0-1 biofilm formation by controlling secretion of the adhesin LapA. *Mol Microbiol* **63**: 656–679.
- Monier, J.M. and Lindow, S.E. (2003) Differential survival of solitary and aggregated bacterial cells promotes aggregate formation on leaf surfaces. *Proc Natl Acad Sci U S A* **100**: 15977–15982.
- Musk, D.J., Banko, D.A., and Hergenrother, P.J. (2005) Iron salts perturb biofilm formation and disrupt existing biofilms of *Pseudomonas aeruginosa*. *Chem Biol* **12**: 789–796.
- Nguyen, T.M., Telek, S., Zicler, A., Andres Martinez, J., Zacchetti, B., Kopp, J., et al. (2021) Reducing phenotypic and genotypic instabilities of microbial population during continuous cultivation based on stochastic switching dynamics. *bioRxiv* 2021.01.13.426484.
- Nikel, P.I., Martínez-García, E., and De Lorenzo, V. (2014) Biotechnological domestication of pseudomonads using synthetic biology. *Nat Rev Microbiol* **12**: 368–379.
- Oh, E., Andrews, K.J., and Jeon, B. (2018) Enhanced biofilm formation by ferrous and ferric iron through oxidative stress in *Campylobacter jejuni*. *Front Microbiol* **9**: 1–9.
- Okshevsky, M., Regina, V.R., and Meyer, R.L. (2015) Extracellular DNA as a target for biofilm control. *Curr Opin Biotechnol* **33**: 73–80.
- Paananen, A., Vuorimaa, E., Torkkeli, M., Penttilä, M., Kauranen, M., Ikkala, O., et al. (2003) Structural hierarchy in molecular films of two class II hydrophobins. *Biochemistry* **42**: 5253–8.
- Platt, R., Drescher, C., Park, S.K., and Phillips, G.J. (2000) Genetic system for reversible integration of DNA constructs and lacZ gene fusions into the *Escherichia coli* chromosome. *Plasmid* **43**: 12–23.
- Qin, Z., Ou, Y., Yang, L., Zhu, Y., Tolker-Nielsen, T., Molin, S., and Qu, D. (2007) Role of autolysin-mediated DNA release in biofilm formation of *Staphylococcus epidermidis*. *Microbiology* **153**: 2083–2092.
- Raneri, M., Pinatel, E., Peano, C., Rampioni, G., Leoni, L., Bianconi, I., et al. (2018) *Pseudomonas aeruginosa* mutants defective in glucose uptake have pleiotropic phenotype and altered virulence in non-mammal infection models. *Sci Rep* **8**: 1–15.
- Ren, D., Madsen, J.S., de la Cruz-Perera, C.I., Bergmark, L., Sørensen, S.J., and Burmølle, M.

- (2014) High-Throughput Screening of Multispecies Biofilm Formation and Quantitative PCR-Based Assessment of Individual Species Proportions, Useful for Exploring Interspecific Bacterial Interactions. *Microb Ecol* **68**: 146–154.
- Rosenberg, M., Azevedo, N.F., and Ivask, A. (2019) Propidium iodide staining underestimates viability of adherent bacterial cells. *Sci Rep* **9**: 1–14.
- Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., and Eliceiri, K.W. (2017) ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics* **18**.
- Sassi, H., Nguyen, T.M., Telek, S., Gosset, G., Grünberger, A., and Delvigne, F. (2019) Segregostat: a novel concept to control phenotypic diversification dynamics on the example of Gram-negative bacteria. *Microb Biotechnol* **12**: 1064–1075.
- Schleheck, D., Barraud, N., Klebensberger, J., Webb, J.S., McDougald, D., Rice, S.A., and Kjelleberg, S. (2009) *Pseudomonas aeruginosa* PAO1 Preferentially Grows as Aggregates in Liquid Batch Cultures and Disperses upon Starvation. *PLoS One* **4**: e5513.
- Schreiber, F., Littmann, S., Lavik, G., Escrig, S., Meibom, A., Kuypers, M.M.M., and Ackermann, M. (2016) Phenotypic heterogeneity driven by nutrient limitation promotes growth in fluctuating environments. *Nat Microbiol* **1**: 1–7.
- Shi, L., Günther, S., Hübschmann, T., Wick, L.Y., Harms, H., and Müller, S. (2007) Limits of propidium iodide as a cell viability indicator for environmental bacteria. *Cytom Part A* **71**: 592–598.
- Sousa, A.M., Machado, I., and Pereira, M.O. (2011) Phenotypic switching: an opportunity to bacteria thrive. *Sci against Microb Pathog Commun Curr Res Technol Adv* 252–262.
- Sriramulu, D.D., Lünsdorf, H., Lam, J.S., and Römling, U. (2005) Microcolony formation: A novel biofilm model of *Pseudomonas aeruginosa* for the cystic fibrosis lung. *J Med Microbiol* **54**: 667–676.
- Stoodley, P., Sauer, K., Davies, D.G., and Costerton, J.W. (2002) Biofilms as complex differentiated communities. *Annu Rev Microbiol* **56**: 187–209.
- Thomas, P., Popovic, N., and Grima, R. (2014) Phenotypic switching in gene regulatory networks. *Proc Natl Acad Sci U S A* **111**: 6994–6999.
- Trappetti, C., Gualdi, L., Di Meola, L., Jain, P., Korir, C.C., Edmonds, P., et al. (2011) The impact of the competence quorum sensing system on *Streptococcus pneumoniae* biofilms varies depending on the experimental model. *BMC Microbiol* **11**: 75.
- Vilain, S., Pretorius, J.M., Theron, J., and Brözel, V.S. (2009) DNA as an adhesin: *Bacillus cereus* requires extracellular DNA to form biofilms. *Appl Environ Microbiol* **75**: 2861–2868.
- Volke, D.C., Friis, L., Wirth, N.T., Turlin, J., and Nikel, P.I. (2020) Synthetic control of plasmid replication enables target- and self-curing of vectors and expedites genome engineering of *Pseudomonas putida*. *Metab Eng Commun* **10**: e00126.
- Volke, D.C. and Nikel, P.I. (2018) Getting Bacteria in Shape: Synthetic Morphology Approaches for the Design of Efficient Microbial Cell Factories. *Adv Biosyst* **2**: 1800111.
- Whitchurch, C.B., Tolker-Nielsen, T., Ragas, P.C., and Mattick, J.S. (2002) Extracellular DNA required for bacterial biofilm formation. *Science (80-)* **295**: 1487.
- Wirth, N.T., Kozaeva, E., and Nikel, P.I. (2020a) Accelerated genome engineering of *Pseudomonas putida* by I-SceI—mediated recombination and CRISPR-Cas9 counterselection. *Microb Biotechnol* **13**: 233–249.
- Wirth, N.T.,
- Worsey, M.J. and Williams, A.P. (1975) Metabolism of toluene and xylenes by *Pseudomonas putida* (arvilla) mt 2: evidence for a new function of the TOL plasmid. *J Bacteriol* **124**: 7–13.
- Yang, J.N., Wang, C., Guo, C., Peng, X.X., and Li, H. (2011) Outer membrane proteome and its regulation networks in response to glucose concentration changes in *Escherichia coli*. *Mol Biosyst* **7**: 3087–3093.
- Yang, L., Barken, K.B., Skindersoe, M.E., Christensen, A.B., Givskov, M., and Tolker-Nielsen, T. (2007) Effects of iron on DNA release and biofilm development by *Pseudomonas aeruginosa*. *Microbiology* **153**: 1318–1328.

1
2

Table 1. Bacterial strains and plasmids used in this study.

Strain	Relevant characteristics ^a	Reference or source
<i>E. coli</i> DH5α λ pir	Cloning host; F ⁻ λ ⁻ <i>endA1 glnX44(AS) thiE1 recA1 relA1 spoT1 gyrA96(Nal^R) rfbC1 deoR nupG</i> Φ 80(<i>lacZ</i> Δ <i>M15</i>) Δ(<i>argF-lac</i>) <i>U169</i> <i>hsdR17(r_K⁻ m_K⁺)</i> , λ pir lysogen	(Platt <i>et al.</i> , 2000)
<i>P. putida</i> KT2440	Wild-type strain, derived from <i>P. putida</i> mt-2 (Worsey and Williams, 1975) cured of the TOL plasmid pWW0	(Bagdasarian <i>et al.</i> , 1981)
LapA	Derivate of <i>P. putida</i> KT2440 with a clean deletion of <i>lapA</i> (<i>PP_0168</i>)	This study
DGC	Derivate of <i>P. putida</i> KT2440 harboring the plasmid pS638::DGC-244	This study
pSEVA638	<i>oriV</i> (pBBR1); <i>XylS/Pm</i> →multiple cloning site (MCS); Gm ^R	(Martínez-García <i>et al.</i> , 2015)
pS638::DGC-244	Derived from pSEVA638 with insertion of DGC-244 into the MCS	This study
pGNW2	<i>oriV</i> (R6K); <i>P_{EM14g}BCD</i> → <i>msfGFP</i> ; Km ^R	(Wirth <i>et al.</i> , 2020a)
pGNW2·Δ <i>lapA</i>	Derived from pGNW2 with homologous flanking region to <i>lapA</i> (<i>PP_0168</i>)	This study
pGNW2·Δ <i>oprB-I</i>	Derived from pGNW2 with homologous flanking region to <i>oprB-I</i> (<i>PP_1019</i>)	This study
pGNW2·Δ <i>oprB-II</i>	Derived from pGNW2 with homologous flanking region to <i>oprB-II</i> (<i>PP_01445</i>)	This study
pGNW2·Δ <i>oprB-III</i>	Derived from pGNW2 with homologous flanking region to <i>oprB-III</i> (<i>PP_3570</i>)	This study

Table 2. Oligonucleotides used in this study^a.

Oligonucleotide	Sequence (5'→3')	Use
P1	AAA <i>GAG CTC</i> TTA <u>GGA GGA AAA ACA</u> TAT GAA AAT CTC AGG CGC CCG GAC	Amplification of <i>dgcA0240</i>
P2	AAA <i>GGA TCC</i> TCA AGC GCT CCT GCG CTT G	
P3†	AGA TCC UAT TCA TCT ATA GAG TGC GGA TTC	Amplification of genomic regions adjacent to <i>lapA</i>
P4†	ATT GGA CUC TCC GTGTGACCCGATGG	
P5†	AGT CCA AUG TGA CAG ACC ACC GGG GCC	
P6†	AGG TCG ACU TCG ATT GGT CGA CGG GTA CG	
P7†	AGA TCC UCC GCA GCA GAT CTA CAA CG	Amplification of genomic regions adjacent to <i>oprB-I</i>
P8†	ACA TCC CUT TGC GTC CTC TTT	
P9†	AGG GAT GUG AGG CAA CTT GTT GTA AAT TTA CG	
P10†	AGG TCG ACU GCT GTC CTG ATG TTC GGT G	
P11†	AGA TCC UAA ACC CGC CAA CGA AAC C	Amplification of genomic regions adjacent to <i>oprB-II</i>
P12†	ACA TGA GAU AGC GCT ATC TTT TGATT	
P13†	ATC TCA TGU AAC CCC TTT TTT GAC CTG ACG	
P14†	AGG TCG ACU GTA GGC CTG CCA TTC GC	
P15†	AGA TCC UAT GCC GTG AAC AAG AAC CG	Amplification of genomic regions adjacent to <i>oprB-III</i>
P16†	ATT ACA GAA UCT CGG GTT GTC TTT G	
P17†	ATT CTG TAA UGT GAA CCG CCG GGG CCG C	
P18†	AGG TCG ACU GCT TCG ATC CAC CGT TCT C	

^a Oligonucleotides designed for *USER* cloning are indicated with a † symbol. Restriction sites are shown in italics, ribosomal binding site is underlined and start codon is shown in bold.

Figure 1. Flow cytometry analysis illustrating the effects of DNase treatment on PI-positive fraction of biofilm sample. **A** Negative control shows non-stained cells according to R1 region on the FC profiles. **B** Sample showing high PI uptake, before treated by the DNase enzyme, the majority of microbial cells were located in the R1 region (62.6%). **C** Sample After treating with DNase enzyme, the PI-positive fraction decreases significantly (32.0%), as a result of eDNA hydrolysis.

Figure 2. Confocal laser scanning microscopy (CSLM) images of *P. putida* KT2440 (planktonic, aggregation and biofilm) samples. **A-B** Planktonic sample co-stained with propidium iodide (PI) and SYTO9. **C** Biofilm cells (co-stained with TOTO-1 and SYTO60). **D** Cell aggregates stained with a combination of TOTO-1 and SYTO60. **E** Biofilm cells stained with PI before treatment with DNase enzyme. **F** Biofilm cells stained with PI after being treated with DNase enzyme.

Figure 3. Effect of iron concentration on biofilm formation, extracellular DNA and the PI-positive subpopulation. **A** Comparison of biofilm formation ability of *P. putida* KT2440 in the presence of different iron concentrations. **B** eDNA quantity in the presence of different iron concentrations. **C** Comparison of PI-positive percentage among DGC, *P. composti*, *P. putida* KT2440 and $\Delta lapA$ in the presence of high (100 μM) and low iron concentration (5 μM).

Figure 4. Comparison of PI-positive subpopulation percentage and biofilm formation capability among DGC, *P. composti*, *P. putida* KT2440, $\Delta lapA$. **A** Comparison of PI-positive fraction (%) cells during batch phase, $N \geq 3$. **B** Biofilm formation ability after 48h, $N \geq 5$. **C** Flow cytometry comparison of PI-positive subpopulation for planktonic, aggregated and biofilm sample from DGC strain.

Figure 5. Effects of deletion porins group B on outer membrane permeability in *P. putida* KT2440 mutant, $N \geq 3$.

Figure 6. **A** Gating the aggregation subpopulation on flow cytometry based on FSC and SSC signals. The blue dotted box separates the aggregation fraction from single cells. **B** Comparison of

the degree of aggregation for *P. composti*, *P. putida* KT2440 and its $\Delta lapA$ derivative by FC, $N \geq 2$.

Figure 7. Dynamics of phenotypic diversification in pulsed continuous cultivation system in *P. putida* KT2440. **A** PI-positive dynamic in continuous feast and famine nutrient cycle during 50 h. **B** Aggregation fluctuation dynamic during 50 h in automated pulsed continuous-culture bioreactor. **C** Macroscopic view of biofilm formation layer on the wall of the 2-L bioreactor after 50 h of cultivation.

Supplemental information

Table S1: comparison of auto-aggregation ability of different strains: *P. putida* DGC (KT2440 background), *P. putida* $\Delta lapA$ (KT2440 background), *P. putida* KT2440 and *P. composti*. Values in a row that are not preceded by the same letter are significantly different ($p \leq 0.01$)

Phenotypes	Strains	Autoaggregation (%)
Agg+ (Strongly autoaggregation)	DGC	100.00±0 ^a
Agg- (non-autoaggregation)	$\Delta lapA$	20.89±1.84 ^d
Agg +/- (Moderate autoaggregation)	<i>P. putida</i> KT2440	37.67±2.96 ^c
	<i>P. composti</i>	55.67±1.53 ^b

Figure S1: effect of different DNase I concentrations on the PI-positive fraction.

1

2

3 **Figure S2:** macroscopic view of biofilm formation on the wall of mini bioreactor vessel among
4 $\Delta lapA$, *P. composti*, *P. putida* KT2440 and DGC at the end of the batch phase.

5

6 **Figure S3:** visual inspection the auto-aggregation ability of different strains: *P. putida* DGC (KT2440
7 background), *P. putida* $\Delta lapA$ (KT2440 background), *P. putida* KT2440 and *P. composti*. Auto-
8 aggregation ability has been determined over a period of 24h at room temperature based on their
9 sedimentation characteristics

1
2
3
4
5
6
7
8
9
10

Movie S1: evolution of PI-positive and aggregates subpopulations, based on individual dotplots with forward scatter (FSC) as x-axis and PI fluorescence as y-axis. These dotplots were acquired at an interval of 12 minutes by on-line flow cytometry and were further assembled into a single .avi movie file