Molecular effects of mannanase-hydrolyzed coprameal to intestinal immunity in the Pacific white shrimp (*Litopenaeus vannamei*)

Wanilada Rungrasamee^{1*}, Sopacha Arayamethakorn¹, Nitsara Karoonuthaisiri¹, Shih-Chu Chen², Eric Chang³, Samu Chan⁴, Yasuhiko Yoshida⁵, Motohiro Maebuchi⁵, Masahisa Ibuki⁶

¹Microarray Research Team, National Center for Genetic Engineering and Biotechnology, Pathum Thani, Khlong Luang, Thailand 12120

²Department of Veterinary Medicine and Graduate Institute of Animal Vaccine Technology, National Pingtung University of Science and Technology, 1 Shuehfu Rd, Neipu, Pingtung, Taiwan, ROC

³Terminalia Garden Aquatic Development, No.73-6 Guanghe Rd., Mituo Dist., Kaohsiung City 827, Taiwan

⁴Fukar International Co., Ltd., No.112, Sec2 Chang-an E. Rd., Zhongshan dist., Taipei, Taiwan

⁵Fuji Oil Co., Ltd., 1 Sumiyoshi-Cho, Izumisano-Shi, Osaka, 598-8540 Japan

⁶Japan Nutrition Co., Ltd., 1-1-1 Minamiaoyama, Minatoku, Tokyo, 107-0062 Japan

*Correspondence to WR

Email: wanilada.run@biotec.or.th

Running title: Shrimp immunity enhanced by mannanase-hydrolyzed copra meal

Abstract

To mitigate disease outbreak, an alternative approach through enhancing shrimp immunity was explored. Mannan oligosaccharides (MOS) have been previously reported to enhance shrimp immune system. Here, coprameal samples were digested with mannanase to yield MOS, namely, mannanase-hydrolyzed coprameal (MCM) and feasibility of MCM as shrimp immunostimulant in grow-out ponds was determined. Pacific white shrimp (*Litopenaeus vannamei*) were fed with the commercial diet containing 1% MCM as the MCM-supplemented group and compared to the non-MCM control diet. There was no significant difference in survival rates between the MCM-supplemented and the control groups throughout the 4-month-period of the trial ($p > 0.05$). Gene expression analysis in shrimp intestines revealed that the transcript levels of antimicrobial peptides (*anti-lipopolysaccharide factor isoform 1 (alf1)*, *penaeidin (pen3a)* and *crustin (crus)*) and lysozymes (*lyz*) were not significantly different in the MCM-supplemented group. Meanwhile, *C-type lectin* and *toll-like receptor* transcript levels, whose gene products play roles as pattern recognition proteins, were significantly higher in a group fed with MCM for 2- and 4-month periods than those of the control group ($p < 0.05$). The increased transcript levels of *C-type lectin* and *toll-like receptor* provide evidence for potential implementation of MCM as feed supplement to modulate shrimp immune system.

Keywords: copra meal; mannan oligosaccharides; Pacific white shrimp; immunostimulant

56 Introduction

57 Aquaculture production play important role to provide food sources for
58 increasing global population. The global demand for shrimp and prawn production is
59 approximately 70% of crustaceans (FAO 2018). This results in the rapid growth of
60 cultivated shrimp production, in particularly, an expanding of Pacific white shrimp
61 farming throughout many Asian countries. However, the shrimp industry continues to
62 suffer from production losses due to the higher disease outbreak incidents. To mitigate
63 shrimp diseases, much attention has been paid to development of feed additives with
64 potential to enhance growth and immune performance as a mean to prevent disease
65 outbreaks in aquaculture.

66 In mammals, prebiotics, which are non-digestible carbohydrates, are shown
67 to selectively stimulate growth of beneficial bacteria in host digestive systems, and
68 consequently, providing health benefits to their host animals (Al-Sheraji et al. 2013;
69 Davani-Davari et al. 2019; Manning & Gibson 2004). Recently, prebiotics as
70 immunostimulants have been applied in aquaculture as a promising approach for growth
71 and immune enhancement (Hasan et al. 2019; Luna-González et al. 2012; Sang et al.
72 2010; Staykov et al. 2007). Among those, mannan oligosaccharides (MOS), derived
73 from yeast cell wall (*Saccharomyces cerevisiae*), have been shown to improve growth
74 performance and enhance immune system of many aquatic animals (Dimitroglou et al.
75 2010; Salem et al. 2015; Zhang et al. 2012; Zhang et al. 2020). In crustaceans,
76 applications of MOS as feed additives has been reported to significantly improve the
77 survival rate of crayfish (*Cherax tenuimanus*) (Sang et al. 2010), and enhance growth
78 rates in tropic spiny lobster (*Panulirus ornatus*) (Do Huu & Jones 2014) and black tiger
79 shrimp (*Penaeus monodon*) (Sang et al. 2014; Sang & Fotedar 2010) and freshwater
80 crayfish (*Astacus leptodactylus*) (Mazlum et al. 2011). The dietary MOS from yeast can
81 significantly increase intestinal microvilli length, growth and percent survival rates in
82 the Pacific white shrimp (Gainza & Romero 2020; Zhang et al. 2012)

83 While MOS used in animal feed additives are mostly derived from yeast, MOS
84 can also be obtained from other natural sources with high mannan sugar contents such
85 as copra meal (Hossain et al. 1996; Kraikaew et al. 2020; Pangsri et al. 2015). Copra
86 meal is a by-product from the coconut milk industry, and it is an alternative source for
87 MOS in a form of β -1, 4 mannoooligosaccharides (Ariandi et al. 2015). The potential

88 applications of MOS from copra meal as immunostimulants have been explored in
89 various animals such as broilers (Prayoonthien et al. 2017; Sundu et al. 2012) and the
90 Pacific white shrimp (Chen et al. 2020; Li et al. 2018; Rungrassamee et al. 2014).
91 Particularly, MOS from copra meal has been reported to improve immune levels and
92 survival rates in the Pacific white shrimp juvenile under a pathogen exposure to *Vibrio*
93 *harveyi* (Rungrassamee et al. 2014), providing a piece of evidence for MOS as a
94 promising shrimp feed additive. Here, we further explored feasibility of MOS through
95 mannanase-hydrolysis of copra meal (MCM) as feed additives in the grow-out pond
96 systems and determined molecular effects on immune gene expression. These findings
97 will lead to further development of MCM (MOS from copra meal) as effective
98 prebiotics for shrimp to maintain their intestinal immunity in the grow-out phase.
99

100 **Materials and methods**

101 *Preparation of mannanase-hydrolyzed coprameal (MCM)*

102 The copra meals were hydrolyzed using mannanase enzyme to yield mannanase-
103 hydrolyzed coprameal (MCM). Briefly, the defatted copra meals were incubated with
104 0.2-0.6% mannanase solution (15,000unit/g minimum, Shin-Nihon Chemical CO., Ltd.,
105 Aichi, Japan) at 40-70 °C for 18-30 h, and then, dried 2-6 h to obtain MCM product.
106 MCM content analysis was carried out using following methods. Mannobiose was
107 determined by high-performance anion-exchange chromatography coupled with a
108 pulsed amperometric detection (HPAEC-PAD) system, and constituent sugars were
109 analyzed by sulfuric hydrolysis. Ashes were determined by a dry-ashing procedure
110 at 55 ± 50°C for 2 h and moisture was determined by oven drying method at 105±1 °C
111 for 4 h. Fat was analyzed by a diethyl ether extraction method (AOAC Method 920.39).
112 Protein content was determined with a Kjeldahl method (Lynch & Barbano 1999).

113

114 *Animal facility and feeding trials*

115 A group of 16-day-old Pacific white shrimp postlarvae was cultivated in the grow-out
116 ponds at the animal facility (Terminalia Garden Aquatic development, Taiwan) (Fig.1).
117 Shrimp were reared in brackish water with 0.07-0.15% salt concentrations and divided
118 into two groups: a control feed with commercial feed pellets (a control diet) and a
119 MCM-supplemented group (a test diet) (Table 1). The test diet was formulated to
120 contain 1% of MCM in the commercial feed pellets. Shrimp were fed under these diets
121 and intestine samples ($N_{\text{pooled}} = 6$ with 5 replicates) were aseptically collected at 2- and
122 4- month of the feeding trial period. All tissue samples were stored in RNeasy[®] RNA
123 Stabilization Solution (Ambion, USA) to preserve tissue integrity for RNA extraction
124 for gene expression analysis. Water quality in ponds was measured every other day for
125 temperature, pH, and dissolved oxygen and weekly for ammonia–nitrogen, nitrite–
126 nitrogen, and alkalinity levels and maintained at the standard ranges for rearing Pacific
127 white shrimp (Rungrassamee et al. 2013).

128

129 *Total RNA extraction and cDNA synthesis*

130 Intestine tissues were homogenized in TRI REAGENT[®] (Molecular Research
131 Center, USA) for RNA extraction according to supplier's instruction. RNA pellets were

132 washed twice with 500 μ L of 75% ice-cold ethanol, air-dried for 5 min, and dissolved in
133 50 μ L diethylpyrocarbonate (DEPC)-treated water. To remove potential genomic DNA
134 contamination, all RNA samples were treated with DNaseI enzyme (RQ1 RNase-free
135 DNaseI, Promega, USA) for 60 min at 37 °C. The DNaseI treated-RNA samples were
136 purified via phenol:chloroform extraction and precipitated with 1/10 volume of 3M
137 sodium acetate and 1 volume of isopropanol as previously done (Rungrassamee et al.
138 2010). All treated-RNA samples were confirmed to be free from genomic DNA via
139 PCR reaction. RNA concentration was quantified with a Nanodrop ND-8000
140 spectrophotometer (NanoDrop, UK). An ImProm-II™ Reverse Transcriptase System kit
141 (Promega, USA) was used to synthesize the first strand cDNA using 1.5 μ g of the
142 DNA-free RNA sample as a template and each reaction was carried out according to the
143 supplier's protocol. The concentration of cDNA was measured by a Nanodrop ND-8000
144 spectrophotometer.

145

146 *Realtime PCR analysis of immune related genes transcript levels in response to MCM* 147 *supplement diet*

148 To determine MCM effects on expression levels of host immune genes, cDNA
149 from each shrimp group after 2- and 4-month of the feeding trial periods were used as
150 templates in realtime PCR analysis using a CFX96™ realtime system (Bio-rad
151 Laboratories, USA). The transcript levels of immune-related genes chosen were anti-
152 lipopolysaccharide factor1 (*alf1*), crustin (*crus*), c-type lectin (*c-lec*), lysozyme (*lyz*),
153 penaeidin3 (*pen3a*) and Toll1 (*Toll1*) and the elongation factor-1 alpha (*EF1a*) was used
154 as an internal control. Each realtime PCR reaction contained IQ™ SYBR Green
155 Supermix (Bio-Rad, USA), gene-specific primer pair (1.25 μ M, Table 2) and cDNA
156 (100 ng). The realtime PCR cycling parameters used were 95 °C for 3 min, 40 cycles of
157 at 95 °C for 30 sec, 57 °C for 20 sec and 72 °C for 30 sec. The fluorescent signal
158 intensities were recorded at the end of each cycle. Melting curve analysis was
159 performed from 55°C to 95 °C with continuous fluorescence measurement at every 0.5
160 °C increment. Relative abundance of the target immune genes in intestines of the
161 Pacific white shrimp was determined by the $\Delta\Delta$ ct method (Livak & Schmittgen 2001).
162 The relative fold change for each target gene was compared between the MCM-
163 supplement to the control groups at the same time point.

164

165 *Statistical analysis*

166 Statistical analysis was conducted using SPSS of Windows version 15.0 to
167 perform Student's t test analysis for significant differences in shrimp weights or gene
168 expression analysis (Landau & Everitt 2004).

169

170 **Results and discussion**

171 *MCM component analysis*

172 In this work, MOS supplement was derived in a form of mannanase-hydrolyzed
173 coprameal (MCM). MCM yield was ~95% of the copra meal. The content analysis of
174 MCM revealed high composition of carbohydrates (55.0%), followed by crude proteins
175 (22.9%), fats (9.3%), moisture (7.0%) and ashes (5.8%) (Table 3). The free sugar
176 contents in MCM were mainly mannobiose sugars (10.3%) and others were glucose,
177 mannose, fructose, sucrose and mannotriose. Among those sugars, mannobiose has been
178 reported for their important role as an immune modulator (Patel & Goyal 2012; Tiwari
179 et al. 2020). For instance, mannobiose is able to activate innate immune response in
180 chicken under *in vivo* and *in vitro* studies (Agunos et al. 2007; Duarte et al. 2014; Ibuki
181 et al. 2010; Ibuki et al. 2011). This suggests that MCM supplementation in this study
182 could potentially be used to enhance disease protection in animals.

183

184 *Expression analysis of the immune related in response to MCM supplementary diet*

185 Since shrimp digestive system, especially its intestine can be prone to pathogen
186 invasion (Aguirre Guzman et al. 2010; Soonthornchai et al. 2015), it is important to
187 determine approaches to modulate intestinal immune level. Hence, we selected genes
188 encoding the antimicrobial peptides (*alf1*, *pen3a* and *crus*), lysozyme (*lyz*), C-type
189 lectin (*c-lectin*), and Toll-like receptor protein (*Toll1*) as our genes of interest due to
190 their important roles in digestive tract of the shrimp as part of host defense mechanisms
191 against pathogens (Tassanakajon et al. 2013). To evaluate molecular effects of MCM
192 supplement on host immune system, the relative expression levels of the
193 aforementioned transcripts were compared in shrimp intestines at 2-month and 4-month
194 after feeding with the MCM-supplement diet to those fed with the non-MCM as a
195 control diet (Fig. 2). Our result showed that the MCM as dietary supplement did not

196 affect the transcript levels of *alfl*, *pen3a* and *crus*. Additionally, the group fed with
 197 MCM showed a significant decreased in transcript level of *lyz* at the 2-month of the
 198 MCM-feeding period ($p < 0.05$), however the transcript level of *lyz* was not
 199 significantly different to the non-MCM diet at 4-month-period. This suggests that MCM
 200 did not have a direct effect on modulating antimicrobial peptides including lysozyme. In
 201 contrast, the *c-lectin* and *Toll1* expression levels were significantly higher than the
 202 control diet by ~2-fold when given MCM supplement diet at 2- and 4-month periods (p
 203 < 0.05). This suggests that MCM supplement specifically increases expression levels of
 204 *c-lectin* and *Toll1*, whose gene products play role as pattern recognition proteins. The
 205 encoded C-type lectin is a member of lectin family proteins that has been reported to
 206 bind to carbohydrate in a calcium-dependent manner (Weis et al. 1998). It plays
 207 important role in pathogen recognition by detecting conserved pathogen cell wall
 208 components of Gram-negative bacteria and further induces immune response to
 209 invaders (Bi et al. 2020; Medzhitov & Janeway 2002; Wang & Wang 2013). Lectins
 210 show strong bacterial-agglutination and opsonic activity, which facilitate phagocytosis
 211 (Luo et al. 2006). In the Chinese white shrimp, C-type lectin has been demonstrated *in*
 212 *vivo* to mediate in *V. anguillarum* clearance (Wang et al. 2009). Thus, the induction of
 213 *C-lectin* could provide a local immune response and protection in Pacific white shrimp
 214 intestine against bacterial pathogen invasion. Similarly, *Toll1*, encodes for Toll-like
 215 receptor protein, was also significantly increased at 2- and 4-month of the MCM
 216 supplemented group in comparison to the control group. Toll-like receptor proteins have
 217 been reported in mammals to play roles in viral pathogen detection and mediate
 218 responses to those viruses (Takeuchi & Akira 2009). Interestingly, the function of *Toll1*
 219 in the Pacific white shrimp has not yet been linked to viral diseases protection in shrimp
 220 (Labreuche et al. 2009), but it has been shown to involve in responses to bacterial
 221 pathogen (Liu et al. 2016). This suggests that MCM is a promising candidate to increase
 222 disease resistance to bacterial pathogens in Pacific white shrimp farming. Further
 223 investigation on optimal dosages and formulation for the MCM supplement should be
 224 conducted for future application.

225 Here, we report that the feed supplement containing mannanase-hydrolyzed
 226 coprameal (MCM) was able to induce transcript levels of pathogen recognition proteins
 227 in the Pacific white shrimp. The significant increase was observable on Month 2 of the

228 feeding trial and the level remained similar on Month 4. Although future shrimp
229 experiments fed with a range of MCM concentration might be necessary to fully
230 elucidate beneficial effects of MCM. Our results provide a promising evidence for
231 further implementation of MCM as feed supplements or immunostimulants to enhance
232 shrimp health. Additionally, a comparison of shrimp survival rates under pathogen
233 challenge between the MCM supplement shrimp group to non-MCM supplement
234 control will strengthen the potential application of MCM as shrimp immunostimulant.

235

236 **Conflict of interest**

237 Yasuhiko Yoshida, Motohiro Maebuchi and Masahisa Ibuki are employees of Fuji Oil
238 Co., Ltd. or Japan Nutrition Co., Ltd., whom kindly supplied mannanase-hydrolysis of
239 copra meal (MCM) used in this study. They had no roles in experimental design and
240 data analysis.

241

242 **Acknowledgements**

243 We thank Dr. Wonnop Visessanguan from National Center for Genetic Engineering and
244 Biotechnology (BIOTEC, Thailand) for advice and insights on evaluation of shrimp
245 feed additive. This project was supported the Thailand Research Fund (RSA5780022).

246
247

References

- Aguirre Guzman G, Sánchez-Martínez J, Pérez-Castañeda R, Palacios A, Trujillo-Rodríguez T, and Ivan N. 2010. Pathogenicity and infection route of *Vibrio parahaemolyticus* in American white shrimp, *Litopenaeus vannamei*. *Journal of The World Aquaculture Society* 41:464-470. 10.1111/j.1749-7345.2010.00388.x
- Agunos A, Ibuki M, Yokomizo F, and Mine Y. 2007. Effect of dietary beta1-4 mannanose in the prevention of *Salmonella enteritidis* infection in broilers. *British Poultry Science* 48:331-341. 10.1080/00071660701370442
- Al-Sheraji SH, Ismail A, Manap MY, Mustafa S, Yusof RM, and Hassan FA. 2013. Prebiotics as functional foods: A review. *Journal of Functional Foods* 5:1542-1553. <https://doi.org/10.1016/j.jff.2013.08.009>
- Ariandi, Yopi, and Meryandini A. 2015. Enzymatic hydrolysis of copra meal by mannanase from *Streptomyces* sp. BF3.1 for the production of mannooligosaccharides. *HAYATI Journal of Biosciences* 22:79-86. <https://doi.org/10.4308/hjb.22.2.79>
- Bi J, Ning M, Xie X, Fan W, Huang Y, Gu W, Wang W, Wang L, and Meng Q. 2020. A typical C-type lectin, perlucin-like protein, is involved in the innate immune defense of whiteleg shrimp *Litopenaeus vannamei*. *Fish & Shellfish Immunology* 103:293-301. <https://doi.org/10.1016/j.fsi.2020.05.046>
- Chen M, Chen X-Q, Tian L-X, Liu Y-J, and Niu J. 2020. Beneficial impacts on growth, intestinal health, immune responses and ammonia resistance of pacific white shrimp (*Litopenaeus vannamei*) fed dietary synbiotic (mannan oligosaccharide and *Bacillus licheniformis*). *Aquaculture Reports* 17:100408. <https://doi.org/10.1016/j.aqrep.2020.100408>
- Davani-Davari D, Negahdaripour M, Karimzadeh I, Seifan M, Mohkam M, Masoumi SJ, Berenjian A, and Ghasemi Y. 2019. Prebiotics: definition, types, sources, mechanisms, and clinical applications. *Foods (Basel, Switzerland)* 8:92. 10.3390/foods8030092
- Dimitroglou A, Davies S, Sweetman J, Divanach P, and Chatzifotis S. 2010. Dietary supplementation of mannan oligosaccharide on white sea bream (*Diplodus sargus* L.) larvae: Effects on development, gut morphology and salinity tolerance. *Aquaculture Research* 41:e245-e251. 10.1111/j.1365-2109.2010.02513.x
- Do Huu H, and Jones CM. 2014. Effects of dietary mannan oligosaccharide supplementation on juvenile spiny lobster *Panulirus homarus* (Palinuridae). *Aquaculture* 432:258-264. <https://doi.org/10.1016/j.aquaculture.2014.05.013>
- Duarte K, Ibuki M, Fukui K, Kato M, Santos E, and Junqueira O. 2014. Effect of hydrolyzed copra meal separately or in combination with *Bacillus cereus* var. *toyoi* on growth performance of broiler chickens. *Acta Scientiarum - Animal Sciences* 36:373-377. 10.4025/actascianimsci.v36i4.22829
- FAO. 2018. *The state of world fisheries and aquaculture 2018 - meeting the sustainable development goals*. Rome: FAO.
- Gainza O, and Romero J. 2020. Effect of mannan oligosaccharides on the microbiota and productivity parameters of *Litopenaeus vannamei* shrimp under intensive cultivation in Ecuador. *Scientific Reports* 10:2719. 10.1038/s41598-020-59587-y

295 Hasan MT, Jang WJ, Lee J, Lee B-J, Lim S, Kang W, Han H-S, and Kong I-S. 2019.
 296 Effects of immunostimulants, prebiotics, probiotics, synbiotics, and potentially
 297 immunoreactive feed additives on Olive flounder (*Paralichthys olivaceus*): A
 298 Review. *Reviews in Fisheries Science & Aquaculture* 27.
 299 10.1080/23308249.2019.1622510
 300 Hossain MZ, Abe J-i, and Hizukuri S. 1996. Multiple forms of β -mannanase from
 301 *Bacillus* sp. KK01. *Enzyme and Microbial Technology* 18:95-98.
 302 [https://doi.org/10.1016/0141-0229\(95\)00071-2](https://doi.org/10.1016/0141-0229(95)00071-2)
 303 Ibuki M, Kovacs-Nolan J, Fukui K, Kanatani H, and Mine Y. 2010. Analysis of gut
 304 immune-modulating activity of beta-1,4-mannobiose using microarray and real-
 305 time reverse transcription polymerase chain reaction. *Poult Sci* 89:1894-1904.
 306 10.3382/ps.2010-00791
 307 Ibuki M, Kovacs-Nolan J, Fukui K, Kanatani H, and Mine Y. 2011. β 1-4 mannanose
 308 enhances Salmonella-killing activity and activates innate immune responses in
 309 chicken macrophages. *Vet Immunol Immunopathol* 139:289-295.
 310 10.1016/j.vetimm.2010.10.011
 311 Kraikaew J, Morakul S, and Keawsompong S. 2020. Nutritional improvement of copra
 312 meal using mannanase and *Saccharomyces cerevisiae*. *3 Biotech* 10:274.
 313 10.1007/s13205-020-02271-9
 314 Labreuche Y, O'Leary NA, de la Vega E, Veloso A, Gross PS, Chapman RW, Browdy
 315 CL, and Warr GW. 2009. Lack of evidence for *Litopenaeus vannamei* Toll
 316 receptor (IToll) involvement in activation of sequence-independent antiviral
 317 immunity in shrimp. *Developmental & Comparative Immunology* 33:806-810.
 318 10.1016/j.dci.2009.02.005
 319 Landau S, and Everitt B. 2004. *A handbook of statistical analyses using SPSS*. Boca
 320 Raton, Florida: Chapman and Hall/ CRC Press LLC.
 321 Li Y, Liu H, Dai X, Li J, and Ding F. 2018. Effects of dietary inulin and mannan
 322 oligosaccharide on immune related genes expression and disease resistance of
 323 Pacific white shrimp, *Litopenaeus vannamei*. *Fish & Shellfish Immunology*
 324 76:78-92. 10.1016/j.fsi.2018.02.034
 325 Liu Y, Song L, Sun Y, Liu T, Hou F, and Liu X. 2016. Comparison of immune response
 326 in Pacific white shrimp, *Litopenaeus vannamei*, after knock down of Toll and
 327 IMD gene *in vivo*. *Developmental & Comparative Immunology* 60:41-52.
 328 10.1016/j.dci.2016.02.004
 329 Livak KJ, and Schmittgen TD. 2001. Analysis of relative gene expression data using
 330 real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*
 331 25:402-408. 10.1006/meth.2001.1262
 332 Luna-González A, Almaraz-Salas JC, Fierro-Coronado JA, Flores-Miranda MdC,
 333 González-Ocampo HA, and Peraza-Gómez V. 2012. The prebiotic inulin
 334 increases the phenoloxidase activity and reduces the prevalence of WSSV in
 335 whiteleg shrimp (*Litopenaeus vannamei*) cultured under laboratory conditions.
 336 *Aquaculture* 362-363:28-32. <https://doi.org/10.1016/j.aquaculture.2012.07.022>
 337 Luo T, Yang H, Li F, Zhang X, and Xu X. 2006. Purification, characterization and
 338 cDNA cloning of a novel lipopolysaccharide-binding lectin from the shrimp
 339 *Penaeus monodon*. *Developmental & Comparative Immunology* 30:607-617.
 340 10.1016/j.dci.2005.10.004

341 Lynch JM, and Barbano DM. 1999. Kjeldahl nitrogen analysis as a reference method
342 for protein determination in dairy products. *Journal of AOAC International*
343 82:1389-1398.

344 Manning TS, and Gibson GR. 2004. Microbial-gut interactions in health and disease.
345 Prebiotics. *Best Practice & Research Clinical Gastroenterology* 18:287-298.
346 10.1016/j.bpg.2003.10.008

347 Mazlum Y, Yılmaz E, Genç MA, and Guner O. 2011. A preliminary study on the use of
348 mannan oligosaccharides (MOS) in freshwater crayfish, *Astacus leptodactylus*
349 Eschscholtz, 1823 juvenile diets. *Aquaculture International* 19:111-119.
350 10.1007/s10499-010-9345-4

351 Medzhitov R, and Janeway CA, Jr. 2002. Decoding the patterns of self and nonself by
352 the innate immune system. *Science* 296:298-300. 10.1126/science.1068883

353 Pangsi P, Piwpankaew Y, Ingkakul A, Nitisinprasert S, and Keawsompong S. 2015.
354 Characterization of mannanase from *Bacillus circulans* NT 6.7 and its
355 application in mannooligosaccharides preparation as prebiotic. *Springerplus*
356 4:771. 10.1186/s40064-015-1565-7

357 Patel S, and Goyal A. 2012. The current trends and future perspectives of prebiotics
358 research: a review. *3 Biotech* 2:115-125. 10.1007/s13205-012-0044-x

359 Prayoonthien P, Nitisinprasert S, and Keawsompong S. 2017. *In vitro* fermentation of
360 copra meal hydrolysate by chicken microbiota. *3 Biotech* 8:41. 10.1007/s13205-
361 017-1058-1

362 Rungrassamee W, Kingcha Y, Srimarut Y, Maibunkaew S, Karoonuthaisiri N, and
363 Visessanguan W. 2014. Mannooligosaccharides from copra meal improves
364 survival of the Pacific white shrimp (*Litopenaeus vannamei*) after exposure to
365 *Vibrio harveyi*. *Aquaculture* 434:403-410.
366 <https://doi.org/10.1016/j.aquaculture.2014.08.032>

367 Rungrassamee W, Leelatanawit R, Jiravanichpaisal P, Klinbunga S, and Karoonuthaisiri
368 N. 2010. Expression and distribution of three heat shock protein genes under
369 heat shock stress and under exposure to *Vibrio harveyi* in *Penaeus monodon*.
370 *Developmental & Comparative Immunology* 34:1082-1089. S0145-
371 305X(10)00125-4 [pii] 10.1016/j.dci.2010.05.012 [doi]

372 Rungrassamee W, Maibunkaew S, Karoonuthaisiri N, and Jiravanichpaisal P. 2013.
373 Application of bacterial lipopolysaccharide to improve survival of the black
374 tiger shrimp after *Vibrio harveyi* exposure. *Developmental & Comparative*
375 *Immunology* 41:257-262. <http://dx.doi.org/10.1016/j.dci.2013.05.021>

376 Salem M, Gaber M, Zaki M, and Nour A. 2015. Effects of dietary mannan
377 oligosaccharides on growth, body composition and intestine of the sea bass
378 (*Dicentrarchus labrax* L.). *Aquaculture Research* 47. 10.1111/are.12801

379 Sang H, Fotedar R, and Filer K. 2010. Effects of dietary mannan oligosaccharide on the
380 survival, growth, immunity and digestive enzyme activity of freshwater crayfish,
381 *Cherax destructor* Clark (1936). *Aquaculture Nutrition* 17:e629-e635.
382 10.1111/j.1365-2095.2010.00812.x

383 Sang H, Kien NT, and Nguyen Thi Thanh T. 2014. Effects of dietary mannan
384 oligosaccharide on growth, survival, physiological, immunological and gut
385 morphological conditions of black tiger prawn (*Penaeus monodon* Fabricius
386 1798). *Aquaculture Nutrition* 20. 10.1111/anu.12083

387 Sang HM, and Fotedar R. 2010. Effects of mannan oligosaccharide dietary
388 supplementation on performances of the tropical spiny lobsters juvenile

389 (Panulirus ornatus, Fabricius 1798). *Fish & Shellfish Immunology* 28:483-489.
390 10.1016/j.fsi.2009.12.011
391 Soonthornchai W, Chaiyapechara S, Jarayabhand P, Söderhäll K, and Jiravanichpaisal
392 P. 2015. Interaction of *Vibrio spp.* with the inner surface of the digestive tract of
393 *Penaeus monodon*. *PLoS One* 10:e0135783. 10.1371/journal.pone.0135783
394 Staykov Y, Spring P, Denev S, and Sweetman J. 2007. Effect of a manna
395 oligosaccharide on the growth performance and immune status of rainbow trout
396 (*Oncorhynchus mykiss*). *Aquaculture International* 15:153–161 15.
397 10.1007/s10499-007-9096-z
398 Sundu B, Hutta U, and Chaudhry A. 2012. Potential use of beta mannan from copra
399 meal as a feed additive for broilers. *World's Poultry Science Journal* 68:707-
400 716. 10.1017/S0043933912000839
401 Takeuchi O, and Akira S. 2009. Innate immunity to virus infection. *Immunological*
402 *Reviews* 227:75-86. 10.1111/j.1600-065X.2008.00737.x
403 Tassanakajon A, Somboonwiwat K, Supungul P, and Tang S. 2013. Discovery of
404 immune molecules and their crucial functions in shrimp immunity. *Fish &*
405 *Shellfish Immunology* 34:954-967. <https://doi.org/10.1016/j.fsi.2012.09.021>
406 Tiwari UP, Fleming SA, Abdul Rasheed MS, Jha R, and Dilger RN. 2020. The role of
407 oligosaccharides and polysaccharides of xylan and mannan in gut health of
408 monogastric animals. *Journal of nutritional science* 9:e21-e21.
409 10.1017/jns.2020.14
410 Wang XW, and Wang JX. 2013. Diversity and multiple functions of lectins in shrimp
411 immunity. *Developmental & Comparative Immunology* 39:27-38.
412 10.1016/j.dci.2012.04.009
413 Wang XW, Zhang XW, Xu WT, Zhao XF, and Wang JX. 2009. A novel C-type lectin
414 (FcLec4) facilitates the clearance of *Vibrio anguillarum in vivo* in Chinese white
415 shrimp. *Developmental & Comparative Immunology* 33:1039-1047.
416 10.1016/j.dci.2009.05.004
417 Weis WI, Taylor ME, and Drickamer K. 1998. The C-type lectin superfamily in the
418 immune system. *Immunological Reviews* 163:19-34. 10.1111/j.1600-
419 065x.1998.tb01185.x
420 Zhang J, Liu Y, Tian L, Yang H, Liang G, and Xu D. 2012. Effects of dietary mannan
421 oligosaccharide on growth performance, gut morphology and stress tolerance of
422 juvenile Pacific white shrimp, *Litopenaeus vannamei*. *Fish & Shellfish*
423 *Immunology* 33:1027-1032. <https://doi.org/10.1016/j.fsi.2012.05.001>
424 Zhang P, Yang F, Hu J, Han D, Liu H, Jin J, Yang Y, Yi J, Zhu X, and Xie S. 2020.
425 Optimal form of yeast cell wall promotes growth, immunity and disease
426 resistance in gibel carp (*Carassius auratus gibelio*). *Aquaculture Reports*
427 18:100465. <https://doi.org/10.1016/j.aqrep.2020.100465>
428
429

Table 1. Composition of feed diets used in this study.

Raw material	% Feed composition	
	Control diet	MCM supplement diet
Wheat flour	35	35
Fish meal	20	20
Squid meal	10	10
Water	30	29
Vitamin and yeast mix	5	5
MCM	0	1

435 **Table 2.** Primer sequences used in realtime PCR analysis
436

Gene	GenBank Accession Number	Primer	Sequence (5'-3')
Anti-lipopolysaccharide factor1 (<i>alf1</i>)	EW713395	alf1_F alf1_R	AGGACCTCATCCCTTCGCTAGT GGACACCACATCCTGCCATTGA
Crustin (<i>crus</i>)	AF430074	crus_F crus_R	CCACAACCTGTTCCAACGACTACAA AAACCTGCGATCCGAAGAATGAGG
C-type lectin (<i>c-lec</i>)	JF834160	lec_F lec_R	CGCTGATGGCTCGGATGAGA AGGCTGAGTTCGGTGGCAATAG
Lysozyme (<i>lyz</i>)	AF425673	lyz_F lyz_R	GGAGTTCGAGTCGTCCTTCAACA GTAGTCGCTTCCGCACCAGTA
Penaedin3 (<i>pen3a</i>)	Y14926	pen3a_F pen3a_R	CGTGGTCTGCCTGGTCTTCTT CAAAGGTCTCACGAAGGGTGGT
Toll1 (<i>Toll1</i>)	DQ923424	Toll1_F Toll1_R	CGAGAGCGAGTTGGACGAGAAG CCTGTGGGTGTGGCATGATGTA
Elongation factor-1 alpha (<i>EF1a</i>)	GU136229	EF1a_F EF1a_R	CGTCGCTTCCGACTCGAAGAA TGGCAATCAAGCACAGGTGAGTAG

437
438
439

440 **Table 3.** Content analysis of mannanase-hydrolyzed coprameal (MCM).
441

		Component in MCM (%)
General composition	Crude protein	22.9
	Crude fat	9.3
	Ash	5.8
	Moisture	7.0
	Carbohydrate	55.0
Free sugar contents	Glucose	1.5
	Mannose	2.6
	Fructose	1.8
	Sucrose	4.6
	Mannobiose	10.3
	Mannotriose	2.1

442
443
444

Figure Legends

Figure 1. Experimental overview for evaluating effects of mannanase-hydrolysis of copra meal (MCM) as immunostimulant in Pacific white shrimp, *Litopenaeus vannamei* in a grow-out pond. The Pacific white shrimp postlarvae were divided into two groups: a control feed with commercial feed pellets and a 1% MCM-supplemented group. The intestine samples ($N_{\text{pooled}} = 6$ with 5 replicates) were collected at 2- and 4- month of the feeding trial period for gene expression analysis.

Figure 2. Relative gene expression analysis of immune-related genes in intestines of Pacific white shrimp after 2-month and 4-month feeding period with the mannanase-hydrolyzed coprameal-supplemented diet group (MCM). The fold changes of transcript of antimicrobial peptides (*alf1*, *pen3a* and *crus*), lysozyme (*lyz*), C-type lectin (*c-lectin*), and Toll-like receptor protein (*Toll1*) were determined by real-time PCR in relative to the control diet. An error bar represents standard deviation, which was calculated from five replicates for each sample. Asterisk indicates significant difference in fold-changes between groups fed with MCM and non-MCM containing diets ($p < 0.05$).

Fig 1

Fig. 2