The IL-10^{GFP} (VeRT-X) mouse strain is not suitable for the detection of IL-10 production by granulocytes during lung inflammation

Müge Özkan¹, Yusuf Cem Eskiocak², Gerhard Wingender^{2,3*}

¹ Department of Genome Sciences and Molecular Biotechnology, Izmir International Biomedicine and Genome Institute, Dokuz Eylul University, 35340 Balçova/Izmir, Turkey;

² Izmir Biomedicine and Genome Center (IBG), 35340 Balçova/Izmir, Turkey.

³ Department of Biomedicine and Health Technologies, Izmir International Biomedicine and Genome Institute, Dokuz Eylul University, 35340 Balçova/Izmir, Turkey.

* Correspondence: gerhard.wingender@ibg.edu.tr (G.W.)

Keywords:

Reporter mice; IL-10^{GFP}; lung inflammation; granulocytes

Abbreviations:

AAD: allergic airway disease; AM: alveolar macrophage; COPD: chronic obstructive pulmonary disease; DC: dendritic cell; HDM: house dust mite; ICCS: intracellular cytokine staining; ILC: innate lymphoid cell; IM: interstitial macrophages; GFP: green fluorescent protein; LPS: lipopolysaccharide; mLN: mediastinal lymph node.

Abstract:

The clear and unequivocal identification of immune effector functions is essential to understand immune responses. The cytokine IL-10 is a critical immune regulator and was shown, for example, to limit pathology during various lung diseases. However, the clear identification of IL-10-producing cells is challenging and, therefore, reporter mouse lines were developed to facilitate their detection. Several such reporter lines utilize GFP, including the IL-10^{GFP} (VeRT-X) reporter strain studied here. In line with previous reports, we found that this IL-10^{GFP} line faithfully reports on the IL-10 production of lymphoid cells. However, we show that the IL-10^{GFP} reporter is not suitable to analyse IL-10 production of myeloid cells during inflammations. During inflammation, the autofluorescence of myeloid cells increased to an extent that entirely masked the IL-10-specific GFP-signal. Our data illustrate a general and important technical caveat using GFP-reporter lines for the analysis of myeloid cells and suggest that previous reports on effector functions of myeloid cells using such GFP-based reporters might require re-evaluation.

Introduction:

Regulatory cytokines are important players during pulmonary inflammations and their right balance is critical to avoid e.g. hampering pathogen clearance, while still limiting immunopathology (1). The regulatory cytokine IL-10 is involved in various lung diseases, like asthma, allergic airway disease (AAD), chronic obstructive pulmonary disease (COPD), and pulmonary infections (1). IL-10 producing cells in the lung; FoxP3⁺ and FoxP3⁻ regulatory T cells (1) , CD8⁺ T cells (2), B cells, alveolar macrophages (AM) (3), interstitial macrophages (IM) (4), airway and alveolar epithelial cells (1) , and airway-associated dendritic cells (DCs) (1) were shown to limit disease pathology. However, due to the low percentage of IL-10⁺ cells within most cell populations and due to the low intensity of the flow cytometric staining of intracellular IL-10 the clear identification of IL-10 producing cells is challenging(5). Therefore, various reporter mouse lines were developed to avoid the need of intracellular cytokine staining to identify IL-10 producing cells(6). The IL-10^{GFP} (VeRT-X) strain studied here, expresses an (IRES)-enhanced green fluorescent protein (eGFP) fusion protein downstream of the exon 5 of the *il10* gene(7), and was reported to enable the identification of IL-10⁺ lymphoid (7), but also myeloid cells. Several myeloid cells possess an autofluorescence around 525 nm (8), which coincides with the emission maxima of GFP around 530 nm (9). However, the impact of this myeloid autofluorescence on the detection of GFP-reporter signals has not been fully clarified. Here, using an LPS-induced lung inflammation model, we demonstrate that the autofluorescence of myeloid cells conceals the IL10^{GFP}-specific signal. This was mainly due to a large increase in the myeloid autofluorescence during

60 the inflammation. These data demonstrate that not all GFP-reporter mouse strains are
61 suitable to analyse effector functions of myeloid cell during inflammation.

62

Results and Discussion:

Neutrophils were suggested to be the primary source of IL-10 in the lung following *Klebsiella pneumoniae* ST258 infection (10). Furthermore, long-term exposure to house-dust mite (HDM) extracts suppressed allergic airway diseases via the IL-10 production by presumably FoxP3⁺ T cells and alveolar macrophages (3). In both studies, the ‘shift’ in the GFP-signal in the experimental IL-10^{GFP} group was compared to either the uninfected IL-10^{GFP} group or the naïve C57BL/6 group as control. When we analysed the IL-10 production during lung inflammation, we noticed that the background signal in the GFP-channel (505 nm - 550 nm) of leukocytes from lung, spleen, or mediastinal lymph node (mLN) was comparable in non-inflamed controls of either C57BL/6 (WT) or IL-10^{GFP} mice (**Figure 1A**). Upon LPS-induced lung inflammation, the GFP-signal clearly increased, in particular in the lungs (**Figure 1B**). Nevertheless, the GFP-signal from WT and IL-10^{GFP} mouse-derived cells remained comparable for the percentage (**Figure 1A, B**) or the mean fluorescence intensity (MFI, **Figure 1C**), irrespective for the organ analysed and the inflammatory condition. These data suggest that the GFP-signal detected in the IL-10^{GFP}-derived cells was actually a background signal. To clarify which cell types could produce IL-10, we directly compared the IL-10 signal derived from the GFP-signal or from intracellular cytokine staining (ICCS) for CD4⁺ and CD8⁺ T cells, neutrophils, macrophages, Siglec-F⁺ cells (eosinophils, alveolar macrophages), monocytes, and ILCs. Cells from the lungs, spleens, and mLN of PBS (control) and LPS-challenged C57BL/6 and IL-10^{GFP} mice were analysed (**Supplementary figure 1**). Although intracellular IL-10 staining is challenging, we obtained a clear IL-10 signal with the commercial antibody (**Supplementary figure 2**). With this side-by-side comparison we found that the IL-10-

86 signal derived from GFP or ICCS correlated well for CD4⁺ (**Figure 2A**) and CD8⁺ T cells
87 (**Figure 2B**). These data indicate that the IL-10^{GFP} - signal faithfully reports on the IL-10
88 production of lymphoid cells. However, the GFP-signal from neutrophils (**Figure 2C**),
89 macrophages (**Figure 2D**), and Siglec-F⁺ (**Figure 2E**) cells was significantly higher than
90 the ICCS-derived signal. Neither the arithmetic subtraction of GFP-signals of control (non-
91 stimulated) C57BL/6 samples (data not shown) nor the staining with a secondary α GFP-
92 AF488-antibody (**Supplementary figure 3**) was able to improve the specificity of the IL-
93 10-signal. Interestingly, the GFP-signal of monocytes (**Figure 2F**) and ILCs (**Figure 2G**)
94 was substantially higher only in the lung tissues, indicating that the increase in the GFP-
95 channel autofluorescence is organ-specific for some cell types. An overlay of the GFP-
96 signals of CD4⁺ T cells, CD8⁺ T cells, neutrophils, macrophages, Siglec-F⁺ cells,
97 monocytes, and ILCs from control C57BL/6 lung (**Figure 3A**) and spleen (**Figure 3B**)
98 indicated that the background GFP-signal was mainly derive from granulocytes.
99 According to the background GFP-signal, cell types ranked as Siglec-F > macrophage >
100 neutrophil > monocyte > ILC. A similar increase in the GFP-channel autofluorescence
101 upon inflammation was noted when alveolar macrophages were analysed (**Figure 4**).
102 In conclusion, our data indicate that the IL-10^{GFP} (VeRT-X) reporter strain is not suitable
103 to analyse IL-10 production of myeloid cells during inflammations, due to the strong
104 increase in the autofluorescence which masks the IL-10-specific GFP-signal. Although
105 we only analysed lung, spleen, and mLN of the VeRT-X IL-10^{GFP} reporter strain, it appears
106 likely that this problem will also be relevant to other organs. Importantly, the strong
107 increase of the granulocytic autofluorescence in the GFP-channels was independent of
108 the strain analysed. Therefore, it is expected that also other fluorescent reporter lines that

utilize the GFP-channel would face similar problems at distinguishing the reporter-specific signal from the autofluorescence signal when analysis granulocytes. Our data indicate that granulocytes are the source of the background GFP-signal, in line with previous publications reporting high levels of autofluorescence (11), although other sources, like collagen deposition [10,11], cannot be excluded. This large increase in false-positive GFP-signals for myeloid cells during inflammations could have been missed previously due to a focus on lymphoid cells or due to a lack of the WT treated controls. Our data illustrates a general and important technical caveat using GFP-reporter lines for the analysis of myeloid cells and suggest that previous reports on effector functions of myeloid cells using such GFP-based reporters might require re-evaluation.

Material and Methods

Mice

C57BL/6J mice and IL-10^{GFP} (also known as Vert-X) B6(Cg)-II10^{tm1.1Karp}/J mice were originally obtained from Jackson Laboratory (Bar Harbor, USA). All mice were housed in the vivarium of the Izmir Biomedicine and Genome Center (IBG, Izmir, Turkey) in accordance with the respective institutional animal care committee guidelines. All mouse experiments were performed with prior approval by the institutional ethic committee ('Ethical Committee on Animal Experimentation'), in accordance with national laws and policies. All the methods were carried out in accordance with the approved guidelines and regulations.

Reagents and monoclonal antibodies

Monoclonal antibodies against the following mouse antigens were used in this study: CD3 ϵ (145.2C11), CD4 (RM4-5), CD8 α (53-6.7), CD11b (M1/70), CD11c (N418), CD45 (30-F11), CD45R/B220 (RA3-6B2), CD90.2 (30-H12), CD127 (A7R34), CD170/Siglec-F (E50-2440), F4/80 (BM8), IL-10 (JESS-16E3), Ly6C (HK1.4), Ly6G (1A8). Antibodies were purchased from BD Biosciences (San Diego, USA), BioLegend (San Diego, USA), eBioscience (San Diego, USA) or ThermoFisher Scientific (Carlsbad, USA). Antibodies were biotinylated or conjugated to Pacific Blue, eFluor 450, V450, Brilliant Violet 421, V500, Brilliant Violet 510, Brilliant Violet 570, eFluor 650, Brilliant Violet 650, Brilliant Violet 711, Brilliant Violet 785, Brilliant Violet 786, FITC, Alexa Fluor 488, PerCP-Cy5.5, PerCP-eFluor 710, PE, PE-CF594, PE-Cy7, APC, Alexa Fluor 647, eFluor 660, Alexa Fluor 700, APC-Cy7, APC-eFluor 780 or APC-Fire750. Anti-mouse CD16/32 antibody

(2.4G2) and Zombie UV Dead Cell Staining Kit were obtained from Tonbo Biosciences (San Diego, USA) and from BioLegend, respectively. Unconjugated mouse and rat IgG antibodies were purchase from Jackson ImmunoResearch (West Grove, USA).

LPS-induced lung inflammation

Lung inflammation was induced by three daily administrations (d0, d1, d2) with 10 µg LPS (#L6386, Sigma-Aldrich, St. Louis, USA) per mouse via pharyngeal/laryngeal aspiration. 16-18 hours after the last challenge, the mice were sacrificed and lungs, spleens, and mediastinal lymph nodes were collected.

Cell preparation

Lungs were removed and minced into smaller pieces in a 6-well plate (Greiner, Germany). The digestion mixture, composed of 1 mg/mL collagenase D and 0.1 mg/mL DNase I (both from Roche, Switzerland) in complete RPMI medium (Gibco, USA), was added to the samples and incubated for 45 min at 37°C on a lateral shaker. The lung samples were filtered through 100 µm mesh with PBS, washed twice, and the red blood cells were eliminated by ACK lysis buffer (Lonza, USA). Spleens and mediastinal lymph nodes were homogenized by filtering through a 76 µm mesh with ice cold PBS (Lonza), washed twice and red blood cells were eliminated by ACK lysis buffer (Lonza).

***In vitro* stimulation**

Lymphocytes were stimulated *in vitro* with PMA (50 ng/mL) and ionomycin (1 µg/mL) (both Sigma-Aldrich, St. Louis, MO) for four hours at 37°C in the presence of both Brefeldin A (GolgiPlug) and Monensin (GolgiStop) in complete RPMI medium (RPMI 1640 medium (Life Technologies); supplemented with 10% (v/v) fetal calf serum (FCS), 1% (v/v) Pen-Strep-Glutamine (10.000 U/ml penicillin, 10.000 µg/ml streptomycin, 29.2 mg/ml L-glutamine (Life Technologies)) and 50 µM β-mercaptoethanol (Sigma)). As GolgiPlug and GolgiStop (both BD Biosciences) were used together, half the amount recommended by the manufacturer were used.

Flow Cytometry

Flow cytometry were performed as described (14). In brief, for staining of cell surface molecules, cells were suspended in staining buffer (PBS, 1% BSA, 0.01% NaN₃) and stained with fluorochrome-conjugated antibodies (0.1-1 µg/10⁶ cells, or according to the manufacturer's recommendations) for 15 min in a total volume of 50 µl at 4°C for 30 min. FcεR-blocking antibody αCD16/32 (2.4G2) and unconjugated rat and mouse IgG (Jackson ImmunoResearch) were added to prevent non-specific binding. If biotin-conjugated antibodies were used, cell bound antibodies were detected with streptavidin conjugates (1:200, or according to the manufacturer's recommendations) in a second incubation step. Dead cells were labelled with commercially available Zombie UV Dead Cell Staining Kit (BioLegend). For the analysis of intracellular cytokines, cells were fixed and permeabilized with Cytofix/Cytoperm (BD Biosciences) for 10 min at 37°C. Cells were washed twice and incubated overnight at 4°C with the fluorochrome-conjugated antibodies and unconjugated rat and mouse IgG in Perm/Wash solution (BD

Biosciences), which was followed by additional 5 min incubation in Perm/Wash solution without antibodies. For the analysis of the-GFP signal, the cells were fixed with freshly prepared 2% formaldehyde solution for 40 minutes on ice as described (15). This fixation allows the detection of the GFP-signal together with other intracellular proteins (15). Then, the cells were permeabilized using the BD Perm/Wash solution. Cells were washed twice and incubated overnight at 4°C with the fluorochrome-conjugated antibodies and unconjugated rat and mouse IgG in Perm/Wash solution, which was followed by additional 5 min incubation in Perm/Wash solution without antibodies. Cells were analyzed with FACSCanto or LSR-Fortessa (BD Biosciences) and data were processed with CellQuest Pro (BD Biosciences) or Flow Jo (Tree Star) software.

Statistical analysis

Data are presented as mean ± standard error of the mean (SEM). The statistical analysis was performed with GraphPad Prism 7.0 software (GraphPad Software, San Diego, USA). One-way ANOVA followed by Holm-Sidak posthoc test are used to compare p-values regarded as *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001.

Authorship

Conceived and designed the experiments: MO, GW. Performed the experiments and analyzed the data: MO, YCE. Wrote the paper: MO, GW. All authors reviewed and approved the manuscript.

Acknowledgments

This work was funded by grants from the Scientific and Technological Research Council of Turkey (TUBITAK, www.tubitak.gov.tr; #116Z272, GW), the Dokuz Eylul University (Izmir, Turkey; www.deu.edu.tr; #2020.KB.SAG.031, GW), and the European Molecular Biology Organization (EMBO, www.embo.org; IG3073; GW). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of Interest

The authors declare no competing financial interests.

References

1. Branchett WJ, Lloyd CM. Regulatory cytokine function in the respiratory tract. *Mucosal Immunol* [Internet]. 2019 Mar 15;12(3):589–600. Available from: <http://dx.doi.org/10.1038/s41385-019-0158-0>
2. Sun J, Madan R, Karp CL, Braciale TJ. Effector T cells control lung inflammation during acute influenza virus infection by producing IL-10. *Nat Med*. 2009;15(3):277–84.
3. Bracken SJ, Adami AJ, Szczepanek SM, Ehsan M, Natarajan P, Guernsey LA, et al. Long-Term Exposure to House Dust Mite Leads to the Suppression of Allergic Airway Disease Despite Persistent Lung Inflammation. *Int Arch Allergy Immunol*. 2015;166(4):243–58.
4. Kawano H, Kayama H, Nakama T, Hashimoto T, Umemoto E, Takeda K. IL-10-producing lung interstitial macrophages prevent neutrophilic asthma. *Int Immunol*. 2016;28(10):489–501.
5. Sag D, Özkan M, Kronenberg M, Wingender G. Improved Detection of Cytokines Produced by Invariant NKT Cells. *Sci Rep*. 2017;7(1).
6. Bouabe H. Cytokine Reporter Mice: The Special Case of IL-10. *Scand J Immunol* [Internet]. 2012 Jun;75(6):553–67. Available from: <http://doi.wiley.com/10.1111/j.1365-3083.2012.02695.x>
7. Madan R, Demircik F, Surianarayanan S, Allen JL, Divanovic S, Trompette A, et al. Nonredundant Roles for B Cell-Derived IL-10 in Immune Counter-Regulation. *J Immunol*. 2009;183(4):2312–20.
8. Mitchell AJ, Pradel LC, Chasson L, Van Rooijen N, Grau GE, Hunt NH, et al.

- Technical Advance: Autofluorescence as a tool for myeloid cell analysis. *J Leukoc Biol* [Internet]. 2010 Sep;88(3):597–603. Available from: <http://doi.wiley.com/10.1189/jlb.0310184>
9. Telford WG, Hawley T, Subach F, Verkhusha V, Hawley RG. Flow cytometry of fluorescent proteins. *Methods* [Internet]. 2012;57(3):318–30. Available from: <http://dx.doi.org/10.1016/j.ymeth.2012.01.003>
10. Peñaloza HF, Noguera LP, Ahn D, Vallejos OP, Castellanos RM, Vazquez Y, et al. Interleukin-10 produced by Myeloid-Derived suppressor cells provides protection to Carbapenem-Resistant *klebsiella pneumoniae* sequence type 258 by enhancing its clearance in the airways. *Infect Immun*. 2019;87(5).
11. Tighe RM, Misharin A V., Jakubzick C V., Brinkman R, Curtis JL, Duggan R, et al. Improving the quality and reproducibility of flow cytometry in the lung. *Am J Respir Cell Mol Biol*. 2019;61(2):150–61.
12. Black PN, Ching PST, Beaumont B, Ranasinghe S, Taylor G, Merrilees MJ. Changes in elastic fibres in the small airways and alveoli in COPD. *Eur Respir J*. 2008;31(5):998–1004.
13. Kretschmer S, Pieper M, Hüttmann G, Bölke T, Wollenberg B, Marsh LM, et al. Autofluorescence multiphoton microscopy for visualization of tissue morphology and cellular dynamics in murine and human airways. *Lab Investig*. 2016;96(8):918–31.
14. Wingender G, Birkholz a. M, Sag D, Farber E, Chitale S, Howell a. R, et al. Selective Conditions Are Required for the Induction of Invariant NKT Cell Hyporesponsiveness by Antigenic Stimulation. *J Immunol* [Internet].

Figure legends

Figure 1: The myeloid-derived GFP signal is comparable between IL10^{GFP} and C57BL/6 mice and increases during lung inflammation. C57BL/6 and IL-10^{GFP} mice were subjected to three daily administrations (d0, d1, d2) of either PBS or 10 µg LPS per mouse with via aspiration. 16-18 hours after the last administration, the single cell suspension from lungs, spleens, and mLNs were stained for CD4⁺ T cells (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁺ CD4⁺ CD8 α ⁻), CD8⁺ T cells (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁺ CD4⁻ CD8 α ⁺), macrophages (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁺), Siglec-F⁺ cells (eosinophils, alveolar macrophages; CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁺ F4/80^{-/+}), neutrophils (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁻ Ly6G⁺ CD11b⁺), monocytes (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁻ Ly6G⁻ Ly6C⁺ CD11b⁺), and ILCs (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁻ Ly6G⁻ CD90.2⁺ CD127^{lo/-}). The expression of IL-10 was measured either by GFP⁺ or by intracellular IL-10 as indicated. **(A, B)** The frequency of GFP⁺ cells (live CD45⁺) in the lungs of (A) PBS or (B) LPS challenged C57BL/6 (red) and IL-10^{GFP} (green) mice are shown. The graph shows combined data from three independent experiments (PBS: n = 9 mice/group, LPS: n =13-15 mice/group). **(C)** Representative histogram showing the shift in the GFP-signal (505 nm - 550 nm) of live CD45⁺ lung cells upon LPS administration in C57BL/6 and IL-10^{GFP} mice strains.

Figure 2: The myeloid-derived IL-10^{GFP} signal is masked by autofluorescence in granulocytes. C57BL/6 and IL-10^{GFP} mice were subjected to three daily administrations (d0, d1, d2) of 10 µg LPS per mouse with via aspiration. 16-18 hours after the last administration, the single cell suspension from lungs, spleens, and mLNs were stained

for CD4⁺ T cells (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁺ CD4⁺ CD8 α ⁻), CD8⁺ T cells (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁺ CD4⁻ CD8 α ⁺), macrophages (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁺), Siglec-F⁺ cells (eosinophils, alveolar macrophages; CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁺ F4/80^{-/+}), neutrophils (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁻ Ly6G⁺ CD11b⁺), monocytes (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁻ Ly6G⁻ Ly6C⁺ CD11b⁺), and ILCs (CD45⁺ CD45R/B220⁻ CD3 ϵ ⁻ Siglec-F⁻ F4/80⁻ Ly6G⁻ CD90.2⁺ CD127^{lo/-}). The expression of IL-10 was measured either by GFP⁺ or by intracellular IL-10 as indicated.

(A-G) Comparison of the IL-10 signal derived from intracellular cytokine staining (ICCS) or from the GFP-signal (IL10^{GFP}) of IL10^{GFP} mice in (D) CD4⁺ T cells, (E) CD8⁺ T cells, (F) neutrophils, (G) macrophages, (H) Siglec-F⁺ cells (eosinophils, alveolar macrophages), (I) monocytes, and (J) ILCs from indicated organs. The graph shows combined data from two independent experiments (PBS: n = 6 mice/group, LPS: n = 7-9 mice/group).

Figure 3: The cell-type-specific autofluorescence in C57BL/6 mice in control conditions. Representative histograms from the **(A)** lung and the **(B)** spleen of PBS challenged C57BL/6 mice, demonstrating the increase in autofluorescence in the GFP-channel for myeloid cells. The graphs show representative data from three independent experiments (PBS: n = 9 mice/group, LPS: n = 13-15 mice/group).

Figure 4: The autofluorescence of alveolar macrophages increases during lung inflammations. C57BL/6 and IL-10GFP mice were subjected to three daily administrations (d0, d1, d2) of 10 μ g LPS per mouse via aspiration. 16-18 hours after the last administration, the single-cell suspension from the lungs. Representative histogram

showing the GFP-signal from alveolar macrophages (CD45⁺ CD45R/B220⁻ CD3⁻ Siglec-F⁺ F4/80⁺ CD11b⁻) from C57BL/6 and IL10GFP mice as indicated. The data shown are representative of three independent experiments (PBS: n = 9 mice/group, LPS: n = 13-15 mice/group).

Supplementary figure 1:

A graphic outline of the gating strategy to identify CD4⁺ T cells (CD45⁺ CD45R/B220⁻ CD3⁺ CD4⁺ CD8^α⁻ CD127^{lo}), CD8⁺ T cells (CD45⁺ CD45R/B220⁻ CD3⁺ CD4⁻ CD8^α⁺), macrophages (CD45⁺ CD45R/B220⁻ CD3⁻ Siglec-F⁻ F4/80⁺), Siglec-F⁺ cells (eosinophils, alveolar macrophages; CD45⁺ CD45R/B220⁻ CD3⁻ Siglec-F⁺ F4/80^{-/+}), neutrophils (CD45⁺ CD45R/B220⁻ CD3⁻ Siglec-F⁻ F4/80⁻ Ly6G⁺ CD11b⁺), monocytes (CD45⁺ CD45R/B220⁻ CD3⁻ Siglec-F⁻ F4/80⁻ Ly6G⁻ Ly6C⁺ CD11b⁺), and ILCs in the inflamed lung.

Supplementary Figure 2: Efficiency of intracellular IL-10 cytokine staining.

Representative flow cytometric dot plots showing the IL-10 staining (ICCS) in splenic CD4⁺ T cells. **(C)** Representative histograms from the spleen of PBS (left) or LPS (right) challenged C57BL/6 and IL-10^{GFP} mice, demonstrating the inefficiency of secondary αGFP-AF488 antibody to improve the GFP-signal.

Supplementary Figure 3: The staining with a secondary αGFP-AF488 antibody does

not improve the resolution of the myeloid IL10GFP signal. Representative histograms from the spleen of PBS (left) or LPS (right) challenged C57BL/6 and IL-10GFP

344 mouse, demonstrating the overall GFP-signal detected with or without labelling with
345 secondary α GFP-AF488.

346

347

348

349

Figure 1

Figure 1

Figure 2

Figure 2

Figure 3

Figure 3

Figure 4

Figure 4