Conservation and Convergence of Genetic Architecture in the Adaptive Radiation of *Anolis* lizards

Joel W. McGlothlin,^{1,*} Megan E. Kobiela,² Helen V. Wright,³ Jason J. Kolbe,⁴ Jonathan B. Losos,⁵ and Edmund D. Brodie III⁶

1. Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia 24061; email:

joelmcg@vt.edu

2. School of Biological Sciences, University of Nebraska, Lincoln, Nebraska 68588; email:

mkobiela2@unl.edu

3. Computing Community Consortium, Computing Research Association, Washington, DC

20036; email: helenvwright2@gmail.com

4. Department of Biological Sciences, University of Rhode Island, Kingston, Rhode Island

02881; email: jjkolbe@uri.edu

5. Department of Biology, Washington University, Saint Louis, Missouri 63130; email:

losos@wustl.edu

6. Department of Biology and Mountain Lake Biological Station, University of Virginia,

Charlottesville, Virginia 22904; email: edb9j@virginia.edu

* Corresponding author

Short running title: *Conservation and Convergence of G*

ABSTRACT: The **G** matrix, which quantifies the genetic architecture of traits, is often viewed as an evolutionary constraint. However, **G** can evolve in response to selection and may also be viewed as a product of adaptive evolution. The evolution of similar **G** matrices in similar environments would suggest that **G** evolves adaptively, but it is difficult to disentangle such effects from phylogeny. Here, we use the adaptive radiation of *Anolis* lizards to ask whether convergence of **G** accompanies the repeated evolution of habitat specialists, or ecomorphs, across the Greater Antilles. We measured **G** in seven species representing three ecomorphs (trunk-crown, trunk-ground, and grass-bush). We found that the overall structure of **G** does not converge. Instead, the structure of **G** is well conserved and displays a phylogenetic signal. However, several elements of **G** showed signatures of convergence, indicating that some aspects of genetic architecture have been shaped by selection. Most notably, genetic correlations between limb traits and body traits were weaker in long-legged trunk-ground species, suggesting effects of recurrent selection on limb length. Our results demonstrate that common selection pressures may have subtle but consistent effects on the evolution of **G**, even as the overall pattern of genetic architecture remains conserved.

Keywords: adaptive radiation, *Anolis* lizards, constraint, **G** matrix, genetic correlation, quantitative genetics

Introduction

Genetic variation translates natural selection into evolutionary change (Falconer and MacKay 1996; Roff 1997). Understanding the nature of such genetic variation and the processes that shape it has been a goal of evolutionary biology since the early days of population genetics (Dobzhansky 1937). In evolutionary quantitative genetics, the pattern of genetic variation in a population is described by the genetic variance-covariance matrix $\mathbf{G}$, which predicts the multivariate response to phenotypic selection (Lande 1979). The $\mathbf{G}$ matrix can be used to make accurate predictions of short-term evolutionary change (Grant and Grant 1995), but its utility for making long-term predictions is more suspect because $\mathbf{G}$ itself may evolve during adaptive evolution (Turelli 1988; Steppan et al. 2002). Although early theory argued for the stability of $\mathbf{G}$ (Lande 1980), both more recent theoretical (Agrawal et al. 2001; Jones et al. 2003; 2004, 2012, 2014; Revell 2007; Arnold et al. 2008) and empirical results (Steppan et al. 2002; Cano et al. 2004; Doroszuk et al. 2008; Hine et al. 2009; Eroukhmanoff and Svensson 2011; Björklund et al. 2013; Careau et al. 2015) suggest that $\mathbf{G}$ can and does evolve, sometimes rapidly. Given enough time, selection is expected to align $\mathbf{G}$ with the adaptive landscape (Cheverud 1984; Arnold et al. 2001, 2008; Jones et al. 2003, 2014; Revell 2007), potentially making $\mathbf{G}$ as much a product of adaptive evolution as a constraint upon it (Merilä and Björklund 2004).

Because stability and evolutionary lability of $\mathbf{G}$ are both plausible theoretical outcomes, the relative importance of history and adaptation in shaping $\mathbf{G}$ is largely an empirical question (Arnold et al. 2008). Several studies have shown a relationship between the shape of $\mathbf{G}$ and divergence, suggesting the importance of genetic constraints in channeling evolutionary outcomes (Bégin and Roff 2003; 2004; Blows and Hoffmann 2005; McGuigan et al. 2005;

McGuigan 2006; Hansen and Houle 2008; Walsh and Blows 2009; Chenoweth et al. 2010; Bolstad et al. 2014; Houle et al. 2017; McGlothlin et al. 2018; Walter et al. 2018). The early stages of adaptive radiation are expected to be aligned with the “genetic line of least resistance” describing the direction of greatest genetic variation within a population (Schluter 1996). Although natural selection should be able to push phenotypes away from this line given enough time, recent results suggest that evolutionary change may be predicted by axes of genetic variation for tens of millions of years (Houle et al. 2017; McGlothlin et al. 2018). Conversely, it is well established that both directional and nonlinear selection may alter aspects of **G**. For example, correlational selection, which occurs when certain combinations of traits are favored over others, can directly alter the strength of genetic correlations each generation (Phillips and Arnold 1989; Jones et al. 2003; Revell 2007). Patterns of genetic correlation are often congruent with axes of correlational selection in the wild (Brodie 1989; 1992; McGlothlin et al. 2005; Roff and Fairbairn 2012), and genetic correlations can evolve in response to artificial correlational selection (Delph et al. 2011; Steven et al. 2020).

Although comparative studies of **G** have become more common in recent years (Steppan et al. 2002; Bégin and Roff 2003; Hine et al. 2009; Eroukhmanoff and Svensson 2011; Walter et al. 2018), none have been able to disentangle the effects of shared ancestry from similar selection pressures in determining the evolution of **G**. Convergent evolution of **G** or its elements in similar environments would provide strong evidence that changes in **G** represent adaptation of genetic architecture (Losos 2011). The adaptive radiation of West Indian *Anolis* lizards provides an ideal testing ground for hypotheses about the evolution of **G** because the effects of phylogenetic history and ecological selection are largely decoupled (Losos 1994; 2009; 2011). In the Greater Antilles, anoles have diversified into 120 species, 95 of which can be classified as one of six

habitat specialists, or ecomorphs, each of which has evolved multiple times throughout the *Anolis* radiation (Williams 1972; Losos et al. 1998; Beuttell and Losos 1999; Losos 2009). Species with dissimilar morphology on the same island tend to be more closely related than are those with similar morphology on different islands, indicating that the characteristic morphology of ecomorphs is due to convergent evolution (Losos et al. 1998; Harmon et al. 2005; Mahler et al. 2013). This repeated adaptive radiation leads to explicit predictions for the evolution of **G**. If **G** responds predictably to similar selection pressures, **G** should show signatures of convergence among the independent origins of the same ecomorph. Conversely, if **G** evolves relatively slowly and does not respond predictably to similar selection pressures, **G** or its elements should be more similar within lineages than within ecomorph classes.

Previous work in *Anolis* using phenotypic variance-covariance matrices (**P**) as proxies for **G** (Cheverud 1988) suggests that selection may indeed lead to convergence in (co)variance structure. A study comparing **P** in eight *Anolis* species showed significant variation in covariance structure across the radiation and demonstrated convergent changes in **P** in three distantly related species from the same ecomorph class (Kolbe et al. 2011). In a separate study, **P** showed significant alignment with the matrix of nonlinear selection (γ) in *A. cristatellus*, suggesting that contemporary stabilizing and correlational selection may act to shape the pattern of phenotypic (co)variance within species (Revell et al. 2010). These results suggest that selection plays a role in shaping genetic architecture in anoles, but patterns of phenotypic covariance do not necessarily mirror patterns of genetic covariance (Hadfield et al. 2007). Thus, comparative studies that directly estimate **G** are necessary to test whether its structure is more influenced by phylogenetic history or convergent evolution.

In this study, we compare **G** matrices in seven *Anolis* species reared in a controlled laboratory environment. We chose species from lineages originating on three different islands, Puerto Rico, Jamaica, and Cuba, and included three ecomorphs, trunk-crown (three species), trunk-ground (three species), and grass-bush (one species), which are distinguished by their habitat use, coloration, and skeletal morphology (Williams 1972; Beuttell and Losos 1999; Harmon et al. 2005; Losos 2009). Trunk-crown lizards are typically found high in trees and are usually green with relatively short legs for climbing and clinging to narrow perches. Trunk-ground lizards tend to be found on low perches or on the ground and are typically brown with long hindlimbs that aid in running quickly and jumping far (Losos and Sinervo 1989; Losos 1990; Irschick and Losos 1998; Beuttell and Losos 1999). The third ecomorph, grass-bush, has a slender body that matches its narrow perches and long hindlimbs that allow it to both run and jump well (Losos 1990; Beuttell and Losos 1999). The three trunk-crown species are distantly related to one another, as are the trunk-ground species. Both ecomorphs may have evolved three separate times, although it cannot be ruled out that one of these ecomorphs represents the ancestral state for the anole radiation (Losos 2009). Because of the importance of skeletal morphology, and limb length in particular, to the evolution of these ecomorphs, our estimates of **G** focus on skeletal traits.

Our previous results have shown that **G** varies substantially across these *Anolis* species, while retaining conserved axes of genetic variation (McGlothlin et al. 2018). Specifically, **G** matrices varied most in size (overall genetic variance), and the major axis of genetic variance remained similar in orientation across all species. This major axis of genetic variance was similar in orientation to the major axis of morphological divergence, suggesting that divergence has occurred along a genetic line of least resistance even though **G** has not remained constant. The

largest evolutionary changes in **G** were also aligned with the major axes of both genetic variance and morphological divergence. This triple alignment may have been caused by deep genetic constraints underlying both the evolution of **G** and morphological divergence, alignment of **G** with the adaptive landscape, genetic drift, or some combination of the three (McGlothlin et al. 2018).

Here, we explicitly consider the role of selection in shaping **G**-matrix evolution across the *Anolis* radiation by testing whether aspects of **G** show patterns of convergence that mirror the repeated evolution of ecomorphs. To do so, we use two types of comparisons. First, we test for convergence of the overall structure of **G** by asking whether random skewers correlations, which are estimates of pairwise similarity in the predicted multivariate response to selection, are better predicted by shared evolutionary history or shared ecology. Second, we conduct similar tests for individual elements of **G** (i.e., variances and covariances of individual traits) to test for signatures of convergence on a finer scale. Although many processes, including both selection and drift, could lead to similarities in **G** among more closely related species, convergence in the structure of **G** among distantly related species of the same ecomorph would provide strong evidence that **G** may be predictably shaped by common selection pressures.

Methods

*Estimation of **G***

Detailed methods for estimation of the **G** matrices used here are reported elsewhere (McGlothlin et al. 2018). Briefly, adults from seven *Anolis* species, representing independent origins of trunk-

crown (*A. evermanni*, Puerto Rico; *A. grahami*, Jamaica; *A. smaragdinus*, a Bahamian species descended from *A. porcatus* on Cuba), trunk-ground (*A. cristatellus*, Puerto Rico; *A. lineatopus*, Jamaica; *A. sagrei*, Cuba), and grass-bush ecomorphs (*A. pulchellus*, Puerto Rico), were collected from the wild (Fig. 1). Due to travel restrictions, species from Cuban lineages were collected from South Bimini, The Bahamas, where they occur naturally. These seven species shared a common ancestor approximately 41.5–43.5 million years ago, and the most recent phylogenetic split (between *A. cristatellus* and *A. pulchellus*) dates is estimated at 19.8–22.5 million years ago (fig. 1, Zheng and Wiens 2016; Poe et al. 2017).

Adults were housed in individual cages in the laboratory except when paired for breeding and held at controlled photoperiod (12L:12D for Puerto Rican and Jamaican adults and 13L:11D for Bahamian adults), temperature (28°C during the day and 25°C at night), and relative humidity (65%). Lizards were provided with a perch, a mesh hammock for basking near an adjacent UVB bulb, and a carpet substrate. Adults were mated in a paternal half-sib breeding design (average of 47 sires and 69 dams per species) to produce offspring (2,904 total, see McGlothlin et al. 2018 for more sampling details). Laying females were provided with potted plants, which were checked weekly for eggs, which were placed in individual cups with a 1:1 mixture of water and vermiculite and held in an incubator at 28°C and 80% humidity until hatching.

Juveniles were reared in individual cages until 6 months of age and were provided with crickets and water daily. At 0, 1, 3, and 6 months of age, we X-rayed juveniles by chilling them for 10 min at 5°C in small plastic bags. The bags were then secured with masking tape to a film cartridge (Kodak Biomax XAR) for imaging in a Faxitron 43805N radiography system. Developed radiographs were digitized using a flatbed scanner. Using ImageJ (NIH), we measured snout-vent length (SVL) and eight other skeletal traits, jaw length (JL), head width (HW),

pectoral width (PECT), pelvis width (PELV), humerus (HUM), ulna (UL), femur (FEM), and tibia (TIB; see Fig. 1A for location of measured traits). In all, 9,369 individual X-ray images were measured (McGlothlin et al. 2018). We used multivariate repeated measures animal models in ASReml 3.0 (Gilmour et al. 2009) to estimate **G** matrices for natural-log transformed traits, with size (natural-log SVL) as a covariate to correct for age and growth. These models included two random animal effects, one linked to the pedigree to estimate additive genetic (co)variance and a second unlinked effect to estimate effects of permanent environment. All species could not be reliably sexed as juveniles; therefore, we did not correct for sex in our models. In one species that has been studied extensively in the laboratory, *A. sagrei*, sexual size dimorphism is not noticeable at hatching and only becomes elaborated after 6 months of age with the maturation of testes in males (Cox et al. 2017). Genetic correlations are shown (along with heritabilities) in table A1 and visualized in fig. 1; full **G** matrices, reprinted from McGlothlin et al. (2018), are also shown in table A1. Permanent environment (co)variances, which were generally at least an order of magnitude smaller than genetic (co)variances, and residual (co)variances are not presented here but were used in the calculation of total phenotypic variance for calculating heritabilities. As reported previously, in all but two species, all eight traits we measured were significantly heritable (mean h^2 across species $\pm$ s.d.: JL, $.40 \pm .150$; HW, $.22 \pm .084$; PECT, $.21 \pm .073$; PELV, $.22 \pm .046$; HUM, $.16 \pm .047$; UL, $.15 \pm .042$; FEM, $.45 \pm .143$, TIB, $.54 \pm .091$, table A1; McGlothlin et al. 2018). In general, genetic correlations were strong and positive for pairs of limb traits and both weaker and more variable across species for other trait combinations (fig. 1, table A1; McGlothlin et al. 2018).

Figure 1: (A) Traits measured in this study: JL = jaw length, HW = head width, PECT = pectoral width, PELV = pelvis width, HUM = humerus, UL = ulna, FEM = femur, TIB = tibia. (B) Schematic of a genetic correlation matrix showing the location of each trait. Elements are color-coded based on morphological modules (head, body, and limbs; see Methods), showing within-module correlations in primary colors (blue = head, red = body, yellow = limbs) and between module correlations in secondary colors (violet = head-body, green = head-limb, orange = body-limb). (C) Graphical representation of genetic correlation matrices for each species. Positive correlations are shown in green and negative correlations are shown in yellow, with brighter colors signifying stronger correlations. See table A1 for values. Photographs by J.B.L. (*A. pulchellus*, *A. lineatopus*, and *A. grahami*), and E.D.B. III (all other species).

Statistical Analyses

All statistical analyses using **G** matrices were performed in R 3.5.3 (R Core Team 2019). To incorporate estimation error in **G** into our analyses, we used the restricted maximum-likelihood multivariate normal (REML-MVN) method developed by Houle and Meyer (2015). We used point estimates of **G** and the sampling (co)variance matrix of (co)variance components, which is calculated by ASReml as the inverse of the average information matrix (Gilmour et al. 2009), to define a multivariate normal distribution. From this distribution, we generated 10,000 samples of **G** for each species using the function rmvn from the R package mgcv (Wood 2012). All further analyses were run using both the point estimates of **G**, from which we estimated parameters and test statistics, and the set of 10,000 REML-MVN samples per species, from which we calculated 95% confidence intervals using the 2.5% and 97.5% quantiles from the parameter distribution. A parameter was considered to be statistically supported when its 95% confidence interval did not overlap the expected null value (often zero). We clarify the specific ways REML-MVN samples were used for each analysis below.

To quantify pairwise similarity in the overall structure of **G**, we used random skewers analysis, which compares the response to selection predicted by a pair of **G** matrices (Marroig and Cheverud 2001; Cheverud and Marroig 2007; Revell 2007; Aguirre et al. 2014). Random skewers analysis has the advantage of providing an evolutionarily relevant comparison of **G** using a single metric. We used the function RandomSkewers in the R package evolQG (Melo et al. 2015) to apply 10,000 random selection gradients (β) to all seven **G** matrices. Each element of these skewers was drawn from a normal distribution, after which each skewer was normalized to unit length and used to calculate the predicted multivariate selection response ($\Delta\bar{z}$) from the

multivariate breeder's equation, $\Delta\bar{\mathbf{z}} = \mathbf{G}\boldsymbol{\beta}$ (Lande 1979). The correlation in response to selection for two species, or the random skewers correlation (r_{RS}), was calculated as the pairwise vector correlation of the resultant 10,000 estimates of $\Delta\bar{\mathbf{z}}$. To incorporate estimation error, r_{RS} was recalculated for 10,000 sets of seven $\mathbf{G}$ matrices each using our REML-MVN samples. We report the mean and 95% confidence intervals of this distribution in addition to our point estimate. Incorporating error pairs of $\mathbf{G}$ matrices leads to negatively skewed distributions of the REML-MVN estimates of r_{RS} . Thus, means of the REML-MVN estimates of r_{RS} tend to be lower than the point estimates.

To determine whether more closely related species had more similar $\mathbf{G}$ matrices, we used a Mantel test to compare a matrix of r_{RS} to a patristic distance matrix, which was calculated using a tree pruned from a dated squamate phylogeny (Zheng and Wiens 2016); dates from a tree of all *Anolis* species (Poe et al. 2017), which had identical topology for our species, were similar. Although Mantel tests are not useful for comparing (co)variance matrices directly (Steppan et al. 2002) and are not recommended for use in the phylogenetic comparative method when they can be avoided, they are the only option when comparative analyses involve data that can only be expressed as either a correlation or a distance between species (Harmon and Glor 2010). A negative Mantel correlation ($r_{\text{M}} < 0$) between r_{RS} and patristic distance would indicate that the similarity in the structure of $\mathbf{G}$ is explained by phylogenetic similarity. To determine whether unrelated species of the same ecomorph displayed convergence in the structure of $\mathbf{G}$, we generated a matrix consisting of zeros (different ecomorph) and ones (same ecomorph) and compared it to a matrix of r_{RS} using a partial Mantel test that used the patristic distance matrix as the control matrix. The single grass-bush species, *A. pulchellus*, was excluded from this and all

other analyses that included an ecomorph effect. A positive correlation ($r_M < 0$) between the r_{RS} matrix and the ecomorph matrix would indicate convergence.

Mantel correlations were calculated in R using the base function `corr` (simple Mantel test) or the function `pcor` (partial Mantel test) from the package `ppcor` (Kim 2015). We did not perform permutation tests to assess statistical significance as is the typical practice when performing Mantel tests. Rather, we conducted separate Mantel tests for each of 10,000 sets of r_{RS} matrices calculated using our REML-MVN samples and report the mean and 95% confidence intervals of this distribution.

Evolutionary patterns in **G** may involve changes that are subtler than can be detected in analyses of its overall structure. Therefore, we also tested for the effects of shared evolutionary history and shared ecology on the individual elements of **G**. We compared genetic variances (diagonal elements of **G**) and genetic correlations (off-diagonal elements of **G** standardized by the square root of the product of the variances) across **G** matrices, testing for both phylogenetic signal and differences between ecomorphs (trunk-crown vs. trunk-ground). We present comparisons of genetic correlations rather than genetic covariances so that tests for associations between traits would be independent of differences in variance across species. However, we note that analyses using covariances gave nearly identical results (not shown).

To test for the effects of shared evolutionary history, we used Blomberg's K as an estimate of phylogenetic signal (Blomberg et al. 2003; implented in the R package `phytools`, Revell 2012). Values of $K = 1$ indicate phylogenetic signal consistent with a Brownian motion model of evolution. Values of $K < 1$ indicate weaker phylogenetic signal, and $K > 1$ indicate stronger signal than would be predicted from a Brownian motion model. We tested for effects of shared ecology on genetic variance and genetic correlations using phylogenetic generalized least

squares (Martins and Hansen 1997) with ecomorph as a predictor (coding trunk-crown as 0 and trunk-ground as 1; *A. pulchellus* was excluded), our dated tree, and an assumption of Brownian motion evolution (implemented in the R package APE, Paradis et al. 2004). Alternative models that assumed an Ornstein-Uhlenbeck evolutionary model provided similar results (not shown). For each test, we report the evolutionary correlation (r_e) between the element of **G** and ecomorph to remove effects of scale. For a given element of **G**, $r_e > 0$ indicates a larger value for trunk-ground species, and $r_e < 0$ indicates a larger value for trunk-crown species. Both of these analyses were conducted for both our point estimates and our REML-MVN samples to incorporate estimation error. Using the 95% confidence intervals from the distributions of REML-MVN estimates, we note deviations from both $K = 0$ and $K = 1$ for phylogenetic signal and $r_e = 0$ for the evolutionary correlation.

For visualization, we grouped our estimates of phylogenetic signal and evolutionary correlation into six groups based on phenotypic module. Three groups consisted of elements of **G** within modules, head (JW, HL), body (PECT, PELV), and limbs (HUM, UL, FEM, and TIB), and three contained elements of **G** between modules, head-limb, head-body, and body-limb.

Results

*Overall Structure of **G***

Predicted responses to selection were highly correlated between all pairs of species (mean $r_{RS} = .84$, range .75 – .92, REML-MVN mean $r_{RS} = 0.73$, range 0.61 – 0.86; table A2, fig. 2), suggesting that the species have **G** matrices with similar overall structure despite being separated

for 20–44 million years. **G**-matrix similarity was negatively correlated with phylogenetic distance ($r_M = -.44$; REML-MVN estimate: r_M [95% CI] = $-.41$ [$-.59, -.15$]), indicating that closely related species have **G** matrices that predict a more similar evolutionary response. More distantly related pairs of species also displayed more variable values of r_{RS} , with some distantly related pairs of species showing highly similar **G** matrices and others showing highly dissimilar **G** (fig. 2). Overall similarity of **G** was not predicted by ecomorph ($r_M = -.12$; REML-MVN estimate: r_M [95% CI] = $.04$ [$-.22, .32$]). However, the two distantly related trunk-ground species *A. cristatellus* and *A. sagrei* did have highly similar **G** matrices (figs. 2 & 3).

Individual Elements of G

All individual elements of **G** showed phylogenetic signal significantly higher than $K = 0$ but indistinguishable from $K = 1$, which is the null expectation of the Brownian motion model (mean $K = .91$, REML-MVN mean $K = .94$; fig. 4A, table A3). Phylogenetic signal did not exhibit any detectable patterns across trait groups (fig. 4A).

Six elements of **G** differed significantly between trunk-crown and trunk-ground ecomorphs. Specifically, genetic correlations between pelvis width and both jaw length and all four limb bones were significantly lower in trunk-ground species than in trunk-crown species (fig. 4B; table A4). In addition, genetic correlations between jaw length and femur length were higher in trunk-ground species than in trunk-crown species (fig. 4B; table A4). In some cases, these effects involved changes in the sign of genetic correlations from positive in trunk-crown species to negative in trunk-ground species (fig. 1). Although not a statistically significant effect, genetic correlations both among limb bones and between pectoral width and all four limb bones

Figure 2: Correlations between species in the overall structure of the **G** matrix as measured via random skewers correlations (r_{RS}). Stronger correlations are represented by darker shading. Within-island comparisons are shown with a solid border, and within-ecomorph comparisons are shown with a dashed border (TC = trunk-crown, TG = trunk-ground). Islands of origin are represented by their shapes on the phylogeny (Puerto Rico, Jamaica, and Cuba, from top to bottom) and for the three within-island comparisons. Note that despite originating on Cuba, *A. sagrei* is actually more closely related to Jamaican species than to *A. smaragdinus*. The phylogeny is pruned from a dated tree of squamates (Zheng and Wiens 2016).

Figure 4 (previous page): Element-by-element comparisons of genetic variances (solid circles) and correlations (open circles). Elements are split into three within-module groups (head, body, and limbs) and three between-module groups (head/body, body/limb, and limb/head) and are labeled with abbreviations as in Fig. 1. Bars show means and 95% confidence intervals for a set of point estimates within a group and are presented for visualization purposes only. (A) Strength of phylogenetic signal, as estimated using Blomberg's K . Estimates were clustered around $K = 1$, suggesting phylogenetic signal consistent with Brownian motion. (B) Ecomorph effects from phylogenetic least squares, given as the evolutionary correlation (r_e). Points above the midline indicate that trunk-ground species had higher values of a given element of **G** than did trunk-crown species; the converse is true below the midline. Six genetic correlations showed a significant correlation with ecomorph (trunk-ground vs. trunk-crown; $p < .05$, denoted by *).

tended to be weaker in trunk-ground species than in trunk-crown species, indicating a trend toward weaker integration among limb and body traits in trunk-ground species.

Although we did not conduct formal tests for the lone grass-bush species, *A. pulchellus*, inspecting its genetic correlation matrix (fig. 1, table A1) shows that this species displays some similarities to trunk-ground lizards. In particular, this species had weak (and occasionally negative) genetic correlations between the pelvis and all limb bones, similar to all trunk-ground species. *A. pulchellus* also showed genetic correlations between hindlimb and forelimb bones that were noticeably weaker than the other two species in the Puerto Rican lineage (fig. 1, table A1).

Discussion

Here we present two significant findings about the evolution of quantitative genetic architecture within the adaptive radiation of West Indian *Anolis* lizards. First, when viewed in terms of its

effects on multivariate response to selection, we show that the overall structure of the *Anolis* **G** matrix retains a phylogenetic signal. Although the seven species we studied have been separated for 20–44 million years, all **G** matrices predicted a very similar multivariate evolutionary response, and more closely related species had more similar **G** matrices. Second, despite this phylogenetic signal in the overall structure of **G**, pairwise genetic correlations between limb traits and body traits showed consistent signatures of convergence, suggesting that they have been adaptively shaped by similar selection pressures resulting from each ecomorph's niche. In particular, longer-limbed trunk-ground lizards show a decoupling of limb length and pelvis width relative to shorter-limbed trunk-crown lizards, demonstrating that convergent changes in genetic architecture may accompany repeated morphological adaptation. Taken together, our results show that selection may alter **G** in predictable and evolutionarily consequential ways without leading to major changes in its overall structure.

From the perspective of overall multivariate response to selection, **G** is remarkably similar across the *Anolis* radiation, with predicted responses to selection in random directions showing strong positive correlations ranging from .76 to .92. Over the span of a few generations, then, morphological divergence of species with the **G** matrices estimated here should be constrained to lie along directions defined by quantitative genetic architecture. Indeed, previous work has shown that divergence of these species remains aligned with the major axis of genetic variation, $\mathbf{g}_{\max}$, even after ~44 million years of divergence (McGlothlin et al. 2018).

The similarity of the overall structure of **G** declined with greater phylogenetic distance. This pattern suggests that the overall structure of **G** changes relatively slowly, remaining conserved over millions of years. While similarity of **G** in closely related species may in part reflect comparable patterns of contemporary selection in species in the same lineage, most

similarity is likely attributable to shared evolutionary history. In addition to this phylogenetic trend, we found that more distantly related species also show greater variance in random skewers correlations. Although many distantly related species have dissimilar **G** matrices, some pairs have highly congruent **G**, a pattern that is not explained by convergent morphology. This increased variance emphasizes the unpredictability of the evolution of **G** structure over longer timescales.

When considering more subtle changes in **G**—shifts in trait-specific genetic variances and correlations—we found evidence that **G** can change repeatedly and predictably in response to similar selection pressures. The strongest convergence occurred in genetic correlations between limb bones and pelvis width, which were significantly reduced in trunk-ground species relative to trunk-crown species. In some cases, genetic correlations differed in sign between ecomorphs. When this was the case, genetic correlations were usually negative in trunk-ground species and positive in trunk-crown species. These results indicate that the pattern of genetic integration was subtly remodeled in the transition from trunk-crown to trunk-ground ecomorphs (or vice versa), most notably in the relationship between limb length and pelvis width, for which trunk-ground species showed weaker genetic correlations when compared to trunk-crown species.

The ecomorph difference in the genetic correlations between limb length and pelvis width is likely to result from a combination of directional selection and correlational selection acting on these traits. Ecomorph differences in limb length have clear links to performance within their characteristic habitats (Losos and Sinervo 1989; Losos 1990; Irschick and Losos 1998), suggesting that they have been driven apart by divergent directional selection. The longer hindlimbs of trunk-ground lizards facilitate running faster on flatter surfaces, whereas the shorter hindlimbs of trunk-crown lizards are suited for a wider variety of perches (Losos 1990; Losos and

Irschick 1996; Irschick and Losos 1998). Strong directional selection acting only on limb length could reduce its genetic correlations with other traits; however, it is also likely that correlational selection has played a role. The evolution of trunk-ground anoles from a hypothetical trunk-crown ancestor would require evolution of longer legs without a concomitant increase in the pelvis, which may lead to negative correlational selection to decouple the two traits. In contrast, correlational selection might favor a positive correlation between the two traits in trunk-crown anoles, perhaps because a matching limb and pelvic morphology would facilitate agility on branches.

Differences in hindlimb length between trunk-crown and trunk-ground ecomorphs are apparent at hatching and appear to emerge mostly via changes in developmental patterning early in embryonic development rather than differences in growth (Sanger et al. 2012). The developmental genetic networks underlying limb growth and development are well understood (Rabinowitz and Vokes 2012), and comparative genomic evidence indicates that genes expressed in these networks experienced enhanced positive selection during the radiation of anoles (Tollis et al. 2018). Limb-development networks share some genes in common with the network underlying development of the pelvic girdle (Sears et al. 2015). Therefore, it is likely that changes in genetic correlations between limb length and pelvis involve evolutionary changes in the expression of some of these shared genes. Future work should explore remodeling of these networks to understand the developmental genetic underpinnings of convergent morphological evolution in anoles.

Weaker genetic correlations between limbs and body traits likely facilitated the evolution of longer hindlimbs in trunk-ground anoles without correlated changes in the rest of the body. Such genetic decoupling of limbs and body may help explain the remarkable adaptability of the

trunk-ground ecomorph. Trunk-ground anoles seem to be especially capable of colonizing new habitats. Trunk-ground species have successfully colonized the Bahamas and the Virgin Islands (*A. sagrei* and *A. cristatellus*, respectively) and have become established invaders in a number of locations following introduction by humans (Kolbe et al. 2004; Eales et al. 2008; Losos 2009). Trunk-ground anoles also have greater species richness within islands (Losos 2009), in part because they have radiated into ecologically distinct macrohabitats (Glor et al. 2003). Some of this adaptability is likely due to rapid evolution in limb length, such as has been demonstrated in experimental populations of *A. sagrei* (Losos et al. 1997, 2001; Kolbe et al. 2012). Although comparable studies have not been conducted using trunk-crown ecomorphs, their stronger genetic correlations between limb traits and body traits suggest that rapid, independent evolution of limb length would not be as likely in trunk-crown species.

The genetic correlations between pelvis and limb length in the single grass-bush anole we examined resemble those of trunk-ground anoles, suggesting either similar selection or common ancestry, or a combination of the two. Grass-bush anoles have long hindlimbs relative to their body width, suggesting that a combination of directional and correlational selection may have reduced these correlations. However, *A. pulchellus* is likely to have evolved from a trunk-ground ancestor (Poe et al. 2017), which suggests that both this species and the closely related *A. cristatellus* may have inherited weakened genetic correlations between pelvis and limbs from a common ancestor. In other respects, however, the genetic correlation structure of *A. pulchellus* is dissimilar to that of *A. cristatellus*. In contrast to *A. cristatellus*, *A. pulchellus* appears to have attained differences between the lengths of its hindlimbs and forelimbs via a reduction in the genetic correlations between the two, a feature it shares with the distantly related trunk-ground lizard *A. lineatopus*.

Conclusion

The evolution of **G** in West Indian anoles illustrates how the complex interplay between selection and history influences genetic architecture. **G** reflects neither an irresistible pattern of constraint nor an easily adapted phenotype responding quickly to environmental pressures. Patterns of genetic covariation have potentially influenced the pathways that selection may follow, as evidenced by evolution along a deeply conserved genetic line of least resistance in this radiation (McGlothlin et al. 2018). At the same time, as we show here, selection leaves an imprint, if subtle, upon **G** as species diverge. We did not observe a full-scale overhaul of **G** as ecomorphs evolved. However, small-scale differences in the elements of **G** involving critical morphological modules arose predictably between ecomorphs and may have had substantial evolutionary consequences. Our results emphasize that while genetic constraints may change as adaptation proceeds, these changes need not be large to facilitate phenotypic diversification. Rather, the convergent changes we observed in the individual elements of **G**, particularly between the limbs and the body, demonstrate that consistent selection pressures can alter underlying genetic constraints in subtle ways that facilitate adaptation and influence future evolutionary potential.

Acknowledgments

We thank Simon Pearish and Michelle Sivilich for managing the lizard colony. Dozens of undergraduates at the University of Virginia provided animal care and assistance with data collection; special thanks is due to Margo Adler, Tyler Cassidy, Brian Duggar, Maridel Fredericksen, Casey Furr, Jessie Handy, Bryan Hendrick, Uma Pendem, Jeff Wright, and

Elizabeth Zipperle, Leleña Avila, Chris Feldman, Vince Formica, Tonia Hsieh, Melissa Losos, Ashli Moore, Liam Revell, and Matt Sanford assisted with field collections. We thank Brooke Bodensteiner, Vincent Farallo, Kerry Gendreau, Angela Hornsby, and Josef Uyeda for comments on the manuscript, and Luke Harmon, Gabriel Marroig, Emília Martins, and Liam Revell for helpful discussions. This work was supported by the National Science Foundation (grant numbers DEB 0519658 and 0650078 to E.D.B. III and DEB 0519777 and 0722475 to J.B.L.), University of Virginia, and Virginia Tech. The authors declare that there are no conflicts of interest.

Statement of Authorship

J.B.L. and E.D.B. III conceived the study; J.W.M. and J.J.K. contributed to study design; J.W.M., J.J.K., J.B.L., and E.D.B. III performed field collections; J.W.M. and E.D.B. III oversaw the breeding experiment; J.W.M., M.E.K., and H.V.W. collected data; J.W.M. analyzed data; J.W.M. drafted the manuscript and all authors contributed to the final version of the manuscript.

Data and Code Availability

Raw data for estimating **G** matrices are available in a Dryad Data Repository from a previous publication (<https://datadryad.org/stash/dataset/doi:10.5061/dryad.pt2g084>). All code and processed data are available at <https://github.com/joelmcg/AnolisG> and will be archived in Dryad upon manuscript acceptance.

Literature Cited

- Agrawal, A. F., E. D. Brodie, III, and L. H. Rieseberg. 2001. Possible consequences of genes of major effect: transient changes in the **G**-matrix. *Genetica* 112:33-43.
- Aguirre, J. D., E. Hine, K. McGuigan, and M. W. Blows. 2014. Comparing **G**: multivariate analysis of genetic variation in multiple populations. *Heredity* 112:21-29.
- Arnold, S. J., R. Bürger, P. A. Hohenlohe, B. C. Ajie, and A. G. Jones. 2008. Understanding the evolution and stability of the **G**-matrix. *Evolution* 62:2451-2461.
- Arnold, S. J., M. E. Pfrender, and A. G. Jones. 2001. The adaptive landscape as a conceptual bridge between micro- and macroevolution. *Genetica* 112-113:9-32.
- Bégin, M., and D. A. Roff. 2003. The constancy of the **G** matrix through species divergence and the effects of quantitative genetic constraints on phenotypic evolution: a case study in crickets. *Evolution* 57:1107-1120.
- . 2004. From micro- to macroevolution through quantitative genetic variation: positive evidence from field crickets. *Evolution* 58:2287-2304.
- Beuttell, K., and J. B. Losos. 1999. Ecological morphology of Caribbean anoles. *Herpetological Monographs*:1-28.
- Björklund, M., A. Husby, and L. Gustafsson. 2013. Rapid and unpredictable changes of the **G**-matrix in a natural bird population over 25 years. *Journal of Evolutionary Biology* 26:1-13.
- Blomberg, S. P., T. Garland, and A. R. Ives. 2003. Testing for phylogenetic signal in comparative data: Behavioral traits are more labile. *Evolution* 57:717-745.

- Blows, M. W., and A. A. Hoffmann. 2005. A reassessment of genetic limits to evolutionary change. *Ecology* 86:1371-1384.
- Bolstad, G. H., T. F. Hansen, C. Pelabon, M. Falahati-Anbaran, R. Perez-Barrales, and W. S. Armbruster. 2014. Genetic constraints predict evolutionary divergence in *Dalechampia* blossoms. *Philosophical Transactions of the Royal Society B-Biological Sciences* 369.
- Brodie, E. D., III. 1989. Genetic correlations between morphology and antipredator behaviour in natural populations of the garter snakes *Thamnophis ordinoides*. *Nature* 342:542-543.
- . 1992. Correlational selection for color pattern and antipredator behavior in the garter snake *Thamnophis ordinoides*. *Evolution* 46:1284-1298.
- Cano, J. M., A. Laurila, J. Palo, and J. Merilä. 2004. Population differentiation in **G** matrix structure due to natural selection in *Rana temporaria*. *Evolution* 58:2013-2020.
- Careau, V., M. E. Wolak, P. A. Carter, and T. Garland. 2015. Evolution of the additive genetic variance-covariance matrix under continuous directional selection on a complex behavioural phenotype. *Proceedings of the Royal Society B-Biological Sciences* 282.
- Chenoweth, S. F., H. D. Rundle, and M. W. Blows. 2010. The contribution of selection and genetic constraints to phenotypic divergence. *American Naturalist* 175:186-196.
- Cheverud, J. M. 1984. Quantitative genetics and developmental constraints on evolution by selection. *Journal of Theoretical Biology* 110:155-171.
- . 1988. A comparison of genetic and phenotypic correlations. *Evolution* 42:958-968.
- Cheverud, J. M., and G. Marroig. 2007. Comparing covariance matrices: random skewers method compared to the common principal components model. *Genetics and Molecular Biology* 30:461-469.

499 Cox, R. M., C. L. Cox, J. W. McGlothlin, D. C. Card, A. L. Andrew, and T. A. Castoe. 2017.
500 Hormonally mediated increases in sex-biased gene expression accompany the breakdown
501 of between-sex genetic correlations in a sexually dimorphic lizard. *American Naturalist*
502 189:315-332.

503 Delph, L. F., J. C. Steven, I. A. Anderson, C. R. Herlihy, and E. D. Brodie, III. 2011. Elimination
504 of a genetic correlation between the sexes via artificial correlational selection. *Evolution*
505 65:2872-2880.

506 Dobzhansky, T. 1937, *Genetics and the Origin of Species*, Columbia Univ. Press.

507 Doroszuk, A., M. W. Wojewodzic, G. Gort, and J. E. Kammenga. 2008. Rapid divergence of
508 genetic variance-covariance matrix within a natural population. *American Naturalist*
509 171:291-304.

510 Eales, J., R. S. Thorpe, and A. Malhotra. 2008. Weak founder effect signal in a recent
511 introduction of Caribbean *Anolis*. *Molecular Ecology* 17:1416-1426.

512 Eroukhmanoff, F., and E. I. Svensson. 2011. Evolution and stability of the **G**-matrix during the
513 colonization of a novel environment. *Journal of Evolutionary Biology* 24:1363-1373.

514 Falconer, D. S., and T. F. C. MacKay. 1996, *Introduction to Quantitative Genetics*. Harlow,
515 England, Prentice Hall.

516 Gilmour, A. R., B. J. Gogel, B. R. Cullis, and R. Thompson. 2009, *ASReml User Guide Release*
517 3.0. Hemel Hempstead, UK, VSN International Ltd.

518 Glor, R. E., J. J. Kolbe, R. Powell, A. Larson, and J. B. Losos. 2003. Phylogenetic analysis of
519 ecological and morphological diversification in Hispaniolan trunk-ground anoles (*anolis*
520 *cybotes* group). *Evolution* 57:2383-2397.

521 Grant, P. R., and B. R. Grant. 1995. Predicting microevolutionary responses to directional
522 selection on heritable variation. *Evolution* 49:241-251.

523 Hadfield, J. D., A. Nutall, D. Osorio, and I. P. F. Owens. 2007. Testing the phenotypic gambit:
524 phenotypic, genetic and environmental correlations of colour. *Journal of Evolutionary*
525 *Biology* 20:549-557.

526 Hansen, T. F., and D. Houle. 2008. Measuring and comparing evolvability and constraint in
527 multivariate characters. *Journal of Evolutionary Biology* 21:1201-1219.

528 Harmon, L. J., and R. E. Glor. 2010. Poor statistical performance of the Mantel test in
529 phylogenetic comparative analyses. *Evolution* 64:2173-2178.

530 Harmon, L. J., J. J. Kolbe, J. M. Cheverud, and J. B. Losos. 2005. Convergence and the
531 multidimensional niche. *Evolution* 59:409-421.

532 Hine, E., S. F. Chenoweth, H. D. Rundle, and M. W. Blows. 2009. Characterizing the evolution
533 of genetic variance using genetic covariance tensors. *Philosophical Transactions of the*
534 *Royal Society B-Biological Sciences* 364:1567-1578.

535 Houle, D., G. H. Bolstad, K. van der Linde, and T. F. Hansen. 2017. Mutation predicts 40 million
536 years of fly wing evolution. *Nature* 548:447-+.

537 Houle, D., and K. Meyer. 2015. Estimating sampling error of evolutionary statistics based on
538 genetic covariance matrices using maximum likelihood. *Journal of Evolutionary Biology*
539 28:1542-1549.

540 Irschick, D. J., and J. B. Losos. 1998. A comparative analysis of the ecological significance of
541 maximal locomotor performance in Caribbean *Anolis* lizards. *Evolution* 52:219-226.

542 Jones, A. G., S. J. Arnold, and R. Bürger. 2003. Stability of the **G**-matrix in a population
543 experiencing pleiotropic mutation, stabilizing selection, and genetic drift. *Evolution*
544 57:1747-1760.

545 —. 2004. Evolution and stability of the **G**-matrix on a landscape with a moving optimum.
546 *Evolution* 58:1639-1654.

547 Jones, A. G., R. Bürger, and S. J. Arnold. 2014. Epistasis and natural selection shape the
548 mutational architecture of complex traits. *Nature Communications* 5.

549 Jones, A. G., R. Bürger, S. J. Arnold, P. A. Hohenlohe, and J. C. Uyeda. 2012. The effects of
550 stochastic and episodic movement of the optimum on the evolution of the **G**-matrix and
551 the response of the trait mean to selection. *Journal of Evolutionary Biology* 25:2210-
552 2231.

553 Kim, S. 2015. ppcor: An R package for a fast calculation to semi-partial correlation coefficients.
554 *Commun Stat Appl Methods* 22:665-674.

555 Kolbe, J. J., R. E. Glor, L. R. G. Schettino, A. C. Lara, A. Larson, and J. B. Losos. 2004. Genetic
556 variation increases during biological invasion by a Cuban lizard. *Nature* 431:177-181.

557 Kolbe, J. J., M. Leal, T. W. Schoener, D. A. Spiller, and J. B. Losos. 2012. Founder effects
558 persist despite adaptive differentiation: a field experiment with lizards. *Science* 335:1086-
559 1089.

560 Kolbe, J. J., L. J. Revell, B. Székely, E. D. Brodie, and J. B. Losos. 2011. Convergent evolution
561 of phenotypic integration and its alignment with morphological diversification in
562 Caribbean *Anolis* ecomorphs. *Evolution* 65:3608-3624.

563 Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain:body
564 size allometry. *Evolution* 33:402-416.

- 565 —. 1980. The genetic covariance between characters maintained by pleiotropic mutations.
566 Genetics 94:203-215.
- 567 Losos, J. B. 1990. Ecomorphology, performance capability, and scaling of West Indian *Anolis*
568 lizards: an evolutionary analysis. Ecological Monographs 60:369-388.
- 569 —. 1994. Integrative approaches to evolutionary ecology: *Anolis* lizards as model systems.
570 Annual Review of Ecology and Systematics 25:467-493.
- 571 —. 2009, Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles. Berkeley,
572 Univ. of California Press.
- 573 —. 2011. Convergence, adaptation, and constraint. Evolution 65:1827-1840.
- 574 Losos, J. B., and D. J. Irschick. 1996. The effect of perch diameter on escape behaviour of *Anolis*
575 lizards: laboratory predictions and field tests. Animal Behaviour 51:593-602.
- 576 Losos, J. B., T. R. Jackman, A. Larson, K. de Queiroz, and L. Rodríguez-Schettino. 1998.
577 Contingency and determinism in replicated adaptive radiations of island lizards. Science
578 279:2115-2118.
- 579 Losos, J. B., T. W. Schoener, K. I. Warheit, and D. Creer. 2001. Experimental studies of adaptive
580 differentiation in Bahamian *Anolis* lizards. Genetica 112:399-415.
- 581 Losos, J. B., and B. Sinervo. 1989. The effects of morphology and perch diameter on sprint
582 performance of *Anolis* lizards. Journal of Experimental Biology 145:23-30.
- 583 Losos, J. B., K. I. Warheit, and T. W. Schoener. 1997. Adaptive differentiation following
584 experimental island colonization in *Anolis* lizards. Nature 387:70-73.
- 585 Mahler, D. L., T. Ingram, L. J. Revell, and J. B. Losos. 2013. Exceptional convergence on the
586 macroevolutionary landscape in island lizard radiations. Science 341:292-295.

587 Marroig, G., and J. M. Cheverud. 2001. A comparison of phenotypic variation and covariation
588 patterns and the role of phylogeny. Ecology, and ontogeny during cranial evolution of
589 new world monkeys. *Evolution* 55:2576-2600.

590 Martins, E. P., and T. F. Hansen. 1997. Phylogenies and the comparative method: a general
591 approach to incorporating phylogenetic information into the analysis of interspecific data.
592 *American Naturalist* 149:646-667.

593 McGlothlin, J. W., M. E. Kobiela, H. V. Wright, D. L. Mahler, J. J. Kolbe, J. B. Losos, and E. D.
594 Brodie, III. 2018. Adaptive radiation along a deeply conserved genetic line of least
595 resistance in *Anolis* lizards. *Evolution Letters* 2:310-322.

596 McGlothlin, J. W., P. G. Parker, V. Nolan, Jr., and E. D. Ketterson. 2005. Correlational selection
597 leads to genetic integration of body size and an attractive plumage trait in dark-eyed
598 juncos. *Evolution* 59:658-671.

599 McGuigan, K. 2006. Studying phenotypic evolution using multivariate quantitative genetics.
600 *Molecular Ecology* 15:883-896.

601 McGuigan, K., S. F. Chenoweth, and M. W. Blows. 2005. Phenotypic divergence along lines of
602 genetic variance. *American Naturalist* 165:32-43.

603 Melo, D., G. Garcia, A. Hubbe, A. P. Assis, and G. Marroig. 2015. EvolQG - An R package for
604 evolutionary quantitative genetics. *F1000Res* 4:925.

605 Merilä, J., and M. Björklund. 2004. Phenotypic integration as a constraint and adaptation, Pages
606 107-129 in M. Pigliucci, and K. Preston, eds. *Phenotypic Integration: Studying the*
607 *Ecology and Evolution of Complex Phenotypes*. Oxford, Oxford Univ. Press.

608 Paradis, E., J. Claude, and K. Strimmer. 2004. APE: Analyses of Phylogenetics and Evolution in
609 R language. *Bioinformatics* 20:289-290.

610 Phillips, P. C., and S. J. Arnold. 1989. Visualizing multivariate selection. *Evolution* 43:1209-
611 1222.

612 Poe, S., A. Nieto-Montes de Oca, O. Torres-Carvajal, K. de Queiroz, J. A. Velasco, B. Truett, L.
613 N. Gray et al. 2017. A phylogenetic, biogeographic, and taxonomic study of all extant
614 species of *Anolis* (Squamata; Iguanidae). *Systematic Biology* 66:663-697.

615 R Core Team. 2019. R: A language and environment for statistical computing. . Vienna, Austria,
616 R Foundation for Statistical Computing.

617 Rabinowitz, A. H., and S. A. Vokes. 2012. Integration of the transcriptional networks regulating
618 limb morphogenesis. *Developmental Biology* 368:165-180.

619 Revell, L. J. 2007. The **G** matrix under fluctuating correlational mutation and selection.
620 *Evolution* 61:1857-1872.

621 —. 2012. phytools: an R package for phylogenetic comparative biology (and other things).
622 *Methods in Ecology and Evolution* 3:217-223.

623 Revell, L. J., D. L. Mahler, J. R. Sweeney, M. Sobotka, V. E. Fancher, and J. B. Losos. 2010.
624 Nonlinear selection and the evolution of variances and covariances for continuous
625 characters in an anole. *Journal of Evolutionary Biology* 23:407-421.

626 Roff, D. A. 1997, *Evolutionary Quantitative Genetics*. New York, Chapman & Hall.

627 Roff, D. A., and D. J. Fairbairn. 2012. A test of the hypothesis that correlational selection
628 generates genetic correlations. *Evolution* 66:2953-2960.

629 Sanger, T. J., L. J. Revell, J. J. Gibson-Brown, and J. B. Losos. 2012. Repeated modification of
630 early limb morphogenesis programmes underlies the convergence of relative limb length
631 in *Anolis* lizards. *Proceedings of the Royal Society B-Biological Sciences* 279:739-748.

632 Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. *Evolution* 50:1766-
633 1774.

634 Sears, K. E., T. D. Capellini, and R. Diogo. 2015. On the serial homology of the pectoral and
635 pelvic girdles of tetrapods. *Evolution* 69:2543-2555.

636 Stepan, S. J., P. C. Phillips, and D. Houle. 2002. Comparative quantitative genetics: evolution of
637 the **G** matrix. *Trends in Ecology & Evolution* 17:320-327.

638 Steven, J. C., I. A. Anderson, E. D. Brodie III, and L. F. Delph. 2020. Rapid reversal of a
639 potentially constraining genetic covariance between leaf and flower traits in *Silene*
640 *latifolia*. *Ecol Evol* 10:569-578.

641 Tollis, M., E. D. Hutchins, J. Stapley, S. M. Rupp, W. L. Eckalbar, I. Maayan, E. Lasku et al.
642 2018. Comparative genomics reveals accelerated evolution in conserved pathways during
643 the diversification of anole lizards. *Genome Biology and Evolution* 10:489-506.

644 Turelli, M. 1988. Phenotypic evolution, constant covariances, and the maintenance of additive
645 variance. *Evolution* 42:1342-1347.

646 Walsh, B., and M. W. Blows. 2009. Abundant genetic variation plus strong selection =
647 multivariate genetic constraints: a geometric view of adaptation. *Annual Review of*
648 *Ecology Evolution and Systematics* 40:41-59.

649 Walter, G. M., J. D. Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018. Evolution of genetic
650 variance during adaptive radiation. *American Naturalist* 191:E108-E128.

651 Williams, E. E. 1972. The origin of faunas. Evolution of lizard congeners in a complex island
652 fauna: a trial analysis. *Evolutionary Biology* 6:47-89.

653 Wood, S. 2012. mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML smoothness
654 estimation.

655 Zheng, Y. C., and J. J. Wiens. 2016. Combining phylogenomic and supermatrix approaches, and
656 a time-calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes
657 and 4162 species. *Molecular Phylogenetics and Evolution* 94:537-547.

Appendix

Supplemental Tables

Table A1: **G** matrices and matrices of heritabilities (h^2 , diagonal) and genetic correlations (r_g , off-diagonal) for seven *Anolis* species. Approximate standard errors for are shown below each matrix. **G** matrices and standard errors are reprinted from McGlothlin et al. (2018). Although we did not conduct formal likelihood-ratio tests, parameters that exceeded their standard errors by a factor of two are shown in bold, which provides a guide to statistical significance. Traits are abbreviated as follows: JL = jaw length, HW = head width, PECT = pectoral width, PELV = pelvic width, HUM = humerus, UL = ulna, FEM = femur, and TIB = tibia. All traits were natural-log transformed and size-corrected for analysis.

Anolis cristatellus (Trunk-ground, Puerto Rico)

G ($\times 10^{-3}$)

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.329							
HW	.094	.449						
PECT	-.083	.349	1.426					
PELV	-.089	.240	.487	.546				
HUM	.293	.071	-.371	-.168	1.441			
UL	.079	.075	-.098	.017	1.008	1.087		
FEM	.208	.030	-.053	-.051	.904	.731	.809	
TIB	.268	.009	-.172	-.081	.945	.780	.783	.949

SE ($\times 10^{-3}$)

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.052							
HW	.029	.048						
PECT	.053	.066	.170					
PELV	.056	.049	.093	.113				
HUM	.057	.067	.128	.090	.200			
UL	.053	.063	.119	.087	.144	.175		
FEM	.035	.042	.077	.058	.093	.085	.071	
TIB	.037	.043	.081	.060	.096	.089	.067	.076

Anolis cristatellus (continued)

h^2/r_g

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.458							
HW	.245	.353						
PECT	-.122	.435	.306					
PELV	-.210	.484	.552	.196				
HUM	.425	.088	-.259	-.190	.231			
UL	.132	.108	-.078	.022	.805	.185		
FEM	.403	.049	-.049	-.076	.837	.779	.482	
TIB	.480	.014	-.148	-.113	.808	.768	.894	.561

h^2/r_g SE

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.057							
HW	.068	.028						
PECT	.079	.073	.029					
PELV	.130	.093	.092	.037				
HUM	.081	.082	.084	.101	.028			
UL	.091	.090	.094	.111	.057	.027		
FEM	.066	.068	.072	.087	.049	.058	.025	
TIB	.063	.066	.069	.085	.051	.057	.017	.023

Anolis evermanni (Trunk-crown, Puerto Rico)

$G (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.251							
HW	-.013	.314						
PECT	.016	.411	1.062					
PELV	-.017	.183	.435	.623				
HUM	.143	.022	.051	.139	.866			
UL	.141	-.026	.097	.145	.884	.971		
FEM	.070	.040	.228	.279	.850	.922	1.224	
TIB	.121	.047	.178	.258	.971	1.064	1.427	1.751

$SE (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.029							
HW	.027	.049						
PECT	.048	.069	.163					
PELV	.040	.054	.099	.146				
HUM	.049	.061	.110	.091	.160			
UL	.049	.062	.113	.093	.135	.156		
FEM	.041	.054	.098	.083	.108	.110	.117	
TIB	.048	.064	.114	.097	.124	.128	.125	.174

h^2/r_g

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.415							
HW	-.047	.244						
PECT	.031	.712	.280					
PELV	-.043	.414	.535	.227				
HUM	.306	.042	.053	.189	.194			
UL	.286	-.046	.096	.186	.964	.212		
FEM	.126	.064	.200	.319	.825	.846	.599	
TIB	.182	.064	.131	.247	.788	.816	.975	.704

$h^2/r_g SE$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.033							
HW	.097	.032						
PECT	.094	.077	.035					
PELV	.101	.108	.106	.048				
HUM	.096	.117	.117	.122	.032			
UL	.092	.113	.111	.117	.044	.029		
FEM	.072	.087	.083	.089	.066	.053	.030	
TIB	.069	.085	.083	.089	.071	.057	.011	.034

Anolis grahami (Trunk-crown, Jamaica)

$G (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.246							
HW	.218	.296						
PECT	.127	.122	.351					
PELV	.193	.126	.173	.704				
HUM	.152	.067	.167	.264	1.003			
UL	.129	.131	.122	.130	.856	1.171		
FEM	.126	-.005	.081	.186	.625	.559	.510	
TIB	.146	.034	.171	.269	.732	.724	.549	.648

$SE (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.049							
HW	.051	.082						
PECT	.064	.081	.151					
PELV	.068	.084	.113	.169				
HUM	.140	.186	.238	.176	.465			
UL	.103	.133	.177	.186	.323	.405		
FEM	.052	.099	.139	.111	.230	.191	.170	
TIB	.056	.102	.145	.104	.230	.181	.163	.179

h^2/r_g

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.421							
HW	.807	.240						
PECT	.434	.377	.133					
PELV	.465	.277	.349	.277				
HUM	.307	.123	.281	.315	.168			
UL	.240	.222	.191	.143	.790	.159		
FEM	.355	-.013	.191	.311	.873	.723	.453	
TIB	.367	.076	.358	.399	.908	.831	.955	.544

$h^2/r_g SE$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.056							
HW	.092	.056						
PECT	.190	.229	.054					
PELV	.134	.173	.210	.054				
HUM	.266	.341	.392	.192	.072			
UL	.188	.221	.281	.199	.155	.050		
FEM	.136	.249	.323	.171	.165	.160	.122	
TIB	.127	.231	.289	.134	.165	.124	.037	.110

Anolis lineatopus (Trunk-ground, Jamaica)

G ($\times 10^{-3}$)

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.442							
HW	.286	.263						
PECT	.385	.290	.750					
PELV	.193	.132	.343	.916				
HUM	.321	.240	.223	-.016	.968			
UL	.412	.297	.308	-.200	.942	1.047		
FEM	.423	.180	.547	.228	.530	.540	1.013	
TIB	.368	.145	.528	.182	.464	.495	.864	.825

SE ($\times 10^{-3}$)

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.103							
HW	.059	.090						
PECT	.101	.100	.309					
PELV	.158	.139	.254	.352				
HUM	.127	.103	.213	.200	.291			
UL	.106	.104	.186	.174	.240	.270		
FEM	.168	.130	.233	.256	.255	.248	.346	
TIB	.147	.117	.212	.227	.226	.222	.295	.272

h^2/r_g

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.536							
HW	.837	.177						
PECT	.668	.652	.151					
PELV	.303	.269	.414	.253				
HUM	.491	.475	.262	-.017	.161			
UL	.605	.566	.348	-.204	.936	.169		
FEM	.633	.349	.627	.237	.535	.525	.645	
TIB	.610	.311	.672	.209	.519	.533	.945	.587

h^2/r_g SE

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.089							
HW	.114	.056						
PECT	.158	.206	.059					
PELV	.217	.265	.263	.087				
HUM	.167	.196	.251	.223	.044			
UL	.118	.178	.223	.177	.071	.040		
FEM	.143	.237	.202	.232	.201	.196	.155	
TIB	.146	.238	.188	.233	.206	.196	.029	.143

Anolis pulchellus (Grass-bush, Puerto Rico)

$G (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.084							
HW	.080	.157						
PECT	.222	.215	.895					
PELV	.070	.065	.300	.766				
HUM	.148	.040	.153	-.036	.745			
UL	.180	.080	.214	-.152	.773	.982		
FEM	.092	.040	.156	.126	.475	.524	.701	
TIB	.089	.030	.136	.134	.403	.524	.648	.723

$SE (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.050							
HW	.040	.059						
PECT	.075	.070	.268					
PELV	.066	.061	.192	.244				
HUM	.072	.068	.163	.127	.204			
UL	.082	.076	.155	.145	.197	.254		
FEM	.054	.053	.110	.083	.133	.118	.156	
TIB	.054	.045	.087	.082	.141	.143	.148	.168

h^2/r_g

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.076							
HW	.701	.124						
PECT	.810	.574	.211					
PELV	.276	.187	.363	.184				
HUM	.594	.118	.188	-.047	.106			
UL	.629	.204	.228	-.175	.905	.116		
FEM	.382	.122	.196	.172	.658	.632	.420	
TIB	.362	.090	.170	.181	.549	.622	.911	.476

$h^2/r_g SE$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.044							
HW	.218	.044						
PECT	.237	.153	.057					
PELV	.259	.176	.191	.054				
HUM	.299	.201	.196	.167	.027			
UL	.295	.192	.164	.160	.066	.028		
FEM	.211	.158	.134	.113	.127	.113	.074	
TIB	.214	.134	.108	.112	.159	.130	.038	.086

Anolis sagrei (Trunk-ground, Bahamas)

$G (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.292							
HW	.112	.302						
PECT	-.002	.166	.286					
PELV	-.077	-.018	.035	.463				
HUM	.156	-.073	.027	-.109	.603			
UL	.027	-.142	-.047	.082	.462	.646		
FEM	.192	-.033	-.027	-.011	.409	.315	.368	
TIB	.177	-.111	-.036	-.003	.527	.507	.434	.606

$SE (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.023							
HW	.021	.033						
PECT	.024	.030	.048					
PELV	.050	.055	.043	.073				
HUM	.035	.050	.050	.068	.092			
UL	.037	.044	.053	.062	.080	.116		
FEM	.024	.037	.042	.041	.051	.051	.054	
TIB	.026	.044	.051	.042	.058	.061	.056	.073

h^2/r_g

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.469							
HW	.377	.269						
PECT	-.007	.566	.127					
PELV	-.211	-.049	.095	.147				
HUM	.372	-.171	.065	-.206	.133			
UL	.063	-.321	-.110	.149	.740	.130		
FEM	.585	-.100	-.085	-.027	.869	.647	.270	
TIB	.421	-.259	-.087	-.006	.873	.811	.919	.449

$h^2/r_g SE$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.023							
HW	.059	.024						
PECT	.087	.083	.020					
PELV	.134	.146	.114	.021				
HUM	.077	.116	.118	.126	.019			
UL	.084	.097	.123	.112	.069	.022		
FEM	.059	.112	.131	.097	.055	.083	.034	
TIB	.054	.097	.122	.084	.048	.060	.026	.041

Anolis smaragdinus (Trunk-crown, Bahamas)

$G (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.348							
HW	.054	.161						
PECT	.251	.065	.830					
PELV	.102	-.002	.442	.560				
HUM	.268	.151	.130	.122	.395			
UL	.238	.094	.249	.300	.331	.452		
FEM	.205	.061	.136	.115	.285	.287	.344	
TIB	.296	.008	.221	.278	.351	.423	.420	.609

$SE (\times 10^{-3})$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.129							
HW	.091	.139						
PECT	.171	.174	.413					
PELV	.072	.068	.144	.142				
HUM	.130	.093	.163	.122	.280			
UL	.098	.134	.176	.175	.181	.286		
FEM	.114	.085	.184	.127	.149	.149	.165	
TIB	.125	.072	.205	.141	.120	.132	.158	.197

h^2/r_g

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.455							
HW	.227	.121						
PECT	.468	.176	.254					
PELV	.230	-.007	.648	.251				
HUM	.722	.600	.228	.259	.100			
UL	.601	.348	.406	.596	.783	.090		
FEM	.592	.257	.255	.262	.773	.728	.283	
TIB	.644	.025	.311	.477	.717	.806	.919	.454

$h^2/r_g SE$

	JL	HW	PECT	PELV	HUM	UL	FEM	TIB
JL	.137							
HW	.353	.101						
PECT	.248	.450	.116					
PELV	.151	.254	.179	.053				
HUM	.298	.421	.302	.261	.068			
UL	.235	.481	.296	.306	.290	.055		
FEM	.207	.348	.337	.274	.307	.311	.122	
TIB	.154	.223	.279	.205	.269	.243	.094	.118

Table A2: Random skewers correlations (r_{RS}) for each species pair. Point estimates, REML-MVN estimates, and 95% confidence intervals are shown. Point estimates occasionally lie outside the REML-MVN confidence interval because incorporating estimation error from two **G** matrices lead to negatively skewed distributions.

species 1	species 2	r_{RS}	REML-MVN	r_{RS}	2.5%	97.5%
CRIS	EVER	.89		.86	.80	.91
CRIS	GRAH	.84		.74	.59	.84
CRIS	LINE	.78		.70	.55	.81
CRIS	PULC	.87		.81	.71	.88
CRIS	SAGR	.88		.85	.79	.89
CRIS	SMAR	.81		.66	.43	.82
EVER	GRAH	.83		.74	.60	.84
EVER	LINE	.81		.73	.55	.83
EVER	PULC	.90		.84	.74	.91
EVER	SAGR	.84		.81	.75	.87
EVER	SMAR	.82		.68	.47	.81
GRAH	LINE	.85		.70	.52	.83
GRAH	PULC	.88		.75	.60	.86
GRAH	SAGR	.87		.76	.62	.86
GRAH	SMAR	.79		.61	.38	.77
LINE	PULC	.92		.79	.64	.89
LINE	SAGR	.76		.67	.52	.78
LINE	SMAR	.81		.63	.39	.79
PULC	SAGR	.82		.76	.66	.83
PULC	SMAR	.84		.66	.44	.82
SAGR	SMAR	.76		.63	.42	.77

Table A3: Estimates of phylogenetic signal (Blomberg's K) for all genetic variances and correlations. Point estimates, REML-MVN estimates, and 95% confidence intervals are shown. All estimates of K were significantly different from 0 and indistinguishable from 1.

Variances

Trait	K	REML-MVN K	2.5%	97.5%
JL	.81	.86	.65	1.11
HW	.69	.77	.61	1.05
PECT	1.16	1.13	.79	1.53
PELV	.90	.90	.69	1.18
RAD	.87	.90	.69	1.20
UL	1.15	1.03	.77	1.31
FEM	.99	1.00	.83	1.31
TIB	.90	.91	.77	1.13

Correlations

Trait 1	Trait 2	K	REML-MVN K	2.5%	97.5%
JL	HW	.92	.96	.74	1.26
JL	PECT	.71	.76	.64	1.00
JL	PELV	.88	.91	.67	1.25
JL	HUM	.91	.93	.69	1.20
JL	UL	.78	.84	.67	1.12
JL	FEM	1.04	1.00	.73	1.32
JL	TIB	.95	.92	.73	1.17
HW	PECT	.97	.97	.74	1.19
HW	PELV	1.04	1.01	.71	1.39
HW	HUM	1.04	1.01	.82	1.20
HW	UL	.97	.99	.84	1.14
HW	FEM	.91	.97	.78	1.20
HW	TIB	.97	.97	.82	1.16
PECT	PELV	1.02	.98	.74	1.30
PECT	HUM	.85	.92	.65	1.30
PECT	UL	.92	.94	.71	1.19
PECT	FEM	.91	.95	.77	1.18
PECT	TIB	1.02	1.01	.80	1.24
PELV	HUM	.90	.92	.75	1.15
PELV	UL	1.03	1.00	.77	1.19
PELV	FEM	.81	.89	.71	1.11
PELV	TIB	.93	.94	.72	1.18
HUM	UL	.90	.99	.76	1.22
HUM	FEM	.78	.92	.72	1.17
HUM	TIB	.76	.87	.71	1.13
UL	FEM	.87	.98	.71	1.31
UL	TIB	.80	.91	.71	1.17
FEM	TIB	.79	.93	.72	1.14

Table A4: Ecomorph effects given as evolutionary correlations (r_e) for all genetic variances and correlations. A positive value indicates that trunk-ground species have a higher value than trunk-crown species. Point estimates, REML-MVN estimates, and 95% confidence intervals are shown. Significant values are indicated with boldface.

Variances

Trait	r_e	REML-MVN r_e	2.5%	97.5%
JL	.62	.48	-.17	.88
HW	.52	.39	-.26	.84
PECT	.23	.19	-.27	.56
PELV	.03	-.04	-.85	.58
RAD	.44	.36	-.24	.78
UL	.09	.08	-.59	.70
FEM	.01	-.02	-.49	.34
TIB	-.35	-.34	-.65	.02

Correlations

Trait 1	Trait 2	r_e	REML-MVN r_e	2.5%	97.5%
JL	HW	.28	.25	-.13	.52
JL	PECT	-.23	-.21	-.60	.22
JL	PELV	-.56	-.49	-.84	-.03
JL	HUM	.09	.07	-.64	.79
JL	UL	-.20	-.17	-.64	.32
JL	FEM	.63	.55	.05	.90
JL	TIB	.49	.43	-.06	.84
HW	PECT	.24	.11	-.73	.57
HW	PELV	.03	.02	-.60	.58
HW	HUM	-.13	-.09	-.63	.52
HW	UL	-.01	.00	-.47	.50
HW	FEM	.08	.04	-.68	.64
HW	TIB	-.07	-.09	-.75	.48
PECT	PELV	-.37	-.27	-.79	.36
PECT	HUM	-.55	-.35	-.86	.39
PECT	UL	-.44	-.33	-.85	.35
PECT	FEM	-.11	-.10	-.70	.42
PECT	TIB	-.24	-.21	-.68	.28
PELV	HUM	-.96	-.78	-.98	-.32
PELV	UL	-.68	-.58	-.89	-.06
PELV	FEM	-.82	-.62	-.97	-.01
PELV	TIB	-.87	-.74	-.97	-.25
HUM	UL	-.16	-.14	-.82	.46
HUM	FEM	-.37	-.21	-.72	.59
HUM	TIB	-.33	-.21	-.70	.59
UL	FEM	-.64	-.35	-.86	.41
UL	TIB	-.58	-.37	-.82	.45
FEM	TIB	-.65	-.45	-.94	.20