Thermally tolerant symbionts may explain Caribbean octocoral resilience to heat stress

Jessie Pelosi^{1, #, *}, Katherine M. Eaton^{1, ^, *}, Samantha Mychajliw², Casey P. terHorst³, and Mary Alice Coffroth^{4, +}

¹Department of Biological Sciences, University at Buffalo, Buffalo, NY 14260, ²Department of Environment and Sustainability, University at Buffalo, Buffalo, NY 14260, ³Department of Biology, California State University, Northridge, Northridge, CA 91330, ⁴Department of Geology, University at Buffalo, Buffalo, NY 14260

*These authors contributed equally to this work.

Current addresses:

[#]Department of Biology, University of Florida, Gainesville, FL 32608

[^]Department of Biological Sciences, Auburn University, Auburn, AL 36849

⁺Corresponding author: coffroth@buffalo.edu

Keywords: symbiosis, Symbiodiniaceae, octocorals, thermal tolerance

Abstract

Coral reef ecosystems are under threat from the frequent and severe impacts of anthropogenic climate change, particularly rising sea surface temperatures. The effects of thermal stress may be ameliorated by adaptation and/or acclimation of the host, symbiont, or holobiont (host + symbiont) to increased temperatures. We examined the role of the symbiont in promoting thermal tolerance of the holobiont, using *Antillologorgia bipinnata* (octocoral host) and *Breviolum antillologorgium* (symbiont) as a model system. We identified five distinct genotypes of *B. antillologorgium* from symbiont populations isolated from *Antillologorgia* colonies in the Florida Keys. Three symbiont genotypes were cultured and maintained at 26°C (ambient historical temperature) and two were cultured and maintained at 30°C (elevated historical temperature) for two years. Following culturing, we analyzed the growth rate and carrying capacity of each symbiont genotype at both ambient and elevated temperatures in culture (*in vitro*). All genotypes grew well at both temperatures, indicating thermal tolerance among these *B. antillologorgium* cultures. Prior culturing at the elevated temperature, however, did not result in increased thermal tolerance. We then inoculated juvenile *A. bipinnata* polyps with each of the five symbiont genotypes, and reared these polyps at both ambient and elevated temperatures (*in hospite* experiment). All genotypes were able to establish symbioses with polyps in both temperature treatments. Survivorship of polyps at 30°C was significantly lower than survivorship at 26°C, but all treatments had surviving polyps at 56 days post-infection, suggestive of broad thermal tolerance in *B. antillologorgium*, which may play a part in the increased resilience of Caribbean octocorals during heat stress events.

Introduction

In recent years, anthropogenic climate change has severely impacted coral reefs, which comprise some of the most biodiverse and economically important ecosystems on Earth (Hoegh-Guldberg 2011; Hoegh-Guldberg et al. 2017). The success of reef ecosystems in oligotrophic tropical waters is due to their obligate endosymbioses with dinoflagellate algae in the family Symbiodiniaceae (Muscattine and Porter 1977). In this mutualistic relationship, the coral host relies on photosynthate produced by its intracellular symbionts to thrive (Muscattine and Cernichiaro 1969; Muscattine and Porter 1977). In exchange for these metabolites, the symbiotic algae recover host nitrogenous waste. This relationship, which is vital to the survival of the reef ecosystem, is currently threatened by anthropogenic environmental changes.

Coral bleaching, or the loss of algal symbionts, occurs due to a variety of environmental stressors including changes in salinity, pH, light intensity, and, most commonly, water temperature (Hoegh-Guldberg 1999). Most reef ecosystems now routinely encounter temperatures at or above their thermal maximum each year (Hoegh-Guldberg et al. 2007; Eakin et al. 2009), with bleaching events becoming more frequent and severe (Heron et al. 2016; van Hooidonk et al. 2016; Hughes et al. 2018; Oliver et al. 2018). As ocean temperatures continue to increase, it is essential to understand if and how corals and their symbionts can respond to the present threat.

The holobiont (host + symbiont) can respond to rising seawater temperatures via changes in the symbiont community within a single host, with the basic premise that hosts associating with more thermally tolerant symbionts will be more likely to survive bleaching events (i.e., shuffling and/or switching, *sensu* Baker 2003). Although most coral species host a single

dominant symbiont type (Goulet 2006; LaJeunesse 2020), many hosts also harbor background or cryptic symbionts that are often undetected by traditional techniques (Mieog et al. 2007; Correa et al. 2009; Silverstein et al. 2012). During thermal stress, a more thermally tolerant background symbiont can become the dominant symbiont type, potentially conferring thermal tolerance to the holobiont (Berkelmans and van Oppen 2006; Jones et al. 2008; LaJeunesse et al. 2009; Keshavmurthy et al. 2012; Silverstein et al. 2015). In the aforementioned studies, the background symbionts often differed at the genus level and were typically replaced by the previously dominant symbiont type once the stress had subsided.

Thermally tolerant holobionts may arise through adaptation of symbiont populations, even in the absence of evolutionary changes in the host. Large population sizes, short generation times, and high mutation rates could greatly augment the standing genetic variation within symbiont populations (van Oppen et al. 2011). With greater standing variation, natural selection is more likely to produce populations with greater thermal tolerance (van Oppen et al. 2011; Torda et al. 2017; Quigley et al. 2018; Buerger et al. 2020). Numerous studies have established that more thermally tolerant symbionts can be selected from within a single symbiont genus or species (Howells et al. 2011; Chakravarti et al. 2017; Buerger et al. 2020). By sampling symbionts in the genus *Cladocopium* (ITS type C1) from reefs with different temperature regimes, Howells et al. (2011) showed that distinct symbiont genotypes had different thermal tolerances. Symbionts from the warmer reef displayed increased performance and survival at elevated temperatures as compared to those from the cooler environment. Furthermore, juvenile hosts inoculated with symbionts from the warmer reef performed better (grew faster, had less bleaching, and had lower mortality) than those inoculated with symbionts from the cooler region.

Chakravarti et al. (2017) and Buerger et al. (2020) presented evidence for selection within a species by examining the adaptive capabilities of the symbionts within a monoclonal culture of *Cladocopium goreau*. Symbionts selected for thermal tolerance (i.e., those maintained at 31°C for 2.5 to 4 years) performed better at 31°C than the original monoclonal culture that had not undergone this selection treatment. Juvenile polyps infected with symbiont cells from the elevated temperature grew at 27°C and 31°C, while polyps infected with the original culture showed negative growth at 31°C. Infection with some strains of thermally selected cells, however, did not reduce the rate or severity of bleaching compared to those infected with the original culture (Buerger et al. 2020). These studies establish that selection for thermal tolerance can occur over ecologically relevant time scales within at least some symbiont populations. Given sufficient standing genetic variation, this is likely the case for other species within Symbiodiniaceae and requires further inquiry (Chakravarti et al. 2017).

By starting with a single monoclonal culture, all genetic variation in the symbionts used by Chakravarti et al. (2017) and Buerger et al. (2020) likely arose from *de novo* mutation. Intraspecific markers have revealed numerous instances where host colonies harbor multiple genotypes within a symbiont species (Santos et al. 2003; Pettay and LaJeunesse 2007; Howells et al. 2009; Kirk et al. 2009), suggesting the possibility for selection on pre-existing genetic variation within *in hospite* populations, resulting from both mutation and recombination (Baums et al. 2014; Chi et al. 2014; Wilkinson et al. 2015; Thornhill et al. 2017). Indeed, many studies have found variation in thermal tolerance within a single symbiont species (Parkinson and Baums 2014; Karim et al. 2015; Díaz-Almeyda et al. 2017; Grégoire et al. 2017; Bayliss et al. 2019). For example, Bayliss et al. (2019) identified genetic and trait variation within and among *Breviolum* species isolated from the octocoral *Antilloporgia bipinnata*.

Here we ask if there are thermally tolerant genotypes present in a symbiont species within a single host colony. We utilized the octocoral *A. bipinnata* and its host-specialist endosymbiotic algae, *Breviolum antillogorgium*, as a model to study the role of symbiont genetic variation in the thermal response of the holobiont. We first selected for thermally tolerant symbionts from *in hospite* populations to capitalize on pre-existing genetic variation. We then examined symbiont growth at two temperatures with an *in vitro* experiment using these cultured isolates of *B. antillogorgium*. Subsequently, we examined whether thermal tolerance in the symbiont could be conferred to the host using an *in hospite* experiment. The presence of thermally tolerant symbionts within a population contained in a single host colony expands the potential for the holobiont to ameliorate the consequences of increased temperatures, potentially slowing the loss of these invaluable ecosystems.

Methods

Symbiont isolation

Although culturing of Symbiodiniaceae is highly selective and rarely recovers the symbionts dominant within the host (Santos et al. 2001; LaJeunesse 2002), *Breviolum antillogorgium* (ITS2-type B1) has been successfully cultured and associates specifically with octocorals in the genus *Antillogorgia* (Santos et al. 2004; Parkinson et al. 2015). Symbiont cells were isolated from sixteen adult *Antillogorgia* colonies collected from 12-18 m depth at sites in Elbow and Pickles Reefs, Florida Keys in September 2016 (Table 1). A sample of each host colony was preserved in 95% ethanol for comparison of dominant host symbionts with those isolated. An approximately 2 cm piece was sampled from each colony and homogenized in 2 mL of filtered seawater (FSW). Each homogenate was passed through a series of mesh filters (250,

120, 70 and 20 μm), followed by several 1 mL 0.22 μm FSW rinses for a final volume of 3-6 mL. Homogenates were spun at 800 rpm for five minutes. Supernatants were decanted, and the pellets were resuspended in 10 mL of FSW. This was repeated three additional times, followed by resuspending each pellet in 1 mL of f/2 media (Guillard and Ryther 1962). For each colony, 50 μL of the resultant cell suspension was added to six flasks containing 30 mL of f/2 media. Three flasks of the symbionts from each colony were maintained at 26°C, mimicking typical thermal conditions, and three were immediately placed at 30°C to select for thermally tolerant genotypes that can survive at an elevated temperature. An elevated temperature of 30°C was chosen based on the historical thermal regime of the collection site during November/December, when newly settled *A. bipinnata* polyps acquire symbionts (Williams and Miller 2015; Online Resource 1). Initial growth, observed as a small dot at bottom of the flask, was immediately transferred into a new flask with fresh media. This was repeated each time new growth was observed in any flask over the first month, resulting in 131 cultures. All cultures were transferred to fresh media monthly.

Genotype identification

After two years (approximately 735-1084 asexual generations, assuming a doubling time of 1.01-1.97 days, based on growth rates from this study), total symbiont genomic DNA from 106 cultures and the host source colonies was extracted using the 2X CTAB method (Coffroth et al. 1992) and diluted to 5-10 ng μL^{-1} , and the flanking region of the B7Sym15 microsatellite (Pettay and LaJeunesse 2007) was sequenced (TACGEN, Richmond, California, USA). Sequences were compared to available sequences on NCBI using BLASTn for symbiont species-level identification.

Cultures identified as *B. antillogorgium* (n=24) and samples from the source colonies of these cultures were amplified with five microsatellite loci following published protocols: CA 6.38 (Santos et al. 2003), GV_1C (Andras et al. 2009), B7Sym8, B7Sym34, and B7Sym36 (Pettay and LaJeunesse 2007). Amplicons were labeled with fluorescently tagged primers and visualized on a 6.5% polyacrylamide gel with the LI-COR NEN Global IR2 DNA Sequencer (LI-COR Biotechnology Division) following Santos et al. (2003). Allele sizes were scored by eye relative to size standards between 50 and 350 bp. Based on the microsatellite results, 21 *B. antillogorgium* cultures were unialgal and contained one of five genotypes. Ultimately, this resulted in the occurrence of three distinct genotypes (referred to here as G1, G2, and G3) from one or more of the cultures reared at 26°C and two genotypes (referred to here as G4 and G5) from one or more of the cultures reared at 30°C (see Results). One of each of these five genotypes were used in the experiments outlined below (Table 1).

***In vitro* growth experiments**

Six replicates of each of the five cultures identified above were started by inoculating 18 mL of f/2 media with 10,000 cells mL⁻¹. Three of the replicates for each culture were subsequently grown at 25.53 ± 0.89°C (mean ± SD; referred to as “26°C”) and three were grown at 29.84 ± 0.48°C (mean ± SD; referred to as “30°C”) on a 12:12 h light:dark cycle. Light levels were maintained at 120.58 ± 11.14 μmol m⁻²s⁻¹ (mean ± SD) (Philips Energy Advantage T5 HO 49W) and 103.79 ± 37.3 μmol m⁻²s⁻¹ (mean ± SD) (Philips T5 HO 24W/840) for the 26°C and 30°C incubators, respectively (Welch’s Two Sample t-test, p = 0.13). Cell densities were measured using six replicate hemocytometer cell counts every three days. Experiments were continued until cell density peaked and exhibited a consistent decline in abundance. We used the ‘growthrates’ package ver. 0.8.1 (Petzoldt 2019) in R ver. 4.0.2 (R Core Team 2020) to fit

logistic growth curves to the cell counts over time, up to the maximum cell count for each culture. We extracted values of r and K from the ‘growthrates’ package and used these as estimates of maximum population growth rate and carrying capacity, respectively. Samples of each replicate were then preserved in 95% ethanol, from which DNA was extracted and amplified following the methods detailed above to verify symbiont genotype at the end of the experiment.

***In hospite* polyp experiments**

Branches from adult colonies of *A. bipinnata*, a gonochoric surface brooder (Kahng et al. 2011), were collected from Tennessee Reef, Florida Keys in November 2018 (N 24° 45.150' W 81° 45.275'). These branches were transported to flow-through seawater tanks at Keys Marine Laboratory, where they were monitored daily for release of embryos. Aposymbiotic brooded larvae were collected from the colony surface on 16, 17, 22, 23, 25, and 26 November, washed three times in 0.22 μ m filtered instant ocean (FIO), and transferred to Buffalo, NY. Following transport, larvae (4-14 days old) were again washed in FIO to remove environmental contaminants and groups of 25 larvae were transferred to pre-conditioned polypropylene containers (volume = approximately 115 mL) with 50 mL of FIO. For pre-conditioning, the bottom and sides of each container were roughened with sandpaper and then soaked for four weeks in 1.2 μ m filtered seawater (FSW) to establish a biofilm. Once larvae were added, water in the containers (FIO) was changed daily for one week, then every other day until larvae settled and metamorphosed into polyps (approximately 4 weeks). Containers of newly settled polyps (approximately 1-2 weeks post-settlement) were then divided into five treatments (G1 through G5, Online Resource 2) and each treatment was inoculated with 1000 cells mL⁻¹ of one of the five genotypes of *B. antillogorgium* utilized in the *in vitro* experiments (Table 1). For each

genotype, 9-10 containers were placed at 26°C ($25.51 \pm 1.10^\circ\text{C}$; mean $\pm$ SD) and 6-8 at 30°C ($29.79 \pm 0.28^\circ\text{C}$; mean $\pm$ SD) and maintained in a 12:12 h light:dark cycle. Initial inoculations occurred twice over a period of one week. Polyp locations in the containers were mapped and survival for each individual polyp was recorded every 3-4 days with water changes. Visual infection, indicated by a light brown coloration in the polyp, was limited within the first month following initial introduction of symbionts. To increase the likelihood of the establishment of the symbiosis, inoculations were repeated with each water change during the second month and combined with feeding with Reefs-Roids (Polyplab, Quebec, Canada) (2 mL of Reef-Roids solution at a concentration of 1 mg/mL FIO) to induce symbiont uptake. Inoculations continued throughout the second month, and establishment of the symbiosis through visual observation was recorded throughout the experiment. An unidentified green alga, first observed in three containers on Day 42, but subsequently found in several additional containers on Days 56-59, led to extensive polyp mortality over time. After 69 days, all surviving polyps were collected, symbiont cell density was determined (as detailed above), and tissue was preserved in 95% ethanol. Symbiont DNA was extracted from preserved tissue and symbiont genotype was verified using the five microsatellite loci described above.

Statistical analyses

We analyzed the effects of temperature treatment and genotype on *in vitro* symbiont maximum population growth rates and carrying capacity using separate two-way ANOVAs for each dependent variable (r and K), including the interaction term between these factors (temperature and genotype). Data were log-transformed to meet assumptions of homoscedasticity for the *in vitro* analyses. For the *in hospite* experiment, we quantified polyp survivorship as the proportion of surviving polyps after 31 and 52 days post-initial inoculation.

These time points were prior to the observation of widespread contamination by the unidentified green alga, to avoid the possibility that this affected survivorship. Containers in which the contaminating algae was observed prior to Day 52 (n=5) were excluded from survivorship analyses beginning at the first observation of contamination. We analyzed survivorship over the first 31 days and then from Day 32 to Day 52 as described above; with survivorship for Days 1-31 arcsine transformed to meet assumptions of normality (survivorship data for Days 32-52 were not transformed as the raw data fit the assumptions of the model without transformation). We analyzed temperature and genotype effects on visually observed infection at Days 31 and 52 as well as on symbiont cell densities at the end of the experiment (Day 69) as above. Symbiont cell densities were square-root transformed to meet assumptions of the model. Assumptions of normality and homogeneity were checked using residuals and Q-Q probability plots. All post-hoc pairwise comparisons were carried out using Tukey's HSD (honest significance test) (Tukey 1949). *In vitro* statistical analyses were conducted in R ver. 4.0.2 and *in hospite* analyses in R ver. 3.6.1 (R Core Team 2020). R scripts are available at www.github.com/jessiepelosi/symbiont_tolerance and raw data are available at <https://www.bco-dmo.org/award/658940>.

Results

Sequencing and genotyping

Following two years of growth at either 26°C or 30°C, there were 131 viable cultures. The majority of cultures (106 of 131) either could not be classified or aligned most closely with *B. minutum*, a species commonly recovered from a range of species where it is undetected by normal techniques (LaJeunesse et al. 2012; Parkinson et al. 2015). Twenty-four of the cultures

were identified as *B. antillogorgium* based on a comparison of B7Sym15 microsatellite flanking region sequences to available sequences on NCBI. Three of these cultures contained multiple genotypes. Each of the remaining 21 *B. antillogorgium* cultures were unialgal and contained one of five genotypes. Ten *B. antillogorgium* cultures reared at 26°C were either genotype G1, G2 or G3, while 11 *B. antillogorgium* cultures reared at 30°C were genotype G4 or G5. These five genotypes were used in the *in vitro* and *in hospite* experiments. Based on microsatellite genotyping, one of the five genotypes (G3) was the same as the dominant genotype in the source colony; the other four genotypes represented background symbionts as is common in culturing from host tissue (Santos et al. 2001; LaJeunesse 2002; LaJeunesse et al. 2012; Parkinson et al. 2015).

***In vitro* growth experiments**

Cultures reached their maximum cell density in 28-40 days and maintained this density for at least several days afterward (Fig 2a, Online Resource 3). The effect of temperature on maximum population growth rate was dependent on the symbiont genotype (Temperature*Genotype: $F_{4,20} = 30.20$, $p < 0.001$). At 30°C the mean growth rate for two of the five genotypes (G1 and G2) was significantly lower than that at 26°C; there was no significant effect of temperature for the remaining three genotypes (Fig. 2a, b). The effect of temperature on carrying capacity also depended on the symbiont genotype (Temperature*Genotype: $F_{4,20} = 17.90$, $p < 0.001$). One genotype (G3) showed a decrease in carrying capacity at the elevated temperature, but the remaining genotypes showed non-significant responses to temperature (Fig. 2c). The time for populations to reach their peak abundance was also dependent on the interaction between temperature and symbiont genotype (Temperature*Genotype: $F_{4,20} = 6.57$, $p = 0.002$).

In hospite polyp experiments

For the first 31 days, we did not find a significant impact of genotype ($F_{4,73} = 1.84$, $p = 0.130$), temperature ($F_{1,73} = 0.03$, $p = 0.861$), nor Temperature*Genotype ($F_{4,73} = 0.83$, $p = 0.508$) on the proportion of visually infected polyps (Fig. 3a). Similarly, between 32 and 52 days, we did not find a significant impact of genotype ($F_{4,56} = 0.66$, $p = 0.623$), temperature ($F_{1,56} = 3.77$, $p = 0.057$), nor Temperature*Genotype ($F_{4,56} = 1.22$, $p = 0.312$) on the proportion of visually infected polyps (Fig. 3b). We found a significant interaction between the effects of temperature and genotype on the symbiont cell densities in polyps at the end of the experiment, when the vast majority of polyps (range 66-100% of polyps per treatment) were infected (69 days post first inoculation, $F_{4,264} = 6.34$, $p < 0.001$). Average symbiont densities in polyps varied between genotypes, depending on temperature at Day 69 (termination of the experiment, Fig. 3c). Polyp survivorship at the end of the first 31 days was not significantly affected by symbiont genotype ($F_{4,74} = 0.81$, $p = 0.522$), nor by treatment temperature ($F_{1,74} = 3.05$, $p = 0.085$), and there was not a significant interaction term ($F_{4,74} = 0.76$, $p = 0.574$) (Fig. 4a, b). Polyp survivorship at the end of 52 days decreased dramatically in both temperature treatments (Fig. 4a, c), with those polyps at 30°C having significantly lower survivorship than those reared at 26°C ($F_{1,68} = 15.64$, $p < 0.001$). There was not a significant effect of genotype ($F_{4,68} = 2.11$, $p = 0.089$) nor the interaction between temperature and genotype on survival for the second period (Temperature*Genotype, $F_{4,68} = 0.13$, $p = 0.970$). Including the five containers removed after Day 42 (due to algal contamination) did not change the results. Although survivorship results are only reported prior to the appearance of an unidentified green alga, all treatments were maintained for 69 days. At that time, all treatments except G4 at 30°C had surviving polyps (range 2-70 polyps per treatment).

Discussion

Cultured symbionts isolated from standing genetic variation within host colonies revealed the presence of multiple genotypes capable of growth at 30°C, with three of the five unique genotypes used here isolated from a single host colony (Table 1). However, there was no clear evidence of selection for increased thermal tolerance in cultures reared at 30°C for over 700 generations (2 years) compared to those reared at 26°C over this time (Figs. 2-4). Genotypic variability was clear as genotypes differed in their response to thermal stress (i.e., growth rate and carrying capacity, Fig. 2). Regardless of this variability, all tested genotypes showed growth at 30°C, suggesting that *Breviolum antillogorgium*, as a species, is thermally tolerant.

In vitro growth experiments

By isolating genotypes that were able to grow at elevated temperatures from standing genetic variation among *in hospite* populations, we established that thermally tolerant symbiont genotypes are present within the host. We maintained these cultures at 30°C for two years to select for long-term growth at elevated temperatures, anticipating that these cultures would adapt and outperform other genotypes at elevated temperatures. Unlike previous studies that found evidence for thermal adaptation in Symbiodiniaceae after one to 2.5 years of exposure to increased temperatures (*Cladocopium* - Chakravarti et al. 2017; *Fugacium*, *Gerakladium*, *Symbiodinium* - Chakravarti and van Oppen 2018), a history of long-term growth at 30°C did not yield better performance for *B. antillogorgium* at 30°C (as compared to 26°C). Our results suggest that local adaptation to this temperature did not occur over this time period. It is possible that all examined genotypes possessed pre-existing adaptations to temperatures at or above 30°C, as over the past decade and a half, if not longer, this threshold has been frequently exceeded during the summer months in the Florida Keys (Online Resource 1).

Growth rate serves as a good proxy for fitness of symbiont populations. Carrying capacity may serve as a proxy for efficiency of resource use because it suggests that more organisms can be sustained at the same level of resource. We found variation among genotypes in carrying capacity, but interestingly, also found variation in the response of carrying capacity to increased temperature. One genotype (G3) showed a significant decrease in carrying capacity at higher temperature (Fig. 2c), but the remaining four genotypes showed non-significant responses to temperature. The ability of all cultures to survive and grow positively at 30°C for several months, regardless of historical temperature, suggests that these genotypes of *B. antillogorgium* possess pre-existing, broad thermal tolerance, possibly due to plastic acclimatory responses.

Prior studies examining the potential for symbionts to adapt to elevated temperatures have focused on a single genotype (Chakravarti et al. 2017; Chakravarti and van Oppen 2018), but in the present study we used several genotypes within the same species, selected from pre-existing genetic variation within *in hospite* populations. Variability in response to elevated temperatures observed among the genotypes used in our experiments highlights the potential role of intraspecific genetic variation in responding to thermal stress and further underscores the importance of examining both intrageneric and intraspecific variation in thermal responses. Other *in vitro* studies (Chakravarti et al. 2017; Chakravarti and van Oppen 2018; Buerger et al. 2020) have explored how dinoflagellate symbionts in other genera respond to temperatures higher (31°C-34°C) than used in our experiments. It is possible that the rapid adaptation seen in previous studies might also occur in *B. antillogorgium* if these genotypes were grown at higher temperatures. It is clear that elevated temperatures in excess of 30°C are becoming frequent during the summer months in the Florida Keys (Online Resource 1) and further work should explore the limits of thermal tolerance in *B. antillogorgium* to temperatures greater than 30°C.

***In hospite* polyp experiments**

There were no significant differences in the percent of visually infected polyps between genotypes and temperature treatments through Day 52. In general, the percent of infected polyps on Day 52 was largely congruent with the average number of symbiont cells per polyp at the end of the experiment; those genotypes that had a greater proportion of visibly infected polyps also had a greater average number of symbiont cells per polyp (Fig. 3b, c). Final symbiont cell density varied with genotype and temperature, indicating that symbionts were either taken up or grew at different rates within hosts. The lack of a consistent effect of temperature on symbiont density within polyps suggests that the establishment of the symbiosis at 30°C may be as stable as those at ambient temperatures (26°C). Given the significant interaction between treatment temperature and symbiont genotype on symbiont density within polyps at the end of the experiment (Day 69), the ability of the symbiont to infect and thrive in the host likely depends on multiple factors, with potential for variation in the establishment and maintenance of the symbiosis (Davy et al. 2012; Hawkins et al. 2016). It is clear that biotic interactions (e.g., recognition and phagocytosis of algal symbionts, Davy et al. 2012) drive the establishment, while abiotic factors, such as temperature, may dictate the maintenance of the symbiosis (e.g., Rädcker et al. 2021).

The overall thermal tolerance recorded *in vitro* was observed in the polyp experiment, even though performance *in vitro* was not directly mirrored *in hospite* (i.e., those symbiont genotypes that had the highest growth rates *in vitro* did not necessarily confer the highest survivorship under the same conditions *in hospite*). All treatments had relatively high polyp survival early in the experiment, with a more rapid decline in survivorship towards the end (Fig. 4). Polyps showed sustained survivorship over an extended period of heat exposure, with some

surviving polyps in all treatments at 56 days post-inoculation. Furthermore, after 69 days, all treatments except G4 at 30°C had surviving polyps. This is a substantially longer exposure to heat stress than has been previously reported for *in hospite* experiments (i.e., Chakravarti et al. 2017: 28 days; Buerger et al. 2020: 7 days). Our results suggest that the observed thermal tolerance exhibited by our symbiont types may have contributed to or enhanced the thermal tolerance of the holobiont, although the long-term value of this has yet to be determined.

In this study, the host-symbiont pairs exhibited thermal tolerance. At the elevated temperature, however, hosts infected with *Breviolum* cultures that had been reared at 30°C did not consistently outperform their counterparts (i.e., hosts infected with cultures reared at 26°C). This indicates that selection for thermal tolerance of certain genotypes likely occurred prior to the isolation of these symbionts. Increasingly elevated temperatures are becoming more commonplace on coral reefs (Heron et al. 2016; van Hooidonk et al. 2016; Hughes et al. 2018; Oliver et al. 2018), and repeated exposure to these elevated temperatures may have selected for symbiont genotypes capable of surviving such extremes. Ultimately, our results indicate that there exists standing genetic variation within symbiont populations, which in this case, contained genotypes that were thermally tolerant. These symbionts may be vital for the holobiont's survival, although the basis for this thermal tolerance requires further investigation.

Role of the symbiont and implications for Caribbean reefs

Most previous studies of the capacity for rapid thermal adaptation in symbionts have focused on Pacific strains of Symbiodiniaceae. Our results shed new light on the existence of thermally tolerant Caribbean coral endosymbionts in genera in addition to those thermally tolerant species within the genus *Durussdinium* (LaJeunesse et al. 2014). We provide key evidence that even in the absence of rapid adaptation to elevated temperatures, thermal tolerance

may be widespread in some *Breviolum* species. It is likely that broad thermal tolerance is an inherent trait among these genotypes and perhaps among other species within *Breviolum*, particularly those within the B1 (ITS2-type) lineage (Pleistocene (B1) radiation, *sensu* LaJeunesse 2005; Parkinson et al. 2015) such as *B. antillogorgium*. Under induced bleaching and/or thermal stress, for example, the octocoral *Briareum asbestinum* lost its dominant *Breviolum* ITS2 type B19 symbiont, which was replaced by a *Breviolum* ITS2 type B1 species (Lewis and Coffroth 2004; Poland and Coffroth 2019). Transitioning from a dominant symbiont species with lower thermal tolerance to a more thermally tolerant cryptic species during bleaching events may buffer Caribbean octocorals from increasing temperatures.

Coral reefs are in decline worldwide and some regions, such as the Caribbean, are experiencing a phase shift from scleractinian-dominated to octocoral-dominated ecosystems (Ruzicka et al. 2013; Lenz et al. 2015; Lasker et al. 2020). This pattern has been attributed, in part, to high levels of octocoral recruitment (Lasker et al. 2020). However, bleaching in octocorals is rarely observed during mass bleaching events (Lasker et al. 1984; Lasker 2003; Prada et al. 2010) and if it is observed, most species show a high level of resilience (Prada et al. 2010). The fact that species within *Breviolum* are the dominant symbionts in Caribbean octocorals (LaJeunesse 2002; van Oppen et al. 2005; Goulet et al. 2008), and that some symbiont species within this genus appear to be thermally tolerant, as shown here, may contribute to this resilience. Additional studies are needed to determine the prevalence of these thermally tolerant symbionts, which may exist at background level, and their role in mitigating the effects of increasing temperatures on reef cnidarians. The survival of Caribbean reefs may be reliant on evolutionary rescue (Gomulkiewicz and Holt 1995) perpetrated by selection for these more thermally tolerant symbiont genotypes within a given species; a process that could be

particularly important for specialist hosts that are not able to harbor different species of
symbionts.

Acknowledgements

We would like to thank H. Franklin, undergraduate volunteers, staff of the KML, the FKNMS
and the FWC (Permit FKNMS-2014-018-A2 and SAL-18-2052-SR, respectively). This work
was supported by NSF-OCE-1559286 (MAC) and NSF-OCE-1559105 (CPt), a UB Honors
Research & Creativity Grant (JAP), and a UB CURCA Award (KME). We are grateful to D.
Williams for providing the NOAA temperature data and to H. Lasker, A. Martinez, C. Wells, and
two anonymous reviewers for helpful and insightful comments on previous drafts of this
manuscript.

Conflict of Interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- Andras JP, Kirk NL, Coffroth MA, Harvell CD (2009) Isolation and characterization of
microsatellite loci in *Symbiodinium* B1/B184, the dinoflagellate symbiont of the Caribbean
sea fan coral, *Gorgonia ventalina*. Mol. Ecol. Resour. 9:989–993
- Baker AC (2003) Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and
biogeography of *Symbiodinium*. Annu. Rev. Ecol. Evol. Syst. 34:661–689
- Baums IB, Devlin-Durante MK, LaJeunesse TC (2014) New insights into the dynamics between
reef corals and their associated dinoflagellate endosymbionts from population genetic
studies. Mol. Ecol. 23:4203–4215
- Bayliss SLJ, Scott ZR, Coffroth MA, terHorst CP (2019) Genetic variation in *Breviolum*
antilogorgium, a coral reef symbiont, in response to temperature and nutrients. Ecol. Evol.
9:2803–2813
- Berkelmans R, van Oppen MJH (2006) The role of zooxanthellae in the thermal tolerance of

430 corals: a “nugget of hope” for coral reefs in an era of climate change. *Proc. Biol. Sci.*
431 273:2305–2312

432 Buerger P, Alvarez-Roa C, Coppin CW, Pearce SL, Chakravarti LJ, Oakeshott JG, Edwards OR,
433 van Oppen MJH (2020) Heat-evolved microalgal symbionts increase coral bleaching
434 tolerance. *Sci. Adv.* 6:eaba2498

435 Chakravarti LJ, Beltran VH, van Oppen MJH (2017) Rapid thermal adaptation in
436 photosymbionts of reef-building corals. *Glob. Chang. Biol.* 23:4675–4688

437 Chakravarti LJ, van Oppen MJH (2018) Experimental evolution in coral photosymbionts as a
438 tool to increase thermal tolerance. *Front. Mar. Sci.* 5:227

439 Chi J, Parrow MW, Dunthorn M (2014) Cryptic sex in *Symbiodinium* (Alveolata, Dinoflagellata)
440 is supported by an inventory of meiotic genes. *J. Eukaryot. Microbiol.* 61:322–327

441 Coffroth MA, Lasker HR, Diamond ME, Bruenn JA, Bermingham E (1992) DNA fingerprints of
442 a gorgonian coral: a method for detecting clonal structure in a vegetative species. *Mar. Biol.*
443 114:317–325

444 Correa AMS, McDonald MD, Baker AC (2009) Development of clade-specific *Symbiodinium*
445 primers for quantitative PCR (qPCR) and their application to detecting clade D symbionts
446 in Caribbean corals. *Mar. Biol.* 156:2403–2411

447 Davy SK, Allemand D, Weis VM (2012) Cell biology of cnidarian-dinoflagellate symbiosis.
448 *Microbiol. Mol. Biol. Rev.* 76:229–261

449 Díaz-Almeyda EM, Prada C, Ohdera AH, Moran H, Civitello DJ, Iglesias-Prieto R, Carlo TA,
450 LaJeunesse TC, Medina M (2017) Intraspecific and interspecific variation in
451 thermotolerance and photoacclimation in *Symbiodinium* dinoflagellates. *Proc. Biol. Sci.*
452 284:20171767

453 Eakin CM, Lough JM, Heron SF (2009) Climate variability and change: monitoring data and
454 evidence for increased coral bleaching stress. In: van Oppen M.J.H., Lough J.M. (eds) *Coral*
455 *Bleaching*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 41–67

456 Gomulkiewicz R, Holt RD (1995) When does evolution by natural selection prevent extinction?
457 *Evolution* 49:201

458 Goulet TL, Simmons C, Goulet D (2008) Worldwide biogeography of *Symbiodinium* in tropical
459 octocorals. *Mar. Ecol. Prog. Ser.* 355:45–58

460 Goulet TL (2006) Most corals may not change their symbionts. *Mar. Ecol. Prog. Ser.* 321:1–7

461 Grégoire V, Schmacka F, Coffroth MA, Karsten U (2017) Photophysiological and thermal
462 tolerance of various genotypes of the coral endosymbiont *Symbiodinium* sp. (Dinophyceae).
463 *J. Appl. Phycol.* 29:1893–1905

464 Guillard RR, Ryther JH (1962) Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt,

and *Detonula confervacea* (Cleve) Gran. Can. J. Microbiol. 8:229–239

Hawkins TD, Hagemeyer JCG, Warner ME (2016) Temperature moderates the infectiousness of two conspecific *Symbiodinium* strains isolated from the same host population. Environ. Microbiol. 18:5204–5217

Heron SF, Maynard JA, van Hooidonk R, Eakin CM (2016) Warming trends and bleaching stress of the world’s coral reefs 1985–2012. Sci. Rep. 6:38402

Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, Hatzitolos ME (2007) Coral reefs under rapid climate change and ocean acidification. Science 318:1737–1742

Hoegh-Guldberg O, Poloczanska ES, Skirving W, Dove S (2017) Coral reef ecosystems under climate change and ocean acidification. Front. Mar. Sci. 4:158

Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world’s coral reefs. Mar. Freshwater Res. 50:839

Hoegh-Guldberg O (2011) Coral reef ecosystems and anthropogenic climate change. Reg. Environ. Change 11:215–227

Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH (2011) Coral thermal tolerance shaped by local adaptation of photosymbionts. Nat. Clim. Chang. 2:116–120

Howells EJ, van Oppen MJH, Willis BL (2009) High genetic differentiation and cross-shelf patterns of genetic diversity among Great Barrier Reef populations of *Symbiodinium*. Coral Reefs 28:215–225

Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H, Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Wilson SK (2018) Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. Science 359:80–83

Jones AM, Berkelmans R, van Oppen MJH, Mieog JC, Sinclair W (2008) A community change in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field evidence of acclimatization. Proc. Biol. Sci. 275:1359–1365

Kahng SE, Benayahu Y, Lasker HR (2011) Sexual reproduction in octocorals. Mar. Ecol. Prog. Ser. 443:265–283

Karim W, Nakaema S, Hidaka M (2015) Temperature effects on the growth rates and photosynthetic activities of *Symbiodinium* cells. J. Mar. Sci. Eng. 3:368–381

Keshavmurthy S, Hsu C-M, Kuo C-Y, Meng P-J, Wang J-T, Chen CA (2012) Symbiont communities and host genetic structure of the brain coral *Platygyra verweyi*, at the outlet of a nuclear power plant and adjacent areas. Mol. Ecol. 21:4393–4407

- 501 Kirk NL, Andras JP, Harvell CD, Santos SR, Coffroth MA (2009) Population structure of
502 *Symbiodinium* sp. associated with the common sea fan, *Gorgonia ventalina*, in the Florida
503 Keys across distance, depth, and time. Mar. Biol. 156:1609–1623
- 504 LaJeunesse TC, Parkinson JE, Reimer JD (2012) A genetics-based description of *Symbiodinium*
505 *minutum* sp. nov. and *S. psygmophilum* sp. nov. (Dinophyceae), two dinoflagellates
506 symbiotic with cnidaria. J. Phycol. 48:1380–1391
- 507 LaJeunesse TC, Smith RT, Finney J, Oxenford H (2009) Outbreak and persistence of
508 opportunistic symbiotic dinoflagellates during the 2005 Caribbean mass coral “bleaching”
509 event. Proc. Biol. Sci. 276:4139–4148
- 510 LaJeunesse TC, Wham DC, Pettay DT, Parkinson JE, Keshavmurthy S, Chen CA (2014)
511 Ecologically differentiated stress-tolerant endosymbionts in the dinoflagellate
512 genus *Symbiodinium* (Dinophyceae) Clade D are different species. Phycologia 53:305–319
- 513 LaJeunesse TC (2005) “Species” radiations of symbiotic dinoflagellates in the Atlantic and Indo-
514 Pacific since the Miocene-Pliocene transition. Mol. Biol. Evol. 22:570–581
- 515 LaJeunesse T (2002) Diversity and community structure of symbiotic dinoflagellates from
516 Caribbean coral reefs. Mar. Biol. 141:387–400
- 517 LaJeunesse T (2020) Zooxanthellae. Curr Biol 30:R1110–R1113
- 518 Lasker HR (2003) Zooxanthella densities within a Caribbean octocoral during bleaching and
519 non-bleaching years. Coral Reefs 22:23–26
- 520 Lasker HR, Martínez-Quintana Á, Bramanti L, Edmunds PJ (2020) Resilience of octocoral
521 forests to catastrophic storms. Sci. Rep. 10:4286
- 522 Lasker HR, Peters EC, Coffroth MA (1984) Bleaching of reef coelenterates in the San Blas
523 Islands, Panama. Coral Reefs 3:183–190
- 524 Lenz EA, Bramanti L, Lasker HR, Edmunds PJ (2015) Long-term variation of octocoral
525 populations in St. John, US Virgin Islands. Coral Reefs 34:1099–1109
- 526 Lewis CL, Coffroth MA (2004) The acquisition of exogenous algal symbionts by an octocoral
527 after bleaching. Science 304:1490–1492
- 528 Mieog JC, van Oppen MJH, Cantin NE, Stam WT, Olsen JL (2007) Real-time PCR reveals a
529 high incidence of *Symbiodinium* clade D at low levels in four scleractinian corals across the
530 Great Barrier Reef: implications for symbiont shuffling. Coral Reefs 26:449–457
- 531 Muscatine L, Cernichiari E (1969) Assimilation of photosynthetic products of zooxanthellae by a
532 reef coral. Biol. Bull. 137:506–523
- 533 Muscatine L, Porter JW (1977) Reef corals: mutualistic symbioses adapted to nutrient-poor
534 environments. Bioscience 27:454–460
- 535 Oliver JK, Berkelmans R, Eakin CM (2018) Coral bleaching in space and time. Coral Bleaching.

- Springer, pp 27–49
- Parkinson JE, Baums IB (2014) The extended phenotypes of marine symbioses: ecological and evolutionary consequences of intraspecific genetic diversity in coral-algal associations. *Front. Microbiol.* 5:445
- Parkinson JE, Coffroth MA, LaJeunesse TC (2015) New species of Clade B *Symbiodinium* (Dinophyceae) from the greater Caribbean belong to different functional guilds: *S. aenigmaticum* sp. nov., *S. antillogorgium* sp. nov., *S. endomadracis* sp. nov., and *S. pseudominutum* sp. nov. *J. Phycol.* 51:850–858
- Pettay DT, LaJeunesse TC (2007) Microsatellites from clade B *Symbiodinium* spp. specialized for Caribbean corals in the genus *Madracis*. *Mol. Ecol. Notes* 7:1271–1274
- Petzoldt T (2019) growthrates. URL: <https://github.com/tpetzoldt/growthrates>.
- Poland DM, Coffroth MA (2019) Host growth and survivorship varies with endosymbiotic algal partner in developing cnidarians. *Mar. Ecol. Prog. Ser.* 612:87–100
- Prada C, Weil E, Yoshioka PM (2010) Octocoral bleaching during unusual thermal stress. *Coral Reefs* 29:41–45
- Quigley KM, Baker AC, Coffroth MA, Willis BL, van Oppen MJH (2018) Bleaching resistance and the role of algal endosymbionts. Ch. 6 pp. 111-151. In: *Coral Bleaching: Patterns, processes, causes and consequences* eds; J. Lough and M. van Oppen. Springer Ecological Series. Cham, Switzerland
- R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>.
- Rädecker N, Pogoreutz C, Gegner HM, Cárdenas A, Roth F, Bougoure J, Guagliardo P, Wild C, Pernice M, Raina J-B, Meibom A, Voolstra CR (2021) Heat stress destabilizes symbiotic nutrient cycling in corals. *Proc Natl Acad Sci USA* 118(5):e2022653118
- Ruzicka RR, Colella MA, Porter JW, Morrison JM, Kidney JA, Brinkhuis V, Lunz KS, Macaulay KA, Bartlett LA, Meyers MK, Colee J (2013) Temporal changes in benthic assemblages on Florida Keys reefs 11 years after the 1997/1998 El Niño. *Mar. Ecol. Prog. Ser.* 489:125–141
- Santos SR, Gutiérrez-Rodríguez C, Lasker HR, Coffroth MA (2003) *Symbiodinium* sp. associations in the gorgonian *Pseudopterogorgia elisabethae* in the Bahamas: high levels of genetic variability and population structure in symbiotic dinoflagellates. *Mar. Biol.* 143:111–120
- Santos SR, Shearer TL, Hannes AR, Coffroth MA (2004) Fine-scale diversity and specificity in the most prevalent lineage of symbiotic dinoflagellates (*Symbiodinium*, Dinophyceae) of the Caribbean. *Mol. Ecol.* 13:459–469
- Santos SR, Taylor DJ, Coffroth MA (2001) Genetic comparisons of freshly isolated versus

cultured symbiotic dinoflagellates: implications for extrapolating to the intact symbiosis. *J. Phycol.* 37:900–912

Silverstein RN, Correa AMS, Baker AC (2012) Specificity is rarely absolute in coral-algal symbiosis: implications for coral response to climate change. *Proc. Biol. Sci.* 279:2609–2618

Silverstein RN, Cunning R, Baker AC (2015) Change in algal symbiont communities after bleaching, not prior heat exposure, increases heat tolerance of reef corals. *Glob. Chang. Biol.* 21:236–249

Thornhill DJ, Howells EJ, Wham DC, Steury TD, Santos SR (2017) Population genetics of reef coral endosymbionts (*Symbiodinium*, Dinophyceae). *Mol. Ecol.* 26:2640–2659

Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L, Berumen ML, Bourne DG, Cantin N, Foret S, Matz M, Miller DJ, Moya A, Putnam HM, Ravasi T, van Oppen MJH, Thurber RV, Vidal-Dupiol J, Voolstra CR, Watson S-A, Whitelaw E, Munday PL (2017) Rapid adaptive responses to climate change in corals. *Nat. Clim. Chang.* 7:627–636

Tukey JW (1949) Comparing individual means in the analysis of variance. *Biometrics* 5:99–114

van Hooidonk R, Maynard J, Tamelander J, Gove J, Ahmadi G, Raymundo L, Williams G, Heron SF, Planes S (2016) Local-scale projections of coral reef futures and implications of the Paris Agreement. *Sci. Rep.* 6:39666

van Oppen MJH, Mieog JC, Sánchez CA, Fabricius KE (2005) Diversity of algal endosymbionts (zooxanthellae) in octocorals: the roles of geography and host relationships. *Mol. Ecol.* 14:2403–2417

van Oppen MJH, Souter P, Howells EJ, Heyward A, Berkelmans R (2011) Novel genetic diversity through somatic mutations: fuel for adaptation of reef corals? *Diversity (Basel)* 3:405–423

Wilkinson SP, Fisher PL, van Oppen MJH, Davy SK (2015) Intra-genomic variation in symbiotic dinoflagellates: recent divergence or recombination between lineages? *BMC Evol. Biol.* 15:325

Williams DE, Miller MW (2015) Water temperature data from reef sites off the upper Florida Keys from 2003-09-18 to 2019-11-25 (NCEI Accession 0126994). [Water temperature data from reef sites off the upper Florida Keys from 2003-09-18 to 2019-11-25 (NCEI Accession 0126994)]. NOAA National Centers for Environmental Information. Dataset. <https://www.ncei.noaa.gov/archive/accession/0126994>. Accessed [7 Jan 2021.]

Figure Captions

Figure 1. Experimental design for a) *in vitro* growth experiment and b) *in hospite* experiment.

Figure colors refer to historical temperature treatment (blue, maintained at 26°C; red, maintained at 30°C), with tip figures colored based on experimental treatment (not historical temperature).

Figure 2. a) Mean logistic growth fits to *in vitro* cell densities of each genotype of *Breviolum antillogorgium* over time. Blue represents growth at 26°C and red at 30°C. b) Estimated mean (+/- SE) maximum symbiont population growth rates (r) and c) estimated mean (+/- SE) carrying capacities (K). Colors for genotype names (x-axis) refer to the historical temperature at which they were grown, blue indicating cultures maintained at 26°C and red indicating cultures maintained at 30°C two years prior to the start of the experiment. Letters above bars represent differences among groups based on Tukey post-hoc tests.

Figure 3. Mean percent of visually infected polyps at the end of a) Day 31 and b) Day 52. c) Symbiont cell densities within *Antillogorgia bipinnata* polyps inoculated with five symbiont genotypes at 26°C and 30°C at the end of the experiment (Day 69). Colors for genotype names (x-axis) refers to historical temperature, blue indicating cultures maintained at 26°C and red indicating cultures maintained at 30°C two years prior to the start of the experiment. Color of boxes refers to experimental temperature (blue- 26°C, red- 30°C). Boxes represent interquartile range and whiskers the range of the data. Points indicate outliers. Letters above boxes represent differences among groups based on Tukey post-hoc tests.

Figure 4. a) Mean polyp survivorship over time for *A. bipinnata* polyps inoculated with five symbiont genotypes at 26°C and 30°C. Error bars are $\pm$ SE. Blue lines refer to treatments at 26 °C and red lines refer to treatments at 30°C. Differences in average survivorship at b) Day 31 and c) Day 52. Colors for genotype names (x-axis) refer to historical temperature, blue indicating cultures maintained at 26°C and red indicating cultures maintained at 30°C two years prior to the start of the experiment. Color of boxes refers to experimental temperature (blue- 26°C, red- 30°C). Boxes represent interquartile range and whiskers the range of the data. Points indicate outliers. Letters above boxes represent differences among groups based on Tukey post-hoc tests.

Online Resource 1. Average water temperature at Elbow Reef A at 3 m depth from 2005 to 2015. Data from Williams and Miller (2015). Horizontal lines are at 30°C (red) and 26°C (blue) for reference.

Online Resource 2. Experimental set up for the *in hospite* study. Letter color refers to historical temperature, blue indicating cultures reared at 26°C and red indicating cultures reared at 30°C two years prior to the start of the experiment.

Online Resource 3. Mean logistic growth fits to *in vitro* cell densities of each genotype of *Breviolum antillogorgium* over time for each replicate for each genotype and temperature treatment.

a Day 31

b Day 52

c Day 69