RELATIONSHIP BETWEEN PARENTAL AGE AND SEVERITY OF OROFACIAL CLEFTS

Olawale Adamson¹, Abimibola V. Oladugba², Azeez Alade^{3,4}, Waheed O. Awotoye⁵, Tamara Busch⁵, Mary Li⁵, Joy Olotu⁶, Veronica C. Sule⁷, Azeez Fashina¹, James Olutayo¹, Mobolanle O. Ogunlewe¹, Wasiu L. Adeyemo¹, Azeez Butali^{4,5}

¹ Department of Oral and Maxillofacial Surgery, University of Lagos, Lagos, Nigeria

² Department of Biostatistics, University of Nigeria, Nsukka, Nigeria.

³ Department of Epidemiology, College of Public Health, University of Iowa, Iowa City, IA, USA.

⁴Department of Oral Pathology, Radiology and Medicine, College of Dentistry, University of Iowa,

⁵Iowa Institute of Oral Health Research, College of Dentistry, University of Iowa, Iowa City, IA, USA
Iowa City, IA, USA

⁶Department of Anatomy, University of Port Harcourt, Port Harcourt, Nigeria.

⁷Department of Operative Dentistry, College of Dentistry, University of Iowa.

Corresponding author

A. Butali, Department of Oral Pathology, Radiology and Medicine / Iowa Institute for Oral Health Research, College of Dentistry, University of Iowa, Butali Laboratory, ML2198, 500 Newton Road, Iowa City, IA 52242, USA. Email: azeez-butali@uiowa.edu Fax: 319-384-1169. Tel: 319-335-8980.

23 ABSTRACT

24 **OBJECTIVES:** This study aims to investigate the relationship between paternal age, maternal age, and
25 both on the severity of orofacial clefts.

26 **DESIGN:** This was a retrospective study of cases which were subjects clinically diagnosed with non-
27 syndromic cleft lip and/or palate (CL/P). Data was obtained from the AFRICRAN project database on
28 Nigerian non-syndromic orofacial cleft cases.

29 **SETTING:** The samples for cases in this study were obtained at the Cleft clinic of Oral and
30 Maxillofacial surgery at the Lagos University Teaching Hospital, Lagos.

31 **OUTCOME:** Primary outcome measure is severity of orofacial clefts and secondary outcome measure
32 is to evaluate the effect of parental age in determining the incidence of left or right sided orofacial clefts.

33 **RESULTS:** There is no statistical significant association between type of CL $\pm$ P and parental age in
34 young fathers ($p=0.93$). When old fathers are considered, percentage of complete (more severe) CL $\pm$ P
35 cases increases especially in old mothers and this was statistically significant ($p=0.036$). In old fathers,
36 the risk of CL $\pm$ P is increased (OR: 2.66, CI: 1.04-6.80) and also there is increased risk of developing
37 right sided CL $\pm$ P (OR: 1.61, CI: 1.0-2.59). There is reduced risk of isolated cleft palate in young
38 fathers (OR: 0.36, CI: 0.07-1.71) but the risk increases when considering complete types (more severe)
39 of isolated cleft palates (OR: 1.63, CI: 0.71-3.7)

40 **CONCLUSION:** The study shows a higher risk of CL $\pm$ P is associated with increase father's age.

41

42 **Keywords:** parental age, severity, orofacial cleft

43

44

45 INTRODUCTION

46 Birth defects are reported to contribute significantly to infant morbidity and mortality globally (1).
 47 Orofacial clefts (OFC) are amongst the most common craniofacial birth defects with a prevalence of
 48 1:700 live births (2). OFC can be syndromic or non-syndromic with syndromic accounting for 70% of
 49 all OFC (Dixon et al., 2011). The phenotypic presentation of OFC differs and ranges from cleft lip
 50 (CL), cleft lip and palate (CLP) and cleft palate only (CPO). The aetiology of OFC is considered to be
 51 multifactorial with polygenic, environmental, epigenetics and interaction between genetics and
 52 environmental factors playing a role (3). Environmental factors implicated in aetiology of OFC include
 53 smoking, alcohol, metabolic syndromes such as diabetes mellitus and maternal obesity as well as
 54 parental age.

55 Parental age has been proposed as a possible risk factor for OFC.(4) Previous studies conducted on the
 56 association between parental age and incidence of birth defects have yielded inconsistent results(5)(6,7).
 57 It is generally reported that advanced age may predispose chromosomes to irreversible changes and
 58 genetic alterations. In a study by Sartorelli et al.,(8) the frequency of numerical and structural
 59 chromosomal aberrations (acentric fragments and complex radial figures) was significantly greater in
 60 chromosomes of older donors when compared with those of the younger group. Many autosomal
 61 dominant diseases have been shown to be associated with increasing paternal age.(9) Crouzon
 62 syndrome, Apert syndrome and Pfeiffer syndrome are all autosomal dominant craniosynostosis disorders
 63 that can be caused by mutations in the FGFR2 gene occurring in a normal father's germ line. All the
 64 FGFR2 mutations were associated with increased paternal age and molecularly proven to be of paternal
 65 origin.(10) A Danish population-based study of 1,920 OFC affected births of 1,489,014 live births
 66 concluded that paternal age is associated with CLP, independently of maternal age.(11). It is worthy to

note that the fetal congenital anomalies attributed to advanced paternal age is low in absolute terms and though there is a relationship, it is not causal in effect.(9)

There are studies suggesting that maternal age and parity might play an important role in the development of certain isolated birth defects.(12). Kim et al (13) reported that risk of trisomy 21, trisomy 18, triple X syndrome, and all aneuploidies showed a significant increase related to increase in maternal age. For Down syndrome, the risk of maternal age did not change when controlling for paternal age. On the other hand, paternal age effects changed from very large risk to a small sparing risk when controlling for maternal age.

There is no clear consensus on the effect of parental age regarding the risk of orofacial clefts though many studies have reported associations between advanced maternal or paternal age and risk of orofacial clefts. A study by Bille et al(4) using the population-based Danish Facial Cleft Database, reported that the influence of maternal and paternal ages on the risk of Cleft lip and/or palate (CL ± P) increases with the advancing age of the other parent, and that the influence vanishes if the other parent is young. In contrast, the risk of having a child with cleft palate is influenced only by father's age, not mother's age. In a study of Brazilians with OFC, Martelli et al(14) reported an association between maternal age and increased risk for CLP while paternal age risk is not significant.

In addition to the fact that the association between parental age and risk of orofacial clefts has been inconsistent, there is sparse literature on the influence of parental age on the severity of orofacial clefts. This study aims to investigate the relationship between paternal age, maternal age, and both on the severity of orofacial clefts. In addition, we plan to evaluate the effect of parental age in determining the incidence of left or right sided orofacial clefts.

MATERIAL AND METHODS

Study design: This was a retrospective study in which cases were subjects clinically diagnosed with non-syndromic cleft lip and/or palate (CL $\pm$ P). Selection of cases was based on standardized examination performed by trained Surgeons who participated in the Pan-African Association of Cleft lip and palate network for repair of orofacial clefts in Africa. Clinical information including detailed description of the phenotype, parental age and clinical photographs were recorded in the database.

Study location: The samples for cases in this study were obtained at the Cleft clinic of Oral and Maxillofacial surgery at the Lagos University Teaching Hospital, Lagos. The Research and ethics committee of Lagos University Teaching Hospital was informed, and ethical approval obtained before commencing the study.

Method:

Data was obtained from the AFRICRAN project database on Nigerian non-syndromic orofacial cleft cases. All infants born with orofacial clefts were clinically examined with the overall goal to measure and characterize the craniofacial morphology and development, and data on parental age were also included. The infants were classified according to whether they were unilateral (left [L] or right [R] sided) or bilateral, as well as the severity of their cleft graded.

For the current analysis regarding the influence of parental age on cleft severity, the groups with CL $\pm$ P and isolated cleft palate were considered two separate populations because of their different embryological origins. The CL $\pm$ P population comprised: unilateral incomplete cleft lip (UICL), bilateral incomplete cleft lip (BICL), unilateral complete cleft lip (UCCL), bilateral complete cleft lip (BCCL), unilateral incomplete cleft lip and palate (UICLP), bilateral incomplete cleft lip and palate (BICLP), unilateral complete cleft lip and palate (UCCLP) and BCCLP (bilateral complete cleft lip and palate).

In the CL $\pm$ P population, the data were grouped by the analysis of the influence of severity. For this purpose, the previously described subgroups were combined as follows: IC (incomplete/less severe clefts = UICL+BICL+UICLP+BICLP) vs CC (complete/ more severe clefts = UCCL+BCCL+UCCLP+BCCLP), as well as L vs R-sided cleft (for this analysis, only unilateral clefts (UCL $\pm$ P = UICL+UCCL+UICLP+UCCLP) were included).

The CP population comprised of: incomplete cleft palate (ICP) and complete cleft palate (CCP)

The parental age was classified into young father, old father or young mother, old mother based on the median ages of the parents. The risk of orofacial clefts was analyzed based on these groups.

Statistical analysis: For the primary analysis, a binary outcome variable was defined with two values (0 = IC, 1 = CC). Pearson's Chi-square test was applied to analyze the association between parental age and the severity of orofacial clefts. Based on logistic regression, the relative risk with confidence interval was calculated between severity of orofacial clefts and parental age.

RESULTS: The total number of non-syndromic orofacial cleft cases analyzed was 267 with 202 CL $\pm$ P and 65 CP cases. Table 1 shows the parental age distribution of the cleft cases. Young fathers are categorized as those below 35 years while old fathers are greater than or equals to 35 years old while young mothers are categorized as those below 30 years while old mothers are greater than or equals to 30 years old.

Table 1: Summary statistics of the age distribution of the father and mother in the data

		CL $\pm$ P		CP	
		fathers age	mothers age	Father's age	Mother's age
	Minimum	20	17	25	19
	1 st Quartile	32.00	26.00	32.00	26.00
	Median	35.00	30.00	35.00	29.00
	Mean	35.48	29.51	35.52	29.09
	3 rd Quartiles	38.00	32.00	39.00	32.00
	Maximum	54	43	48	42

CLEFT LIP AND PALATE

Generally, there are more complete cleft CL $\pm$ P than incomplete cases (Table 2). There is no statistical significant association between type of CL $\pm$ P and parental age in young fathers (p=0.93). When old fathers are considered, percentage of complete CL $\pm$ P cases increases especially in old mothers and this was statistically significant at (p=0.036). These findings indicate that old father-old mother combination is more associated with more severe CL $\pm$ P.

161 **Table 2: Shows the relationship of parental age to cleft lip and palate cases.**

Fathers age based on median				Type of Cleft lip		Total	p-value
				IC	CC		
< 35	Mothers age based on median	< 30	Count	13	42	55	0.930
			% of Total	16.9%	54.5%	71.4%	
		> 30	Count	5	17	22	
			% of Total	6.5%	22.1%	28.6%	
		Total		Count	18	59	
				% of Total	23.4%	76.6%	
	Total			Count	18	59	
				% of Total	23.4%	76.6%	
≥35	Mothers age based on median	< 30	Count	12	32	44	0.036
			% of Total	9.6%	25.6%	35.2%	
		> 30	Count	10	71	81	
			% of Total	8.0%	56.8%	64.8%	
		Total		Count	22	103	
				% of Total	17.6%	82.4%	
	Total			Count	22	103	
				% of Total	17.6%	82.4%	

162

163

164 Severity of CL ± P

165 There is no increased risk of CL ± P in young fathers (OR: 1.05, CI: 0.3-3.4) and there is no increased
 166 risk for any subtype of CL ± P (table 3). In old fathers, the risk of CL ± P is increased (OR: 2.66, CI:
 167 1.04-6.80). There is also increased risk for incomplete CL ± P in old fathers (OR: 2.209, CI: 1.04-4.70)
 168 but the risk reduces when complete CL ± P was considered (OR: 0.83, CI: 0.68-1.01). These show that
 169 the risk of CL ± P increases with paternal age which is higher in less severe form of incomplete CL ± P
 170 (Table 3).

171 **Table 3: Relative risk of severity of CL ± P in relation to parental age.**

Risk Estimate				
Fathers age based on median		Value	95% Confidence Interval	
			Lower	Upper
< 35	Odds Ratio for Mothers age based on median (< 30 / > 30)	1.052	.325	3.409
	For cohort Type of Cleft lip = IC	1.040	.421	2.571
	For cohort Type of Cleft lip = CC	.988	.754	1.295
≥ 35	Odds Ratio for Mothers age based on median (< 30 / > 30)	2.663	1.043	6.797
	For cohort Type of Cleft lip = IC	2.209	1.039	4.699
	For cohort Type of Cleft lip = CC	.830	.680	1.012
Total	Odds Ratio for Mothers age based on median (< 30 / > 30)	1.982	.974	4.035
	For cohort Type of Cleft lip = IC	1.734	.973	3.090
	For cohort Type of Cleft lip = CC	.875	.761	1.006

Risk of left or right sided cleft in unilateral CL ± P

There is no associated increase in risk of unilateral CL ± P for either left or right side in young fathers (Table 4). In old fathers, there is increased risk of developing right sided CL ± P (OR: 1.61, CI: 1.0-2.59) and the risk of developing left sided clefts reduces indicating that mother's age is a more associated with left-sided clefts in old fathers.

Table 4: Relative risk of Left or right sided Unilateral CL ± P

Risk Estimate				
Fathers age based on median		Value	95% Confidence Interval	
			Lower	Upper
< 35	Odds Ratio for Mothers age based on median (< 30 / > 30)	1.045	.352	3.104
	For cohort Cleft Lip Details = Left	1.026	.548	1.919
	For cohort Cleft Lip Details = Right	.981	.618	1.558
≥ 35	Odds Ratio for Mothers age based on median (< 30 / > 30)	.442	.193	1.011
	For cohort Cleft Lip Details = Left	.714	.493	1.033
	For cohort Cleft Lip Details = Right	1.614	1.006	2.588
Total	Odds Ratio for Mothers age based on median (< 30 / > 30)	.508	.273	.943
	For cohort Cleft Lip Details = Left	.733	.549	.978
	For cohort Cleft Lip Details = Right	1.443	1.028	2.025

Severity of bilateral CL $\pm$ P

In bilateral CL $\pm$ P, there is a slight risk of bilateral CL $\pm$ P in young fathers (OR: 1.14, CI: 0.7-16.94) (Table 5). There was two-fold increase in risk of bilateral CL $\pm$ P in old fathers (OR: 2.0, CI: 0.11-36.9) and this was more predominant in incomplete bilateral CL $\pm$ P (OR: 1.87, CI: 0.13-26.1).

Table 5: Relative risk of bilateral CL $\pm$ P

Risk Estimate				
Fathers age based on median		Value	95% Confidence Interval	
			Lower	Upper
< 35	Odds Ratio for Mothers age based on median (< 30 / > 30)	1.143	.077	16.947
	For cohort Type of Cleft lip = IC	1.091	.184	6.476
	For cohort Type of Cleft lip = CC	.955	.382	2.387
$\geq$ 35	Odds Ratio for Mothers age based on median (< 30 / > 30)	2.000	.108	36.954
	For cohort Type of Cleft lip = IC	1.875	.134	26.161
	For cohort Type of Cleft lip = CC	.938	.698	1.259
Total	Odds Ratio for Mothers age based on median (< 30 / > 30)	2.857	.477	17.110
	For cohort Type of Cleft lip = IC	2.368	.524	10.698
	For cohort Type of Cleft lip = CC	.829	.605	1.135

188 CLEFT PALATE

189 There is reduced risk of isolated cleft palate in young fathers (OR: 0.36, CI: 0.07-1.71) but the risk
 190 increases when considering complete cleft palates (OR: 1.63, CI: 0.71-3.7) though this was not
 191 statistically significant (Table 6). This indicates that maternal age is more associated with less severe
 192 cleft palate while paternal age is associated with more severe cleft palate.

193

194 **Table 6: Risk of parental age and severity of cleft palate only**

Risk Estimate				
Fathers age based on median		Value	95% Confidence Interval	
			Lower	Upper
<35	Odds Ratio for Mothers age based on median (<30 / >30)	.359	.075	1.714
	For cohort Populations = ICP	.583	.267	1.276
	For cohort Populations = CCP	1.625	.713	3.706
≥35	Odds Ratio for Mothers age based on median (<30 / >30)	.593	.149	2.365
	For cohort Populations = ICP	.781	.395	1.545
	For cohort Populations = CCP	1.316	.646	2.680
Total	Odds Ratio for Mothers age based on median (<30 / >30)	.445	.165	1.200
	For cohort Populations = ICP	.663	.398	1.106
	For cohort Populations = CCP	1.492	.904	2.462

195

DISCUSSION

This study evaluates the relationship between parental age and severity of cleft using data derived from Nigerian patients with cleft lip and palate. To our knowledge, this is the first of such study to be conducted in an African population. This study attempts to go further than just linking parental age with risk of OFC, but highlighting the effects on severity and also on left or right selection of unilateral cleft lip and palate cases.

This study shows that increased parental age is associated with more severe CL $\pm$ P cases as combination of older parents produce more severe cases. This aligns with various studies that have reported increased congenital malformations in older parents.(6,15). A population-based study on Danish Facial Cleft Database, reported the influence of maternal and paternal ages on the risk of cleft lip with or without cleft palate increases with the advancing age of the other parent, and that the influence vanishes if the other parent is young.(7) Though there has been varying reports on whether the maternal or paternal chromosomes are culpable. The exact mechanism of this occurrence has not been elucidated, though single gene mutations are suggested mechanisms.(11)

Also from this study, advanced paternal age is associated with increased risk of less severe unilateral and bilateral CL $\pm$ P. This is in agreement with a similar study by Herman et al(16) that reported that paternal age increases risk of CL $\pm$ P which is more pronounced with advanced maternal age. The paternal age seems to have a great deal of influence on the prevalence of CL $\pm$ P in any population. The influence of paternal age also spills over in to cleft palate where many studies have reported association between paternal age and cleft palate.(Martelli *et al.*, 2010; Hoda Badr *et al.*, 2011) In this study, paternal age is associated with increased risk of more severe cleft palate.

Though maternal age has been associated with chromosomal abnormalities in some studies but paternal age is usually associated with birth defects.(12,13) It is reported in some literature that the risk of birth

defects such as heart malformation, other musculoskeletal anomalies, tracheo-oesophageal fistula/oesophageal atresia, Down's syndrome and other chromosomal anomalies, increases slightly with advancing paternal age.(5,14). Association between younger fathers and several selected birth defects like neural tube defects has also been published (18) The association of paternal age with birth defects has been attributed to accumulation of chromosomal aberrations and mutations during the maturation of male germ cells.(10,19). The amount of DNA damage in sperm of men aged 36–57 is three times that of men 35 years and less.(11)

Prevalence and pattern of occurrence of OFC in a given population is expected to fluctuate as the average parental ages change. Increase occurrence of more severe cleft is expected with advanced parental ages and this may take a toll on available resources.

Strengths and Limitations

The strength of this study is that it is a population-based investigation of a genetically homogeneous population who has similar environmental exposures. Furthermore, only parents of children with non-syndromic cleft were included. The limitation of this study is the small sample size and other environmental factors like socio-economic status of the parents, maternal intake of alcohol and smoking were not considered

Conclusion

Increased parental age is associated with increased risk of OFC. In this study, advanced paternal age is associated with increased risk of less severe unilateral and bilateral CL ± P but a more severe cleft palate. Future prospective studies on different populations and also considering other socio-economic factors may provide more insights into the influence of parental age on occurrence and severity of OFC.

ACKNOWLEDGEMENTS

We are grateful to the families who voluntarily participated in this study in Nigeria. We are also grateful to all the administrative and research staffs, students, nurses and resident doctors who assisted with participant recruitment, consent, and data collection. This research is supported by the National Institute of Dental and Craniofacial Research (R00 DE022378 and R01DE028300; A.B.).

REFERENCES

1. Butali A, Adeyemo WL, Mossey PA, Olasoji HO, Onah II, Adebola A, et al. Prevalence of orofacial clefts in Nigeria. *Cleft Palate-Craniofacial J.* 2014;
2. Stone C. Cleft Lip and Palate: Etiology, Epidemiology, Preventive and Intervention Strategies. *Anat Physiol.* 2013;04(03):2–6.
3. Wong FK, Hägg U. An update on the aetiology of orofacial clefts. *Hong Kong Med J.* 2004;10(5):331–6.
4. Bille C, Skytthe A, Vach W, Knudsen LB, Andersen AMN, Murray JC, et al. Parent’s age and the risk of oral clefts. *Epidemiology.* 2005;16(3):311–6.
5. Yang Q, Wen SW, Leader A, Chen XK, Lipson J, Walker M. Paternal age and birth defects: How strong is the association? *Hum Reprod.* 2007;22(3):696–701.
6. de Carvalho PHP, Machado RA, Reis SR de A, Martelli DRB, Dias VO, Martelli-Júnior H. Parental age is related to the occurrence of cleft lip and palate in Brazilian populations. *Brazilian J Oral Sci.* 2016;15(2):167–70.
7. Leth Jensen B, Kreiborg S, Dahl E, Fogh-Andersen P. Cleft lip and palate in Denmark, 1976–1981: Epidemiology, variability, and early somatic development. *Cleft Palate J.* 1988;25(3):258–69.
8. Sartorelli EMP, Mazzucatto LF, De Pina-Neto JM. Effect of paternal age on human sperm chromosomes. *Fertil Steril.* 2001;76(6):1119–23.
9. Nybo Andersen AM, Urhoj SK. Is advanced paternal age a health risk for the offspring? *Fertil Steril.* 2017;107(2):312–8.
10. Wiener-Megnazi Z, Auslender R, Dirnfeld M. Advanced paternal age and reproductive outcome. *Asian J Androl.* 2012;14(1):69–76.

- 286 11. Bray I, Gunnell D, Smith GD. Advanced paternal age: How old is too old? *J Epidemiol*
287 *Community Health*. 2006;60(10):851–3.
- 288 12. Luo YL, Cheng YL, Gao XH, Tan SQ, Li JM, Wang W, et al. Maternal age, parity and isolated
289 birth defects: A population-based case-control study in Shenzhen, China. *PLoS One*.
290 2013;8(11):4–9.
- 291 13. Kim YJ, Lee JE, Kim SH, Shim SS, Cha DH. Maternal age-specific rates of fetal chromosomal
292 abnormalities in Korean pregnant women of advanced maternal age. *Obstet Gynecol Sci*.
293 2013;56(3):160.
- 294 14. Martelli DRB, da Cruz KW, de Barros LM, Silveira MF, Swerts MSO, Júnior HM. Maternal and
295 paternal age, birth order and interpregnancy interval evaluation for cleft lip/palate. *Braz J*
296 *Otorhinolaryngol*. 2010;76(1):107–12.
- 297 15. Harville EW, Wilcox AJ, Lie RT, Vindenes H, Åbyholm F. Cleft lip and palate versus cleft lip
298 only: Are they distinct defects? *Am J Epidemiol*. 2005;162(5):448–53.
- 299 16. Hermann N V., Darvann TA, Munch A, Kreiborg S. Parental age in relation to the severity of
300 cleft lip and/or palate. *Orthod Craniofac Res*. 2018;21(4):236–41.
- 301 17. Hoda Badr, Cindy L. Carmack, Deborah A. Kashy, Massimo Cristofanilli and TAR. 基因的改变
302 NIH Public Access. *Bone*. 2011;23(1):1–7.
- 303 18. Zhu JL, Madsen KM, Vestergaard M, Olesen A V., Basso O, Olsen J. Paternal age and congenital
304 malformations. *Hum Reprod*. 2005;20(11):3173–7.
- 305 19. Thompson JA. Disentangling the roles of maternal and paternal age on birth prevalence of down
306 syndrome and other chromosomal disorders using a Bayesian modeling approach. *BMC Med Res*
307 *Methodol*. 2019;19(1):1–8.

308