An evolutionarily conserved coreceptor gene is essential for CLAVATA signaling in *Marchantia polymorpha*

**Go Takahashi^{1†}, Shigeyuki Betsuyaku^{2†}, Natsuki Okuzumi¹, Tomohiro Kiyosue¹,
and Yuki Hirakawa^{1†*}**

¹ Graduate School of Science, Gakushuin University, Toshima-ku, Tokyo, Japan

² Faculty of Agriculture, Ryukoku University, Otsu, Shiga, Japan

[†]These authors have contributed equally to this work.

*** Correspondence:**

Yuki Hirakawa

Email: yuki.hirakawa@gakushuin.ac.jp

Running title: CLAVATA coreceptor in *Marchantia*

Keywords: CLAVATA, coreceptor, *Marchantia*, meristem, LRR-RLK

Abstract

Growth and development of land plants are controlled by CLAVATA3/EMBRYO SURROUNDING REGION-related (CLE) family of peptide hormones. In contrast to the genetic diversity of CLE family in flowering plants, the liverwort *Marchantia polymorpha* possesses a minimal set of CLE, MpCLE1(TDIF homolog) and MpCLE2 (CLV3 homolog). MpCLE1 and MpCLE2 peptides exert distinct function at the apical meristem of *M. polymorpha* gametophyte via specific receptors, MpTDIF RECEPTOR (MpTDR) and MpCLAVATA1 (MpCLV1), respectively, both belonging to the subclass XI of leucine-rich repeat receptor-like kinases (LRR-RLKs). Biochemical and genetic studies in *Arabidopsis* have shown that TDR/PXY family and CLV1/BAM family recognize the CLE peptide ligand in a heterodimeric complex with a member of subclass-II coreceptors. Here we show that three LRR-RLK genes of *M. polymorpha* are classified into subclass II, representing three distinct subgroups evolutionarily conserved in land plants. To address the involvement of subclass-II coreceptors in *M. polymorpha* CLE signaling, we performed molecular genetic analysis on one of them, MpCLAVATA3 INSENSITIVE RECEPTOR KINASE (MpCIK). Two knockout alleles for MpCIK formed narrow apical meristems marked by *prom*MpYUC2:*GUS* marker, which were not expanded by MpCLE2 peptide treatment, phenocopying *Mpclv1*. Loss of sensitivity to MpCLE2 peptide was also observed in gemma cup formation in both *Mpclv1* and *Mpcik*. Biochemical analysis using a *Nicotiana benthamiana* transient expression system revealed weak association between MpCIK and MpCLV1, as well as MpCIK and MpTDR. While MpCIK may also participate in MpCLE1 signaling, our data show that the conserved CLV3-CLV1-CIK module functions in *M. polymorpha*, controlling meristem activity for development and organ formation for asexual reproduction.

Introduction

CLAVATA3/EMBRYO SURROUNDING REGION-related (CLE) peptides are a family of peptide hormones in land plants, mediating cell-to-cell communication in the plant body (Murphy et al., 2012; Hirakawa and Sawa, 2019; Fletcher, 2020). CLE peptides are genetically encoded as a precursor protein possessing a conserved CLE domain(s) at or near the C-terminus. Biosynthesis of CLE peptide hormone from the CLE domain involves post-translational events including proteolytic cleavage, post-translation modifications and secretion to the apoplast (Ito et al., 2006; Kondo et al., 2006; Ohyama et al., 2009; Tamaki et al., 2013; Matsubayashi, 2014). In flowering plants, a large number of *CLE* genes are encoded in the genome, which have been extensively studied for the past two decades (Cock and McCormick 2001; Oelkers et al., 2008; Jun et al., 2010; Fletcher, 2020). The function of CLE genes cover a wide range of physiological processes including stem cell homeostasis in meristems, vascular cell differentiation, stomata differentiation and responses to various environmental cues (Fletcher et al., 1999; Suzaki et al., 2008; Okamoto et al., 2009; Stahl et al., 2009; Mortier et al., 2010; Etchells and Turner 2010; Hirakawa et al. 2010; Kondo et al., 2011; Fiume and Fletcher, 2012; Depuydt et al., 2013; Endo et al., 2013; Araya et al., 2014; Czyzewicz et al., 2015; Gutierrez-Alanis et al. 2017; Rodríguez-Leal et al., 2017; Qian et al., 2018; Takahashi et al., 2018; Ma et al., 2020). In bryophytes, which diverged early in the land plant lineage with respect to flowering plants (Puttick et al., 2018; Morris et al., 2018), relatively low number of *CLE* genes are encoded in the genome, providing simplified models to study the function of *CLE* genes (Bowman et al., 2017; Whitewoods et al. 2018). The minimal set of *CLE* genes, Mp*CLE1* (Mp6g07050) and Mp*CLE2* (Mp5g18050), are encoded in the genome of the liverwort *Marchantia polymorpha* (Bowman et al., 2017; Hirakawa et

al., 2019; Montgomery et al., 2020; Figure 1A). MpCLE1 and MpCLE2 are the orthologs of TDIF (tracheary element differentiation inhibitor factor) and CLV3 (CLAVATA3) of *Arabidopsis thaliana*, respectively, representing the two distinct subgroups of CLE peptide family. In *Arabidopsis*, specific bioactivities of TDIF and CLV3 are attributed to the difference in a few amino acids between them, which are mediated by two distinct groups of receptors, TDIF RECEPTOR/PHLOEM INTERCALATED WITH XYLEM (TDR/PXY) and CLAVATA1/BARELY ANY MERISTEMs (CLV1/BAMs), respectively (Fletcher et al., 1999; DeYoung et al., 2006; Fisher and Turner, 2007; Ogawa et al., 2008; Hirakawa et al., 2008; Rodriguez-Villalon et al., 2014; Shimizu et al., 2015; Shinohara and Matsubayashi, 2015; Hirakawa et al., 2017; Crook et al., 2020). Since the ligand-receptor pairs are conserved among flowering plants and bryophytes and no CLE homologs were found in sister streptophyte algae, the specific CLE peptide-receptor pairs may have originated in the common ancestor of land plants (Whitewoods et al., 2018; Hirakawa et al., 2019; Hirakawa et al., 2020). In *M. polymorpha*, CLE genes regulate the activity of the apical meristem located at the apical notch of the thalloid gametophyte body. MpCLE1-MpTDR signaling acts as a negative regulator of cell proliferation at the apical notch, while MpCLE2-MpCLV1 signaling functions as a positive regulator of stem cell activity in the apical notch (Hirakawa et al., 2019; Hirakawa et al., 2020).

Both TDR/PXY and CLV1/BAM belong to the subclass XI of leucine-rich repeat receptor-like kinase (LRR-RLK) family. In addition to CLE peptides, a number of peptide ligands have been shown to bind to specific members of subclass-XI receptors, which possess a long extracellular domain (ECD) composed of more than 20 LRRs (Shiu and Bleecker, 2001; Yamaguchi et al., 2006; Tabata et al., 2014; Ou et al., 2016; Shinohara et al., 2016; Song et al., 2016; Doblas et al., 2017; Nakayama et al., 2017; Doll et al., 2020).

Accumulating evidence indicates that subclass-II receptors, such as SOMATIC EMBRYOGENESIS RECEPTOR KINASE/BRASSINOSTEROID INSENSITIVE1-ASSOCIATED KINASE1 (SERK/BAK1) family, participate in the peptide hormone perception by forming a heterodimeric complex with subclass-XI receptors (Hohmann et al., 2017; Gou and Li, 2020). Structural studies have revealed that SERK coreceptors have a short ECD containing five LRRs. The ECD of subclass-II receptors do not interact strongly or at all to the peptide ligand by themselves and rather recognize the ligand-receptor complex (Santiago et al., 2013; Sun et al., 2013; Wang et al., 2015; Santiago et al., 2016; Okuda et al., 2020). In line with this scheme, PXY/TDR and SERK2 are reported to form a heterodimeric complex for TDIF recognition, and multiple knockout mutants for Arabidopsis *SERK* genes show reduced TDIF sensitivity in vascular development (Morita et al., 2016; Zhang et al., 2016a; Zhang et al., 2016b).

In contrast to TDIF, involvement of SERK family has not been observed in CLV3-type CLEs. Instead, another group of subclass-II receptors, CLV3 INSENSITIVE RECEPTOR KINASEs (CIKs), have been implicated in CLV3 peptide perception. CIK proteins can form protein complexes with CLV1/BAM receptors (Hu et al., 2018; Cui et al., 2018). Quadruple mutants for Arabidopsis *CIK1-4* genes develop enlarged shoot apical meristems, which is similar to those of *clv* mutants. The growth from the enlarged meristems is not arrested by treatment with CLV3 peptide, a negative regulator of stem cells in Arabidopsis (Hu et al., 2018). Furthermore, full activity of Arabidopsis CLE26/CLE45 peptides in root phloem cell differentiation requires *CLE-RESISTANT RECEPTOR KINASE (CLERK)/CIK2* although biochemical interaction is not detected between the ECDs of CLERK and the subclass-XI receptor BAM3 (Anne et al., 2018).

In this study, we searched for the homologs of *CIK* genes in *M. polymorpha* and analyzed their involvement in CLE peptide signaling by molecular genetic approach.

Results

A single *CIK* ortholog in *M. polymorpha*

In the *M. polymorpha* genome, three LRR-RLK genes (Mp7g09160/Mapoly0068s0069, Mp7g14210/Mapoly0009s0106, Mp7g15980/Mapoly0560s0001) have been classified into subclass II (Sasaki et al., 2007; Bowman et al., 2017; Montgomery et al., 2020).

To better understand the evolutionary relationships, we performed phylogenetic analysis of the subclass-II genes from land plants (*Arabidopsis thaliana*, *Amborella trichopoda*, *Picea abies*, *Selaginella moellendorffii*, *Physcomitrium* (*Physcomitrella*) *patens*, *Sphagnum fallax*, *Marchantia polymorpha*) and charophycean algae (*Spirogyra pratensis*, *Coleochaete orbicularis*) based on the amino acid sequence of the kinase domain using a Bayesian method (Figure 1B). The tree inferred three subgroups diverged in the land plant lineage, each of which contains a single *M. polymorpha* gene. Mp7g14210, designated as Mp*CIK*, was grouped into a single subgroup with all *CIK* genes from *Arabidopsis*. Likewise, Mp7g09160/Mp*SERK* was grouped into the *SERK* subgroup with all *Arabidopsis* *SERK* genes. Thus, *M. polymorpha* genome may lack redundancy in CLE ligand/receptor/coreceptor orthologs (Figure 1A).

CRISPR-Cas9 editing of Mp*CIK* does not affect overall growth of gametophyte

To analyze the physiological function of the *CIK* coreceptor gene in *M. polymorpha*, we generated loss-of-function alleles for Mp*CIK* using CRISPR-Cas9 editing (Sugano et al., 2018). Sanger sequencing revealed that two independent transgenic lines, Mp*ciik*-I^{ge} and

Mpcik-2^{ge}, possess different mutations at the CRISPR/Cas9 target site, both predicted to result in gene knockout due to premature termination of translation (Figure 1C). We could not find significant differences in the overall morphology of thalli in 10-day-old *Mpcik-1^{ge}* and *Mpcik-2^{ge}* plants grown from gemmae, compared to any of wild-type (Tak-1), *Mpcle2-2^{ge}* and *Mpchl1-3^{ge}* genotypes (Figure 1D).

Figure. 1 Analysis of LRR-RLK subclass II in *Marchantia polymorpha*

(A) The number of CLE-receptor homologs. (B) A phylogenetic tree of subclass-II LRR-RLKs, generated with a Bayesian method based on the conserved kinase domain. The posterior probabilities of trees are shown at the nodes. Coleochaete sequence was used as an outgroup. Land plant sequences form a monophyletic clade, which can be divided into three subgroups as indicated on the right. Inset shows the list of species with their phylogenetic relationships. (C) Gene/protein structures and genome editing alleles of MpCIK. (top) Protein structure of MpCIK. (middle) Structure of MpCIK/Mp7g14210 locus with the position of a designed guide RNA (gRNA). (bottom) Genotyping of genome editing alleles. Target guide sequence is in bold and PAM sequence is in blue. Deleted bases are indicated with hyphens in magenta. Exon and intron are indicated in capital and small letters, respectively. N-terminal region of WT and mutant proteins deduced from the genomic DNA sequences are indicated below. Asterisks indicate translational termination. (D) Overall morphology of 10-day-old plants grown from gemmae. Scale bars represent 0.5 mm.

Figure 2 Phenotypes of MpCIK knockout alleles in the apical notch

(A) Morphology of 4-day-old gemmalings grown with or without MpCLE2 peptide as indicated above. Width of apical notch is indicated by white lines. (B) Quantification of apical notch width. Asterisks indicate a significant difference in a two-tailed Welch's *t* test, $p < 0.05$; ns, not significant; $n = 30$. (C) *proMpYUC2:GUS* marker in 4-day-old gemmalings. Genetic background is indicated above each panel. Scale bars represent 500 μm in (A) and 100 μm in (C).

MpCIK is necessary for MpCLE2 signaling to control apical notch expansion

To analyze the involvement of MpCIK in MpCLE2 peptide signaling, we examined the apical notch morphology in 4-day-old gemmalings grown on liquid M51C medium supplemented with or without 3 μM MpCLE2 peptide. In wild-type gemmalings, apical notches were expanded by treatment with MpCLE2 peptide, as reported previously (Figures 2A and 2B; Hirakawa et al., 2020). By contrast, apical notches in both *Mpcik-1* and *Mpcik-2* were insensitive to MpCLE2 peptide, which is similar to those in *Mpclv1-3* (Figures 2A and 2B). Importantly, apical notches of the *Mpcik* and *Mpclv1* alleles were narrower than those of wild type in the growth without MpCLE2 peptide, indicating that MpCIK is involved in intrinsic MpCLE2-MpCLV1 signaling. Consistently, *Mpcle2-2^{ge}*

developed narrow apical notches but it was sensitive to the treatment with the MpCLE2 peptide as reported previously (Figure 2B; Hirakawa et al., 2020). *pro*MpYUC2(*YUCCA2*):*GUS* is a marker for the tip of apical notch, and *pro*MpYUC2:*GUS*-positive (MpYUC2⁺) region is affected by MpCLE2-MpCLV1 signaling (Eklund et al., 2015; Hirakawa et al., 2020). Compared to wild-type (Tak-1) background, MpYUC2⁺ region was reduced in *Mpcik* backgrounds, phenocopying *Mpclv1-3* (Figure 2C). These data support that MpCIK is an essential component of the MpCLE2 peptide perception.

Figure. 3 MpCIK knockout suppresses gain-of-function phenotypes of MpCLE2

(A) Overall morphology

of 16-day-old plants grown from gemmae.

Genotypes are indicated above the panels. The upper middle panel shows the magnification of Tak-1 image in which arrows indicate gemma

cups. Note that *pro*MpYUC2:MpCLE2 (*CLE2ox*) exhibits multichotomy and produces no gemmae cups at this age. Scale bars represent 1 cm. (B) Number of gemmae cups (mean and SD; n=9-12). *CLE2ox* showed a significant delay of gemmae cup formation. Data are obtained at 4, 7, 14, 15, 16, 17, 18, 21 days after gemmae plating. This experiment was repeated twice with similar results.

MpCLE2 signaling affects thallus branching and gemma cup formation via MpCLV1 and MpCIK

To further address the involvement of MpCIK in MpCLE2 signaling, we used *pro*MpYUC2:MpCLE2, a gain of function allele of MpCLE2, which stably develops supernumerary branching from the expanded apical meristems (Hirakawa et al., 2020).

We generated *Mpcik* and *Mpclv1* knockout alleles in *pro*MpYUC2:MpCLE2-2 background

by CRISPR/Cas9-mediated genome editing (Figure S1). Both *Mpcik-3^{ge}* and *Mpclv1-4^{ge}* suppressed the supernumerary branching phenotype in 16-day-old plants (Figure 3A), which is consistent with the results in peptide treatment assay (Figure 2B). Furthermore, the number of gemma cup formed on thalli was reduced in *proMpYUC2:MpCLE2* and it was suppressed in both *Mpcik-3^{ge}* and *Mpclv1-4^{ge}* (Figure 3A). Time-course analysis showed that the gemma cup formation was significantly reduced and delayed in *proMpYUC2:MpCLE2* plants compared to wild type (Figure 3B). Meanwhile, all *Mpcik* and *Mpclv1* alleles showed minor increase in gemma cup formation compared to wild type (Figure 3B).

Biochemical interaction of MpCIK and MpCLV1 proteins

Since the genetic analysis suggests that MpCIK may function as a coreceptor for MpCLV1 for the perception of MpCLE2 peptide, we examined the biochemical interaction between MpCIK and MpCLV1 proteins expressed in a *Nicotiana benthamiana* transient expression system, which has been utilized to analyze the interaction of CLV and CIK receptors of Arabidopsis (Kinoshita et al., 2010; Betsuyaku et al. 2011; Hu et al. 2018). MpCIK, MpCLV1 and MpTDR were expressed under the control of 35S promoter in *N. benthamiana* as proteins C-terminally fused to 3× HAs-single StrepII or 3× FLAG (MpCIK-3HS, MpCLV1-3FLAG, MpTDR-3FLAG), respectively (Figure 4A). MpTDR, a receptor for MpCLE1, was also included in this interaction assay (Hirakawa et al. 2019). In co-immunoprecipitation experiments using an anti-HA affinity matrix, MpCLV1-3FLAG was detected not strongly but reproducibly in the immunoprecipitates containing MpCIK-3HS, suggesting a weak or transient interaction between MpCIK-3HS and MpCLV1-3FLAG (Figure 4B). Similarly, MpTDR-3FLAG was also shown to associate

weakly with MpCIK-3HS (Figure 4B). Thus, MpCLV1 and MpTDR are capable of interacting with MpCIK.

Figure 4. MpCIK weakly associates with MpCLV1 and MpTDR in *N. benthamiana*

(A) Schematic illustration of expressed receptors.

(B) Co-immunoprecipitation experiment using anti-HA affinity matrix. The indicated combinations of MpCIK-3HS, MpCLV1-3FLAG and MpTDR-3FLAG constructs, together with p19 silencing suppressor, were transiently expressed in *N. benthamiana*. Total proteins were extracted and immunoprecipitated with anti-HA affinity matrix. Immunoblot analyses were performed using anti-HA or anti-FLAG antibody. In the presence of MpCIK-3HS, MpCLV1-3FLAG and MpTDR-3FLAG were co-precipitated with anti-HA affinity matrix. This experiment was repeated twice with similar results.

MpCIK is sensitive to MpCLE1 and TDIF

Our biochemical data suggests that MpCIK could also function as a coreceptor for MpTDR upon the perception of MpCLE1, a TDIF-type CLE peptide. Previously, we showed that synthetic TDIF-type peptides cause slight reduction of overall growth and twisted lobes in *M. polymorpha* thalli (Hirakawa et al., 2019). In order to address the possible involvement of MpCIK in MpCLE1 signaling, we first examined the effects of TDIF, the strongest analog among known TDIF-type CLE peptides including MpCLE1 peptide. In 14-day-old plants grown from gemmae on growth medium supplemented with 3 μ M TDIF, both Tak-1 and MpCIK-1 showed slight reduction of overall growth and twist in the thallus lobes (Figure 5A), indicating that MpCIK is not necessary for the perception of TDIF. We further analyzed the effects of MpCLE1 overexpression using

proMpYUC2:MpCLE1 transformants. In the wild-type (Tak-1) background, *proMpYUC2:MpCLE1* resulted in small, twisted thalli in 14-day-old plants. This phenotype was not observed in *proMpYUC2:MpCLE1¹⁻⁴¹⁷*, a truncated version of *MpCLE1* lacking an essential asparagine residue in the CLE peptide motif (Figure 5B; Hirakawa et al., 2019). In the *Mpcik-1^{ge}* background, *proMpYUC2:MpCLE1* also resulted in small twisted thalli although the effects on growth was mild compared to the Tak-1 background, as judged from the ground cover area in 14-day-old plants (Figures 5B and 5C). These data suggests that *MpCIK* could be partially involved in *MpCLE1* perception.

Figure 5. Growth of *Mpcik* thalli is sensitive to *MpCLE1*/TDIF activity.

(A) Overall morphology of 14-day-old plants grown from gemmae on medium supplemented with or without TDIF peptide as indicated above. Side view panels show the images taken from the right in the top view panels as indicated by red arrowheads. Note that thalli are twisted in plants treated with TDIF, resulting in the uplift of the thalli from the medium as indicated by arrows.

(B) Overall morphology of 14-day-old plants grown from gemmae. *MpCLE1ox* indicates overexpression under *MpYUC2* promoter (*proMpYUC2:MpCLE1*) in Tak-1 and *Mpcik-1^{ge}* background. *MpCLE1(1-417)ox*, indicates a truncate version of *MpCLE1*. **(C)** Ground cover in 14-day-old plants (mean and SD; n= 11-14). Means sharing the superscripts are not significantly different from each other in Tukey's HSD test, $p < 0.05$. Scale bars represent 1 cm in (A) and (B).

Discussion

In this study, we performed functional analysis of MpCIK, the sole *M. polymorpha* ortholog of Arabidopsis CIK genes, and showed that MpCIK is essential for MpCLE2 peptide signaling to regulate the apical meristem activity in gametophyte. Biochemical analysis in *N. benthamiana* supports the idea that the MpCLV1-MpCIK is a receptor-coreceptor pair for MpCLE2 peptide perception. Since the same ligand-receptor-coreceptor relationship of CLV3-CLV1-CIK is consistently observed for multiple paralogs in Arabidopsis (Hu et al., 2018; Cui et al., 2018; Anne et al., 2018), this system is likely an evolutionarily conserved mechanism for the perception of CLV3-type CLE peptide in land plants. In addition, we suggested partial involvement of MpCIK in signaling of TDIF-type CLE peptide, MpCLE1, although MpCIK is not necessary for MpCLE1 signaling. Studies in Arabidopsis have suggested that subclass-II receptors in the SERK subgroup function as coreceptors for TDIF-PXY/TDR signaling (Zhang et al., 2016b). Further studies on MpSERK would clarify the contribution of different coreceptors. The phylogenetic analysis inferred the divergence of CIK and SERK subgroups in the common ancestor of land plants, which coincides with the appearance of subclass-XI genes. Recent studies have also shown that SERK-interacting receptors from subclass X, such as BRI1/BRL and EMS1, are also encoded in bryophytes (Ferreira-Guerra et al., 2020; Furumizu and Sawa 2021). Interestingly, the two sequences from *Spirogyra pratensis* (Figure 1B) showed high similarity to the subclass-II genes from land plants. Studies on Zygnematales algae would provide a clue to understand the evolution of these receptors.

Gemma cups are specialized structures for vegetative propagation, found in certain species of Marchantiopsida (Yasui et al., 2019; Kato et al., 2020). Gemma cup formation

initiates at the cells in the dorsal epidermis close to the apical meristem (Suzuki et al., 2020). We show that gain-of-function of MpCLE2 results in the delay of gemma cup formation although it is still unclear if the phenotypes in gemma cup formation can be uncoupled from the defects in the apical meristem. More detailed experiments will clarify the role for CLE in gemma cup formation. It is known that hormonal and environmental cues affect the formation of gemma cup (Flores-Sandoval et al., 2015; Aki et al., 2019; Li et al., 2020; Rico-Reséndiz et al., 2020). A possible role for MpCLE2 peptide signaling would be to mediate certain environmental cues to control the timing of gemma cup formation and thus clonal propagation, cooperatively with other hormonal inputs. In addition, involvement of CIK subgroup members into antiviral responses has been suggested in Arabidopsis (Fontes et al., 2004). Further studies of Mpcik knockout plants under various environmental conditions would provide a new insight into signals that allowed plants to survive on land.

Our biochemical data reveals that MpCIK is capable of associating with MpCLV1 or MpTDR in an ectopic and transient expression system of *N. benthamiana*. Furthermore, weak associations observed for MpCIK-MpCLV1 as well as MpCIK-MpTDR indicates possible requirement of other components in MpCIK-containing complex formation. For instance, in a ligand-induced dimerization model, receptor-coreceptor interaction can be induced by the perception of ligand at their ectodomains, which in turn allows for their kinase domains to transphosphorylate and activate signaling (Jaillais et al., 2011; Hohmann et al., 2018; Perraki et al., 2018). Thus, ligand and/or other membrane receptors could be required for strong MpCIK-MpCLV1/MpTDR association. With its genetic simplicity in CLE signaling, *M. polymorpha* will be a nice experimental system to address this point in future studies.

Materials and Methods

Phylogenetic analysis

Gene sequences of land plants were retrieved from Phytozome v12.1 database (<https://phytozome.jgi.doe.gov/pz/portal.html>) except for those of *Arabidopsis thaliana* (<https://www.arabidopsis.org/>), *Picea abies* (<http://congenie.org/>) and *Marchantia polymorpha* (<https://marchantia.info/>). Sequences of charophycean algae were reported in Bowman et al. 2017, obtained from transcriptome databases for *Spirogyra pratensis* (<http://www.ncbi.nlm.nih.gov/Traces/wgs/wgsviewer.cgi?val=GBSM01&search=GBSM01000000&display=scaffolds>) and *Coleochaete orbicularis* (<http://www.ncbi.nlm.nih.gov/Traces/wgs/wgsviewer.cgi?val=GBSL01&search=GBSL01000000&display=scaffolds>). Gene IDs and the protein sequences are listed in Table S1. Predicted protein sequences were aligned in Clustal W (<https://www.genome.jp/tools-bin/clustalw>). We excluded ambiguously aligned sequence to produce an alignment of 297 amino acid characters in the conserved cytosolic domain. Bayesian analysis was performed using MrBayes 3.2.7 (Ronquist et al., 2012). Two runs with four chains of Markov chain Monte Carlo (MCMC) iterations were performed for 1,500,000 generations, keeping one tree every 100 generations. The first 25% of the generations were discarded as burn-in and the remaining trees were used to calculate a 50% majority-rule tree. The standard deviation for the two MCMC iteration runs was below 0.01, suggesting that it was sufficient for the convergens of the two runs. Convergence was assessed by visual inspection of the plot of the log likelihood scores of the two runs calculated by MrBayes (Gelman and Rubin, 1992). Character matrix used for the Bayesian phylogenetic analysis is provided in Data Sheet S1.

Plant materials and growth conditions

Marchantia polymorpha male Takaragaike-1 (Tak-1) accession was used as wild type in this study. *M. polymorpha* plants were grown on half-strength Gamborg B5 medium (pH 5.5) solidified with 1.4% agar at 22 °C under continuous white light. *Nicotiana benthamiana* seeds were grown on BM2 soil (Berger) in a growth room at 23 °C under continuous LED light.

Peptide treatment

Synthetic peptides used in this study were analytically pure and dissolved in 0.1% TFA (trifluoroacetic acid) solution as stock solutions. For MpCLE2 peptide treatment, approximately 20 mature gemmae were floated on 2 mL liquid M51C medium containing 2% sucrose supplemented with 3 µM MpCLE2 peptide (KEVHypNGHypNPLHN) or mock (TFA) solution, in 12-well plates as described previously (Hirakawa et al., 2020). For the TDIF treatment, gemmae were plated on half-strength B5 agar plates supplemented with 3 µM TDIF (HEVHypSGHypNPISN) or mock (TFA) solution as described previously (Hirakawa et al., 2019).

Constructs

Primers and plasmids are listed in Tables S2 and S3. All plant transformation vectors were generated using the Gateway cloning system (Thermo Fisher Scientific, MA, USA). Gateway destination vectors are described in Kinoshita et al., 2010, Ishizaki et al., 2015 and Sugano et al., 2018, except for pMpGWB301-YUC2p, which was generated in this study. A 3032 bp DNA fragment of MpYUC2 promoter sequence flanking the translation initiation site was PCR amplified from pENTR-proMpYUC2 vector (Hirakawa et al.

2020) with a primer pair of MpYUC2prom3k_F_InFusion_XbaI and MpYUC2prom_R_InFusion_XbaI, and cloned into the *XbaI* digestion site of pMpGWB301 using In-Fusion HD Cloning Kit (Takara Bio, Shiga, Japan). For construction of *proMpYUC2:MpCLE1*, entry clones, pENTR-MpCLE1 and pENTR-MpCLE(1-417) (Hirakawa et al., 2019), were transferred to the pMpGWB301-YUC2p vector using Gateway LR Clonase II Enzyme mix (Thermo Fisher Scientific). For genome editing of *MpCIK*, a guide RNA was designed at the first exon/intron junction of Mp7g14210 using CRISPRdirect (<https://crispr.dbcls.jp/>) (Naito et al., 2015) and the plasmid for genome editing was constructed according to Sugano et al. 2018. For the expression of epitope-tagged receptors in *N. benthamiana*, coding sequences of *MpCIK*, *MpCLV1* and *MpTDR* were PCR amplified from *M. polymorpha* cDNA and cloned into pENTR/D-TOPO vector. Resultant entry clones (pENTR-MpCIK, pENTR-MpCLV1 and pENTR-MpTDR) were transferred to pXCSG-3FLAG or pXCSG-3HS vector using Gateway LR Clonase II Enzyme mix (Thermo Fisher Scientific).

Production of transgenic *M. polymorpha*

Transgenic *M. polymorpha* plants are listed in Table S3. *Agrobacterium*-mediated transformation of *M. polymorpha* was performed using regenerating thalli according to Kubota et al. 2013. CRISPR/Cas9-based genome editing was performed according to Sugano et al. 2018 and mutations in the guide RNA target loci were examined by direct sequencing of PCR product amplified from genome DNA samples with primers listed in Table S2. Genome editing of *MpCLV1* was performed as described previously (Hirakawa et al., 2020). Nomenclature of genes and mutants are according to Bowman et al. 2016.

Plant imaging and phenotypic measurement

Overall morphology of plants was observed under a digital microscope (DMS1000, Leica Microsystems, Wetzlar, Germany) or under a digital camera (TG-6, Olympus, Tokyo, Japan). For the quantification of ground cover area in plant images, blue color was extracted and quantified using ImageJ (Schneider et al., 2012). For the measurement of apical notch width, plants grown on liquid medium were individually transferred onto agar medium and imaged under a digital microscope (DMS 1000, Leica Microsystems). To quantify the apical notch width, distance between the rims of apical notch was measured on the obtained images using ImageJ (Schneider et al., 2012).

Promoter GUS assay

Individual plants were stained separately in 30-50 μ L GUS staining solution (50 mM sodium phosphate buffer pH 7.2, 1 mM potassium-ferrocyanide, 1 mM potassium-ferricyanide, 10 mM EDTA, 0.01% Triton X-100 and 1 mM 5-bromo-4-chloro-3-indolyl- β -D-glucuronic acid) at 37 °C in dark. GUS-stained samples were washed with water, cleared with ethanol and mounted with clearing solution for imaging under a light microscope (BX51, Olympus).

Transient expression in *N. benthamiana*

Agrobacterium tumefaciens strains GV3101 MP90RK carrying expression constructs were grown in YEB medium with appropriate antibiotics, harvested by centrifugation at 4,500 rpm for 10 min, and resuspended in infiltration buffer [10 mM MES (pH 5.7), 10 mM MgCl₂, 150 μ M acetosyringone]. The cultures were adjusted to an OD 600 of 1.0 and incubated at room temperature for at least 3 h prior to infiltration. Equal volumes of

cultures of different constructs were mixed for co-infiltration, and then mixed with agrobacterial cultures (OD 600 of 1.0) carrying the p19 silencing suppressor in a 1:1 ratio (Voinnet et al., 1999). The resulting cultures were infiltrated into leaves of 3- to 4-week-old *N. benthamiana*. The leaf samples were harvested 3 d after infiltration for subsequent protein extraction (Betsuyaku et al., 2011).

Protein extraction

Total protein was extracted from the infiltrated *N. benthamiana* leaves with IP extraction buffer (1 : 1 w/v, 50 mM Tris-HCl pH 8.0, 150 mM NaCl, 10 % glycerol, 1 % Triton X-100, 1×Proteinase inhibitor cocktail SIGMA P9599 and 1 mM EDTA) and incubate the extract at 4 °C for 30 min. The lysates were centrifuged at 20,000×g for 20 min at 4 °C and the supernatants were then centrifuged again at 20,000×g for 5 min at 4 °C. The resultant supernatants were used as total protein lysates.

Co-immunoprecipitation

For immunoprecipitation, 1 ml of the lysates prepared with IP extraction buffer from 0.5 g of leaves was incubated with anti-HA Affinity Matrix (Roche 11815016) for o/n in a rotary shaker at 4 °C. The beads were collected and washed three times with 1 ml of the extraction buffer. Immunoprecipitated proteins were eluted from the beads by boiling in SDS sample buffer at 95 °C and analyzed by Western blot using the corresponding antibodies. We used the following antibodies; Anti-HA-Peroxidase High Affinity (3F10) (Roche 12013819001) and Monoclonal ANTI-FLAG M2-Peroxidase (HRP) (SIGMA A8592).

Data Availability Statement

The original contributions presented in this study are included in the article/Supplementary Material.

Author Contributions

YH conceived the study, designed the work with input from all authors, and prepared the manuscript draft. GT, SB, NO and YH performed the experiments and analyzed the data. All authors contributed to the article and approved the submitted version.

Funding

This work was supported by grants from Japan Science and Technology Agency ERATO (JPMJER1502 to S.B.) and JSPS KAKENHI (JP19K06727 to Y.H.).

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgments

We thank colleagues in the peptide/protein center of WPI-ITbM for peptide synthesis, Eriko Betsuyaku and Ikuko Nakanomyo for technical assistance.

References

Aki, S.S., Mikami, T., Naramoto, S., Nishihama, R., Ishizaki, K., Kojima, M., et al. (2019). Cytokinin Signaling Is Essential for Organ Formation in *Marchantia polymorpha*. *Plant*

1 *Cell Physiol.* **60**:1842-1854.

2 Anne, P., Amiguet-Vercher, A., Brandt, B., Kalmbach, L., Geldner, N., Hothorn, M., et al.

3 (2018) CLERK is a novel receptor kinase required for sensing of root-active CLE peptides

4 in Arabidopsis. *Development* **145**:dev162354.

5 Araya, T., Miyamoto, M., Wibowo, J., Suzuki, A., Kojima, S., Tsuchiya, Y.N., et al. (2014).

6 CLE-CLAVATA1 peptide-receptor signaling module regulates the expansion of plant root

7 systems in a nitrogen-dependent manner. *Proc. Natl. Acad. Sci. U.S.A.* **111**:2029-2034.

8 Betsuyaku, S., Takahashi, F., Kinoshita, A., Miwa, H., Shinozaki, K., Fukuda, H., et al.

9 (2011). Mitogen-activated protein kinase regulated by the CLAVATA receptors

10 contributes to shoot apical meristem homeostasis. *Plant Cell Physiol.* **52**:14-29.

11 Bowman, J.L., Araki, T., Arteaga-Vazquez, M.A., Berger, F., Dolan, L., Haseloff, J., et al.

12 (2016). The Naming of Names: Guidelines for Gene Nomenclature in Marchantia. *Plant*

13 *Cell Physiol.* **57**:257-261.

14 Bowman, J.L., Kohchi, T., Yamato, K.T., Jenkins, J., Shu, S., Ishizaki, K., et al. (2017).

15 Insights into Land Plant Evolution Garnered from the Marchantia polymorpha

16 Genome. *Cell* **171**:287-304.

17 Cock, J.M., and McCormick, S. (2001). A large family of genes that share homology with

18 CLAVATA3. *Plant Physiol.* **126**:939-942.

19 Crook, A.D., Willoughby, A.C., Hazak, O., Okuda, S., VanDerMolen, K.R., Soyars, C.L.,

20 Cattaneo, P., Clark, N.M., Sozzani, R., Hothorn, M., Hardtke, C.S. and Nimchuk, Z.L.

21 (2020). BAM1/2 receptor kinase signaling drives CLE peptide-mediated formative cell

22 divisions in Arabidopsis roots. *Proc. Natl. Acad. Sci. U.S.A.* **117**:32750-32756.

23 Cui, Y., Hu, C., Zhu, Y., Cheng, K., Li, X., Wei, Z., et al. (2018). CIK Receptor Kinases

24 Determine Cell Fate Specification during Early Anther Development in

1 Arabidopsis. *Plant Cell* **30**:2383-2401.

2 Czyzewicz, N., Shi, C.L., Vu, L.D., Van De Cotte, B., Hodgman, C., Butenko, M.A., et
3 al. (2015). Modulation of Arabidopsis and monocot root architecture by
4 CLAVATA3/EMBRYO SURROUNDING REGION 26 peptide. *J. Exp. Bot.* **66**:5229-
5 5243.

6 DeYoung, B.J., Bickle, K.L., Schrage, K.J., Muskett, P., Patel, K., and Clark, S.E. (2006).
7 The CLAVATA1-related BAM1, BAM2 and BAM3 receptor kinase-like proteins are
8 required for meristem function in Arabidopsis. *Plant J.* **45**:1-16.

9 Depuydt, S., Rodriguez-Villalon, A., Santuari, L., Wyser-Rmili, C., Ragni, L., and
10 Hardtke, C.S. (2013). Suppression of Arabidopsis protophloem differentiation and root
11 meristem growth by CLE45 requires the receptor-like kinase BAM3. *Proc. Natl. Acad.*
12 *Sci. U.S.A.* **110**:7074-7079.

13 Doblas, V.G., Smakowska-Luzan, E., Fujita, S., Alassimone, J., Barberon, M., Madalinski,
14 M., et al. (2017). Root diffusion barrier control by a vasculature-derived peptide binding
15 to the SGN3 receptor. *Science* **355**:280-284.

16 Doll, N.M., Royek, S., Fujita, S., Okuda, S., Chamot, S., Stintzi, A., et al. (2020). A two-
17 way molecular dialogue between embryo and endosperm is required for seed
18 development. *Science* **367**:431-435.

19 Eklund, D.M., Ishizaki, K., Flores-Sandoval, E., Kikuchi, S., Takebayashi, Y., Tsukamoto,
20 S., et al. (2015). Auxin Produced by the Indole-3-Pyruvic Acid Pathway Regulates
21 Development and Gemmae Dormancy in the Liverwort *Marchantia polymorpha*. *Plant*
22 *Cell* **27**:1650-1669.

23 Endo, S., Shinohara, H., Matsubayashi, Y., and Fukuda, H. (2013). A novel pollen-pistil
24 interaction conferring high-temperature tolerance during reproduction via CLE45

1 signaling. *Curr. Biol.* **23**:1670-1676.

2 Etchells, J.P., and Turner, S.R. (2010). The PXY-CLE41 receptor ligand pair defines a
3 multifunctional pathway that controls the rate and orientation of vascular cell
4 division. *Development* **137**:767-774.

5 Ferreira-Guerra, M., Marquès-Bueno, M., Mora-García, S., and Caño-Delgado, A.I.
6 (2020). Delving into the evolutionary origin of steroid sensing in plants. *Curr. Opin. Plant*
7 *Biol.* **57**:87-95.

8 Fisher, K., and Turner, S. (2007). PXY, a receptor-like kinase essential for maintaining
9 polarity during plant vascular-tissue development. *Curr. Biol.* **17**:1061-1066.

10 Fletcher, J.C., Brand, U., Running, M.P., Simon, R., and Meyerowitz, E.M. (1999).
11 Signaling of cell fate decisions by CLAVATA3 in Arabidopsis shoot
12 meristems. *Science* **283**:1911-1914.

13 Fletcher, J.C. (2020). Recent Advances in Arabidopsis CLE Peptide Signaling. *Trends.*
14 *Plant. Sci.* **25**:1005-1016.

15 Flores-Sandoval, E., Eklund, D.M., and Bowman, J.L. (2015). A Simple Auxin
16 Transcriptional Response System Regulates Multiple Morphogenetic Processes in the
17 Liverwort *Marchantia polymorpha*. *PLoS Genet.* **11**:e1005207.

18 Fontes, E.P., Santos, A.A., Luz, D.F., Waclawovsky, A.J., and Chory, J. (2004). The
19 geminivirus nuclear shuttle protein is a virulence factor that suppresses transmembrane
20 receptor kinase activity. *Genes. Dev.* **18**:2545-2556.

21 Furumizu, C., and Sawa, S. (2021). Insight into early diversification of leucine-rich repeat
22 receptor-like kinases provided by the sequenced moss and hornwort genomes. *Plant Mol.*
23 *Biol.* Epub ahead of print.

24 Gelman, A., and Rubin, D.B. (1992) Inference from Iterative Simulation Using Multiple

1 Sequences. *Statist. Sci.* **7**:457-472.

2 Gou, X., and Li, J. (2020). Paired Receptor and Coreceptor Kinases Perceive Extracellular
3 Signals to Control Plant Development. *Plant Physiol.* **182**:1667-1681.

4 Gutiérrez-Alanís, D., Yong-Villalobos, L., Jiménez-Sandoval, P., Alatorre-Cobos, F.,
5 Oropeza-Aburto, A., Mora-Macías, J., et al. (2017). Phosphate Starvation-Dependent Iron
6 Mobilization Induces CLE14 Expression to Trigger Root Meristem Differentiation
7 through CLV2/PEPR2 Signaling. *Dev. Cell* **41**:555-570.

8 Hirakawa, Y., Fujimoto, T., Ishida, S., Uchida, N., Sawa, S., Kiyosue, T., et al. (2020).
9 Induction of Multichotomous Branching by CLAVATA Peptide in *Marchantia*
10 *polymorpha*. *Curr. Biol.* **30**:3833-3840.

11 Hirakawa, Y., Kondo, Y., and Fukuda, H. (2010). TDIF peptide signaling regulates
12 vascular stem cell proliferation via the WOX4 homeobox gene in *Arabidopsis*. *Plant*
13 *Cell* **22**:2618-2629.

14 Hirakawa, Y., and Sawa, S. (2019). Diverse function of plant peptide hormones in local
15 signaling and development. *Curr. Opin. Plant. Biol.* **51**:81-87.

16 Hirakawa, Y., Shinohara, H., Kondo, Y., Inoue, A., Nakanomyo, I., Ogawa, M., et al.
17 (2008). Non-cell-autonomous control of vascular stem cell fate by a CLE peptide/receptor
18 system. *Proc. Natl. Acad. Sci. U.S.A.* **105**:15208-15213.

19 Hirakawa, Y., Shinohara, H., Welke, K., Irle, S., Matsubayashi, Y., Torii, K.U., et al.
20 (2017). Cryptic bioactivity capacitated by synthetic hybrid plant peptides. *Nat.*
21 *Commun.* **8**:14318-.

22 Hirakawa, Y., Uchida, N., Yamaguchi, Y.L., Tabata, R., Ishida, S., Ishizaki, K., et al.
23 (2019). Control of proliferation in the haploid meristem by CLE peptide signaling in
24 *Marchantia polymorpha*. *PLoS Genet.* **15**:e1007997.

1 Hohmann, U., Lau, K., and Hothorn, M. (2017). The Structural Basis of Ligand
2 Perception and Signal Activation by Receptor Kinases. *Annu. Rev. Plant Biol.* **68**:109-
3 137.

4 Hohmann, U., Santiago, J., Nicolet, J., Olsson, V., Spiga, F.M., Hothorn, L.A., et al.
5 (2018). Mechanistic basis for the activation of plant membrane receptor kinases by
6 SERK-family coreceptors. *Proc. Natl. Acad. Sci. U.S.A.* **115**:3488-3493.

7 Hu, C., Zhu, Y., Cui, Y., Cheng, K., Liang, W., Wei, Z., et al. (2018). A group of receptor
8 kinases are essential for CLAVATA signalling to maintain stem cell homeostasis. *Nat.*
9 *Plants* **4**:205-211.

10 Ishizaki, K., Nishihama, R., Ueda, M., Inoue, K., Ishida, S., Nishimura,
11 Y., et al. (2015). Development of Gateway Binary Vector Series with Four Different
12 Selection Markers for the Liverwort *Marchantia polymorpha*. *PLoS One* **10**:e0138876.

13 Ito, Y., Nakanomyo, I., Motose, H., Iwamoto, K., Sawa, S., Dohmae, N., et al. (2006).
14 Dodeca-CLE peptides as suppressors of plant stem cell differentiation. *Science* **313**:842-
15 845.

16 Jaillais, Y., Belkhadir, Y., Balsemão-Pires, E., Dangl, J.L., and Chory, J. (2011).
17 Extracellular leucine-rich repeats as a platform for receptor/coreceptor complex
18 formation. *Proc. Natl. Acad. Sci. U.S.A.* **108**:8503-8507.

19 Jun, J., Fiume, E., Roeder, A.H., Meng, L., Sharma, V.K., Osmont, K.S., et al. (2010).
20 Comprehensive analysis of CLE polypeptide signaling gene expression and
21 overexpression activity in *Arabidopsis*. *Plant Physiol.* **154**:1721-1736.

22 Kato, H., Yasui, Y., and Ishizaki, K. (2020). Gemma cup and gemma development in
23 *Marchantia polymorpha*. *New Phytol.* **228**:459-465.

24 Kinoshita, A., Betsuyaku, S., Osakabe, Y., Mizuno, S., Nagawa, S., Stahl, Y., et al. (2010).
RPK2 is an essential receptor-like kinase that transmits the CLV3 signal in

1 *Arabidopsis. Development* **137**:3911-3920.

2 Kondo, Y., Hirakawa, Y., Kieber, J.J., and Fukuda, H. (2011). CLE peptides can
3 negatively regulate protoxylem vessel formation via cytokinin signaling. *Plant Cell*
4 *Physiol.* **52**:37-48.

5 Kubota, A., Ishizaki, K., Hosaka, M., and Kohchi, T. (2013). Efficient *Agrobacterium*-
6 mediated transformation of the liverwort *Marchantia polymorpha* using regenerating
7 thalli. *Biosci. Biotechnol. Biochem.* **77**:167-172.

8 Li, D., Flores-Sandoval, E., Ahtesham, U., Coleman, A., Clay, J.M., Bowman, J.L. et al.
9 (2020). Ethylene-independent functions of the ethylene precursor ACC in *Marchantia*
10 *polymorpha*. *Nat. Plants* **6**:1335-1344.

11 Ma, D., Endo, S., Betsuyaku, S., Shimotohno, A., and Fukuda, H. (2020). CLE2 regulates
12 light-dependent carbohydrate metabolism in *Arabidopsis* shoots. *Plant Mol.*
13 *Biol.* **104**:561-574.

14 Matsubayashi, Y. (2014). Posttranslationally modified small-peptide signals in
15 plants. *Annu. Rev. Plant Biol.* **65**:385-413.

16 Montgomery, S.A., Tanizawa, Y., Galik, B., Wang, N., Ito, T., Mochizuki, T., et al. (2020).
17 Chromatin Organization in Early Land Plants Reveals an Ancestral Association between
18 H3K27me3, Transposons, and Constitutive Heterochromatin. *Curr. Biol.* **30**:573-588.

19 Morita, J., Kato, K., Nakane, T., Kondo, Y., Fukuda, H., Nishimasu, H., et al. (2016).
20 Crystal structure of the plant receptor-like kinase TDR in complex with the TDIF
21 peptide. *Nat. Commun.* **7**:12383-.

22 Morris, J.L., Puttick, M.N., Clark, J.W., Edwards, D., Kenrick, P., Pressel, S., et al. (2018).
23 The timescale of early land plant evolution. *Proc. Natl. Acad. Sci. U.S.A.* **115**:E2274-
24 E2283.

1 Mortier, V., Den Herder, G., Whitford, R., Van de Velde, W., Rombauts, S., D'Haeseleer,
2 et al. (2010). CLE peptides control *Medicago truncatula* nodulation locally and
3 systemically. *Plant Physiol.* **153**:222-237.

4 Murphy, E., Smith, S., and De Smet, I. (2012). Small signaling peptides in Arabidopsis
5 development: how cells communicate over a short distance. *Plant Cell* **24**:3198-3217.

6 Nakayama, T., Shinohara, H., Tanaka, M., Baba, K., Ogawa-Ohnishi, M., and
7 Matsubayashi, Y. (2017). A peptide hormone required for Casparian strip diffusion barrier
8 formation in Arabidopsis roots. *Science* **355**:284-286.

9 Naito, Y., Hino, K., Bono, H., and Ui-Tei, K. (2015). CRISPRdirect: software for
10 designing CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics* **31**:1120-
11 1123.

12 Oelkers, K., Goffard, N., Weiller, G.F., Gresshoff, P.M., Mathesius, U., and Frickey, T.
13 (2008). Bioinformatic analysis of the CLE signaling peptide family. *BMC Plant Biol.* **8**:1.

14 Ogawa, M., Shinohara, H., Sakagami, Y., and Matsubayashi, Y. (2008). Arabidopsis
15 CLV3 peptide directly binds CLV1 ectodomain. *Science* **319**:294.

16 Ohyama, K., Shinohara, H., Ogawa-Ohnishi, M., and Matsubayashi, Y. (2009). A
17 glycopeptide regulating stem cell fate in Arabidopsis thaliana. *Nat. Chem. Biol.* **5**:578-
18 580.

19 Okamoto, S., Ohnishi, E., Sato, S., Takahashi, H., Nakazono, M., Tabata, S., et al. (2009).
20 Nod factor/nitrate-induced CLE genes that drive HAR1-mediated systemic regulation of
21 nodulation. *Plant Cell Physiol.* **50**:67-77.

22 Okuda, S., Fujita, S., Moretti, A., Hohmann, U., Doblas, V.G., Ma, Y., et al. (2020).
23 Molecular mechanism for the recognition of sequence-divergent CIF peptides by the plant
24 receptor kinases GSO1/SGN3 and GSO2. *Proc. Natl. Acad. Sci. U.S.A.* **117**:2693-2703.

1 Ou, Y., Lu, X., Zi, Q., Xun, Q., Zhang, J., Wu, Y., et al. (2016). RGF1 INSENSITIVE 1
2 to 5, a group of LRR receptor-like kinases, are essential for the perception of root
3 meristem growth factor 1 in *Arabidopsis thaliana*. *Cell Res.* **26**:686-698.

4 Perraki, A., DeFalco, T.A., Derbyshire, P., Avila, J., Séré, D., Sklenar, J., et al. (2018).
5 Phosphocode-dependent functional dichotomy of a common co-receptor in plant
6 signalling. *Nature* **561**:248-252.

7 Puttick, M.N., Morris, J.L., Williams, T.A., Cox, C.J., Edwards, D., Kenrick, P., et al.
8 (2018). The Interrelationships of Land Plants and the Nature of the Ancestral
9 Embryophyte. *Curr. Biol.* **28**:733-745.

10 Qian, P., Song, W., Yokoo, T., Minobe, A., Wang, G., Ishida, T., et al. (2018). The
11 CLE9/10 secretory peptide regulates stomatal and vascular development through distinct
12 receptors. *Nat. Plants* **4**:1071-1081.

13 Rico-Reséndiz, F., Cervantes-Pérez, S.A., Espinal-Centeno, A., Dipp-Álvarez, M.,
14 Oropeza-Aburto, A., Hurtado-Bautista, E., et al. (2020) Transcriptional and Morpho-
15 Physiological Responses of *Marchantia polymorpha* upon Phosphate Starvation. *Int. J.*
16 *Mol. Sci.* **21**:8354.

17 Rodríguez-Villalon, A., Gujas, B., Kang, Y.H., Breda, A.S., Cattaneo, P., Depuydt, S., et
18 al. (2014). Molecular genetic framework for protophloem formation. *Proc. Natl. Acad.*
19 *Sci. U.S.A.* **111**:11551-11556.

20 Rodríguez-Leal, D., Lemmon, Z.H., Man, J., Bartlett, M.E., and Lippman, Z.B. (2017).
21 Engineering Quantitative Trait Variation for Crop Improvement by Genome
22 Editing. *Cell* **171**:470-480.

23 Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., et al.
24 (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across

1 a large model space. *Syst. Biol.* **61**:539-542.

2 Santiago, J., Brandt, B., Wildhagen, M., Hohmann, U., Hothorn, L.A., Butenko, M.A., et

3 al. (2016) Mechanistic insight into a peptide hormone signaling complex mediating floral

4 organ abscission. *Elife* **5**:e15075.

5 Santiago, J., Henzler, C., and Hothorn, M. (2013). Molecular mechanism for plant steroid

6 receptor activation by somatic embryogenesis co-receptor kinases. *Science* **341**:889-892.

7 Sasaki, G., Katoh, K., Hirose, N., Suga, H., Kuma, K., Miyata, T., et al. (2007). Multiple

8 receptor-like kinase cDNAs from liverwort *Marchantia polymorpha* and two

9 charophycean green algae, *Closterium ehrenbergii* and *Nitella axillaris*: Extensive gene

10 duplications and gene shufflings in the early evolution of streptophytes. *Gene* **401**:135-

11 144.

12 Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25

13 years of image analysis. *Nat. Methods* **9**:671-675.

14 Shimizu, N., Ishida, T., Yamada, M., Shigenobu, S., Tabata, R., Kinoshita, A., et al. (2015).

15 BAM 1 and RECEPTOR-LIKE PROTEIN KINASE 2 constitute a signaling pathway and

16 modulate CLE peptide-triggered growth inhibition in *Arabidopsis* root. *New*

17 *Phytol.* **208**:1104-1113.

18 Shinohara, H., and Matsubayashi, Y. (2015). Reevaluation of the CLV3-receptor

19 interaction in the shoot apical meristem: dissection of the CLV3 signaling pathway from

20 a direct ligand-binding point of view. *Plant J.* **82**:328-336.

21 Shinohara, H., Mori, A., Yasue, N., Sumida, K., and Matsubayashi, Y. (2016).

22 Identification of three LRR-RKs involved in perception of root meristem growth factor

23 in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* **113**:3897-3902.

24 Shiu, S.H., and Bleecker, A.B. (2001). Receptor-like kinases from *Arabidopsis* form a

1 monophyletic gene family related to animal receptor kinases. *Proc. Natl. Acad. Sci.*
2 *U.S.A.* **98**:10763-10768.

3 Song, W., Liu, L., Wang, J., Wu, Z., Zhang, H., Tang, J., et al. (2016). Signature motif-
4 guided identification of receptors for peptide hormones essential for root meristem
5 growth. *Cell Res.* **26**:674-685.

6 Stahl, Y., Wink, R.H., Ingram, G.C., and Simon, R. (2009). A signaling module
7 controlling the stem cell niche in Arabidopsis root meristems. *Curr. Biol.* **19**:909-914.

8 Sugano, S.S., Nishihama, R., Shirakawa, M., Takagi, J., Matsuda, Y., Ishida, S., et al.
9 (2018). Efficient CRISPR/Cas9-based genome editing and its application to conditional
10 genetic analysis in *Marchantia polymorpha*. *PLoS One* **13**:e0205117.

11 Sun, Y., Han, Z., Tang, J., Hu, Z., Chai, C., Zhou, B., et al. (2013). Structure reveals that
12 BAK1 as a co-receptor recognizes the BRI1-bound brassinolide. *Cell Res* **23**:1326-1329.

13 Suzaki, T., Yoshida, A., and Hirano, H.Y. (2008). Functional diversification of
14 CLAVATA3-related CLE proteins in meristem maintenance in rice. *Plant Cell* **20**:2049-
15 2058.

16 Suzuki, H., Harrison, C.J., Shimamura, M., Kohchi, T., and Nishihama, R. (2020).
17 Positional cues regulate dorsal organ formation in the liverwort *Marchantia*
18 *polymorpha*. *J. Plant Res.* **133**:311-321.

19 Tabata, R., Sumida, K., Yoshii, T., Ohyama, K., Shinohara, H., and Matsubayashi, Y.
20 (2014). Perception of root-derived peptides by shoot LRR-RKs mediates systemic N-
21 demand signaling. *Science* **346**:343-346.

22 Takahashi, F., Suzuki, T., Osakabe, Y., Betsuyaku, S., Kondo, Y., Dohmae, N., et al.
23 (2018). A small peptide modulates stomatal control via abscisic acid in long-distance
24 signalling. *Nature* **556**:235-238.

1 Tamaki, T., Betsuyaku, S., Fujiwara, M., Fukao, Y., Fukuda, H., and Sawa, S. (2013).
2 SUPPRESSOR OF LLP1 1-mediated C-terminal processing is critical for CLE19 peptide
3 activity. *Plant J.* **76**:970-981.

4 Voinnet, O., Pinto, Y.M., and Baulcombe, D.C. (1999) Suppression of gene silencing: a
5 general strategy used by diverse DNA and RNA viruses of plants. *Proc. Natl. Acad. Sci.*
6 *U.S.A.* **96**:14147-14152. Erratum in: *Proc. Natl. Acad. Sci. U.S.A.* (2015)
7 **112**:E4812. Wang, J., Li, H., Han, Z., Zhang, H., Wang, T., Lin, G., et al. (2015). Allosteric
8 receptor activation by the plant peptide hormone phytosulfokine. *Nature* **525**:265-268.

9 Whitewoods, C.D., Cammarata, J., Nemec Venza, Z., Sang, S., Crook, A.D., Aoyama, T.,
10 et al. (2018). CLAVATA Was a Genetic Novelty for the Morphological Innovation of 3D
11 Growth in Land Plants. *Curr. Biol.* **28**:2365-2376.

12 Yamaguchi, Y., Pearce, G., and Ryan, C.A. (2006). The cell surface leucine-rich repeat
13 receptor for AtPep1, an endogenous peptide elicitor in Arabidopsis, is functional in
14 transgenic tobacco cells. *Proc. Natl. Acad. Sci. U.S.A.* **103**:10104-10109.

15 Yasui, Y., Tsukamoto, S., Sugaya, T., Nishihama, R., Wang, Q., Kato, H., et al. (2019).
16 GEMMA CUP-ASSOCIATED MYB1, an Ortholog of Axillary Meristem Regulators, Is
17 Essential in Vegetative Reproduction in *Marchantia polymorpha*. *Curr. Biol.* **29**:3987-
18 3995.

19 Zhang, H., Lin, X., Han, Z., Qu, L.J., and Chai, J. (2016a). Crystal structure of PXY-TDIF
20 complex reveals a conserved recognition mechanism among CLE peptide-receptor
21 pairs. *Cell Res.* **26**:543-555.

22 Zhang, H., Lin, X., Han, Z., Wang, J., Qu, L.J., and Chai, J. (2016b). SERK Family
23 Receptor-like Kinases Function as Co-receptors with PXY for Plant Vascular
24 Development. *Mol. Plant* **9**:1406-1414.