1 **Difference in hoof conformation between shod and barefoot-** 2 **managed hooves**

3 J.N. DE KLERK^{*}, BVetMed (Hons) MSTAHCRCVS

4 ^{*}E-mail for correspondence: johardvet@gmail.com

6 **Authors' declaration of interests**

7 No competing interests have been declared.

9 **Ethical animal research**

10 Ethical approval was granted by the Royal Veterinary College on 19th March 2013 (Reference Number
11 2013/R62).

13 **Source of funding**

14 The research was sole funded by the author.

16 **Acknowledgements**

17 The author wishes to acknowledge Mr Neil Nevison for constructing a photographic rig to the author's design and
18 Ms Jenny Butler-Smith, Miss Emma Delport, Mrs Deborah Mead, Mrs Kirsty Stuart, Mrs Diane Goddard, Mrs
19 Shirley Exall, Miss Eva Cox, Mrs Marianne Cox, Ms Sarah Mollett, Miss Lisa Housden, Miss Harriet Bradford,
20 Mrs Auriol Thorne, Ms Bevni Lyons, Miss Emma Burston, Mr Wayne Upton, Mr Jack Hopson, Ms Teresa Mallia,
21 Ms Teresa Watson, Miss Jo Ricey and Miss Natasha Crawley for aiding in the recruitment of horses for the
22 study.

24 **Authorship**

25 J.N. de Klerk contributed to the study design, data collection, analysis and writing of the study.

30 Abstract

31

32 **Reasons for performing the study:** Hoof conformation is linked to biomechanics of the hoof and injury
33 occurrence. There is no scientific data if conformation differs between shod and barefoot-managed hooves.

34 **Objectives:** To investigate if and how shod and barefoot hooves differ in conformation.

35 **Study design:** Retrospective cohort study.

36 **Methods:** Standardised lateral, dorsopalmar/dorsoplantar and solar photographs of 98 shod and 69 barefoot-
37 managed hooves were included. Thirty-six of the barefoot horses were farrier-managed, 33 were podiatrist-
38 managed. Length and angular measurements produced nine conformation parameters; dorsopalmar/plantar
39 balance, solar symmetry, toe angle, heel angle, heel/toe angle difference, heel width, splaying index, flaring index
40 and frog size.

41 **Results:** Barefoot hooves showed significantly fewer underrun heels, steeper heel angles, wider heels,
42 increased splaying, increased flaring and larger frog size compared to hooves of shod horses. Solar symmetry
43 showed a significant difference in front hooves but not hind hooves ($P=0.038$, $P=0.104$) and toe angle was not
44 significantly different ($P=0.368$, $P=0.425$). There was no significant difference in the conformation of barefoot
45 farrier and podiatrist-managed front hooves, however there was a significant difference in the hind hooves:
46 farrier-managed hooves had longer frogs and shorter toes, compared to podiatrist-managed hooves.

47 **Conclusions:** The significant differences in hoof conformation found should be considered when managing the
48 individual horse, since hoof conformation affects loading of the internal structure of the hoof and hence influences
49 aetiopathogenesis of hoof pathology.

50

51 **Keywords:** horse; barefoot; shoes; hoof, conformation; angles

52

53 Introduction

54

55 Horses are shod for comfort over all surfaces, particularly hard or rough terrains, and to prevent mechanical
56 damage to the hoof capsule, such as splitting, bruising, cracking and excessive wear [1]. Good shoeing
57 technique ensures suitable mediolateral and dorsopalmar/dorsoplantar hoof balance and provides support for
58 internal structures. Shoes can also influence load distribution, for example by fitting extensions to address
59 imbalances [2]. Many acknowledge that horseshoes provide support and can prevent or improve many hoof-
60 related pathologies.

61

62 Some parties challenge shoeing, believing that shoes are cruel [3] and cause discomfort and lameness. In the
63 late 20th century, a “natural” barefoot hoof management was introduced collaboratively between a veterinarian
64 and a farrier [4]. However, these concepts were considered radical, as many horses endured hoof pain and sub-
65 solar abscess formation during the post-shoe removal transition period. Now, less radical trimming programmes
66 have been introduced, advocated by the Equine Podiatry Association and the Institute of Applied Equine Podiatry
67 [5].

68

69 Trimming and shoeing influences hoof conformation which affects the loading of the structures within the hoof
70 and higher up the limb. One degree change in solar angle induces a 4% change in the strain of the digital flexor
71 tendon and hence pressure experienced by the navicular bone [6]. These findings were verified by the results of
72 a study where hoof conformation differed significantly between lame horses suffering from different lesions in the
73 hoof [7]. Hoof growth and management influences locomotor biomechanics [8, 9] and hoof conformation [10].
74 The latter study investigated how hoof conformation changed over a 16-month period once implementing
75 standardised barefoot trimming. This work demonstrated that the toe increased in length, the heel angle
76 steepened and that the frog contact area increased significantly [10]. These changes were considered beneficial,
77 particularly the raised heel height. A 5% increase in heel angle results in significantly less stress and
78 displacement of distal limb structures [11]. Furthermore, an increased contact-area of the frog aids the hoof’s
79 haemodynamic mechanism by assisting healing, growth and energy dissipation [12, 13].

80

81 There is limited scientific evidence about how hoof management influences hoof conformation. This study
82 compared hoof conformation of horses managed barefoot and shod and hypothesised there would be a
83 significant difference in hoof conformation between the two groups. In particular:

- 84 1) Lateromedial asymmetry, e.g. splaying, is more common in barefoot-managed horses.
- 85 2) Dorsopalmar/plantar hoof imbalance such as under-run heels, boxy hooves and long toes are more common
86 in shod horses
- 87 3) Barefoot horses have wider, larger frogs and larger width between heels than shod horses

88 In the UK, trimming of hooves is not regulated and is not limited to farriers, but is routinely performed by
89 podiatrists. Training of farriers and podiatrists differs and we hypothesise:

- 90 4) There will be a significant difference in hoof conformation between barefoot-managed hooves managed by
91 farriers compared to podiatrists.

92

93 **Materials and Methods**

94

95 *Horses*

96 Ninety-eight shod and 69 barefoot hooves were included in the study from 46 horses. Twenty-four horses were
 97 geldings and 22 were mares. All horses included in the study were from a general leisure horse population,
 98 regularly exercised, but not in competition work or heavy training routines. The horses were between 5 and 22
 99 years old with a mean±SD age of 12.1±4.8 years. Heights ranged from 12.3hh to 17.3hh. All horses were
 100 maintained exclusively either shod or barefoot for more than 18 months before the study and on a hoof
 101 management regimen of either shoeing with standard shoes or kept barefoot with trimming performed every 5-7
 102 weeks in both groups. A farrier or podiatrist who had been working full time for no less than 2 years, had received
 103 certified training in their area of work, and had undergone continual professional development since being
 104 awarded their degree, performed the shoeing and hoof trimming. The horses were split into 2 groups; shod,
 105 barefoot and the latter was further subdivided in farrier-managed and podiatrist-managed horses (36 farrier-
 106 managed, 33 podiatrist-managed).

107

108 *Conformation Measurements*

109 Frontal, lateral and solar views were obtained from each front and hind hoof within two days of shoeing or
 110 trimming using a digital camera (Samsung WB250): Frontal and lateral photographs were obtained centring on
 111 the centre of the hoof with the camera set at a 30cm distance away using of a custom-made rig. For the solar
 112 photograph, the camera was set to be 30cm away from the apex of the frog. Linear and angular measurements
 113 were performed in Image J (<http://rsbweb.nih.gov/ij/links.html>) (table 1). Additionally, circumference at the
 114 coronary band (cm) and circumference of the sole (cm) were measured with a measuring tape. Type of horse-
 115 build based on height, weight and stature (light-weight/medium-weight/heavy-weight), gender, age and exercise
 116 regimen was also recorded.

117

118 A repeatability assessment of the measurement technique was performed by measuring each parameter twice
 119 and calculating the limits of agreement [14, 15] (table 2).

120

121

122

123

124

Data analysis

Data were tested for normality using the Shapiro-Wilk test and conformation parameters were compared between the shod and barefoot-managed hooves and between farrier and podiatrist-managed hooves using a T-test or Mann Whitney U test depending on the distribution of the individual conformation parameter. Horses were also grouped according to their stature and stature-type was also included in the analysis using a Kruskal-Wallis or One Way ANOVA test to assess if it was a confounding factor. When analysing stature-type, it was found that it was a confounding factor for dorsopalmar/dorsoplantar balance ($P=0.004$), heel angle ($P=0.013$), and frog size ($P=0.001$) and so it was ensured that the same ratio of light-weight, medium-weight and heavy-weight horses were included in each group for further statistical analysis. Left and right hooves were also analysed to ensure the values could be combined. For all parameters, there was no statistical significance, so left and right hooves were combined. All parameters were analysed in SPSS (IBM, version 22.0.0.1). P was set at 0.05.

Results

Comparison of conformation parameters of shod and barefoot-managed horses

Table 3 shows the hoof conformation parameters of the hooves measured in this study for shod and barefoot-managed hooves.

The toe angle in front and hind hooves did not differ significantly between shod and barefoot-managed hooves. The heel angle however differed significantly between the two groups in both front and hind hooves. Shod hooves had a shallower heel angle compared to barefoot-managed hooves. In front hooves, the range of toe angle was much greater in shod hooves than in the barefoot hooves and the heel angle range was much greater for shod hooves in both, front and hind feet. There was a significant difference between the two groups in the difference in toe and heel angles in both front and hind hooves. Shod hooves had larger differences between toe and heel angles than barefoot hooves. There was a high prevalence of under-run heels, defined by a difference of more than 5 degrees [16], observed in both groups. 75.5% of hooves were under-run in the shod group and 40.5% were under-run in the barefoot group.

Barefoot-managed hooves showed a significantly greater heel width compared to shod hooves in front and hind. There was a significant difference in dorsopalmar/dorsoplantar balance between both groups for both front and hind hooves with barefoot hooves showing significantly shorter toes compared to shod hooves. For both hoof management regimens, the hooves were on average wider laterally. In the front hooves, barefoot hooves were significantly more asymmetrical compared to shod hooves, but no significant difference in asymmetry was found between the two groups in the hind hooves.

There was a significant difference between hoof management regimens for splaying and flaring in both front and hind hooves with barefoot hooves being more splayed and flared than shod hooves.

There was a significant difference between hoof management regimens for frog area in both front and hind hooves with barefoot hooves having significantly larger frogs than shod hooves.

Comparison of conformation parameters of barefoot hooves managed by a podiatrist and managed by a farrier

Table 4 shows the difference in hoof conformation parameters of barefoot hooves managed by a farrier compared to a podiatrist.

There was no significant difference between farrier and podiatrist-managed front and hind hooves for heel angle, solar symmetry, toe angle, difference in toe and heel angles, heel width, splaying, flaring or frog size.

There was no significant difference between farrier and podiatrist-managed front hooves for dorsopalmar balance, however there was a significant difference in the hind hooves. Farrier-managed hooves had longer frogs and shorter toes, compared to podiatrist-managed hooves.

Discussion

This study showed that some hoof conformation parameters differed significantly between shod and barefoot-managed hooves and between farrier and podiatrist-managed barefoot hooves.

In concordance to other recent studies on hoof conformation, it was found that hooves were wider laterally than medially in the forelimbs [17] and that hooves were more asymmetrical without shoes in the hindlimbs [18]. In agreement with previous findings there was a relationship between management regime and palmar angle, dorsal to palmar angle ratio, lateral heel height, sole length, sole length to width ratio, medial hoof width and solar symmetry [17]. Traditionally, in past literature, hoof angles were described to be around 45° in the forelimb and 55° in the hindlimb [1], which would suggest the results of this study were slightly steep in the forelimb and slightly shallow in the hindlimb, however this study's results are consistent with other studies conducted in the past year where hoof angles were on average 50.3° in the forelimb [17] and 51.52°-52.16° in the hindlimb [18], suggesting a need to review hoof conformation of the modern domestic horse.

In the presented study there was no significant difference in toe angle between shod and barefoot-managed hooves, however the heel angle was significantly shallower in shod hooves compared to barefoot hooves. This may be due to the fact that these horses were more likely to be managed with shoes or it may be due to the fact that shoes increase the pressure on the horn in the heel area and over time lead to lowering of the heels. It has been shown previously that when hooves are managed without shoes, the distal half of the heels migrate palmarly/plantarly [10], thus increasing heel height. Lower heels are usually associated with a shallower solar angle of the distal phalanx and it has been shown that for each degree change in angle the strain and thus the stress on the navicular bone increases [6]. The opposite effect is seen when heel wedges are applied to increase heel angle, this results in a decrease of navicular bone stress in comparison to flat shoes. The same study showed that barefoot hooves showed a 14% reduction in the forces acting on the navicular bone, possibly due to the same mechanism as wedges [19]. Shallower heel angles may progress to under-run heels, which decrease the capability of the heels to deform under load thereby negatively impacting shock absorption and possibly also compromising the blood supply to the hoof [20]. This consequently may predispose to hoof pathologies. Hyperextension of the interphalangeal and fetlock joints also occurs with under-run heels, thereby increasing strain on the deep digital flexor tendon and the navicular bone [21].

Our study revealed that barefoot horses had more splayed and flared hooves than shod horses, suggesting that horseshoes might restrict multidirectional expansion of the hoof capsule. Shoes have been shown to decrease the deformation of the hoof wall during loading, by redistributing irregular strains without compromising hoof function, thereby protecting against flare [22]. Flare is undesirable in hooves, as it considerably weakens the hoof wall by bending of horn tubules [23].

Overall, all hooves in all groups were wider laterally. Previous studies have demonstrated that horses preferentially land laterally [17, 24] which corresponds to a more splayed, lower lateral and a steeper, higher medial hoof wall [25]. Slight asymmetry throughout all groups to the lateral side to a degree, is therefore considered normal. In the presented study there was only a significant difference in solar symmetry between hoof management regimens for front hooves, with barefoot hooves being more asymmetrical than shod hooves. The difference between the front and the hind hooves may be attributed to the uneven weight distribution of the horse with 58% of the weight being loaded on the forelimbs, and 42% loaded on the hindlimbs [26], as a greater downwards pressure on the hooves may increase lateral splay. Incorrect trimming, pain, or compensation for poor conformation may result in excessive lateral or medial landing and mediolateral imbalance [27]. Excessive asymmetry is important to address as it affects rotation, abduction and adduction of the distal interphalangeal joint [28], thereby potentially predisposing the distal limb to injuries relating to the articular surfaces and collateral ligaments [28,29], as well as causing chronic heel pain, sheared heels, metacarpophalangeal synovitis and side-bones [30]. The point at which asymmetry is considered excessive has not been quantified, however in this study a wider range of asymmetry was observed in barefoot hooves compared to shod hooves. The potential for barefoot-managed hooves to show more hoof asymmetry and hence for more uneven loading ought to be taken into account when dealing with those horses.

Heel width and frog size were both significantly associated with hoof management regimen. Barefoot hooves had wider heels and larger frogs than shod hooves. A wide heel and large frog is considered an advantageous feature of the hoof since it increases the load-bearing area, thus reducing stress. An increased area in the palmar/plantar aspect of the solar dermis could also potentially increase proprioception of the horse during locomotion, due to the high local concentration of Pacinian corpuscles [31]. It has been suggested that the frog hypertrophies when it becomes a weight-bearing structure [32]. Barefoot frogs have more stimuli for growth as they impact the ground, instead of being raised above it like in shod hooves [10]. This is advantageous for the distal limb as increased frog-ground contact increases blood flow, thereby aiding healing, energy dissipation and growth [10,12, 13]

There was a significant difference in hindlimb, but not frontlimb conformation between farrier-managed and podiatrist-managed barefoot hooves with the farrier-managed horses showing significantly shorter hooves and longer frogs in the hindlimbs. This may be due to different approaches to trimming hindfeet in particular based on the different training members of both groups undergo. No significant differences in the frontlimb were observed despite the proclaimed fundamental difference in approach by the two groups. Further studies on the detailed differences of the individual training programmes in relation to trimming are necessary to investigate this further.

Tables

TABLE 1 Conformation parameters included in this study

Parameter	Unit	Definition
Toe Angle	degrees	Direct measurement
Heel Angle	degrees	Direct measurement
Difference in angles	degrees	Toe angle minus heel angle A negative number indicates a steeper heel angle and a positive number indicates a shallower heel angle compared to the toe angle. 0° was the gold standard and greater than +5° indicated under-run heels.
Flaring	degrees	The average of the lateral angle of deviation and the medial angle of deviation.
Dorsopalmar(plantar) Balance	%	Frog length divided by sole length, multiplied by 100
Heel Width	%	Heel width divided by sole width, multiplied by 100
Frog size	%	Frog area divided by sole area, multiplied by 100
Splaying	ratio	Sole circumference plus heel width, divided by coronary band circumference A higher figure indicates that the hoof was more splayed.
Solar Symmetry Index	ratio	Lateral sole width divided by medial sole width These figures represent mediolateral symmetry of the sole over the widest point, where less than 1 indicates the sole is wider medially and greater than 1 indicates the sole is wider laterally.

TABLE 2 Reliability of conformation measurements: 95% limits of agreement for repeat measurements

Parameter	Lower limit of agreement	Upper limit of agreement
Toe Angle (degrees)	-0.35	0.40
Heel Angle (degrees)	-0.49	0.41
Medial Wall Angle (degrees)	-0.51	0.42
Lateral Wall Angle (degrees)	-0.55	0.36
Heel Width (arb. unit)	-0.52	0.66
Sole Width (arb. unit)	-0.91	0.81
Lateral Sole Width (arb. unit)	-0.66	0.65
Medial Sole Width (arb. unit)	-0.68	0.63
Frog Length (arb. unit)	-0.58	0.60
Frog to Toe Length (arb. unit)	-0.83	0.69
Frog Width (arb. unit)	-0.52	0.51

TABLE 3 Conformation parameters for shod hooves and barefoot hooves for front and hind hooves.

Mean and standard deviation (SD) are displayed for normally distributed parameters, median and interquartile range (IQR) are displayed for not-normally distributed parameters; P-value is for comparison between shod and barefoot-managed hooves.

Parameter	Condition	Hoof	Minimum	Maximum	Median	IQR	Mean	SD	P value
Toe Angle (degrees)	Shod	Front	43.22	58.06	50.22	4.39			0.368
	Barefoot	Front	40.47	55.87	50.03	3.04			
	Shod	Hind	42.73	62.59	50.93	4.37			0.425
	Barefoot	Hind	43.19	62.31	51.88	4.78			
Heel Angle (degrees)	Shod	Front	22.87	57.56			41.65	7.34	<0.001
	Barefoot	Front	35.10	56.43			45.83	5.60	
	Shod	Hind	14.68	51.21			37.21	9.62	<0.001
	Barefoot	Hind	33.02	54.46			43.95	5.98	
Difference in toe and heel angles	Shod	Front	-2.04	31.11	8.55	9.56			<0.001
	Barefoot	Front	-12.30	10.37	2.55	3.59			
	Shod	Hind	2.56	33.05	12.08	14.46			<0.001

(degrees)	Barefoot	Hind	-2.08	16.50	5.02	6.63			
Flaring	Shod	Front	0.46	12.42	4.35	3.91			<0.001
(degrees)	Barefoot	Front	1.92	32.15	11.33	7.28			
	Shod	Hind	0.07	11.32	4.86	2.91			<0.001
	Barefoot	Hind	2.42	17.67	10.44	5.49			
Dorsopalmar	Shod	Front	57.97	72.65			65.71	3.46	0.038
(plantar)	Barefoot	Front	64.31	77.62			69.75	3.77	
Balance (%)	Shod	Hind	58.23	68.69			65.14	2.80	0.006
	Barefoot	Hind	62.07	74.88			68.76	3.35	
Heel Width	Shod	Front	38.03	62.61	49.48	7.66			<0.001
(%)	Barefoot	Front	53.41	71.46	63.08	5.79			
	Shod	Hind	41.07	66.67	52.34	9.11			<0.001
	Barefoot	Hind	55.19	88.80	70.65	10.36			
Frog size (%)	Shod	Front	12.18	23.62	16.57	3.02			<0.001
	Barefoot	Front	18.50	29.29	21.92	4.06			
	Shod	Hind	13.46	27.06	19.05	3.51			<0.001
	Barefoot	Hind	20.54	39.02	26.53	5.80			
Solar	Shod	Front	0.88	1.14	1.03	0.06			0.038
Symmetry	Barefoot	Front	0.81	1.34	1.06	0.09			
Index (ratio)	Shod	Hind	0.93	1.30	1.08	0.08			0.104
	Barefoot	Hind	0.89	1.29	1.04	0.11			
Splaying	Shod	Front	1.11	1.51	1.32	0.07			0.001
(ratio)	Barefoot	Front	1.27	1.51	1.39	0.11			
	Shod	Hind	1.22	1.52	1.34	0.08			0.003
	Barefoot	Hind	1.19	1.51	1.39	0.08			

TABLE 4 Conformation parameters for barefoot hooves managed by a farrier and barefoot hooves managed by a podiatrist for front and hind hooves. Mean and standard deviation (SD) are displayed for normally distributed parameters, median and interquartile range (IQR) are displayed for not-normally distributed parameters; P-value is for comparison between farrier and podiatrist-managed hooves.

Parameter	Farrier/Podiatrist	Hoof	Minimum	Maximum	Median	IQR	Mean	SD	P value
Toe Angle (degrees)	Farrier	Front	40.47	55.87	49.76	4.49			0.618
	Podiatrist	Front	41.70	53.89	50.16	2.70			
	Farrier	Hind	48.42	62.31	52.09	3.90			0.403
	Podiatrist	Hind	43.19	58.89	50.68	5.13			
Heel Angle (degrees)	Farrier	Front	39.64	56.43			47.31	4.15	0.351
	Podiatrist	Front	35.10	59.50			45.75	4.79	
	Farrier	Hind	34.99	59.71			46.69	6.04	0.326
	Podiatrist	Hind	33.02	54.46			44.70	6.41	
Difference in toe and heel angles (degrees)	Farrier	Front	-12.30	7.25	2.33	3.96			0.318
	Podiatrist	Front	-2.57	10.37	3.55	5.13			
	Farrier	Hind	-0.35	16.50	4.64	4.78			0.322
	Podiatrist	Hind	-2.08	11.69	7.97	5.49			
Flaring (degrees)	Farrier	Front	1.92	21.63	13.69	10.12			1.000
	Podiatrist	Front	7.44	32.15	11.06	4.80			
	Farrier	Hind	2.42	17.54	9.49	8.41			0.126
	Podiatrist	Hind	6.31	17.67	11.08	3.94			
Dorsopalmar(plantar) balance (%)	Farrier	Front	53.87	77.62			70.20	5.61	0.080
	Podiatrist	Front	59.50	73.04			66.80	4.41	
	Farrier	Hind	64.35	74.88			69.38	3.29	0.002
	Podiatrist	Hind	58.36	69.97			65.51	3.79	
Heel Width (%)	Farrier	Front	57.83	67.69	63.24	4.63			0.771
	Podiatrist	Front	53.41	71.46	62.77	6.60			
	Farrier	Hind	61.78	83.86	71.21	10.57			0.650
	Podiatrist	Hind	55.19	88.80	69.70	6.53			
Frog size (%)	Farrier	Front	19.18	29.29	23.34	2.31			0.135

	Podiatrist	Front	18.50	26.85	21.66	2.10			
	Farrier	Hind	20.54	33.48	26.20	5.81			0.777
	Podiatrist	Hind	21.14	39.02	26.53	5.94			
Solar	Farrier	Front	0.81	1.34	1.06	0.11			0.589
Symmetry	Podiatrist	Front	0.95	1.14	1.06	0.08			
Index (ratio)	Farrier	Hind	0.94	1.29	1.03	0.11			0.479
	Podiatrist	Hind	0.89	1.25	1.05	0.10			
Splaying	Farrier	Front	1.27	1.51	1.41	0.15			0.759
(ratio)	Podiatrist	Front	1.29	1.49	1.38	0.08			
	Farrier	Hind	1.19	1.51	1.37	0.10			0.590
	Podiatrist	Hind	1.32	1.51	1.39	0.07			

References

1. Karle A.S., Tank P.H., Vedpathak H.S., Mahida H.K., Shah R.G. & Dhami M.A. (2010) Horseshoeing: An Overview. *Vet World* **3**, 148-151
2. Stiller P. (2002) Pros and cons of barefoot horses. *J Equine Vet Sci* **22**, 91.
3. Reilly P. T. (2011) The barefoot paradox. *J Equine Vet Sci* **31**, 591-592
4. Teskey T.G. (2005) The Unfettered Foot: A paradigm of change for equine podiatry. *J Equine Vet Sci* **25**, 77-82
5. La Pierre K.C. (2011) Barefoot Hoof Care – Energizing Holistic Hoof Care, Institute of Applied Equine Podiatry. http://appliedequinepodiatry.org/Barefoot_hoof_care.html. Accessed May 27 2013
6. Eliashar E., Mcguigan M. P. & Wilson A. M. (2004) Relationship of foot conformation and force applied to the navicular bone of sound horses at the trot. *Equine Vet J* **36**, 431–435

7. Holroyd K., Dixon J.J., Mair T., Bolas N., Bolt D.M., David F. & Weller R. (2013) Variation in foot conformation in lame horses with different foot lesions. *The Vet J* **195**, 361-365
8. Van Heel M.C.V., Moleman M., Barneveld A., Van Weeren P. R. & Back W. (2005) Changes in location of centre of pressure and hoof-unrollment pattern in relation to an 8-week shoeing interval in the horse. *Equine Vet J* **37**, 536–540
9. Moleman M., van Heel M.C., van Weeren P.R. & Back W. (2006) Hoof growth between two shoeing sessions leads to a substantial increase of the moment about the distal, but not the proximal, interphalangeal joint. *Equine Vet J* **38**, 170-174
10. Clayton H.M., Gray S., Kaiser L.J. & Bowker R.M. (2011) Effects of barefoot trimming on hoof morphology. *Aus Vet J* **89**, 305-311
11. Hinterhofer C.H., Stanek C.H. & Haider, H. (2000) The effect of flat horseshoes, raised heels and lowered heels on the biomechanics of the equine hoof assessed by finite element analysis (FEA). *J Vet Med. A, Physiology, Pathology, Clinical Medicine* **47**, 37-82
12. Bowker R.M., Van Wulfen K.K., Springer S.E., Linder K.E. (1998) Functional anatomy of the cartilage of the distal phalanx and digital cushion in the equine foot and a hemodynamic flow hypothesis of energy dissipation. *Am J Vet Res* **59**, 961-968
13. Darcy P. (2003) The Sports Injury Process. In: *Fundamentals of Sports Injury Management*, 2nd edn., Ed: M.K. Anderson, Lippincott Williams and Wilkins pp 35-40
14. British Standards Institution. (1975) Precision of test methods 1: Guide for the determination and reproducibility for a standard test method. *BS 597, Part 1*, London: BSI
15. Bland J.M. & Altman D.G. (1986) Statistical methods for assessing agreement between two methods of clinical measurement. *The Lancet* **327**, 307-310
16. Turner T. (1992) The use of hoof measurements for the objective assessment of hoof balance. *Proc Am*

Assoc Equine Prac **38**, 389-395

17. Wilson A., Agass R.F., Sherlock E.C., Vaux S., Day P., Weller R. & Pfau T. (2014) Foot placement of the equine forelimb: relationship between foot conformation, foot placement and motion asymmetry. *Equine Vet J* (online)

18. Agass R.F., Wilson A.M., Sherlock E.C., Vaux S., Day P., Weller R. & Pfau T. (2014) The relationship between foot conformation, foot placement and movement symmetry in the equine hind limb at walk and trot. *Equine Vet J* **46**, 19-20

19. Willemen M.A., Savelberg H.H.C.M. & Barneveld A. (1999) The effect of orthopaedic shoeing on the force exerted by the deep digital flexor tendon on the navicular bone in horses. *Equine Vet J* **31**, 25–30

20. Day P., Butts D., Pfau T., Pardoe C. & Weller R. (2013) Does hoof deformation differ between horses with collapsed heels and horses with non-collapsed heels? *J Equine Vet Sci* **33**, 859

21. Balch O.K., Helman R.G. & Collier M.A. (2001) Underrun heels and toe-grab length as possible risk factors for catastrophic musculoskeletal injuries in Oklahoma race-horses. *Proc Am Assoc Equine Prac* **47**, 334–338

22. Thomason J.J. (1998) Variation in surface strain on the equine hoof wall at the midstep with shoeing, gait, substrate, direction of travel, and hoof shape. *Equine Vet J Supp* **26**, 86-95

23. Balch O.K., Butler D. & Collier M.A. (1997) Balancing the normal foot: hoof preparation, shoe fit and shoe modification in the performance horse. *Equine Vet Educ* **9**, 143-154

24. Van Heel M.C.V., Barneveld A., van Weeren P.R. & Back W. (2004) Dynamic pressure measurements for the detailed study of hoof balance: the effect of trimming. *Equine Vet J* **36**, 778-782

25. Milner P. & Hughes I. (2012) Remedial Farriery Part 5: Principles of foot balance. *Comp Ani* **17**, 10-15

26. Hood D.M., Wagner I.P., Taylor D.D., Brumbaugh G.W. & Chaffin M.K. (2001) Voluntary limb load distribution in horses with acute and chronic laminitis. *Am J Vet Res* **62**, 1393-1398

27. Kaneps A.J. & Turner T.A. (2004) Diseases of the Foot. In: *Equine Sports Medicine and Surgery*. Ed: K.W. Hinchcliff, A.J. Kaneps, R.J. Geor. Saunders Elsevier, London. pp 260-288
28. Chateau H., Degruet C., Jerbi H., Crevier-Denoix N., Pourcelot P., Audigie F., Pasqui-Boutard V. & Denoix J. (2002) Three-dimensional kinematics of the equine interphalangeal joints: articular impact of asymmetric bearing. *Vet Res* **33**, 371-382
29. Clayton H.M. (2011) Hoof Mechanics during Locomotion. In *Care and Rehabilitation of the Equine Foot*. Ed P. Ramey, Hoof Rehabilitation Publishing LLC, p 276
30. Balch O.K., White K., Butler D. & Metcalf S. (1995) Hoof balance and lameness: improper toe length, hoof angle and mediolateral balance. *Comp Cont Ed Prac Vet* **17**, 1275-1283
31. Bowker R.M., Brewer A.M., Vex K.B., Guida L.A., Linder K.E., Sonea I.M. & Stinson A.W. (1993) Sensory receptors in the equine foot. *Am J Vet Res* **54**, 1840–1844
32. J.J., Douglas J.E. & Sears, W. (2001) Morphology of the laminar junction in relation to the shape of the hoof capsule and distal phalanx in adult horses (*Equus caballus*). *Cells Tissues Organs* **168**, 295–311

Heel angle

Toe angle

Lateral angle of
deviation (flare)

Medial angle of
deviation (flare)

Frog width

Heel width

Frog length

Medial sole

Lateral sole

Sole width

Frog to toe distance