Cleavage of Braun lipoprotein Lpp from the bacterial peptidoglycan by a paralog of L,D-transpeptidases, LdtF

Raj Bahadur, Pavan Kumar Chodiseti, and Manjula Reddy*

CSIR-Centre for Cellular and Molecular Biology

Hyderabad India 500007

Running title: LdtF is a murein endopeptidase

Key words: Bacterial cell wall, Peptidoglycan, Lpp, L,D-transpeptidase, LdtF

*Address correspondence to

Manjula Reddy

S-106, CSIR-Centre for Cellular and Molecular Biology

Hyderabad India 500007

Ph: +91-40-27192514

Fax: +91-40-27160310

Email: manjula@ccmb.res.in

Abstract

Gram-negative bacterial cell envelope is made up of an outer membrane (OM), an inner membrane (IM) that surrounds the cytoplasm, and a periplasmic space between the two membranes containing peptidoglycan (PG or murein). PG is an elastic polymer that forms a mesh-like sacculus around the IM protecting cells from turgor and environmental stress conditions. In several bacteria including *E. coli*, the OM is tethered to PG by an abundant OM lipoprotein, Lpp (or Braun lipoprotein) that functions to maintain the structural and functional integrity of the cell envelope. Since its discovery Lpp has been studied extensively and although L,D-transpeptidases, the enzymes that catalyse the formation of PG–Lpp linkages have been earlier identified, it is not known how these linkages are modulated. Here, using genetic and biochemical approaches, we show that LdtF (formerly *yafK*), a newly-identified paralog of L,D-transpeptidases in *E. coli* is a murein hydrolytic enzyme that catalyses cleavage of Lpp from the PG sacculus. LdtF also exhibits glycine-specific carboxypeptidase activity on mucopeptides containing a terminal glycine residue. LdtF is earlier presumed to be an L,D-transpeptidase; however, our results show that it is indeed an L,D-endopeptidase that hydrolyses the products generated by the L,D-transpeptidases. To summarize, this study describes the discovery of a murein endopeptidase with a hitherto unknown catalytic specificity that removes the PG–Lpp cross-links suggesting a role for LdtF in regulation of PG-OM linkages to maintain the structural integrity of the bacterial cell envelope.

Significance statement

Bacterial cell walls contain a unique protective exoskeleton, peptidoglycan, which is a target of several clinically important antimicrobials. In Gram-negative bacteria, peptidoglycan is covered by an additional lipid layer, outer membrane that serves as permeability barrier against entry of toxic molecules. In some bacteria, an extremely abundant lipoprotein, Lpp staples outer membrane to peptidoglycan to maintain the structural integrity of the cell envelope. In this study, we identify a previously unknown peptidoglycan hydrolytic enzyme that cleaves Lpp from the peptidoglycan sacculus and show how the outer membrane-peptidoglycan linkages are modulated in *Escherichia coli*. Overall, this study helps in understanding the fundamental bacterial cell wall biology and in identification of alternate drug targets for development of new antimicrobials.

Introduction

Gram-negative bacterial cell envelope is made up of outer membrane (or OM), an asymmetric bilayered lipid membrane which is surface-exposed and an inner membrane (or IM) consisting of a phospholipid bilayer surrounding the cytoplasm. In between these two membranes is the periplasmic space in which a sac-like molecule, the peptidoglycan (PG or murein) is located (1). PG is an elastic heteropolymer that protects bacterial cells from lysis by internal osmotic pressure and from external stress conditions. It is a single, large macromolecule made up of multiple linear glycan strands that are inter-connected with each other by short peptide chains forming a net-like sacculus around the cytoplasmic membrane (Fig.1). The glycan strands are polymers of alternating β -1,4-linked N-acetylglucosamine (GlcNAc) and N-acetylmuramic acid (MurNAc) disaccharide units in which the D-lactoyl moiety of each MurNAc residue is covalently attached to the first amino acid of the stem peptide. Normally, the peptide chains are of two to five amino acids with a pentapeptide made up of L-alanine (L-ala¹)–D-glutamic acid (D-glu²)–mesodiaminopimelic acid (mDAP³)–D-ala⁴–D-ala⁵ residues. In *E. coli*, approximately 40% of the neighbouring peptide chains are linked to each other, either between the D-ala⁴ and mDAP³ (D-ala–mDAP or 4–3) or two mDAP³ residues (mDAP–mDAP or 3–3) residues. Of these, the 4–3 cross-links are more prevalent and are formed by D,D-transpeptidases whereas 3–3 cross-links are much less abundant and are catalysed by L,D-transpeptidases (LDTs), LdtD and LdtE (2–4).

In several bacteria including *E. coli*, the OM is stapled to PG by a lipoprotein, Lpp or Braun lipoprotein. Lpp is the first OM lipoprotein to be discovered almost five decades ago and has been studied extensively (5–9). It is a small abundant protein ($\sim 10^6$ molecules per cell) of 58 amino acids and is known to exist in two conformations each occupying a distinct subcellular location in the cell envelope (9–11). One third of total Lpp is in the periplasm covalently

attached to the mDAP³ residues of the PG peptides (bound form) whereas the other two third spans the OM (free form) (Fig.1). OM-PG tethering by Lpp has been shown to determine the width of the periplasm by controlling the IM-OM distance and to contribute to the stiffness of cell envelope (12,13). Although Lpp is not essential for viability of *E. coli*, mutants that lack Lpp show several pleiotropic defects such as increased sensitivity to hydrophobic agents, leakage of periplasmic contents, OM blebbing, vesiculation, cell separation defects, as well as deficiency in virulence, highlighting the role of Lpp in maintenance of envelope integrity (8,9,14).

Three redundant LDTs, LdtA, -B, -C catalyse the formation of PG–Lpp cross-links by covalently attaching the extreme C-terminal residue of Lpp, lysine to the mDAP³ residue of a tetrapeptide in the mature PG sacculus (15). In this reaction, the terminal D-ala residue of the tetrapeptide is lost leading to the formation of a tripeptide-Lpp cross-link (Fig. 1). About, 10% of the peptides in a PG sacculus are attached to Lpp and this frequency is presumed to vary during conditions of stress (2,5,16–18).

E. coli encodes six LDTs, LdtA-F belonging to YkuD family of proteins (3,18,19,20). Of these, LdtA,-B,-C catalyze the covalent attachment of Lpp to the PG; though these are redundant, LdtB is physiologically relevant because deleting *ldtB* alone abrogates the attachment of Lpp to a significant extent (15). Ribosome profiling (10) has also shown the abundance of LdtB to be much higher (~5,000 copies per cell) than LdtA and LdtC (~50 and 500 respectively). On the other hand, the 3–3 cross-link formation in the PG sacculus is catalysed by LdtD and LdtE (21).

Apart from their ability to form cross-links, LdtA-E catalyse an amino acid exchange reaction in the periplasm wherein the terminal D-ala⁴ residue of the stem peptides is substituted with either glycine or a variety of noncanonical D-amino acids (NCDAA) such as D-tryptophan,

D-methionine or D-aspartate (2,21,22). The significance of this exchange in *E. coli* is not clear, although, it is believed that these substituted mucopeptides do not participate in further steps of PG polymerization. LDTs are presumed to have a larger role in the maintenance of structural integrity of PG because of their ability to form cross-links de novo in a mature PG sacculus independent of active PG precursor synthesis (18).

LdtF is a recently-identified paralog of LDTs and is implicated in facilitating the formation of 3–3 cross-links; however, its precise function remains unclear (18–20). Here, we show that LdtF (encoded by *yafK*) is a murein L,D-endopeptidase that cleaves Lpp from the PG sacculus. We initially identified *ldtF* because of its genetic interaction with *mepS*, a gene encoding a major PG elongation-specific D,D-endopeptidase (23). Further genetic and biochemical analysis demonstrated the role of LdtF in hydrolysing the products generated by the activity of other LDTs. LdtF cleaves Lpp which is bound to the PG sacculus and in addition, cleaves the terminal glycine residue that is incorporated into the stem peptides due to the periplasmic exchange reaction of LDTs. However, LdtF was not able to cleave the terminal NCDAAs residues from the mucopeptides. To summarize, this study identifies a murein endopeptidase with a previously unknown catalytic specificity having an ability to modulate the Lpp-mediated OM-PG linkages.

Results

Lack of PG-Lpp linkages confer growth advantage to an *E. coli* mutant lacking an elongation-specific D,D-endopeptidase, *mepS*

We showed earlier that absence of 3–3 cross-link forming LDTs (*ldtD* and *ldtE*) confer growth advantage to a mutant lacking an elongation-specific 4–3 cross-link cleaving D,D-endopeptidase, MepS signifying the importance of cleavage of both 4–3 and 3–3 cross-links to make space for incorporation of new PG material during cell elongation (23,24). To

examine whether the tethering of OM to PG by Lpp also affects the process of PG expansion, we introduced a deletion of Lpp into a mutant lacking MepS and examined the growth phenotypes of the double mutant on Nutrient agar (NA) because *mepS* deletion mutant is unable to grow on NA (25). Fig. 2A shows that absence of Lpp restores moderate amount of growth to the *mepS* deletion mutant on NA. In addition, a mutant Lpp allele that lacks the C-terminal lysine residue and hence unable to bind PG (Lpp^{ΔK58}; 11) confers suppression like that of Lpp deletion. Likewise, deletion of LdtB, which catalyses the formation of mDAP–Lpp linkages conferred growth to *mepS* deletion mutant (Fig. 2A; Fig. S1, Fig. S2). However, deletion of LdtA or LdtC which also link Lpp to mDAP did not have any effect on *mepS* growth (Fig. S1). Surprisingly, deletion of LdtF, the newly-identified paralog of LDTs, conferred a very small yet consistent growth defect to *mepS* single mutant (Fig. S1) which was further exacerbated in a *mepS* mutant lacking *mepK*, a gene encoding the 3–3 cross-link cleaving PG hydrolase (Fig. 2B).

As these results intrigued us, we further investigated the role of LdtF by introducing the plasmids encoding each of the LDTs (26) into the *mepS* mutant. As shown in Fig. S3, plasmids encoding *ldtA*, *-B*, *-C*, *-D*, *-E* did not confer growth to *mepS* mutant whereas a plasmid encoding *ldtF* alone was able to moderately suppress the growth defects of *mepS* mutant. Another plasmid derivative carrying cloned *ldtF* downstream to an IPTG-dependent promoter (P_{trc}::*ldtF*) also suppressed the growth defects of *mepS* mutant on NA (Fig. 2C) indicating that LdtF may have a distinct function compared to that of other LDTs. LdtF belongs to YkuD family of proteins and members of this family contain L,D-transpeptidase domain with an invariant cysteine residue at the active site (27). To further validate the role of LdtF, we constructed a mutant derivative with an alanine residue substituted for cysteine (LdtF-C143A) and examined its ability to suppress the *mepS* mutation. Fig. 2C shows that a plasmid encoding LdtF-C143A poorly suppressed the *mepS* deletion mutant whereas another

variant coding for LdtF-H135A behaved like that of WT. However, deletion of *ldtF* alone in a WT strain did not confer any discernible phenotype when grown on LB, LBON (LB without NaCl) or NA plates at 30, 37 or 42°C except a slight reduction in the doubling time during exponential phase of growth (Fig. S4). In addition, *ldtF* deletion mutant did not exhibit significant sensitivity to any of the cell wall-antibiotics such as cephalexin, cefsulodin, mecillinam or vancomycin.

LdtF modulates PG–Lpp linkages *in vivo*

To understand the basis of LdtF’s function, we examined the composition of PG in strains either having a deletion or multiple copies of *ldtF*. PG sacculi from these strains were prepared, digested with a muramidase (mutanolysin) followed by separation of soluble muropeptides by RP-HPLC and identification of the peaks by MS or MS-MS analysis (as described in Materials and Methods). No major difference was observed in the muropeptide profile of *ldtF* deletion derivative compared to that of WT (Fig. S5). In contrast, the PG sacculi of cells carrying additional copies of *ldtF* had considerable alterations (Fig. 3A; Table S3), the most significant being absence of peak 3 (tri-lys-arg) which is a disaccharide tripeptide attached to a lys-arg dipeptide (Fig. 3B, 3C). Tri-lys-arg muropeptides are generated due to the proteolytic activity of pronase which is used during preparation of PG sacculi to remove bound Lpp. Pronase cleaves Lpp at the 56th position leaving the extreme C-terminal lys-arg dipeptide attached to the mDAP residue of the stem peptides resulting in generation of several species of muropeptides bound to lys-arg dipeptide (2). In addition to absence of peak 3, a muropeptide peak eluting at 46 min (labelled ‘Y’) was significantly elevated in the PG sacculi of cells carrying additional copies of LdtF (Fig. 3A). MS-MS analysis indicated this peak to be a tetra-tri dimer linked by 4–3 cross-bridge with a molecular mass of 1794 Da (Fig. 3B, 3C; Fig. S6). Absence of peak 3 with concomitant

increase in peak 1 (a monomer of tri) allowed us to speculate that LdtF may have an ability to modulate the mDAP–Lpp linkages. Though the source of peak Y is not clear, it was not detected in a strain deleted for Lpp suggesting it may have originated by the activity of LdtF on PG–Lpp cross-links (Fig. S7A). Additionally, the incidence of peak Y was not dependent on the presence of functional LdtD and -E (Fig. S7B). Further, all other alterations observed due to overexpression of LdtF disappeared in a strain lacking Lpp reinforcing the suggestion that LdtF functions downstream of Lpp (Fig. S7A). As shown in Fig. 3A, analysis of PG in strains carrying plasmids encoding either LdtF-C143A or LdtF-H135A indicated a direct role for LdtF in modulation of mDAP–Lpp linkages (Table S3).

Because the above results implicated LdtF in regulation of PG–Lpp linkages, we examined the extent of these cross-links in cells lacking LdtF. To perform this experiment, Lpp-bound PG sacculi were prepared from WT and *ldtF* deletion mutant as described in Materials and Methods. PG sacculi from both strains were digested with mutanolysin, soluble muropeptides were collected and normalized amounts (Fig. S8) were electrophoresed using SDS-PAGE followed by western blotting and detection with anti-Lpp antibody. Fig. 3D shows that the PG sacculi derived from LdtF deletion mutant indeed contain a greater abundance of high molecular weight Lpp-bound muropeptides compared to that of WT, although the level of low-molecular weight Lpp-bound muropeptides were unchanged. This observation suggested an interesting possibility of LdtF specifically moderating the larger oligomeric Lpp–cross-linked muropeptides of the PG sacculus and the implications of this result are further discussed below.

LdtF is a murein endopeptidase that cleaves PG–Lpp linkages

To examine the enzymatic activity of LdtF, a signal-less hexa-histidine tagged LdtF (LdtF²⁰⁻²⁴⁶-His₆) was overexpressed and purified as described in Materials and Methods. Treatment

of soluble muropeptides derived from the PG sacculi of WT *E. coli* with purified LdtF yielded muropeptide fraction that totally lacked tri-lys-arg (peak 3) and tetra-tri-lys-arg (peak 6) with concomitant increase in tri- and tetra-tri muropeptides (Fig. 4A). Cleavage of tri-lys-arg and tetra-tri-lys-arg into tri- or tetra-tri muropeptides was also confirmed using purified fractions (Fig. 4B, 4C, 4D).

We next examined the ability of LdtF to cleave the bound Lpp from the intact PG sacculi. To perform this experiment, Lpp-bound PG sacculi from WT and *ldtF* mutant were isolated and equal amounts of each were treated with purified LdtF. The soluble fraction was electrophoresed on SDS-PAGE and Lpp was detected by western analysis using anti-Lpp antibody. As a positive control, PG sacculi treated with pronase were used. Fig. 4E shows that both LdtF and pronase cleave Lpp from the PG sacculi and that the amount of Lpp released from the sacculi of LdtF deletion mutant was considerably higher (approximately 5-fold) than that of the WT (compare lanes 2 and 3 with 5 and 6). The remaining insoluble PG fraction was further analysed by RP-HPLC and as expected, the lys-arg muropeptides were not detected in LdtF-treated PG whereas pronase-treated PG contained the lys-arg muropeptides (Fig. S9). Overall, these results demonstrate the catalytic specificity of LdtF on PG–Lpp or PG–lys-arg substrates.

LdtF is a glycine-specific carboxypeptidase that cleaves terminal glycine residue from the stem peptides

The above experiments clearly demonstrated hydrolytic activity of LdtF on PG–Lpp linkages formed by LdtA, -B and -C. LDTs also exchange the terminal D-ala of stem peptides with a glycine residue. To examine whether LdtF has any activity on glycine-substituted muropeptides, soluble muropeptides of a WT strain grown with glycine-supplementation were used as substrates for LdtF. As expected, growth of WT *E. coli* with exogenously added

glycine resulted in accumulation of a large number of mucopeptides with glycine at position 4 whose identity is determined by MS or MS-MS analysis (Fig. S10). Fig. 5A shows that LdtF effectively removes glycine from a variety of glycine-containing mucopeptides. LdtF also cleaved several glycine-containing mucopeptides prepared from PG sacculi of a strain overexpressing LdtD (Fig. 5B). Of these, three distinct types of glycine-containing mucopeptides were purified to homogeneity and Fig. 5C, 5D, 5E show that LdtF removes glycine residue from all these mucopeptides. However, the abundance of glycine-containing mucopeptides remained the same in both WT and *ldtF* deletion mutant when grown with glycine-supplementation suggesting the existence of alternate carboxypeptidases that cleave the terminal glycine residue. In support of this idea, *ldtF* deletion mutant was not sensitive to addition of glycine and behaved just like that of WT strain.

Considering an earlier report that D,D-carboxypeptidases hydrolyse the terminal glycine from the stem peptides (2,28), we made a quadruple mutant deleted for major D,D-carboxypeptidases, DacA, -B, -C, -D and tested for sensitivity to glycine. As expected, the quadruple mutant formed smaller-sized colonies on glycine-supplemented media (Fig. S11). Introduction of *ldtF* deletion marginally exacerbated the defect of this quadruple mutant whereas multiple copies of *ldtF* moderately improved the growth of this mutant on glycine-containing media (Fig. S11) implicating a role for LdtF in removal of terminal glycine residue from the stem peptides. In sum, the above results demonstrate that LdtF is a glycine-specific carboxypeptidase.

LdtF does not cleave NCDAAs from the stem peptides

To examine whether LdtF also cleaves the terminal NCDAAs residues that are substituted by the exchange reaction of the LDTs, PG sacculi were made from WT *E. coli* grown in the presence of D-methionine, D-tryptophan or D-phenylalanine (22). Soluble mucopeptides of

these PG sacculi were separated and the peaks containing the NCDAAs substitutions were identified by MS analysis (Fig. S12A) and used as substrates for LdtF (Fig. S12B). However, LdtF was not able to cleave the terminal NCDAAs from any of these mucopeptides (Fig. S12B).

LdtF removes Lpp-mediated IM-PG linkages

Lpp is transported from the cytosol into the periplasm by Sec-mediated pathway and is eventually translocated to the OM by lipoprotein translocating machinery, LolABCDE (29). However, in certain transport-defective mutants, Lpp is stalled at the periplasmic face of the IM leading to the formation of IM-PG linkages by LDTs (30,31). It has been shown recently that absence of two small cytoplasmic membrane proteins, DcrB and YciB leads to mislocalization of Lpp at the IM, resulting in lethal IM-PG cross-links, and that this lethality is suppressed by deletion of either Lpp or LdtB (31). To examine the ability of LdtF to cleave the Lpp bound to IM, we made use of this mutant and observed that a deletion of *ldtF* exacerbates the growth defect of *yciB dcrB* double mutant (Fig. 6A). In addition, introduction of a multicopy *ldtF* plasmid ($P_{trc}::ldtF$) partially restored the growth of the *yciB dcrB* double mutant, suggesting that LdtF may also cleave Lpp bound to the IM (Fig. 6B).

Discussion

Here, we report identification of a previously unknown peptidoglycan hydrolase, LdtF that cleaves Lpp (or Braun lipoprotein), an abundant OM lipoprotein which links OM to the PG sacculus in *E. coli*. LdtF is also a glycine-specific carboxypeptidase that removes the terminal glycine residue from the PG mucopeptides. LdtF is a recently-identified member of YkuD family of proteins in *E. coli*; the other paralogs of this family comprise LdtA, -B, -C that catalyse the formation of mDAP–Lpp linkages and LdtD and -E which catalyse the formation

of mDAP–mDAP cross-links. This study also represents an instance wherein members of a paralogous family perform contrasting but not overlapping functions.

Role of LdtF in maintenance of envelope structure and stability

We identified LdtF in this study because its absence enhanced growth defects of a mutant lacking two of the PG elongation-specific endopeptidases, MepS and MepK and additionally, multiple copies of *ldtF* rescued the defects of *mepS* mutant (Fig. 2B, 2C). Moreover, we observed that absence of LdtF increases the PG–Lpp linkages (Fig. 3D) whereas more copies of *ldtF* decrease the level of PG-bound Lpp (Fig. 3A), suggesting a role for LdtF in modulating the degree of PG–Lpp cross-linkages. Subsequent biochemical analysis confirmed LdtF to be a hydrolase having two distinctive enzymatic functions- an L,D-endopeptidase activity that cleaves mDAP–Lpp cross-links and a carboxypeptidase activity that cleaves mDAP–gly linkages (Fig. 4; Fig. 5).

LdtF was earlier identified because a transposon insertion in *ldtF* (*yafK*) caused defective biofilm formation in an enteroaggregative *E. coli* strain (32). LdtF deletion has also been shown to confer additive sickness to a mutant defective in the transport of lipopolysaccharide (18). Although the basis of the above phenotypes is not clear, elevated OM-PG linkages may result in a defective cell envelope leading to such phenotypes. Excess OM-PG linkages may also alter the plasticity of the cell wall resulting in decreased fitness and survival of *E. coli*. Nevertheless, under laboratory conditions, absence of *ldtF* did not result in a large effect on the growth of *E. coli* excepting a small decrease in the doubling time (Fig. S4).

It is interesting to note that although the abundance of lower molecular weight Lpp-bound muropeptides was comparable in both WT and *ldtF* mutant (Fig. 3D, Fig. S5), the amount of bound Lpp is considerably higher in absence of LdtF (Fig. 3D, 4E). Occurrence of larger Lpp-bound oligomeric muropeptides in *ldtF* mutant (Fig. 3D) strongly suggests that LdtF

may preferentially cleave PG–Lpp cross-links of higher order structures in the PG sacculus. Lpp-mediated OM-PG cross-linkages resulting in formation of large oligomers may distort the symmetry, or the organization of cell envelope and LdtF may perhaps work towards eliminating such linkages. In addition, the effect of *ldtF* alleles on the growth of mutants in which Lpp is stalled at the IM (Fig. 6) suggests a role for LdtF in removal of rare IM-PG linkages that may occur during transport of Lpp across the periplasm. LdtF may also facilitate PG turnover as the Lpp-linked muropeptides may not efficiently get recycled unless Lpp is cleaved from the PG sacculi.

Earlier studies implicated LdtF in facilitating the formation of 3–3 cross-links because ectopic expression of LdtF along with coexpression of LdtD or -E (in a strain lacking *ldtABCDEF*) increased the level of 3–3 cross-links (18–20). However, we did not observe increased 3–3 dimers when LdtF is overexpressed in WT; in contrast, we detected increased 4–3 linked dimer (peak Y) whose occurrence was dependent on Lpp but not on LdtD or -E (Fig. S7). The basis of this discrepancy is not clear – it may perhaps be due to the strain background used for these studies and needs to be further investigated.

Regulation of PG–Lpp linkages

Lpp is an abundant OM lipoprotein (5,6,10) with one third of it covalently attached to PG, making almost 10% of the peptides linked to Lpp. However, it is not clear how *E. coli* maintains optimal levels of PG–Lpp linkages. The combined activities of LdtABC and LdtF may control the abundance of PG–Lpp linkages or alternatively structural/ conformational constraints of PG sacculi may limit the extent of PG–Lpp cross-link formation. These linkages are reported to be higher during conditions of stress including stationary-phase of growth and certain mutant conditions (2,16–18). LdtF promoter expression is shown to be higher in stationary-phase (18); however, preliminary experiments done in our laboratory to

examine the endogenous LdtF expression (using a LdtF-FLAG fusion) show that the protein levels fall during stationary-phase suggesting a likely basis for the occurrence of higher amount of PG–Lpp linkages in stationary phase. It would be worthwhile to further examine how cells achieve a dynamic yet balanced level of PG-Lpp linkages to maintain the structural and functional integrity of the cell envelope.

Role of PG–Lpp linkages in PG enlargement

Absence of PG–Lpp linkages either by deleting Lpp, LdtB or increasing the copy number of LdtF, partially rescued the growth defects of a *mepS* deletion mutant (Fig. 2A, 2C); however, the suppression was not very robust (Fig. S1, S2). In addition, other phenotypes of *mepS* such as sensitivity to β -lactam antibiotic, mecillinam or its synthetic lethality with deletion of *mepM* (23) were not suppressed by deletion of Lpp or LdtB suggesting that the absence of Lpp linkages may not significantly affect the process of PG enlargement. Lack of OM-PG tethering may perhaps alter the mechanical properties of the cell envelope and increase the flexibility of the PG sacculus, consequently resulting in a marginal growth advantage to *mepS* mutants in low osmolar conditions such as NA.

Materials and Methods

Media, bacterial strains and plasmids. LB has 0.5% yeast extract, 1% tryptone and 1% NaCl (33). LBON is LB without NaCl. NA (Nutrient Agar) has 0.5% peptone and 0.3% beef extract. Antibiotics were used at the following concentrations ($\mu\text{g/mL}$): ampicillin (Amp)-50, chloramphenicol (Cm)-30, and kanamycin (Kan)-50. Bacterial strains and plasmids used in this study are listed in Tables S1-S2 (SI).

Molecular and genetic techniques. Recombinant DNA and plasmid constructions were performed as per standard methods. MG1655 genomic DNA was used as template and

Phusion HF DNA polymerase was used for PCR amplifications and the plasmid clones were confirmed by sequence analysis. P1-phage mediated transductions and transformations were performed using standard methods (33). All strains are derivatives of MG1655 (Coli Genetic Stock Centre). Deletion mutations are from Keio mutant collection (34).

Determination of PG–Lpp linkages in the PG sacculi

Lpp-bound PG sacculi were isolated from cultures of WT and $\Delta ldtF$ mutant, treated with mutanolysin and the soluble fraction was run on 15% SDS-PAGE followed by western blotting using anti-Lpp antibody.

Determination of enzymatic activity

To examine the activity of LdtF, soluble muropeptides were incubated with either buffer or LdtF (4 μ M) at 30°C for 16 h. Samples were heat inactivated, reduced with sodium borohydride and separated by RP-HPLC. Lpp cleavage was examined by incubating purified LdtF (4 μ M) with the Lpp-bound PG sacculi for 16 h in 25 mM Tris-HCl (pH 8.0) at 30°C followed by electrophoresis of the soluble fraction on SDS-PAGE and western blotting with anti-Lpp antibody.

Supplemental Information

Details of strains, plasmids, additional materials and methods are described in Supplementary Information (SI). Tables S1-S3 and Figures S1-S12 are included in SI.

Acknowledgements

We thank NBRP: *E. coli* for Keio collection and ASKA plasmid library; Thomas Silhavy for Lpp-K58 allele and anti-Lpp antibody; Sujata Kumari for initiating the work; V Krishna Kumari and C Subbalakshmi for HPLC; B Raman and Y Kameshwari for mass spectrometry

analysis; N Madhusudhana Rao for suggestions, and members of Reddy laboratory for critical comments. This work is supported by funds from Council of Scientific and Industrial Research (MLP0141) and Department of Biotechnology (Centre of Excellence in Microbial Biology), Govt of India (to MR). We acknowledge financial support from Department of Biotechnology (to RB) and University Grants Commission of India (to PKC).

Author Contributions

RB and MR designed the study; RB and PKC performed the experiments; RB, PKC and MR analysed the data and wrote the manuscript.

References

1. T. J. Silhavy, D. Kahne, S. Walker, The bacterial cell envelope. *Cold Spring Harb Perspect Biol* **2**:a000414 (2010).
2. B. Glauner, J. V. Höltje, U. Schwarz, The composition of the murein of *Escherichia coli*. *J. Biol. Chem.* **263**, 10088-10095 (1988).
3. A. J. F. Egan, J. Errington, W. Vollmer, Regulation of peptidoglycan synthesis and remodelling. *Nat. Rev. Microbiol.* **18**, 446-460 (2020).
4. S. Garde, P. K. Chodisetti, M. Reddy, Peptidoglycan: Structure, Synthesis and Regulation. *EcoSal Plus*. ESP-0010-2020 (2021).
5. V. Braun, K. Rehn, Chemical characterization, spatial distribution and function of a lipoprotein (murein-lipoprotein) of the *E. coli* cell wall. The specific effect of trypsin on the membrane structure. *Eur. J. Biochem.* **10**, 426-438 (1969).
6. V. Braun, V. Bosch, Sequence of the murein-lipoprotein and the attachment site of the lipid. *Eur. J. Biochem.* **28**, 51-69 (1972).
7. M. Inouye, L. Shaw, C. Shen, The assembly of a structural lipoprotein in the envelope of *Escherichia coli*. *J. Biol. Chem.* **247**, 8154-8159 (1972).
8. V. Braun, K. Hantke, Lipoproteins: Structure, Function, Biosynthesis. *Subcell. Biochem.* **92**, 39-77 (2019).
9. A. T. Asmar, J. F. Collet, Lpp, the Braun lipoprotein, turns 50-major achievements and remaining issues. *FEMS Microbiol. Lett.* **365** (2018).
10. G. W. Li, D. Burkhardt, C. Gross, J. S. Weissman, Quantifying absolute protein synthesis rates reveals principles underlying allocation of cellular resources. *Cell.* **157**, 624-635 (2014).

11. C. E. Cowles, Y. Li, M. F. Semmelhack, I. M. Cristea, T. J. Silhavy, The free and bound forms of Lpp occupy distinct subcellular locations in *Escherichia coli*. *Mol. Microbiol.* **79**, 1168-1181 (2011).
12. A. T. Asmar, *et al.*, Communication across the bacterial cell envelope depends on the size of the periplasm. *PLoS Biol.* **15**, e2004303 (2017).
13. M. Mathelié-Guinlet, A. T. Asmar, J. F. Collet, Y. F. Dufrêne, Lipoprotein Lpp regulates the mechanical properties of the *E. coli* cell envelope. *Nat. Commun.* **11**, 1789 (2020).
14. A. N. Sanders, M. S. Pavelka, Phenotypic analysis of *Escherichia coli* mutants lacking L,D-transpeptidases. *Microbiology.* **159**, 1842-1852 (2013).
15. S. Magnet *et al.*, Identification of the L,D-transpeptidases responsible for attachment of the Braun lipoprotein to *Escherichia coli* peptidoglycan. *J. Bacteriol.* **189**, 3927-3931 (2007).
16. G. A. Pisabarro, M. A. de Pedro, D. Vázquez, Structural modifications in the peptidoglycan of *Escherichia coli* associated with changes in the state of growth of the culture. *J. Bacteriol.* **161**, 238-242 (1985).
17. C. Schwechheimer, D. L. Rodrigue, M. J. Kuehn, NlpI-mediated modulation of outer membrane vesicle production through peptidoglycan dynamics in *Escherichia coli*. *Microbiology* **4**, 375-389 (2015).
18. N. Morè, *et al.*, Peptidoglycan remodeling enables *Escherichia coli* to survive severe outer membrane assembly defect. *mBio.* **10**, e02729 (2019).
19. A. Montón Silva, *et al.*, The fluorescent D-amino acid NADA as a tool to study the conditional activity of transpeptidases in *Escherichia coli*. *Front. Microbiol.* **9**, 2101 (2018).

20. K. Peters, *et al.*, Copper inhibits peptidoglycan LD-transpeptidases suppressing β -lactam resistance due to bypass of penicillin-binding proteins. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 10786-10791 (2018).
21. S. Magnet, L. Dubost, A. Marie, M. Arthur, L. Gutmann, Identification of the L,D-transpeptidases for peptidoglycan cross-linking in *Escherichia coli*. *J. Bacteriol.* **190**, 4782-4785 (2008).
22. M. Caparrós, A. G. Pisabarro, M. A. de Pedro, Effect of D-amino acids on structure and synthesis of peptidoglycan in *Escherichia coli*. *J. Bacteriol.* **174**, 5549-5559 (1992).
23. S. K. Singh, L. SaiSree, R. N. Amrutha, M. Reddy, Three redundant murein endopeptidases catalyse an essential cleavage step in peptidoglycan synthesis of *Escherichia coli* K12. *Mol. Microbiol.* **86**, 1036-1051 (2012).
24. P. K. Chodiseti, M. Reddy, Peptidoglycan hydrolase of an unusual cross-link cleavage specificity contributes to bacterial cell wall synthesis. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 7825-7830 (2019).
25. H. Hara, N. Abe, M. Nakakouji, Y. Nishimura, K. Horiuchi, Overproduction of penicillin-binding protein 7 suppresses thermosensitive growth defect at low osmolarity due to an *spr* mutation of *Escherichia coli*. *Microb. Drug. Resist.* **2**, 63-72 (1996).
26. M. Kitagawa, *et al.*, Complete set of ORF clones of *Escherichia coli* ASKA library (a complete set of *E. coli* K-12 ORF archive): unique resources for biological research. *DNA Res.* **12**, 291-299 (2005).
27. S. Biarrotte-Sorin *et al.*, Crystal structure of a novel beta-lactam-insensitive peptidoglycan transpeptidase. *J. Mol. Biol.* **359**, 533-538 (2006).

28. T. Miyamoto, M. Katane, Y. Saitoh, M. Sekine, H. Homma, Involvement of penicillin-binding proteins in the metabolism of a bacterial peptidoglycan containing a non-canonical D-amino acid. *Amino Acids*. **52**, 487-497 (2020).
29. J. Szewczyk, J. F. Collet, The journey of lipoproteins through the cell: one birthplace, multiple destinations. *Adv. Microb. Physiol.* **69**, 1-50 (2016).
30. T. Yakushi, T. Tajima, S. Matsuyama, H. Tokuda, Lethality of the covalent linkage between mislocalized major outer membrane lipoprotein and the peptidoglycan of *Escherichia coli*. *J. Bacteriol.* **179**, 2857-2862 (1997).
31. A. Mychack, *et al.*, A synergistic role for two predicted inner membrane proteins of *Escherichia coli* in cell envelope integrity. *Mol. Microbiol.* **111**, 317-337 (2019).
32. J. Sheikh, S. Hicks, M. Dall'Agnol, A. D. Phillips, J. P. Nataro, Roles for Fis and YafK in biofilm formation by enteroaggregative *Escherichia coli*. *Mol. Microbiol.* **41**, 983-997 (2001).
33. J. H. Miller, A Short Course in Bacterial Genetics: A Laboratory Manual and Handbook for *Escherichia coli* and Related Bacteria (Cold Spring Harbor Lab Press, Cold Spring Harbor, NY) (1992).
34. T. Baba, *et al.*, Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2**:2006.0008 (2006).
35. K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 6640–6645 (2000).
36. S. Uzzau, N. Figueroa-Bossi, S. Rubino, L. Bossi, Epitope tagging of chromosomal genes in *Salmonella*. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 15264-15269 (2001).

479 **Figures:**

Fig. 1

480

481 **Fig. 1.** Diagrammatic representation of the cell envelope of *E. coli*. Cell envelope consists of
482 three layers- outer membrane (OM), peptidoglycan (PG) and inner membrane (IM). PG is
483 stapled to the OM by Lpp or Braun lipoprotein (red helix) which exists in bound or free form
484 (5–9). In the bound form, the N-terminal end of Lpp is anchored to the OM whereas the C-
485 terminal lysine (purple circle) is covalently attached to an mDAP residue (dark green) of the
486 PG stem peptides. The free form of Lpp spans the OM and is exposed to the surface (11).
487 LdtF is identified in this study as an endopeptidase which cleaves PG–Lpp cross-links and
488 also as a glycine-specific carboxypeptidase.

489

Fig. 2

Fig. 2. Genetic interactions of *mepS* with LDTs. (A) WT and its mutant derivatives carrying either deletion of *lpp*, *ldtB* or *lpp^{ΔK58}*, an allele lacking C-terminal lysine were tested for viability at 37°C. (B) Indicated strains were grown and viability was tested as described above. (C) A *mepS* deletion mutant carrying either vector (pTrc99a; P_{trc}::) or pRB1 (P_{trc}::*ldtF*) or pRB2 (P_{trc}::*ldtF_{H135A}*) or pRB3 (P_{trc}::*ldtF_{C143A}*) were grown in LB broth supplemented with ampicillin and growth was examined on nutrient agar (NA) plates with or without IPTG (0.25 mM) at 37°C.

Fig. 3

499

500 **Fig. 3.** LdtF modulates PG-Lpp linkages. (A) HPLC chromatograms of PG sacculi of WT
501 carrying either vector ($P_{trc}::$), pRB1($P_{trc}::ldtF$), pRB2 ($P_{trc}::ldtF_{H135A}$) or pRB3
502 ($P_{trc}::ldtF_{C143A}$). Cultures were grown to an A_{600} of ~ 1 in LB containing 0.2 mM IPTG
503 followed by isolation and analysis of PG sacculi. (B) Structures of mucopeptides (C) Mass
504 spectra of peaks 3 and Y showing molecular mass $(M+H)^+$ of 1,155.58 Da and 1795.78 Da
505 (D) Determination of PG-Lpp linkages in WT, *ldtF* mutant was done by treating intact PG
506 sacculi (with bound Lpp) with mutanolysin followed by electrophoresing the soluble
507 mucopeptides. Lpp containing mucopeptides were visualized by western blot using anti-Lpp
508 antibody. PG from *ldtABC* mutant was used as negative control. Cell lysates of WT, *ldtF* and
509 *ldtABC* were used as controls (lanes 4, 5 and 6).

510

Fig. 4

511

512 **Fig. 4.** Endopeptidase activity of LdtF. (A) Soluble mucopeptides of WT PG sacculi, (B)
513 purified tri-lys-arg, (C) purified tetra-tri-lys-arg, or (D) purified tetra-tetra dimer were
514 incubated either with buffer or LdtF (4 μ M) for 16 h and separated by RP-HPLC. LdtF
515 cleaved peak 3 (tri-lys-arg) to yield tri (peak 1); and peak 6 (tetra-tri-lys-arg) to yield tetra-tri

516 (peak Y). LdtF showed an extremely weak activity on tetra-tetra dimer (peak 5) (E) Cleavage
 517 of Lpp protein from the PG sacculi (containing bound Lpp) of WT, *ldtF* mutant was tested by
 518 incubating the PG sacculi either with buffer, pronase (0.2 mg/ml) or LdtF (4 μ M) for 16 h at
 519 30°C. Pronase, a non-specific protease is used as a positive control.

520

Fig. 5

Fig. 5. LdtF is a glycine-specific carboxypeptidase. (A) Soluble mucopeptides generated from WT cells grown in minimal A medium (33) supplemented with 50 mM glycine, (B) soluble mucopeptides of WT/ $P_{T5-lac}::ldtD$, (C) purified tetra (gly⁴), (D) tri-tetra (gly⁴), or (E) tetra-tetra (gly⁴) were incubated either with buffer or LdtF (4 μ M) and processed as described

526 above. LdtF cleaved the terminal glycine residue completely from peak ‘a’ (tetra-gly⁴), ‘b’
 527 (tri-tetra-gly⁴) or ‘d’ (tetra-tetra-gly⁴) to yield peak 1 (tri), ‘c’ (tri-tri) or Y (tetra-tri)
 528 respectively. All the mucopeptides were analysed by mass spectrometry (Fig. S10).

529

Fig. 6

Fig. 6. LdtF may also cleave Lpp-mediated IM-PG linkages. (A) *yciB dcrB* mutant and its *ldtF*, *ldtB* or *lpp* deletion derivatives were grown overnight in LB broth, serially diluted, and 5 μ L of each dilution were spotted on indicated plates and tested for viability at 30°C. (B) Viability of *yciB dcrB* mutant carrying either vector (*P_{trc}::*) or pRB1 (*P_{trc}::ldtF*) was tested as described above at 37°C. IPTG was used at 0.1 mM.

537 **Supplemental Information:**

538

539

540

541

542

543 **Cleavage of Braun lipoprotein Lpp from the bacterial peptidoglycan by a paralog of**

544 **L,D-transpeptidases, LdtF**

545

546

547

548 Raj Bahadur, Pavan Kumar Chodiseti, and Manjula Reddy*

549 CSIR-Centre for Cellular and Molecular Biology

550 Hyderabad India 500007

551

552

553

554

555

Supporting Materials and Methods

Plasmid constructions

For PCR amplifications, genomic DNA of MG1655 strain was used as a template unless otherwise indicated. Amplification of DNA for cloning purposes was done using Phusion DNA polymerase (NEB) and clones obtained were confirmed by sequence analysis.

pRB1. The *ldtF* gene along with its native ribosome binding site (RBS) was PCR amplified using forward and reverse primers 5'-GCTCTAGAAGGAATAAGCAGTATGCGTAAA-3' and 5'-CCCAAGCTTTTATTTTGCCTCGGGGAGCGTGT-3' respectively and the resulting amplified DNA fragment was cloned using XbaI and HindIII sites (underlined in the primer sequence) in a cloning vector, pTrc99a to obtain pRB1. The clones were confirmed by sequence analysis and shown to suppress the NA-sensitivity of $\Delta mepS$ mutant at 37°C with 250 μ M IPTG.

pRB2 and pRB3. To create site directed variants of *ldtF* (H135A and C143A), a 3 step PCR was performed. For this procedure, two primers were synthesized that are complementary to each other with desired mutations at the center (in bold and underlined). In the first PCR step, N-terminal fragment of *ldtF* gene was amplified using a common forward primer and a reverse primer containing the desired mismatch. In the second PCR step, C-terminal fragment of *ldtF* gene was amplified using a forward primer containing the desired mismatch and a common reverse primer. Desired mismatches code for an alanine instead of H135, and C143 in the LdtF. Common forward primer (containing its native RBS and an XbaI site), and reverse primer (containing a HindIII site) are used for cloning these variants.

Common forward primer: 5'-GCTCTAGAAGGAATAAGCAGTATGCGTAAA-3' and

579 Reverse primer: 5'-CCCAAGCTTTTATTTTGCCTCGGGGAGCGTGT-3'.

580 The forward and reverse primers with nucleotide substitution are:

581 Histidine to alanine change at codon 135-

582 5'-AAGGGAAATACCTGATGATCGCTGGCGATTGTGTTTCCATCGG-3' and

583 5'-CCGATGGAAACACAATCGCCAGCGATCATCAGGTATTTCCCTT-3'

584 Cysteine to alanine change at codon 143-

585 5'-GCGATTGTGTTTCCATCGGCGCTTACGCAATGACCAATCAGGG-3' and

586 5'- CCCTGATTGGTCATTGCGTAAGCGCCGATGGAAACACAATCGC-3'

587 In the third step, both the PCR products were mixed in 1:1 molar ratio and end filling was
588 done by PCR in 10 cycles at low annealing temperature. After addition of common forward
589 and reverse primer, PCR was resumed for the next 30 cycles. The final PCR product was
590 digested with XbaI-HindIII and cloned into pTrc99a digested with the same enzymes. The
591 recombinant plasmids, pRB2 (*ldtF*-H135A) and pRB3 (*ldtF*-C143A) were confirmed for the
592 presence of mutations by sequencing.

593 **pRB4.** A fragment encoding LdtF²⁰⁻²⁴⁶ was cloned into pET21b vector in between NdeI and
594 XhoI sites using forward and reverse primers

595 5'-GGAATTCCATATGGGTTTGCTGGGCAGCAGTAG-3 and

596 5'-CCGCTCGAGTTTTGCCTCGGGGAGCGTGTAG-3' respectively to generate a C-
597 terminal 6XHis fusion vector. The plasmid was confirmed by sequencing and used for
598 expression and purification of LdtF.

599 **Construction of an *ldtF* deletion mutation (Δ 149*ldtF*::Kan)**

We constructed a partial deletion mutant of *ldtF* lacking N-terminal 1-149 amino acids using recombineering as described earlier (35). The hybrid primers used for constructing this deletion are:

FP: 5'-

GTCCTGGCGTGTGTAACCGTTTTATCAAGGAATAAGCAGTATGTGTAGGCTGGAG
CTGCTTC-3'

RP: 5'-

CACCAGCGCACCAGTAACGAACTGGAATATCTCATCAATACCATATGAATATCCT
CCTTAG-3'

In the first step, the Kan^R cassette of pKD4 vector (35) was amplified with above set of primers and the purified PCR product was electroporated into a strain encoding λ Red-Gam system. Transformants were selected on plates supplemented with 25 μ g/ml kanamycin, followed by confirming the deletion by sequencing and linkage analysis. The deletion was subsequently transferred by P1 transduction into MG1655. This deletion mutant behaved exactly like that of the deletion mutant of Keio collection (in terms of causing sickness to *mepS mepK* double mutant and in PG composition) and hence, Keio deletion mutation was used throughout the study.

Construction of *ldtF*-FLAG fusion

Epitope tagging of LdtF using a 3xFlag was done at the 3' end of the gene at its native chromosomal locus by recombineering as described earlier (36). The sequence of the hybrid primers used for construction of this fusion is given below:

621 FP: 5'-
622 GCAGCCACAACCTGGCATCAAACCTACACGCTCCCCGAGGCAAAAGACTACAAAGA
623 CCATGACGGTG-3' and,
624 RP: 5'-
625 CGGGCAATGAAACCTGGCAAAAGATTATGCCAGGCGAATGGCGCCATATGAATA
626 TCCTCCTTAG-3'

627 The 3xFlag tag along with Kan^R cassette was amplified from plasmid pSUB11 (36) using the
628 above primers. The 5' end of (43 bases) forward primer has homology to the C-terminal of
629 *ldtF* without stop a codon whereas the 3' end (22 bases) has homology to the region encoding
630 Flag epitope of pSUB11. Similarly, the 5' end of reverse primer has a region homologous (42
631 bases) to the downstream sequence of *ldtF* whereas the 3'end has a region (22 bases) which
632 has homology to the sequence of pSUB11 plasmid at the 3' end. The PCR product was
633 electroporated into DY378 and transformants were selected on plates supplemented with 25
634 µg/ml Kanamycin at 30°C on LB. The putative *ldtF*-3xFlag-Kan^R region was transferred into
635 MG1655 by P1 transduction and the construct was confirmed by sequencing and linkage
636 analysis. The expression of the fusion tag was confirmed by western blotting using anti-
637 FLAG antibodies.

638 **Confirmation of deletion mutation (mutant from Keio collection) of *ldtF***

639 The Keio deletion mutant of *ldtF* was confirmed by sequencing the gene-Kan^R junctions. The
640 region encompassing the deletion mutation was amplified using the below primers and
641 sequenced using the same primers.

642 FP: 5'- GACAGGCTTGCGTAAACTC-3 and,
643 RP: 5'- CAGGATGTGGAAATCGACTTCAGC-3

Methods:

Purification of LdtF

LdtF encoding plasmid, pET21b-*ldtF*²⁰⁻²⁴⁶ (pRB4) was transformed into BL21 (λDE3) strain and transformants were selected on LB plates supplemented with ampicillin (Amp). One purified colony was grown overnight and used to dilute 1:100 into 50 ml fresh LB broth containing Amp. Culture was induced with 250 μM IPTG at 0.6 OD and further allowed to grow for 2 h at 37°C. Cells were harvested, and pellet was stored at −30°C. When required, pellet was resuspended in 1 ml of buffer (50 mM Tris-Cl, 300 mM NaCl and 20 mM imidazole, pH 8.0) and lysed by sonication. Cell debris was removed by centrifugation and the supernatant was mixed with 200 μl Ni²⁺-NTA agarose (Qiagen) and mixed for 1 h at 4°C. This mixture was loaded onto pre-washed empty column (Bio-Rad) and washed with 30 ml wash buffer-1 (50 mM Tris, 300 NaCl, 30 mM imidazole, pH 8.0) and 20 ml of wash buffer-2 (50 mM Tris, 300 NaCl, 50 mM imidazole, pH 8.0). The bound proteins were eluted with 5 ml of elution buffer (50 mM Tris, 300 NaCl, 150 mM imidazole, pH 8.0) and concentrated to 2.5 ml using a 3 kDa cut-off centrifugal membrane filter (Millipore). This was then loaded onto a buffer exchange PD-10 column and protein was eluted into 3.5 ml 2 x storage buffer (100 mM Tris, 200 mM NaCl, 2 mM DTT). The fraction was further concentrated to 250 μl using 3 kDa cut-off centrifugal membrane filter and mixed with equal volume of 100% glycerol and stored at −30°C.

Viability assays and Microscopy. Viability of the indicated strains was examined by growing cultures overnight, serially diluting (10^{-2} , 10^{-4} , 10^{-5} and 10^{-6}), and placing 3-5 μl aliquots of each dilution onto the required plates. Plates were normally incubated for 18-24 h at the indicated temperature. To measure growth rates, overnight grown cultures were diluted 1:100 into fresh medium, allowed to grow and OD at 600 nm was determined after every 1 h

interval and data were plotted using Origin software. For microscopy, immobilized cultures on a thin 1% agarose pad were visualized using Zeiss AxioImager.Z2 microscope by DIC.

Preparation of PG sacculi. Isolation of PG was done as described earlier (2,24). Cells grown to A_{600} of 1.0 were collected by centrifugation at $10,000\times g$ for 10 min at 4°C . Cell pellet (from 1000 ml) was resuspended in 6 ml of ice-cold deionized water and added drop wise into 6 ml of boiling 8% SDS with vigorous stirring followed by boiling for another 45 min to solubilize membranes and to destroy high molecular weight DNA. After overnight incubation at room temperature, the PG sacculi were collected by high speed centrifugation ($200,000\times g$, 40 min) and washed thoroughly with deionized water to completely remove SDS. High molecular weight glycogen and covalently bound lipoprotein, Lpp were removed by treating with α -amylase (100 $\mu\text{g/ml}$ in 10 mM Tris-HCl, pH 7.0, 2 h at 37°C) and pre-digested pronase (200 $\mu\text{g/ml}$, 90 min at 60°C). Enzymes were inactivated by boiling with equal volume of 8% SDS for 15 min and pure sacculi were obtained by ultracentrifugation and washed several times with water till SDS was completely removed. Final pellet was resuspended in 0.5 ml of 25 mM Tris-HCl (pH 8.0) and stored at -30°C .

Lpp-bound PG sacculi were prepared from cultures grown up to A_{600} of ~ 6.0 using the above-described protocol except that the treatment with Pronase was not done.

Analysis of PG sacculi. PG analysis was done as previously described (2,24). Essentially, the sacculi were digested with 10 U mutanolysin (Sigma-Aldrich) at 37°C in 25 mM Tris-HCl (pH 8.0) for 16 h and soluble muropeptides were collected after centrifugation. This fraction was reduced with 1 mg of sodium borohydride in 50 mM sodium borate buffer (pH 9.0) for 30 min and excess borohydride was destroyed by addition of 20% phosphoric acid. pH was adjusted to 3–4 and samples were loaded onto a reverse phase C18 column (Zorbax 300 SB; 250×4.6 mm, 5 mm) connected to Agilent technologies RRLC 1200 system. Column

temperature was 55°C and binding was done at a flow rate of 0.5 ml/min with 1% acetonitrile in water containing 0.1% trifluoroacetic acid (TFA) for 10 min. Muropeptides were eluted in a gradient of 1–10% acetonitrile containing 0.1% TFA at a flow rate of 0.5 ml/min for next 60 min (using RRLC online software called Chemstation). Absorbance of muropeptides was detected at 205 nm.

Mass spectrometry (MS) analysis of muropeptides. Muropeptide fractions collected during HPLC were dried and reconstituted into 5% acetonitrile with 0.1% formic acid and loaded onto a reverse phase PepMapTM RSLC - C18 column (3 µm, 100Å, 75µmx, 15cm) connected to Q-ExactiveTM HF Hybrid Quadrupole-OrbitrapTM Mass Spectrometer (Thermo Fisher Scientific, USA). Peaks were analysed by tandem MS and structures were decoded based on molecular mass of the individual fragments.

Determination of Lpp-bound muropeptides

Normalized muropeptides were boiled with laemmli loading dye and separated using 15% SDS-PAGE. Lpp was detected by western blot using rabbit anti-Lpp antibody (rabbit). As a positive control (free form of Lpp) cell lysates of WT, *ΔldtF*, or *ΔldtABC* (equivalent of 0.012 OD) were used.

Cleavage of Lpp from intact PG sacculi

Lpp-bound PG sacculi were isolated from WT and *ΔldtF* (grown upto ~6 OD) without pronase treatment as described. Equal volumes of the PG sacculi were treated with buffer, pronase (as positive control) or LdtF (4 µM) for 16 h at 30°C. Reaction was stopped by heat inactivation and supernatant was collected after centrifugation at 15000xg for 15 min. These fractions were run on 15% SDS-PAGE to detect the Lpp released from intact PG sacculi. The

714 remaining pellet fraction was further digested with mutanolysin and the resulting soluble
715 mucopeptides were separated by RP-HPLC.

716

717 **Supplementary Figures:**

718

Fig. S1

719

720 **Fig. S1.** Genetic interactions of *mepS* with *ldts*. WT (MG1655) and its deletion mutant
721 derivatives were grown overnight in LB broth, serially diluted, and 5 μ L of each dilution
722 were spotted on indicated plates and tested for viability at 37°C.

Fig. S2

Fig. S2. Microscopic images of WT and its various mutant derivatives. Indicated strains were grown overnight and diluted 1:500 into prewarmed Nutrient Broth and grown till A_{600} of 1.0 at 37°C and visualized with DIC microscopy. The scale bar represents 5 μm.

Fig. S3

Fig. S3. Effect of multiple copies of Ldts on growth of *mepS* mutant. $\Delta mepS$ mutant carrying either pCA24N vector (P_{T5-lac}::) or its derivatives (P_{T5-lac}::*ldtA-F*) were grown and viability was tested on NA plates at 37°C with or without 50 μM IPTG.

Fig. S4

Fig. S4. Growth curve of *ldtF* mutant. Overnight grown cultures of WT and *ldtF* were sub-cultured 1:500 into fresh LB and growth was monitored every 30 min at 37°C. In addition, *ldtF* mutant shows comparable decrease in the doubling time compared to that of WT when grown in any media including LBON or Minimal A media with or without glycine-supplementation.

Fig. S5

737

738 **Fig. S5.** PG composition of *ldtF* mutant. HPLC chromatograms of PG sacculi isolated from
739 WT and *ldtF* mutant. Strains were grown to an A_{600} of ~ 1 in LB followed by isolation and
740 analysis of PG sacculi. Data shown is representative of three independent experiments.

744 +2 charges) is shown. The signature peaks indicating the 3–3 cross-links in tri-tetra are
745 highlighted in red.

746

Fig. S7

Fig. S7. PG composition of strains with ectopic expression of LdtF. HPLC chromatograms of PG sacculi isolated from (A) *lpp* or (B) *ldtDE* deletion mutants carrying either vector (pTrc99a; $P_{trc}::$) or pRB1($P_{trc}::ldtF$). Strains were grown to an A_{600} of ~ 1 in LB containing 150 μ M IPTG followed by isolation and analysis of PG sacculi. Note that in absence of Lpp, increased LdtF does not alter the PG composition. On the other hand, the effect of LdtF is independent of LdtD and -E.

Fig. S8

Fig. S8. HPLC chromatograms of Lpp-bound PG sacculi from WT, *ldtF* and *ldtABC* mutant. The data obtained from these chromatograms was used to normalize the amount of sample loaded for performing the experiment described in Fig. 3D.

Fig. S9

Fig. S9. HPLC chromatograms of Lpp-bound PG sacculi from WT and *ldtF* mutant. The PG sacculi were treated either with buffer, pronase (0.2 mg/ml) or LdtF (4 μ M) for 16 h at 30°C and the remaining insoluble fraction was treated with mutanolysin and separated by HPLC. Note that the tri-lys-arg peak is absent in LdtF-treated PG sacculi.

Fig. S10

Fig. S10. Identification of muropeptides using Mass spectrometry analysis. Mass spectra of various muropeptides indicating the molecular mass (both +1 and +2 charges) is shown.

Fig. S11

Fig. S11. Effect of LdtF on glycine-toxicity of *dacABCD* mutant. Cultures of *dacABCD* and *dacABCD ldtF* mutants carrying either vector (pTrc99a; P_{trc}::) or pRB1(P_{trc}::*ldtF*) were grown in LB broth, serially diluted, and 5 µL of each dilution were spotted on indicated plates and tested for viability at 37°C. Glycine and IPTG are used at 25 mM and 100 µM respectively. *dacABCD* deletion mutant was slow-growing on glycine-supplementation and *ldtF* deletion exacerbated the sickness whereas more copies of *ldtF* marginally improved the growth of the quadruple mutant.

Fig. S12 A

774
775

Fig. S12 B

776

777 **Fig. S12.** Effect of LdtF on NCDAAs containing mucopeptides. (A) Mass spectrometry

778 analysis of mucopeptides containing NCDAAs (D-methionine, D-tryptophan or D-

779 phenylalanine) (B) Treatment of soluble mucopeptides from PG sacculi of strains grown in 20

780 mM D-methionine (I), 15 mM D-tryptophan (II), or 15 mM D-phenylalanine (III) were

781 incubated either with buffer or LdtF (4 μ M) for 16 h and separated by RP-HPLC. LdtF did
782 not cleave the terminal D-amino acids from any of the NCDAA-muropeptides.

Table S1. Strains used in this study

Strains	Genotype ^a	Source/Reference
WT (MG1655)	<i>rph1 ilvG rfb-50</i>	Lab collection
DH5α	F ⁻ <i>hsdR17 deoR recA1 endA1phoA supE44 thi-1 gyrA96 relA1 Δ(lac-argF)U169 φ80dlacZ ΔM15</i>	Lab collection
BL21 (λDE3)	<i>ompT rB⁻ mB⁻ (P_{lac}UV5::T7gene1)</i>	Lab collection
DY378	Recombineering strain; W3110 <i>λcI857 Δ(cro-bioA)</i>	Lab collection
	<i>ΔmepS::frt</i>	23
RB11	<i>ΔldtF::Kan</i> (Keio)	34
RB12	<i>ΔI49ldtF::Kan</i>	This study
RB13	<i>Δlpp::Kan</i>	This study
RB14	<i>^blpp^{ΔK58}</i>	This study
RB15	<i>ΔldtABC::frt</i>	This study
RB15	<i>ΔldtDE::frt</i>	This study
	<i>ΔyciB ΔdcrB</i>	31
RB16	<i>ΔdacABCD::frt</i>	This study

^aDeletion alleles used in this study are sourced from Keio collection (34). The deletion mutations were used after testing for their authenticity (by linkage analysis, PCR and sequence analysis) and introduced into different strain backgrounds by P1 phage-mediated transduction (33). MG1655 was used as the wild type strain. The Kan marker from strains was flipped out using pCP20 plasmid encoding a Flp recombinase (35). Flipping creates a ‘*frt*’ scar at the site of deletion.

^blpp^{K58} allele is from Thomas Silhavy’s laboratory. It was transferred by P1 transduction using a 60% linked tetracycline marker.

794 **Table S2. Plasmids used in this study**

Plasmid	Relevant features	Source/Reference
pET21b	ColE1, Amp ^R , <i>lacIq</i> , T7 <i>lac</i>	Lab collection
pTrc99a	ColE1, Amp ^R , <i>lacIq</i> , P _{trc}	Lab collection
pCA24N	Cm ^R , <i>lacIq</i> , P _{T5-lac}	26
pCA24N- <i>ldtA</i>	Cm ^R , <i>lacIq</i> , P _{T5-lac} :: <i>ldtA</i>	26
pCA24N- <i>ldtB</i>	Cm ^R , <i>lacIq</i> , P _{T5-lac} :: <i>ldtB</i>	26
pCA24N- <i>ldtC</i>	Cm ^R , <i>lacIq</i> , P _{T5-lac} :: <i>ldtC</i>	26
pCA24N- <i>ldtD</i>	Cm ^R , <i>lacIq</i> , P _{T5-lac} :: <i>ldtD</i>	26
pCA24N- <i>ldtE</i>	Cm ^R , <i>lacIq</i> , P _{T5-lac} :: <i>ldtE</i>	26
pCA24N- <i>ldtF</i>	Cm ^R , <i>lacIq</i> , P _{T5-lac} :: <i>ldtF</i>	26
pRB1	pTrc99a- <i>ldtF</i>	This study
pRB2	pTrc99a- <i>ldtF</i> -H135A	This study
pRB3	pTrc99a- <i>ldtF</i> -C143A	This study
pRB4	pET21b- <i>ldtF</i> ²⁰⁻²⁴⁶	This study

795

Table S3. Muropeptide composition of strains carrying multiple copies of *ldtF*

Muropeptide (Peak)	% Area of muropeptide peaks ^a			
	WT/ P _{trc} ::	WT/ P _{trc} :: <i>ldtF</i>	WT/ P _{trc} :: <i>ldtF</i> _{H135A}	WT/ P _{trc} :: <i>ldtF</i> _{C143A}
Tri (1)	5.81 ± 0.06	15.74 ± 0.47	9.51 ± 0.47	8.91 ± 1.41
Tetra (2)	38.91 ± 0.37	34.19 ± 1.70	35.14 ± 1.42	37.6 ± 1.47
Tri-lys-arg (3)	4.57 ± 0.53	0.7 ± 0.60	2.2 ± 1.22	1.86 ± 1.06
Tetra-tri (Y)	3.0 ± 0.23	11.75 ± 2.67	5.34 ± 0.06	5.27 ± 0.77
Tri-tetra (4)	3.54 ± 0.35	2.47 ± 0.45	3.99 ± 0.25	4.87 ± 0.18
Tetra-tetra (5)	32.22 ± 0.70	25.15 ± 1.79	27.44 ± 2.6	26.33 ± 2.91

^aMuropeptide analysis was done by calculating the relative percentage area of each muropeptide from the HPLC chromatograms.