Title: Approximate Bayesian computation untangles signatures of contemporary and historical hybridization between two endangered species

Dittberner Hannes¹, Tellier Aurelien² and de Meaux Juliette^{1*}

¹ Institute of Plant Sciences, Zülpicher str. 47b, University of Cologne, Germany.

² Department of Life Science Systems, Technical University of Munich, Freising, Germany

* Correspondance to jdemeaux@uni-koeln.de

ABSTRACT

Contemporary gene flow, when resumed after a period of isolation, can have crucial consequences for endangered species, as it can both increase the supply of adaptive alleles and erode local adaptation. Determining the history of gene flow and thus the importance of contemporary hybridization, however, is notoriously difficult. Here, we focus on two endangered plant species, *Arabis nemorensis* and *A. sagittata*, which hybridize naturally in a sympatric population located on the banks of the Rhine. Using reduced genome sequencing, we determined the phylogeography of the two taxa but report only a unique sympatric population. Molecular variation in chloroplast DNA indicated that *A. sagittata* is the principal receiver of gene flow. Applying classical D-statistics and its derivatives to whole-genome data of 35 accessions, we detect gene flow not only in the sympatric population but also among allopatric populations. Using an Approximate Bayesian computation approach, we identify the model that best describes the history of gene flow between these taxa. This model shows that low levels of gene flow have persisted long after speciation. Around 10 000 years ago, gene flow stopped and a period of complete isolation began. Eventually, a hotspot of contemporary hybridization was formed in the unique sympatric population. Occasional sympatry may have helped protect these lineages from extinction in spite of their extremely low diversity.

INTRODUCTION

Individual taxa do not always evolve in isolation. Interspecific hybridization, when it leads to fertile offspring, can result in the transfer of alleles across species barriers and even allow new species to emerge (Seehausen 2004; Abbott 2017). The footprints of gene flow are detectable in many genera and demes (Seehausen 2004; Marcet-Houben and Gabaldón 2015; Ackermann et al. 2019; Taylor and Larson 2019). Genomic analyses have revealed that species barriers are established progressively, depending on the size and degree of reproductive isolation of the species, with genetic variation being shared over periods of time that are longer than previously thought (Brandvain et al. 2014; Novikova et al. 2016; Edelman et al. 2019; Small et al. 2020). The climatic oscillations of quaternary glaciation cycles have likely contributed to multiple opportunities for hybridization in many taxa (Hewitt 2000).

The evolutionary consequences of hybridization can be manifold, offering a powerful channel for some taxa to capture and quickly fix alleles that have been subjected to selection in other species (Hedrick 2013; Goulet et al. 2017; Suarez-Gonzalez et al. 2018a; Vallejo-Marín and Hiscock 2018). The potent adaptive potential unleashed by hybridization have been confirmed in a number of species (Rieseberg et al. 2003; Baduel et al. 2018; Ma et al. 2019; Marburger et al. 2019). For example, introgression appeared to accelerate local adaptation via the transfer of alleles contributing to the success of a relative of the sunflower (Todesco et al. 2020). In *Heliconius* butterflies, an inversion associating genes that control color patterns has been repeatedly exchanged between species (Edelman et al. 2019). In addition, transgressive phenotypic variation can emerge from the recombination of resident and incoming alleles throughout the genome (Rieseberg et al. 1999; Seehausen 2004). Positive selection is therefore expected to increase introgression rates specifically in and around adaptive loci. Heterogeneous introgression rates, however, can also arise along the genome in the absence of adaptive gene flow (Schumer, Rosenthal, et al. 2018). Covariation between the rate of introgression, recombination and gene density, for example, indicates that selection acts to limit introgression and points to the existence of barriers to gene flow (Brandvain et al. 2014; Schumer, Xu, et al. 2018). In humans, the impact of deleterious alleles was shown to correlate negatively with the frequency of alleles introgressed from the related species *Homo neanderthalis*, whose

population size was low and eventually collapsed (Juric et al. 2016; Steinrücken et al. 2018). Ultimately, high rates of gene flow can introduce maladapted alleles and ultimately impede adaptation (Lenormand 2002; Yeaman 2015; Tigano and Friesen 2016). These alleles, in turn, can select for alleles that will reinforce species barriers to prevent resources from being wasted by producing poorly performing hybrids (Hopkins and Rausher 2012). Gene flow, regardless of whether it promotes or erodes adaptation, is expected to create a heterogeneous pattern of introgression throughout the genome (Martin and Jiggins 2017; Schumer, Rosenthal, et al. 2018).

If gene flow can be either a blessing or a curse, it is particularly crucial to understand its importance for species that have to be protected from extinction. In a context of global climate change, the number of threatened species is expected to increase (Díaz et al. 2019; Eichenberg et al. 2020). At the same time, many species barriers that have so far been maintained by habitat, phenological or behavioral separation are likely to disintegrate, generating unprecedented opportunities for novel episodes of hybridization (Anderson et al. 2012; Chunco 2014). Contemporary hybridization will most affect endangered species if the taxa involved have been separated long enough to acquire distinct ecological and population genomics characteristics. For example, the hybridization of long separated continental subspecies of salmon has been associated with the manifestation of Dobzhansky-Muller incompatibilities (Rougemont and Bernatchez 2018). In *Mimulus*, barriers to gene flow are strong today despite an ancient history of hybridization, but contemporary hybridization has been reported at specific locations (Brandvain et al. 2014; Kenney and Sweigart 2016). The potential of hybridization to create a novel genetic make-up today that may impact future evolution and support the evolutionary rescue of biodiversity depends, therefore, on how much time has elapsed during the period of isolation.

Determining the history of gene flow between taxa is a complex task. Dating gene flow cannot be achieved with a single summary statistic of genomic variation, because effective population sizes, migration rates and/or the time since species formation jointly influence patterns of standing variation, divergence and expected allele sharing. The widely used D-statistics tests for gene flow, while accounting for

incomplete lineage sorting at the time of speciation (Durand et al. 2011). This and related statistics have, for example, revealed the extent of gene flow between species thought to be separated by differences in ploidy (Arnold et al. 2016; Paape et al. 2018; Kryvokhyzha et al. 2019). Yet such statistics do not allow speciation or gene flow to be dated, nor do they evaluate the duration of isolation periods (Schumer, Rosenthal, et al. 2018; Hibbins and Hahn 2019). Failing to understand the correct history of speciation and gene flow can lead to confounding neutral patterns of gene flow with convergent evolution, sympatric speciation or even adaptive divergence (Bierne et al. 2013; Ravinet et al. 2017).

The evolutionary and demographic histories of hybridizing species are often too complex to be determined analytically. Intensive simulation methods, such as Approximate Bayesian Computation (ABC), offer a powerful alternative (Csilléry et al. 2010). It not only makes it possible to choose among competing models for the one best able to explain the data, but also enables population parameter (population sizes, migration rates) and their fluctuation over time to be estimated (Rougemont and Bernatchez 2018). However, ABC approaches have rarely been applied to all genome data and when, then only to infer relatively simple demographic events such as bottlenecks or expansions (Jay et al. 2019). The current limitation stems from the inherent statistical complexity of the approach (high dimensionality of whole-genome data), the necessity for an educated guess regarding the choice of summary statistics to use, and the lack of user-friendly ready-to-use software capable of handling all genome data. The potential of ABC approaches to determine the timing, intensity and duration of hybridization episodes based on all genome data thus remains to be fully leveraged.

Here, we focused on a documented case of contemporary hybridization between two endangered plant species and reconstructed the history of gene flow. The species we examined, *Arabis nemorensis* and *A. sagittata*, are selfing biennial forbs from the Brassicaceae family. *A. nemorensis*, which is strictly confined to floodplain environments in Central and Northern Europe, harbors extremely low levels of nucleotide diversity (Dittberner et al. 2019). Since the 1950s, conversion to arable land and management intensification have reduced the area covered by species-rich floodplain meadows, the ecosystem in which *A. nemorensis* thrives, by more than 80

%. Only small remnant patches have persisted within protected areas (Hölzel 2005). The unique ecology of *A. nemorensis*, its shrinking habitat and its selfing mating system all contribute to the acute danger of extinction this species finds itself in (Hölzel 2005; Burmeier et al. 2010; Mathar et al. 2015). *A. nemorensis*, however, was found to occur in sympatry with its relative, *A. sagittata*, in a small set of pristine habitat patches located on the banks of the Rhine near Mainz, Germany (Dittberner et al. 2019). *A. sagittata*, known to thrive in relatively dry environments, is also endangered, but its presence in floodplain meadows is novel (Hand and Gregor 2006; Dittberner et al. 2019). Interestingly, approximately 10% of the individuals of the sympatric population appeared to be hybrids between these two species (Dittberner et al. 2019).

Using a combination of reduced sequencing and whole-genome sequencing, we asked the following questions: Is hybridization local or is there a large contact zone? Is gene flow symmetrical? Do rates of introgression vary along the genome? Is hybridization contemporary or was gene flow continuous throughout the history of the species? Our study confirms the existence of a contemporary hotspot of hybridization where asymmetric gene flow has resumed after approximately 10 000 years of complete isolation between taxa.

RESULTS

Hybrids detected only in a single sympatric population

We combined previously published and newly generated RAD-sequencing data for a total of 231 accessions to describe the phylogeographic distribution of *A. nemorensis* and assess the distribution of *A. sagittata* and interspecific hybrids in the area of distribution of *A. nemorensis* (Table S1). This collection of genotypes covered all sites where the presence of *A. nemorensis* had been documented (see methods). We performed an admixture analysis (Alexander and Lange 2011a) and identified two genetic clusters that were previously assigned to *A. nemorensis* and *A. sagittata* (Figure S1-5, Dittberner et al. 2019). We found individuals of the two species growing in sympatry in only one of the eleven sites (the Rhine population), indicating that opportunities for natural hybridization are restricted (Figure 1A, Dittberner et al. 2019). Of the 140 individuals sampled in the sympatric population, 75 were *A. sagittata*, 42 were *A. nemorensis* and 23 were individuals we identified as having a hybrid origin because they showed less than 95% purity in the output of the ADMIXTURE analysis. Phenotypic observations, such as low seed number per silique and elongated stems, had suggested that hybridization may occur in some rare instances (Novotná and Czapik 1974; Titz 1979). In contrast, genetic analyses indicate that the two species frequently cross-hybridize within the sympatric population (Dittberner et al. 2019). Interestingly, *A. sagittata* ancestry predominates in most hybrid individuals (Figure 1B), indicating the frequent backcross of hybrids with *A. sagittata* individuals, which are more frequent than *A. nemorensis* individuals in the sympatric population.

Chloroplast DNA indicates that *A. nemorensis* is the maternal parent of hybrids

Chloroplast DNA is maternally inherited and its sequence variation can provide information about the maternal genotype of hybrids (McCauley 1995). To determine the maternal genotypes of hybrid individuals, we focused on RADseq stacks that mapped to

a total of 10 kb (5%) of the chloroplast sequence of the close relative *Arabis hirsuta* (Kawabe et al. 2018). Haplotype network analysis showed that the chloroplast sequences of *A. sagittata* formed four closely related haplotype groups and two for *A. nemorensis* (Figure 3C). As all but one interspecific hybrid had an *A. nemorensis* haplotype, we conclude that *A. nemorensis* is the maternal genotype of most hybrid individuals. This observation confirms that gene flow between the two clusters is asymmetrical. Surprisingly, 24 *A. sagittata* individuals were found to carry the *A. nemorensis* chloroplast haplotype although the admixture analysis found no trace of *A. nemorensis* in their nuclear genome. This suggests that some individuals may have a hybrid ancestry but their *A. nemorensis* ancestry is undetectable in the admixture analysis of RAD-seq data, showing in turn that the examination of whole-genome sequences was necessary.

Figure 1: Hybridization between *A. nemorensis* and *A. sagittata*

A) Map of sampled populations. Each population is represented by a pie chart showing the average ancestry proportions of *A. nemorensis* and *A. sagittata* in the given population, based on RAD-sequencing data. B) Representation of individual ancestry components in the sympatric ("Rhine") population, based on RAD-seq data. Each bar represents one individual and is colored according to its genomic ancestry. Bars are ordered by decreasing *A. sagittata* ancestry. Hybrid admixed individuals are

framed with a purple rectangle. C) Network of chloroplast RAD-seq haplotypes. Each pie-chart represents one haplotype and the fractions of species/populations carrying this haplotype. Haplotypes are connected to their closest relative by a line. Orthogonal dashes represent the number of mutations. Abbreviations in the legend: nemo. - *A. nemorensis*, sag. - *A. sagittata*, allo. - allopatric, symp. - sympatric.

Gene flow between *A. nemorensis* and *A. sagittata* is not restricted to the sympatric population

To quantify interspecific gene flow between the two genetic clusters, we resequenced 35 whole genomes of accessions from both allopatric and sympatric populations for the two species, as well as one individual of the closest diploid relative, *A. androsacea* (Table S2). In order to understand the history of gene flow, we specifically excluded individuals that were predicted to be admixed hybrids (Fig. 1B), which were obviously formed a handful of generations ago. Analyses presented hereafter were all conducted on data generated by whole-genome sequencing in this set of 35 accessions. We first computed Patterson's *D* (ABBA-BABA) statistic (Durand et al. 2011; Green et al. 2010), a statistic that quantifies gene flow after accounting for incomplete allele sorting. This statistic is computed over the whole genome for phylogenies of the form (((P1,P2),P3),O), where O is the outgroup *A. androsacea* and P1 and P2 two populations of the same species. *D* was highest for the phylogeny with P1 = *A. sagittata* allopatric, P2 = *A. sagittata* sympatric and P3 = all *A. nemorensis* individuals with 0.2 ($p < 0.001$; block jackknife test). This significant and positive *D* value shows that gene flow is higher between P3 *A. nemorensis* and P2, the sympatric *A. sagittata* population, than between *A. nemorensis* and P1, the allopatric *A. sagittata* individuals (Figure 2A). Although this test showed elevated rates of gene flow in the sympatric population, it did not rule out introgressions in the allopatric populations. Thus, we also compared the two allopatric *A. sagittata* populations Lob and Adl-1 (Figure 1A) with P1 = *A. sagittata* Lob, P2 = *A. sagittata* Adl-1 and P3 = all *A. nemorensis*, and found *D* was 0.09 ($p < 0.001$; block jackknife test) indicating significant gene flow outside of the sympatric population. Finally, we tested the phylogeny with P1 = *A. nemorensis* sympatric, P2 = *A. nemorensis* allopatric and P3 = *A. sagittata* Lob (allopatric), and found that *D* was 0.17 ($p < 0.001$, block jackknife test).

This positive value indicated that gene flow from *A. sagittata* into *A. nemorensis* was stronger in allopatry than in sympatry, demonstrating that the history of hybridization in this system predates the formation of the sympatric population.

The frequency of introgression is heterogeneous along the genome

The *D*-statistic provides information about genome-wide rates of gene flow but not about its distribution across the genome. Thus, we calculated the f_D -statistic (Martin et al. 2015) across the genome for five phylogenies covering all possible scenarios of interspecific gene flow (Figure 2B). Distributions of f_D were highly zero-inflated for all populations, indicating that introgressions were generally rare across the genome. Yet, in all populations, we found genomic regions with elevated f_D values. We did not find strong differences in the distribution of these regions among chromosomes. The *D* and f_D statistics rely on the assumption that there was no gene flow between P3 and P1 (Green et al. 2010; Durand et al. 2011; Martin et al. 2015). As we found introgressions in all populations, this assumption was violated, meaning any introgression shared by P1 and P2 would remain undetected. Our estimates of gene flow are therefore likely to be conservative.

Introgressed fragments are on average largest in the sympatric population

To locate introgressed fragments and their boundaries in the genome, we calculated the ratio of average intra- and interspecific genetic distance for each individual in 10 kb genomic windows (see Methods). This ratio is expected to be smaller than the ratio throughout the genome, except in introgressed regions. We used a value equal or greater than 2 as a conservative threshold for the ratio value calling introgressed fragments. The average number of introgressions per individual was highest in the sympatric *A. sagittata* population, with 24.7 introgressed fragments per genome (standard deviation (sd) = 3.6), followed by the allopatric *A. sagittata* populations, with an overall mean of 13.7 fragments per genome (sd = 3.5, Table S3). In contrast, in *A. nemorensis*, we observed an average of 9.6 introgressed fragments per genome (sd = 1.47) in the sympatric population and 2.6 (sd = 0.55) in the allopatric. These results confirmed that interspecific gene flow from *A. nemorensis* to *A. sagittata*

was stronger than vice versa. Furthermore, the presence of introgressions in allopatric populations suggests that gene flow is at least partly historical.

The high occurrence of introgression fragments in the sympatric *A. sagittata* population indicates that some of these fragments could have been introduced more recently by contemporary gene flow. We thus assessed whether the introgression was likely to have occurred in the sympatric population (recent gene flow) or whether it predated the separation of local populations (historical gene flow). For this, we determined the individual in the donor species, whose orthologous region was most closely related to the introgressed fragment identified in the receiver species. We then observed that the proportion of introgressed fragments that were most similar to alleles of the sympatric *A. nemorensis* population was almost twice as high in the sympatric *A. sagittata* population (38.5%) as in the allopatric *A. sagittata* population (20%), a difference that was marginally significant (Chi square= 3.76, $p=0.05$, Figure 2C). This trend was reversed in *A. nemorensis*, with 25% of introgressions originating from a sympatric *A. sagittata* lineage in the sympatric *A. nemorensis* population as opposed to 43% in the allopatric *A. sagittata* population. However, as there were only four introgressions in the sympatric population, this reversal is likely due to chance.

Figure 2: Patterns of introgression in sympatric and allopatric populations of parental species.

A) D-statistic results calculated over the whole genome for different phylogenies. Bar color represents the species of the P3 / donor population. Asterisks represent the result of a Jackknife test: *** $p < 0.001$, ** $p < 0.01$. B) Genome-wide distribution of f_D calculated in 50-kbp windows for different phylogenies, as shown on the right. Plots are ordered by decreasing overall genome-wide f_D . C) Fractions of introgression origins for each population, inferred based on minimum genetic distances (see Methods). Numbers on top show the total number of introgressions in each population. This plot does not take into account the frequency of the identified introgression tracts. D) Distribution of introgression size in each population, represented by dots and boxplots. Dots are colored according to the introgression origin. This plot does not take into account the frequency of the identified introgression tracts. The compact letter display indicates significant differences among groups. Abbreviations in the legends: nemo. - A. nemorensis, sag. - A. sagittata, allo. - allopatric, symp. - sympatric

We further reasoned that recent introgressions originating from sympatric donor lineages should be on average larger than introgressions originating from allopatric donor lineages because the latter being more ancient, they should have had more time to be broken down by recombination. Thus, we compared the distributions of

introgression size among populations. In *A. sagittata*, introgressions in the sympatric population were significantly larger than in the allopatric population ($Z=-4.47$, $p<0.001$), with median values of 37,900 bp and 10,000 bp, respectively (Figure 2D). We did not find this difference in the *A. sagittata* allopatric populations (Kruskal-Wallis chi-squared = 0.5174, $df = 1$, $p = 0.472$) or in the two *A. nemorensis* populations (symp.: Kruskal-Wallis chi-squared = 0.6, $df = 1$, $p = 0.4386$; allopat.: Kruskal-Wallis chi-squared = 1.772, $df = 1$, $p = 0.1831$).

The best fit model excludes constant gene flow between the two taxa

Based on the above, we observed i) asymmetrical gene flow between species, ii) evidence for gene flow both within and outside of the sympatric population. Our data thus indicate that gene flow may have been continuous throughout the history of the two species. However, we also observed that gene flow from *A. nemorensis* into *A. sagittata* might have intensified in the sympatric population. Gene flow could thus have occurred in the past and stopped during a phase of complete isolation, only to resume recently in the sympatric population. We took a modelling ABC approach to date gene flow between *A. nemorensis* and *A. sagittata* and to estimate its strength and past fluctuation.

We modelled the history of interspecific gene flow using a random-forest-based approximate Bayesian computation (ABC) approach (Raynal et al. 2019). We started with five demographic models, which explored different modes of intra- and interspecific gene flow (Figure 3A), and generated 50,000 coalescent simulations under each one (see Methods). To choose the model that best fitted the data, we first trained a random-forest classifier on the data all five models use. The accuracy of the trained classifier was evaluated with a confusion matrix that described how many simulations were assigned to the model under which they were generated (Figure 3B). Based on this matrix, we identified two subsets of models that summary statistics could accurately distinguish (max. error 5.2 %). The first subset comprised the two weak-migration models: the model without any migration and the model with early ancestral interspecific migration. The second subset contained the three models with strong interspecific migration: ancestral migration, ancestral and recent sympatric migration, and

continuous migration. Within this last subset, the first model was classified with a high accuracy of 84 %, but the second and third models were only classified with accuracies of 53 % and 61 %, respectively. These results show that i) models with both high and low migration rates could be distinguished easily by the classifier; and ii) determining the exact model within each of the two groups was less straightforward. This result was not surprising because when recent migration is low, the ancestral and recent migration model gives results similar to those generated under the ancestral migration model, which does not allow migration in the sympatric population.

Next, we used the trained random-forest classifier to determine the model that best explained the observed data. To this end, we randomly selected 100 datasets, each consisting of 500 loci of individual length 75 kbp sampled from the observed genotype (full genome) data and summarized with 200 summary statistics (Table S4). Models with non-constant gene flow always received the majority of the votes. The model with ancestral & recent sympatric migration was chosen as the best model for 78 % of the observed genomic subsamples (Figure 3C). This was followed by the model assuming ancestral migration only for 22 % of observed genomic subsamples. The other three models were not selected as the most likely model.

Ancestral and recent sympatric migration separated by a period of isolation most aptly characterize the history of this plant system (Figure 3A). A history of low but uninterrupted gene flow (Model 5) can be clearly ruled out. We further quantified the goodness-of-fit of the model we identified as the best. The distance between observed and simulated datasets was not significantly different, confirming that the observed data lay well within expectations of the model (Figure 3D). Next, we used this model to estimate the timing, strength and direction of gene flow between species and populations as well as its fluctuation across genome sub-samples.

Figure 3: Choosing the best demographic model.

A) Schematic representation of the tested demographic models. Populations are colored according to the bottom-right code. Arrows represent migration rates. Interspecific migration rates can be

asymmetric (even with double-sided arrows) and intraspecific migration is symmetric. Dashed lines are timepoints at which populations split and/or migration rates change. B) Confusion matrix of the model-choice random forest model. For each model, simulated (out-of-bag) samples were classified by the trained random forest. Correct classifications are on the diagonal. Results are represented as percentages. C) Mean model classification results, i.e. random forest votes, for 100 observed datasets, with 500 randomly selected genomic windows each. Error bars represent the standard deviation. Bars are colored according to the percentage of observed datasets for which the respective model was selected as the best fit. D) Evaluation of fit for the best model. For each of the 100 observed datasets, the median normalized summary statistic distance between observed data and the closest 1% of simulated datasets was calculated (orange). As a null distribution, the same calculation was done with 1000 simulated datasets of pseudo-observed data. For better clarity, the x-axis was trimmed at 300, but no observed distance was larger than that.

Estimation of demographic parameters indicates very low N_e in both taxa

To estimate demographic parameters of the best model, we generated a total of 360,000 coalescent simulations under this demographic model. We used two methods for parameter estimation -- random-forest-based ABC (Marin et al. 2019), which also generated confidence intervals, and extreme gradient boosting (XGBoost; Chen et al. 2020), which provided the most accurate point estimates -- and estimated each parameter independently (see Methods for details). We achieved highest accuracy for the estimation of present population sizes and lowest accuracy for the time and population size at speciation (Figure S6-7). Of the two methods, XGBoost tended to have the lowest RMSE and the highest R^2 , indicating that it could best reconstruct the parameters under which simulated data was generated (Figure S8).

We then estimated the demographic parameters of the two species based on observed data. We present in detail the estimates obtained for one of the observed datasets we randomly picked among those assigned to the selected model. The estimated contemporary population sizes for *A. nemorensis* were 6,775 individuals (sd = 1,988) for the allopatric and 2,964 (sd = 615) for the sympatric population (Figure 4, Table S5). For *A. sagittata*, 14,895 individuals (sd = 2,407) made up the effective allopatric population and 6,425 (sd = 1,649) individuals, the effective sympatric population. These results agree with those of the previous report, namely, that in the

sympatric population, *A. sagittata* harbored levels of genetic diversity higher than those of *A. nemorensis* (Dittberner et al. 2019). That the estimated sizes of the allopatric populations was larger in both species could be because the allopatric samples included individuals collected in several sites. We further estimated that the ancestral population in *A. sagittata* was more than twice as large as that in *A. nemorensis*, with 59,031 (sd = 12,529) and 25,206 (sd = 9,777), respectively. The magnitude of the population size declined after the split in both species, with a mean factor of 6.11 for *A. nemorensis*, which was similar to the factor of 6.57 estimated for *A. sagittata*. A decline of effective population sizes in the recent past has also been observed in the closely related species *Arabidopsis thaliana*, which is also selfing (Durvasula et al. 2017). However, contemporary population sizes in this species are still larger by at least an order of magnitude. This difference indicates that *A. nemorensis* and *A. sagittata* have not been particularly successful in their natural landscape.

The period of isolation dates back to the last glaciation

The split between the species occurred approximately 894,801 (sd = 119,548) generations ago, when the effective size of the ancestral population was approximately 150,281 individuals (sd = 39,960). Standard deviations were high, presumably because our model also allowed for migration between populations (Figure S7). Within species, the populations split almost simultaneously: 10,441 (sd = 3,910) generations ago in *A. nemorensis* and 10,858 (sd = 3,417) generations ago in *A. sagittata*, and their split time coincides with the last glacial maximum. Our analysis thus suggests that the genetic landscape of the two taxa was established after the last glaciation, a period during which they became completely isolated.

Estimates of migration rate confirmed gene flow resumed in sympatry

The low but significant estimates of gene flow showed that the model with ancestral and recent migration explained the data better than a model assuming only ancestral migration (Figure 3). We report the migration rates as the log₁₀-transformed fraction of a population migrating per generation. Migration from *A. nemorensis* to *A. sagittata* in the ancestral population was approximately five times higher than vice

versa: -6.34 (sd = 0.36) and -7.00 (sd = 0.85), respectively. This amounts to about one migrant every 33 generations. Point estimates indicated that, in the sympatric population, migration rates were lower than in the ancestral population. Yet, the rate was still approximately ten times higher from *A. nemorensis* to *A. sagittata* than vice versa, with a rate of -7.72 (sd = 1.39) and -8.74 (sd = 1.33). The mode of the posterior distribution for these two parameters differed even more strongly (-6.36 for migration from *A. nemorensis* to *A. sagittata* and -9.80 for the opposite direction). This result aligns well with the asymmetry of gene flow revealed by the analysis of chloroplast genome variation. As expected estimates of intraspecific migration rate were higher than interspecific migration rates. We noted that the estimate was two orders of magnitude higher in *A. nemorensis* compared to *A. sagittata* (-5.61, sd = 1.61 and -7.09, sd = 1.60 in *A. nemorensis* and *A. sagittata*, respectively).

Figure 4: Model parameter estimation

A) Schematic representation of the model and its parameters: N = effective population size, M = migration rate (fraction of migrants per generation), T = Species/population split times. B) Estimation results of the model parameters based on two methods: ABCRF and XGBoost. Points represent the point estimate for each method, and triangles, the mode of the posterior distribution estimated by ABCRF. Solid error bars represent the standard deviation of the point estimate of ABCRF. Dashed error bars represent the 95% prediction interval of the ABCRF posterior distribution.

Variation in the observed genomic sample affects ancestral population size and sympatric migration rates

Since we observed that the introgression rate varies across the genome, we examined whether estimates of gene flow varied depending on the genomic regions included in the observed sample. To test this, we compared the distributions of random-forest-based and xgboost-based point estimates of all demographic parameters for the 100 observed datasets (Table S6). Estimates were fairly robust to genome subsamples because they were always in the same order of magnitude (Figure 5).

Yet, we made two notable observations. First, the abcrf estimate of sympatric migration from *A. nemorensis* to *A. sagittata* was 4.6 times more variable than the estimate of migration in the opposite direction (Figure 5). The distribution of point estimates for recent sympatric migration from *A. nemorensis* to *A. sagittata* ranged from -8.3 to -6.5, with a mean of -7.5 (Figure S9). Since the estimate was positively correlated with the proportion of introgression regions included in each genomic subsample, it partly reflects the heterogeneity of introgression rates along the genome (Spearman correlation coefficient $Rho=0.2$, $p=0.03$). However, migration rates estimated by gradient boosting, a machine learning method that does not take the variance of the coalescent process into account, appeared more robust to genomic heterogeneity in introgression rate. Second, while both methods indicated the same asymmetry in gene flow between *A. nemorensis* and *A. sagittata* in the sympatric population, they yielded opposite conclusions for ancestral gene flow. Drawing a firm conclusion on the direction of ancestral gene flow is therefore not possible.

Figure 5: Distribution of parameter estimates for the 100 observed datasets formed by sub-sampling 500 75kb-long regions in the genome. M_N and M_S : intraspecific migration rate among *A. nemorensis* and *A. sagittata* populations, respectively. M_{NS} and M_{SN} : interspecific migration rate from *A. nemorensis* into *A. sagittata*, respectively. Sym: sympatric population, Anc.: ancient gene flow. N_{N1} : Effective population size of allopatric *A. nemorensis* populations, N_{N2} : Effective population of sympatric *A. nemorensis* population, N_{S1} : Effective population of sympatric *A. sagittata* population, N_{S2} : Effective population size of allopatric *A. sagittata* population, N_{Nanc} : Ancestral population size of *A. nemorensis* N_{Nanc} , N_{Sanc} : Ancestral population size of *A. sagittata*, T_N : time elapsed since split of *A. nemorensis* population, T_S : Time elapsed since split of *A. sagittata* populations, T_{Anc} : time of speciation, N_{Anc} : ancestral population size prior to speciation.

DISCUSSION

ABC modelling untangles contemporary and ancestral gene flow between two taxa

Explorations of both genome-wide and local estimates of interspecific introgressions, as well as allelic distributions within and between species at introgressed loci, indicated that these two *Arabis* species share a history of gene flow that predates the formation of the sympatric population. The potential of ABC approaches had been revealed previously by the study of population isolation in *Salmo salar* (Rougemont and Bernatchez 2018). Patterns of gene flow in *S. salar* populations, however, were

resolved by genotyping more than 2,000 individuals at some 5,000 SNPs (Rougemont and Bernatchez 2018). This study now shows that ABC approaches have the power to resolve complex demographic histories even with a relatively small sample of genotypes. Indeed, we provide compelling evidence that, although low levels of gene flow were maintained long after speciation, interspecific gene flow was not continuous in the history of these two plant species. Models postulating a phase of complete isolation initiated approximately 10,000 generations ago, were markedly better at explaining the data than those assuming constant gene flow.

We can further confirm that the Rhine population forms a contemporary hotspot of hybridization. The migration rates estimated for past gene flow were low in absolute number but they clearly supported a model in which gene flow has resumed specifically in the sympatric population. This inference aligns well with the observation that, in the sympatric population of *A. sagittata*, introgressed fragments are longer than in the allopatric population. We note that the estimate of gene flow in the sympatric population should be considered as the lower bound since the sample used for ABC demographic inference excluded individuals with obvious signatures of admixture. The low but significant rate of gene flow estimated between the two species in the sympatric populations suggests that gene flow might have happened in the past as it happens today: through the occasional formation of sympatric demes leading to localized bursts of interspecific gene flow.

Interspecific gene flow as a chance for evolutionary rescue?

A. nemorensis and *A. sagittata* are both considered endangered in Germany and in many parts of Europe (Dittberner et al. 2019). The demographic model confirms that effective population sizes are particularly small in these two low-diversity taxa (Figure S2-S5). Genetic diversity in both species was approximately an order of magnitude lower than in more common selfing relatives such as *Arabidopsis thaliana* (Alonso-Blanco et al. 2016) or *Capsella rubella* (Onge et al. 2011). These low levels of diversity, however, do not necessarily point to a low adaptive potential that would increase their risk of extinction. Indeed, the hotspot of contemporary gene flow we uncovered has formed after a period of isolation that might have been sufficiently long to allow

functional divergence. Variance in abcrf estimates associated with the sub-sampling of genomic variation indicates that the signature of contemporary gene flow is not homogeneous throughout the genome, in contrast to that of ancient gene flow. Such a heterogeneity suggests that the contemporary contact is not without fitness consequences, and, indeed, we found several high-frequency introgressions that could be a signature of adaptive gene flow (Racimo et al. 2017). Studies in organisms such as yeast or sunflower have shown that hybrid populations can adapt efficiently to changing environmental conditions (Todesco et al. 2020; Mitchell et al. 2019; Stelkens et al. 2014). Moreover, gene flow can also resolve fitness tradeoffs limiting adaptation in the parental species (Walter et al. 2020).

At this stage, however, it is premature to conclude that introgressions will positively affect the receiving taxon (Suarez-Gonzalez et al. 2018; Shumer et al. 2018). First, we do not know whether the two species possess alleles of potential fitness relevance for each other. Second, F1 hybrids display low fertility, indicating that gene flow causes fertility reduction, at least initially (HD pers. com.; Titz 1979). The simple removal of incompatibility alleles could be sufficient to explain the pattern of heterogeneous introgression documented here (Schumer et al. 2014). Introgressions may also carry deleterious alleles, either due to fitness trade-offs between the parental species (Arnegard et al. 2014) or due to slightly deleterious variants in the parental genomes that have not been purged by selection, i.e., genetic load. This phenomenon is known to have shaped the landscape of introgression from Neanderthals to humans (Harris and Nielsen 2016; Juric et al. 2016), resulting in more introgressions outside of either functionally important regions or of regions with low recombination rates (Schumer, Xu, et al. 2018). Novel methods for the detection of adaptive gene flow have been proposed, but this task remains arduous in selfing species, where recombination events are rare (Setter et al. 2020). Further work is required to identify and disentangle the positive and negative fitness effects of gene flow between *A. sagittata* and *A. nemorensis*. Although Having identified a local hotspot of contemporary hybridization, it is now possible to monitor these consequences in situ.

MATERIAL AND METHODS

Plant material and DNA extraction

In 2016 and 2017, we identified 30 sites reported to host *A. nemorensis* populations on the Deutschland Flora Database. Of these, 24 could be visited, and *A. nemorensis*/*A. sagittata* populations were observed in 11 of them. We sampled seeds from at least 10 *A. nemorensis* and/or *A. sagittata* plants per site for a total of 231 accessions (Table S1). Populations were located in Southern Germany and Austria (Figure 1A). One of these sites, referred to as “Rhine,” was previously described in (Dittberner et al. 2019) and consists of multiple proximate pristine habitat patches at one site. Individuals from these patches were assembled in one sympatric population. We extracted DNA for genotyping as previously described (Dittberner et al. 2019).

RAD-seq genotyping for phylogeography

In addition to the 140 accessions originating from the Rhine populations that were previously genotyped in Dittberner et al. (2019), we genotyped 91 accessions using the original RAD-seq protocol (Etter et al. 2011) with the modification described in (Dittberner et al. 2019). Libraries were sequenced at the Cologne Center for Genomics on three Illumina HiSeq 4000 lanes with 2 × 150bp. We used *FastQC* (Andrews 2010) to check the raw reads. We trimmed adapters and removed reads shorter than 100bp using *Cutadapt* (Martin 2011). We removed PCR duplicates based on a 5 bp stretch of random nucleotides at the end of the adapter, using the *clone_filter* module of *Stacks* version 1.37 (Catchen et al. 2013). We de-multiplexed samples using the *process_radtags* module from *Stacks*. We filtered reads with ambiguous barcodes (allowed distance 2) and cut-sites, reads with uncalled bases and low-quality reads (default threshold).

We used our previously described pipeline to call nuclear genotypes in all samples (Dittberner et al. 2019). Briefly, we used BWA (Li and Durbin 2009) to map reads against the high-quality reference genome we had sequenced and assembled for *A. nemorensis* (Dittberner et al. 2019). We filtered mapped reads using *SAMtools* (Li et al. 2009) to remove regions with excessively low (30%) or high (2-fold) coverage,

compared to the mean coverage. We called genotypes using samtools mpileup and VarScan2 (Koboldt et al. 2012). We filtered genotyped loci using VCFtools (Danecek et al. 2011). Knowing that both species are predominantly selfing, we filtered out SNPs with more than 20% overall heterozygosity and SNPs with more than 75% heterozygosity within a population, because heterozygous SNPs tended to cluster on a few single RAD-seq fragments, indicating inaccuracies in mapping that were not picked up by filtering on coverage. In total, we genotyped 2.6 million sites (~ 1% of the genome) (excluding sites with more than 5% missing data in our sample) and identified 25,634 single nucleotide polymorphisms (SNPs).

To obtain SNPs in the chloroplast sequence, we mapped the RAD-seq reads against a chloroplast reference genome of *Arabis hirsuta* (Kawabe et al. 2018), using the *mem* algorithm of BWA (Li and Durbin 2009) with default settings. We called genotypes and SNPs as described above. We set heterozygous genotype calls to missing data, as these calls most likely resulted from mapping errors (the plastome is effectively haploid). Furthermore, we used VCFtools to remove SNPs with either more than 20% missing data or more than two alleles, which resulted in 23 chloroplast SNPs.

Determining hybrids with RAD-seq and chloroplast data

To determine the genomic make-up of each population and identify the presence of hybrids, we first analyzed SNP data using ADMIXTURE (Durand et al. 2011), varying K from 2 to 10. First we converted VCF files to bed format using PLINK (Purcell et al. 2007; Purcell 2009). We ran ADMIXTURE analysis (Alexander and Lange 2011a) for K=1 to K=6, with 10 iterations of cross-validation each. We normalized clusters across runs using CLUMPAK (Kopelman et al. 2015). We used K=2 for further analysis, as we were analyzing two species and the value was well supported by the cross-validation error (Figure S1). Individuals were defined as hybrids if they had less than 95% ancestry from either species. We created plots using the libraries *ggplot2* (Wickham 2009), *ggmap* (Kahle and Wickham 2013), *scatterpie* (Yu 2018) and *ggsn* (Baquero 2017). Finally, we used the library *pegas* (Paradis et al. 2016b), to determine chloroplast haplotypes and build a haplotype network.

Analysis of genetic diversity

We used the RAD-seq data to determine the level of genetic diversity within and between species. Using the *vcfR* package (Knaus and Grünwald 2017), the genotype data were loaded into R and converted to DNAbin format. We used the *pegas* package (Paradis et al. 2016b) to calculate pairwise genetic distances among all individuals. Based on the resulting distance matrix, we calculated average genetic distances within and between both populations and species. We calculated the diversity in the hybrid complex as the average genetic distance within the whole Rhine population.

Whole genome resequencing for gene flow quantification and ABC

To achieve greater resolution for introgression detection, we randomly selected five *A. nemorensis* and 14 *A. sagittata* accessions from the sympatric population, 4 and 6 accessions from two allopatric *A. sagittata* populations and one accession of each of the six allopatric *A. nemorensis* populations. In total, the whole genome of 35 accessions was sequenced (Table S2). To provide an outgroup species for estimating gene flow, we also sequenced one accession of *Arabis androsacea*, provided by Jean-Gabriel Valay (Jardin Alpin du Lautaret, France). DNA for these accessions was extracted as described above. Libraries were prepared using Illumina TruSeq DNA PCR-free kits at the Cologne Center for Genomics. Six samples were sequenced on a HiSeq 4000 with 50 million 2x75 bp reads per sample. The remaining samples were sequenced on a NovaSeq6000 with 25 million 2x150 bp reads per sample, resulting in a depth of ~ 25x.

We used *FastQC* (Andrews 2010) to quality-check the resulting reads. We filtered the reads using the *process_shortreads* module from *Stacks* v2.2 (Catchen et al. 2013), removing reads shorter than 70 bp or 100 bp (threshold was adapted to the sequencing device) and reads with uncalled bases or low-quality scores (default threshold). At this point, we included reads of the *A. nemorensis* reference genome accession from the sympatric population (Dittberner et al. 2019). We mapped the reads against the *A. nemorensis* reference genome (Dittberner et al. 2019) using the *mem* algorithm of BWA (Li and Durbin 2009) with default settings. We filtered out poorly mapped reads using the following criteria: mapping quality < 30, read not mapped in

proper pairs, > 50% of the read is soft-clipped. We called genotypes as described above but with the default strand-filter of *VarScan2* (Koboldt et al. 2012). Genotypes were filtered using *VCFtools* (Danecek et al. 2011). As in the above, filters were set to allow maximum 20% missing data, maximum two alleles and not more than 20% heterozygosity per site (see above) and to remove indels and low quality genotype calls marked “filtered” by *VarScan2*. Additionally, we removed sites with a mean depth greater than 45 (twice the mean depth of all sites) and smaller than 7.5 (mean depth of all sites divided by 3). From the resulting genotype dataset, which comprised 78,976,767 nucleotide positions, we extracted a total of 2,954,526 SNPs using *VCFtools*. To verify sample identity, we performed a PCA using the *adegenet* package (Jombart et al. 2016). Based on this PCA and the RAD-seq data above, we re-assigned one *A. sagittata* mis-labeled sample (174) from one allopatric population to the other.

Estimation of the population recombination rate

In order to take within-locus recombination into account in our ABC estimations of the demographic history, we estimated the population recombination rate R ; we used the sympatric *A. sagittata* population because we had sequenced the most individuals in this population, which should maximize the accuracy of the estimate. To this end, we first phased the reads using the read-aware phasing algorithm implemented in SHAPEIT (Delaneau et al. 2013). We then used FastEPRR (Gao et al. 2016) to estimate the local recombination rate in non-overlapping 75 kbp windows along the genome, with otherwise default settings. We used the median of the distribution ($R=11.5$) as the per locus population recombination rate (i.e. $R=0.15$ per kb).

Detection of interspecific gene flow

To investigate signatures of gene flow among non-hybrid individuals, we analyzed the whole-genome resequencing (WGS) data using two four-taxon statistics: we calculated D (Durand et al. 2011) over the whole genome to detect interspecific gene flow, and we calculated f_D (Martin et al. 2015) in 50kbp sliding windows to analyze the heterogeneity of gene flow along the chromosomes. These tests assumed the general phylogeny: (P1,P2)(P3) outgroup; where P1 was the control population (no

gene flow), P2 was the gene-flow target population and P3 was the donor population of gene flow. We used *A. androsacea* as the outgroup to determine the ancestral allelic state. We calculated f_D in 50 kbp sliding windows with 25 kbp overlap, following Martin et al. (2015). We skipped sites with derived allele frequency of 0 in P3, as these could bias values of f_D . We calculated these statistics for the phylogenies shown in Figure 2.

To complement the four-taxon analysis, we applied a method based on analyses of genetic distance to locate and assign boundaries to putative introgressed fragments in each accession. Our reasoning was as follows: if there is no introgression, the mean genetic distance of a given accession to members of the same species should be smaller than the distance to members of other species; for introgression, the opposite is true. Thus, in 10 kbp windows along the genome, we calculated the mean intra- and interspecific genetic distance for each accession, excluding individuals from the same population for the intraspecific case, as they are likely to carry the same introgression. We added a constant of 0.0001 to all interspecific distances to avoid zero-division and removed windows with zero interspecific distance and non-extreme intraspecific distance ($< 90^{\text{th}}$ percentile), as these could lead to spurious signals of introgression. For each window, we calculated the ratio of intraspecific and interspecific genetic distance (hereafter, distance ratio), which we used to identify introgressions.

For each accession, we identified introgressions by first removing all windows with a distance ratio smaller than or equal to 1. We then classified windows as introgressions if their distance ratio was > 2 or if both of their neighboring windows fulfilled this condition. Adjacent introgression windows were connected to a single introgression. We applied several filters to the resulting set of candidate introgressions: We excluded candidate introgressions from further analysis if they contained too many repetitive elements (i.e those with mean number of genotyped positions less than the 25% quantile of the number of genotyped position in all introgressions). Furthermore, we overlapped candidate introgressions with D and f_D values (in 50 kbp windows) for the respective target population; for introgressions larger than 50 kbp, we calculated the mean of all windows within the introgression. We kept introgressions with values of $D > 0.1$ and $f_D > 0.01$. Subsequently, we removed introgressions with $> 30\%$ mean heterozygosity, as these often gave inaccurate signals: the expectation for this distance

ratio is not the same as for the homozygous case (detecting heterozygous introgressions with this method would require haplotype data). To define the boundaries of each introgression accurately within each of the remaining candidate introgressions, we recalculated the genetic distances in sliding 2 kbp windows in 200 bp steps. We applied the same thresholds as described above to define our final set of introgression regions in all these windows.

Finally, we aimed to determine the likely origin of each introgression by calculating genetic distances of the introgression region between all accessions of the donor species and the target accession. We then determined the accession(s) with the minimum genetic distance to a given introgression. We standardized genetic distances by dividing them with the approximate number of genotyped sites for each pair of accessions (we used an approximation to reduce computational requirements). The approximate number was estimated by taking the mean of all 2 kbp windows (200 bp steps) within an introgression and scaling this value by the length of the introgression. If all accessions originated from the same population (i.e. sympatric or allopatric for the donor species), we inferred that the introgression likely originated from this population (or a closely related one). If donor accessions originated from multiple populations, the inferred origin was ambiguous.

Statistical analysis was performed in R (R Development Core Team 2008). Visualization was done using the ggplot2 library (Wickham 2009). To account for the introgression frequency in statistical analyses of origin and introgression size, we counted each unique introgression (identical start- and end-points) only once, regardless of its frequency. We tested whether introgression size differed among populations using a Kruskal-Wallis test followed by a pairwise Dunn-Test, implemented in the *FSA* package (Ogle et al. 2020).

Modelling the history of interspecific gene flow

We used coalescent simulations and an ABC algorithm with random forest-based Bayesian parameter inference to determine the most likely demographic history of the two species, focusing especially on the history of interspecific gene flow (Raynal et al. 2019). Modelling was performed in two steps: i) we compared data obtained from

each postulated demographic model to our observed genetic data to identify the best-fitting one and ii) we estimated the posterior distribution for each demographic parameter of the best model.

The summary statistics for the observed data were computed from the 35 fully sequenced genomes. We randomly selected 500 genomic windows of 75 kbp each, all of which were at least 150 kbp apart from one another and contained at least 20,000 genotyped sites, which allowed excluding regions that were excessively repetitive. We computed the following summary statistics using a custom python script: overall θ_W , θ_W and θ_π within each population, Tajima's D for each population, d_{XY} and F_{ST} between all pairs of populations, the fraction of shared, private, invariant and fixed SNPs for all pairs of populations, Patterson's D and f_D (ABBA-BABA) for 8 possible phylogenies. All summary statistics, except Patterson's D, were calculated per locus and the resulting distributions were further summarized as the mean and variance as well as the 5% and 95% quantile, resulting in a total of 236 summary statistics. To estimate the variance in model selection and parameter inference due to window choice, we assembled 100 random sets of these 500 genomic windows and considered each as one independent observation.

Simulated data for each of the five model described in Figure 3A were obtained with the software *escrm* (Sellinger et al. 2020), which is a modification of the widely used coalescent simulation software *scrm* (Staab et al. 2015) that accommodates selfing (we set a 90% selfing rate for all populations). Fixed parameter values and prior distributions for all demographic parameters are found in Table 1. For each model, we simulated 50,000 datasets, each consisting of five hundred 75 kbp-long loci. Summary statistics for the observed data are computed using the same custom python script.

Model choice procedure

In the first step, we specified five different demographic models, all of which differed in their mode of intra- and interspecific gene flow (Figure 3A). In the first model, we assumed no migration at all, either intra- or interspecific. In the second model, interspecific migration stopped 100,000 generations after the species split. In the third

model, ancestral interspecific migration continued until the first intraspecific population-split. The fourth model was an extension of the third model that additionally allowed migration between the sympatric populations, after populations in both species had split. In the fifth model, interspecific migration continued throughout the history of the species, allowing a change in intensity after the intraspecific population had split in both species. All interspecific migration rates were allowed to be asymmetrical, while symmetric intraspecific migration was assumed in all models (except model # 1, which assumed no migration at all). Population sizes were constant for each population but were allowed to change at all population split points (Table 1).

Table 1: Overview of demographic parameters (see Figure 3A) and their prior distributions or fixed values. M_{Sym} and M_{Anc} had the same prior distributions for both directions of migration. 500 loci of length 75bp were simulated assuming a per site mutation rate of $7 \cdot 10^{-9}$ (Ossowski et al. 2010) and a population recombination rate of 0.15 per kb.

Parameter	Minimum	Maximum	Distribution
$N_{N1}, N_{N2}, N_{S1}, N_{S4}$	500	100,000	log-uniform
N_{Nanc}, N_{Sanc}	500	100,000	Uniform
N_{Anc}	50,000	200,000	Uniform
T_N, T_S	0	20,000	Uniform
T_{Anc}	500,000	1,500,000	Uniform
M_{Sym}, M_{Anc}	$1e-11$	$1e-3$	log-uniform
M_N, M_S	$1e-11$	$1e-1$	log-uniform

We used *abcrf*, an R package designed for ABC-based demographic modelling based on random forests, to choose the model that best fit our data and to infer their parameters (Marin et al. 2019; Raynal et al. 2019). We trained a random forest classifier

to distinguish between the five models based on a set of summary statistics, i.e., the model was the dependent variable and the 236 summary statistics were independent variables. We trained the random forest algorithm on 50,000 simulated datasets for each of the five models drawing from the priors in Table 1 and using *abcrf* default parameters, except that parameter “lda” was set to false, disabling linear discriminant analysis. To estimate the classification error for model choice, we used the so-called out-of-bag method implemented in *abcrf*. Due to bootstrap aggregating, each sample was used only to build a subset of trees in the random forest, and the remaining trees could be used to predict the best model for this sample. The results were represented as a confusion matrix showing how frequently the predicted model matches (or does not match) the true model (Figure 3 B). All samples were expected to fall on the diagonal of such a matrix. We then used the trained classifier to assign the most likely demographic model to each of the 100 observed samples (see above). The model that was most frequently selected as the best model for the observed samples was selected as the model fitting our data best.

We further quantified the goodness-of-fit for the best model using a method implemented in the R package *abc* (Katalin et al. 2015), which is based on a rejection algorithm. Briefly, the normalized distance of summary statistics between the observed and all 50,000 simulated datasets was calculated; then the median of the lowest 1% of distances was extracted (the rejection step). The same procedure was performed for 1,000 simulated datasets under this model, drawing parameter values from the prior distributions of distances (so-called pseudo-observed distances). If the model fit is good, the observed distance (1% best-fitting simulations to the observed data) should lie well within the distribution of pseudo-observed distances (1% best simulations fit each of the pseudo-observed dataset). We conducted this analysis for all 100 observed datasets. For each dataset, a p-value was calculated as the fraction of pseudo-observed distances, which are higher than the observed distance.

Estimation of demographic parameters

After identifying the best-fitting demographic model, we estimated the posterior distributions of the different parameters. Demographic parameters with log-uniform

prior distributions were log-transformed prior to training to make them uniform. To reduce computational load, we first selected a subset of the most informative summary statistics for each parameter; we then trained a random forest for each demographic parameter based on 115,000 simulated datasets and extracted the top ten most important summary statistics. This approach reduced the 236 summary statistics to a set of 111 unique summary statistics for training the full models.

The accuracy of random-forest predictions can be greatly increased by tuning training parameters. We tuned the following training parameters by training random forests using all possible combinations: the number of variables to possibly split in each node (mtry; search space: 11, 22, 44, 66, 88); and the minimal node size (search space: 5, 10, 30, 50). This training was conducted for a subset of representative demographic parameters (N_{N1} , N_{Nanc} , T_N , $M_{SN,anc}$, M_S , $M_{SN,sym}$). We plotted the out-of-bag prediction error for each combination of training and demographic parameters (Figure S10). Based on these results, we took mtry=44 as the optimal number of randomly sampled variables, as higher values strongly increased training time while increasing accuracy only minimally. The influence of minimal node size was generally small, so we kept it at the default value of 5.

Finally, we trained a random forest for each parameter on 359,000 simulated datasets, keeping 1,000 pseudo-observed datasets out of the training dataset to use as a test. We quantified the estimation error for each model parameter by computing the parameter estimates for each pseudo-observed dataset and calculating R^2 between true and estimated values. Additionally, we computed the root-mean-square error (RMSE)

per model parameter as: $\sqrt{\frac{\sum_{t=1}^T (\hat{y}_t - y_t)^2}{T}}$, where T is the number of pseudo-observed datasets, y_t is the true value and $\hat{y}_t$ is the estimated value. To make RMSE more comparable among parameters, we further converted RMSE to percentages (%RMSE) by scaling it to the prior range of each parameter (Figure S8).

To calculate parameter estimates for the observed data, we first randomly selected one of the 100 genomic sub-samples (observed samples) that was previously classified as fitting the best model. For this dataset, point estimate, standard deviation, posterior mode and the 5% and 95% quantiles of the posterior distributions were

determined for each demographic parameter. Additionally, we determined point estimates for the remaining 99 observed datasets to assess the variation of demographic parameters depending on the chosen genomic sub-sample.

To complement *abcrf*, which estimates a posterior distribution, we also used *XGBoost* (Chen et al. 2020), a widely used tree-based machine-learning method that gives only a point estimate but often achieves greater accuracy than random forests. We included all summary statistics in *XGBoost* models. We split the 50 000 simulated datasets into a test dataset of 20,000 samples and a training dataset with the remaining 30,000 samples. Tree depth (dp; values: 3, 6, 9) and minimum child weight (values: 5, 10), are the two training parameters that control the complexity of the trees. We tuned these parameters for the same demographic parameters as described above, choosing those that minimized RMSE (Figure S11). The algorithm was stopped if the test statistic did not improve for 100 rounds. For tree depth, we chose six as the optimal value for training models for T_S , T_N , N_{Nanc} and N_{Sanc} , and nine for all other model parameters. We chose 10 as the optimal value for minimum child weight. Using these settings, we trained an *XGBoost* model for each demographic parameter, again stopping after 100 rounds if there was no improvement. We assessed accuracy using the test dataset with 20,000 simulated observations, as described for *abcrf*. We then performed the estimation of the demographic parameters for the 100 genomic sub-samples forming the observed datasets.

ACKNOWLEDGMENTS

This research was funded by the European Research Council (ERC) through the “AdaptoSCOPE” grant 648617, and by the Deutsche Forschungsgemeinschaft through DFG Grant ME2742/13-1. HD acknowledges funding by the International Max Planck Research School. We thank Dr. Markus Stetter and Dr. Gregor Schmitz and Prof. N. Hölzel for helpful discussions. Sequence data are available at ENA under ERS5040446-ERS5040483. Markdown scripts are available as supplementary information.

SUPPLEMENTARY MATERIAL

Suppl_Fig.pdf: Supplementary figures S1-S11 with legends.
RADseq.html: R Markdown of RAD-seq genotyping script and analyses.

Introg_Det_final.html: R Markdown of Introgression detection in whole genome data
 Demography_estimation_final2.html: R markdown for ABC modeling
 Suppl_Tables: Table S1 Description of genotypes included in the RAD-seq analysis,
 Table S2: Description of genotypes chosen for whole-genome sequencing –Table S3:
 list of introgressed fragments identified in the genomes of individuals described in Table
 S2. Table S4: Summary statistics describing the 100 observed datasets. Table S5:
 ABCRF parameter distribution and XGBoost point estimate for the representative
 observed dataset shown in Fig. 4. Table S6: Variation in parameter point estimates for
 the 100 observed datasets.

REFERENCES

- Abbott RJ. 2017. Plant speciation across environmental gradients and the occurrence
 and nature of hybrid zones. *J. Syst. Evol.* 55:238–258.
- Ackermann RR, Arnold ML, Baiz MD, Cahill JA, Cortés-Ortiz L, Evans BJ, Grant BR, Grant
 PR, Hallgrímsson B, Humphreys RA, et al. 2019. Hybridization in human evolution: Insights from
 other organisms. *Evol. Anthropol. Issues News Rev.* 28:189–209.
- Alexander DH, Lange K. 2011a. Enhancements to the ADMIXTURE algorithm for
 individual ancestry estimation. *BMC Bioinformatics* 12:246.
- Alexander DH, Lange K. 2011b. Enhancements to the ADMIXTURE algorithm for
 individual ancestry estimation. *BMC Bioinformatics* 12:246.
- Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, Cao J, Chae
 E, Dezwaan TM, Ding W, et al. 2016. 1,135 Genomes Reveal the Global Pattern of
 Polymorphism in *Arabidopsis thaliana*. *Cell* 166:481–491.
- Anderson JT, Panetta AM, Mitchell-Olds T. 2012. Evolutionary and Ecological Responses
 to Anthropogenic Climate Change: Update on Anthropogenic Climate Change. *Plant Physiol.*
 160:1728–1740.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.
 Available from: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>
- Arnegard ME, McGee MD, Matthews B, Marchinko KB, Conte GL, Kabir S, Bedford N,
 Bergek S, Chan YF, Jones FC, et al. 2014. Genetics of ecological divergence during speciation.
Nature 511:307–311.
- Arnold BJ, Lahner B, DaCosta JM, Weisman CM, Hollister JD, Salt DE, Bomblies K, Yant L.

2016. Borrowed alleles and convergence in serpentine adaptation. *Proc. Natl. Acad. Sci. U. S. A.* 113:8320–8325.

Baduel P, Hunter B, Yeola S, Bomblies K. 2018. Genetic basis and evolution of rapid cycling in railway populations of tetraploid *Arabidopsis arenosa*. *PLOS Genet.* 14:e1007510.

Baquero OS. 2017. ggsn: North Symbols and Scale Bars for Maps Created with “ggplot2” or “ggmap.” Available from: <https://CRAN.R-project.org/package=ggsn>

Bierne N, Gagnaire P-A, David P. 2013. The geography of introgression in a patchy environment and the thorn in the side of ecological speciation. *Curr. Zool.* 59:72–86.

Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. 2014. Speciation and Introgression between *Mimulus nasutus* and *Mimulus guttatus*. *PLOS Genet.* 10:e1004410.

Burmeier S, Eckstein RL, Donath TW, Otte A. 2010. Plant Pattern Development during Early Post-Restoration Succession in Grasslands-A Case Study of *Arabis nemorensis*. *Restor. Ecol.* 19:648–659.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. 2013. Stacks: an analysis tool set for population genomics. *Mol. Ecol.* 22:3124–3140.

Chen T, He T, Benesty M, Khotilovich V, Tang Y, Cho H, Chen K, Mitchell R, Cano I, Zhou T, et al. 2020. xgboost: Extreme Gradient Boosting. Available from: <https://CRAN.R-project.org/package=xgboost>

Chunco AJ. 2014. Hybridization in a warmer world. *Ecol. Evol.* 4:2019–2031.

Csilléry K, Blum MGB, Gaggiotti OE, François O. 2010. Approximate Bayesian Computation (ABC) in practice. *Trends Ecol. Evol.* 25:410–418.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics* 27:2156–2158.

Delaneau O, Howie B, Cox AJ, Zagury J-F, Marchini J. 2013. Haplotype Estimation Using Sequencing Reads. *Am. J. Hum. Genet.* 93:687–696.

Díaz S, Settele J, Brondízio ES, Ngo HT, Agard J, Arneth A, Balvanera P, Brauman KA, Butchart SHM, Chan KMA, et al. 2019. Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science* [Internet] 366. Available from:

965 <https://science.sciencemag.org/content/366/6471/eaax3100>

966 Dittberner H, Becker C, Jiao W-B, Schneeberger K, Hölzel N, Tellier A, Meaux J de. 2019.

967 Strengths and potential pitfalls of hay transfer for ecological restoration revealed by RAD-seq

968 analysis in floodplain Arabis species. *Mol. Ecol.* 28:3887–3901.

969 Durand EY, Patterson N, Reich D, Slatkin M. 2011. Testing for Ancient Admixture

970 between Closely Related Populations. *Mol. Biol. Evol.* 28:2239–2252.

971 Durvasula A, Fulgione A, Gutaker RM, Alacakaptan SI, Flood PJ, Neto C, Tsuchimatsu T,

972 Burbano HA, Picó FX, Alonso-Blanco C, et al. 2017. African genomes illuminate the early history

973 and transition to selfing in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci.* 114:5213–5218.

974 Edelman NB, Frandsen PB, Miyagi M, Clavijo B, Davey J, Dikow RB, García-Accinelli G,

975 Belleghem SMV, Patterson N, Neafsey DE, et al. 2019. Genomic architecture and introgression

976 shape a butterfly radiation. *Science* 366:594–599.

977 Eichenberg D, Bowler DE, Bonn A, Bruelheide H, Grescho V, Harter D, Jandt U, May R,

978 Winter M, Jansen F. 2020. Widespread decline in Central European plant diversity across six

979 decades. *Glob. Change Biol.* [Internet] n/a. Available from:

980 <https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.15447>

981 Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA. 2011. SNP discovery and

982 genotyping for evolutionary genetics using RAD sequencing. *Methods Mol. Biol. Clifton NJ*

983 772:157–178.

984 Gao F, Ming C, Hu W, Li H. 2016. New Software for the Fast Estimation of Population

985 Recombination Rates (FastEPRR) in the Genomic Era. *G3 GenesGenomesGenetics* 6:1563–1571.

986 Goudet J, Jombart T. 2015. hierfstat: Estimation and Tests of Hierarchical F-Statistics.

987 Available from: <https://cran.r-project.org/web/packages/hierfstat/index.html>

988 Goulet BE, Roda F, Hopkins R. 2017. Hybridization in Plants: Old Ideas, New Techniques.

989 *Plant Physiol.* 173:65–78.

990 Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W,

991 Fritz MH-Y, et al. 2010. A Draft Sequence of the Neandertal Genome. *Science* 328:710–722.

992 Hand R, Gregor T. 2006. Die Verbreitung von *Arabis sagittata* In Deutschland. Ergebnisse

993 einer Herbarstudie. *Kochia* 1:21–31.

994 Harris K, Nielsen R. 2016. The Genetic Cost of Neanderthal Introgression. *Genetics*
995 203:881–891.

996 Hedrick PW. 2013. Adaptive introgression in animals: examples and comparison to new
997 mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* 22:4606–4618.

998 Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405:907–913.

999 Hibbins MS, Hahn MW. 2019. The Timing and Direction of Introgression Under the
1000 Multispecies Network Coalescent. *Genetics* [Internet]. Available from:
1001 <https://www.genetics.org/content/early/2019/01/21/genetics.118.301831>

1002 Hölzel N. 2005. Seedling recruitment in flood-meadow species: The effects of gaps, litter
1003 and vegetation matrix. *Appl. Veg. Sci.* 8:115–124.

1004 Hopkins R, Rausher MD. 2012. Pollinator-Mediated Selection on Flower Color Allele
1005 Drives Reinforcement. *Science* 335:1090–1092.

1006 Jay F, Boitard S, Austerlitz F. 2019. An ABC Method for Whole-Genome Sequence Data:
1007 Inferring Paleolithic and Neolithic Human Expansions. *Mol. Biol. Evol.* 36:1565–1579.

1008 Jombart T, Kamvar ZN, Lustrik R, Collins C, Beugin M-P, Knaus B, Solymos P, Schliep K,
1009 Ahmed I, Cori A, et al. 2016. adegenet: Exploratory Analysis of Genetic and Genomic Data.
1010 Available from: <https://cran.r-project.org/web/packages/adegenet/index.html>

1011 Juric I, Aeschbacher S, Coop G. 2016. The Strength of Selection against Neanderthal
1012 Introgression. *PLOS Genet.* 12:e1006340.

1013 Kahle D, Wickham H. 2013. ggmap: Spatial Visualization with ggplot2. *R J.* 5:144–161.

1014 Katalin C, Louisiane L, Olivier F, Michael B. 2015. abc: Tools for Approximate Bayesian
1015 Computation (ABC). Available from: <https://CRAN.R-project.org/package=abc>

1016 Kawabe A, Nukii H, Furihata HY. 2018. Exploring the History of Chloroplast Capture in
1017 Arabis Using Whole Chloroplast Genome Sequencing. *Int. J. Mol. Sci.* [Internet] 19. Available
1018 from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5855824/>

1019 Kenney AM, Sweigart AL. 2016. Reproductive isolation and introgression between
1020 sympatric *Mimulus* species. *Mol. Ecol.* 25:2499–2517.

1021 Knaus BJ, Grünwald NJ. 2017. VCFR: a package to manipulate and visualize variant call
1022 format data in R. *Mol. Ecol. Resour.* 17:44–53.

1023 Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding
1024 L, Wilson RK. 2012. VarScan 2: Somatic mutation and copy number alteration discovery in
1025 cancer by exome sequencing. *Genome Res.* 22:568–576.

1026 Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I. 2015. Clumpak: a
1027 program for identifying clustering modes and packaging population structure inferences across
1028 K. *Mol. Ecol. Resour.* 15:1179–1191.

1029 Kryvokhyzha D, Salcedo A, Eriksson MC, Duan T, Tawari N, Chen J, Guerrina M, Kreiner
1030 JM, Kent TV, Lagercrantz U, et al. 2019. Parental legacy, demography, and admixture influenced
1031 the evolution of the two subgenomes of the tetraploid *Capsella bursa-pastoris* (Brassicaceae).
1032 *PLoS Genet.* 15:e1007949.

1033 Lenormand T. 2002. Gene flow and the limits to natural selection. *Trends Ecol. Evol.*
1034 17:183–189.

1035 Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler
1036 transform. *Bioinformatics* 25:1754–1760.

1037 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
1038 R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map
1039 format and SAMtools. *Bioinforma. Oxf. Engl.* 25:2078–2079.

1040 Ma Y, Wang J, Hu Q, Li J, Sun Y, Zhang L, Abbott RJ, Liu J, Mao K. 2019. Ancient
1041 introgression drives adaptation to cooler and drier mountain habitats in a cypress species
1042 complex. *Commun. Biol.* 2:1–12.

1043 Marburger S, Monnahan P, Seear PJ, Martin SH, Koch J, Paajanen P, Bohutínská M,
1044 Higgins JD, Schmickl R, Yant L. 2019. Interspecific introgression mediates adaptation to whole
1045 genome duplication. *Nat. Commun.* 10:5218.

1046 Marcet-Houben M, Gabaldón T. 2015. Beyond the Whole-Genome Duplication:
1047 Phylogenetic Evidence for an Ancient Interspecies Hybridization in the Baker’s Yeast Lineage.
1048 *PLOS Biol.* 13:e1002220.

1049 Marin J-M, Raynal L, Pudlo P, Robert CP, Estoup A. 2019. abcrf: Approximate Bayesian
1050 Computation via Random Forests. Available from: <https://CRAN.R-project.org/package=abcrf>

1051 Martin M. 2011. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal* 17:10–12.

Martin SH, Davey JW, Jiggins CD. 2015. Evaluating the use of ABBA-BABA statistics to locate introgressed loci. *Mol. Biol. Evol.* 32:244–257.

Martin SH, Jiggins CD. 2017. Interpreting the genomic landscape of introgression. *Curr. Opin. Genet. Dev.* 47:69–74.

Mathar W, Kleinebecker T, Hölzel N. 2015. Environmental variation as a key process of co-existence in flood-meadows. Collins B, editor. *J. Veg. Sci.* 26:480–491.

Mattila TM, Tyrmi J, Pyhäjärvi T, Savolainen O. 2017. Genome-Wide Analysis of Colonization History and Concomitant Selection in *Arabidopsis lyrata*. *Mol. Biol. Evol.* 34:2665–2677.

McCauley DE. 1995. The use of chloroplast DNA polymorphism in studies of gene flow in plants. *Trends Ecol. Evol.* 10:198–202.

Mitchell N, Owens GL, Hovick SM, Rieseberg LH, Whitney KD. 2019. Hybridization speeds adaptive evolution in an eight-year field experiment. *Sci. Rep.* 9:6746.

Novikova PY, Hohmann N, Nizhynska V, Tsuchimatsu T, Ali J, Muir G, Guggisberg A, Paape T, Schmid K, Fedorenko OM, et al. 2016. Sequencing of the genus *Arabidopsis* identifies a complex history of nonbifurcating speciation and abundant trans-specific polymorphism. *Nat. Genet.* 48:1077–1082.

Novotná I, Czapik R. 1974. Studies on the Progenies of Hybrids from the *Arabis hirsuta* Complex. *Folia Geobot. Phytotaxon.* 9:341–357.

Ogle D, Wheeler P, Dinno A. 2020. FSA: Simple Fisheries Stock Assessment Methods. Available from: <https://CRAN.R-project.org/package=FSA>

Onge KRS, Källman T, Slotte T, Lascoux M, Palmé AE. 2011. Contrasting demographic history and population structure in *Capsella rubella* and *Capsella grandiflora*, two closely related species with different mating systems. *Mol. Ecol.* 20:3306–3320.

Ossowski S, Schneeberger K, Lucas-Lledó JI, et al. The rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*. *Science*. 2010;327(5961):92–94.

Paape T, Briskine RV, Halstead-Nussloch G, Lischer HEL, Shimizu-Inatsugi R, Hatakeyama M, Tanaka K, Nishiyama T, Sabirov R, Sese J, et al. 2018. Patterns of polymorphism and selection

in the subgenomes of the allopolyploid *Arabidopsis kamchatica*. *Nat. Commun.* 9:3909.

Paradis E, Jombart T, Schliep K, Potts A, Winter D. 2016a. pegas: Population and Evolutionary Genetics Analysis System. Available from: <https://cran.r-project.org/web/packages/pegas/index.html>

Paradis E, Jombart T, Schliep K, Potts A, Winter D. 2016b. pegas: Population and Evolutionary Genetics Analysis System. Available from: <https://cran.r-project.org/web/packages/pegas/index.html>

Purcell S. 2009. PLINK.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW, Daly MJ, et al. 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

R Development Core Team. 2008. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing Available from: <http://www.R-project.org>

Racimo F, Marnetto D, Huerta-Sánchez E. 2017. Signatures of Archaic Adaptive Introgression in Present-Day Human Populations. *Mol. Biol. Evol.* 34:296–317.

Ravinet M, Faria R, Butlin RK, Galindo J, Bierne N, Rafajlović M, Noor M a. F, Mehlig B, Westram AM. 2017. Interpreting the genomic landscape of speciation: a road map for finding barriers to gene flow. *J. Evol. Biol.* 30:1450–1477.

Raynal L, Marin J-M, Pudlo P, Ribatet M, Robert CP, Estoup A. 2019. ABC random forests for Bayesian parameter inference. *Bioinformatics* 35:1720–1728.

Rieseberg LH, Archer MA, Wayne RK. 1999. Transgressive segregation, adaptation and speciation. *Heredity* 83 (Pt 4):363–372.

Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, Durphy JL, Schwarzbach AE, Donovan LA, Lexer C. 2003. Major Ecological Transitions in Wild Sunflowers Facilitated by Hybridization. *Science* 301:1211–1216.

Rougemont Q, Bernatchez L. 2018. The demographic history of Atlantic salmon (*Salmo salar*) across its distribution range reconstructed from approximate Bayesian computations*: ATLANTIC SALMON HISTORY AND LINKED SELECTION. *Evolution* 72:1261–1277.

1110 Schumer M, Cui R, Powell DL, Dresner R, Rosenthal GG, Andolfatto P. 2014. High-
1111 resolution mapping reveals hundreds of genetic incompatibilities in hybridizing fish
1112 species. McVean G, editor. *eLife* 3:e02535.

1113 Schumer M, Rosenthal GG, Andolfatto P. 2018. What do we mean when we talk about
1114 hybrid speciation? *Heredity* 120:379–382.

1115 Schumer M, Xu C, Powell DL, Durvasula A, Skov L, Holland C, Blazier JC, Sankararaman S,
1116 Andolfatto P, Rosenthal GG, et al. 2018. Natural selection interacts with recombination to
1117 shape the evolution of hybrid genomes. *Science* 360:656–660.

1118 Seehausen O. 2004. Hybridization and adaptive radiation. *Trends Ecol. Evol.* 19:198–207.

1119 Sellinger TPP, Awad DA, Moest M, Tellier A. 2020. Inference of past demography,
1120 dormancy and self-fertilization rates from whole genome sequence data. *PLOS Genet.*
1121 16:e1008698.

1122 Setter D, Mousset S, Cheng X, Nielsen R, DeGiorgio M, Hermisson J. 2020.
1123 VolcanoFinder: Genomic scans for adaptive introgression. *PLOS Genetics* 16(6): e1008867

1124 Small ST, Labbé F, Lobo NF, Koekemoer LL, Sikaala CH, Neafsey DE, Hahn MW, Fontaine
1125 MC, Besansky NJ. 2020. Radiation with reticulation marks the origin of a major malaria vector.
1126 *Proc. Natl. Acad. Sci.* 117:31583–31590.

1127 Staab PR, Zhu S, Metzler D, Lunter G. 2015. scrm: efficiently simulating long sequences
1128 using the approximated coalescent with recombination. *Bioinformatics* 31:1680–1682.

1129 Steinrücken M, Spence JP, Kamm JA, Wieczorek E, Song YS. 2018. Model-based
1130 detection and analysis of introgressed Neanderthal ancestry in modern humans. *Mol. Ecol.*
1131 27:3873–3888.

1132 Stelkens RB, Brockhurst MA, Hurst GDD, Greig D. 2014. Hybridization facilitates
1133 evolutionary rescue. *Evol. Appl.* 7:1209–1217.

1134 Suarez-Gonzalez A, Lexer C, Cronk QCB. 2018a. Adaptive introgression: a plant
1135 perspective. *Biol. Lett.* 14:20170688.

1136 Suarez-Gonzalez A, Lexer C, Cronk QCB. 2018b. Adaptive introgression: a plant
1137 perspective. *Biol. Lett.* 14:20170688.

1138 Takou M, Härmälä T, Steige KA, Koch E, Dittberner H, Yant L, Genete M, Sunyaev S,

Castric V, Vekemans X, et al. 2019. Maintenance of adaptive dynamics in a bottlenecked range-edge population that retained outcrossing. *bioRxiv*:709873.

Taylor SA, Larson EL. 2019. Insights from genomes into the evolutionary importance and prevalence of hybridization in nature. *Nat. Ecol. Evol.* 3:170–177.

Tigano A, Friesen VL. 2016. Genomics of local adaptation with gene flow. *Mol. Ecol.* 25:2144–2164.

Titz W. 1979. Die Interfertilitätsbeziehungen europäischer Sippen der *Arabis hirsuta*-Gruppe (Brassicaceae). *Plant Syst. Evol.* 131:291–310.

Todesco M, Owens GL, Bercovich N, Légaré J-S, Soudi S, Burge DO, Huang K, Ostevik KL, Drummond EBM, Imerovski I, et al. 2020. Massive haplotypes underlie ecotypic differentiation in sunflowers. *Nature* 584:602–607.

Vallejo-Marín M, Hiscock SJ. 2018. Hybridization and hybrid speciation under global change. *New Phytol.* 211:1170–1187.

Walter GM, Richards TJ, Wilkinson MJ, Blows MW, Aguirre JD, Ortiz-Barrientos D. 2020. Loss of ecologically important genetic variation in late generation hybrids reveals links between adaptation and speciation. *Evol. Lett.* 4:302–316.

Wickham H. 2009. Ggplot2: elegant graphics for data analysis. New York: Springer

Yeaman S. 2015. Local Adaptation by Alleles of Small Effect*. *Am. Nat.* 186:S74–S89.

Yu G. 2018. scatterpie: Scatter Pie Plot. Available from: <https://CRAN.R-project.org/package=scatterpie>