Circulating miRNAs associated with bone mineral density in healthy adult baboons.

Ellen. E. Quillen¹, Jaydee Foster², Anne Sheldrake², Maggie Stainback¹, Todd L. Bredbenner³

1. Department of Internal Medicine, Section of Molecular Medicine, Wake Forest School of Medicine

2. Department of Genetics, Texas Biomedical Research Institute

3. Department of 3 Mechanical and Aerospace Engineering, University of Colorado Colorado Springs

Abstract

MicroRNAs (miRNAs) regulate gene expression post-transcriptionally and circulate in the blood, making them attractive biomarkers of disease state for tissues like bone that are challenging to interrogate directly. Here we report on five miRNAs – miR-197-3p, miR-320a, miR-320b, miR-331-5p, and miR-423-5p – that are associated with bone mineral density (BMD) in 147 healthy adult baboons. These baboons range in age from 15 to 25 years (45 to 75 human equivalent years) and were 65% female with a broad range of BMDs including a minority of osteopenic individuals. miRNAs were generated via RNA sequencing from buffy coats collected at necropsy and areal BMD evaluated via DXA of the lumbar vertebrae post-mortem. Differential expression analysis controlled for the underlying pedigree structure of these animals to account for genetic variation which may be driving miRNA abundance and BMD values. While many of these miRNAs have been associated with risk of human osteoporosis, this finding is of interest because the cohort represent a model of normal aging and bone metabolism rather than a disease cohort. The replication of miRNA associations with osteoporosis or other bone metabolic disorders in animals with healthy BMD suggests an overlap in normal variation and disease states. We suggest that these miRNAs are involved in the regulation of cellular proliferation, apoptosis, and protein composition in the extracellular matrix throughout life. However, age-related dysregulation of these systems may lead to disease causing associations of the miRNAs among individuals with clinically defined disease.

Keywords: bone mineral density, miRNA, non-human primate

Research interest in microRNAs (miRNAs) has exploded in the past decade with recognition of the potential for these small circulating RNAs to act as biomarkers of disease state or regulators of bone across cell types. As post-transcriptional regulators, these small non-coding RNAs are uniquely able to regulate gene expression in the cytoplasm and may travel among cells to do so. While a number of studies have focused on the potential of miRNAs as biomarkers for osteoporosis and fracture risk, little is known about the miRNAs in the context of normal variation in spinal bone mineral density (BMD) among healthy adults. Here we report on five circulating miRNAs that are associated with variation in BMD within the healthy range for middle-aged and older baboons.

The baboon is unique model for age-related bone loss, as it is closely related phylogenetically to humans, is relatively large bodied, and exhibits intracortical remodeling throughout life, unlike rodent models of skeletal aging. The adult bone remodeling and skeletal fracture properties of baboons are also more similar to humans than are other mid- to large-sized mammals (Wang, Mabrey, and Agrawal 1998; Brommage 2001). Like humans, baboons undergo postmenopausal bone loss and sex and age influence BMD (L Havill et al. 2003) leading to osteopenia in approximately 25% of older females (L. M. Havill et al. 2008). Furthermore, biomechanical properties directly relevant to fracture, including vertebral trabecular bone mechanical properties (LM Havill, Allen, and Bredbenner 2010), femoral cortical bone microstructure (L. Havill et al. 2008), and femoral bone shape (Hansen et al. 2009), are strongly heritable in the baboon, making pedigreed animals ideal for assessing the effects of genetic and non-genetic factors bone biomarkers to bone fragility (Cox et al. 2013).

We leveraged the unique anatomical and genetic features of the baboon to identify circulating miRNAs associated with BMD across a broad range of mildly osteopenic and healthy adult baboons while controlling for underlying genetic factors.

Methods

Study Population

We generated miRNA and BMD data for 147 middle-aged and older baboons (hybrid *Papio hamadryas* species) drawn from a larger pedigree of baboons housed at the Southwest National Primate Center and Texas Biomedical Research Institute (Vagtborg 1973). Animals were 65% female and ranged in age from 15 to 25 years of age which is the equivalent of approximately 45 to 75 human years (Figure 1).

Figure 1. Distribution of animals by age and sex in sample.

During life, all animals were housed outdoors in large social group cages and maintained on commercial monkey chow to which they had ad libitum access. Animals with medical conditions known to influence bone metabolism (e.g. diabetes, chronic renal disease) or a history of traumatic fracture were excluded. No animals were sacrificed for this study, all were euthanized for other reasons and monitored by the Institutional Animal Care and Use Committee. Blood samples were drawn in EDTA tubes at necropsy and processed buffy coat stored at -80°C until RNA was extracted. Lumbar spines were stored at -20°C.

DXA

Dual-energy x-ray absorptiometry (DXA) scans were performed post-mortem on thawed lumbar vertebrae using a Lunar DPX 6529 (General Electric) and areal BMD analyzed with the manufacturer's software for adults. All analyzed values are for DXA measured for the anterior-posterior (AP) axis of L4-L2 vertebrae. Baboons exhibit high levels of sexual dimorphism in weight and body size leading to significant differences in mean BMD (0.22 g/cm^2 , $p = 1 \times 10^{-6}$, Figure 2). The distribution of BMD in these animals is consistent with previously published reference standards from more than 650 baboons (L Havill et al. 2003). Within the 10-year age range of our study, there is no significant decline in BMD. Nevertheless, the distribution of BMD across the sample is broad ranging from 4.3 standard deviations below to 8.4 standard deviations above the healthy, young adult mean for females and -3.5 to 9.3 standard deviation for males. To account for sexual dimorphism in BMD, we performed z-scoring on male and female BMD values separately and combined the scaled values for analysis.

Figure 2. Distribution of BMD in study sample by age and sex.

miRNA Sequencing and Analysis

Total RNA was isolated from buffy coat using TRIzol (Invitrogen) and the Qiagen miRNeasy Mini Kit as described previously (Spradling et al. 2013). RNA quality was assessed using an Agilent Bioanalyzer 2100 and RNA was enriched for small non-coding RNAs (sncRNAs) by using the Ambion mirVana miRNA Isolation Kit. Complementary DNA (cDNA) libraries were generated with the Illumina Small RNA Prep Kit v1.5 following the manufacturer's protocol and sequenced in Illumina's Genome Analyzer (GAIIx) (Karere et al. 2012). mirDeep2 (Mackowiak 2011) was used to align reads to the known human miRbase version 21 (Griffiths-Jones et al. 2008, 2006) mature and hairpin miRNAs and to identify novel miRNAs.

Raw miRNA counts were analyzed using the R statistical package DESeq2 to identify individual miRNAs associated with BMD (Love, Huber, and Anders 2014; R Core Team 2019). DESeq2 has the advantage of incorporating shrinkage estimators for dispersion and fold change which better handles low-abundance miRNAs compared to other methods. After z-scoring, weight and age were no longer associated with BMD and so were not included as covariates. miRNAs significantly associated with BMD at a FDR < 0.05 were further tested to determine if this association was driven by underlying genetic variation in the pedigree. A likelihood ratio test was calculated to compare linear mixed effects models fit with lme4 in the coxme R package (Therneau 2020) with and without miRNA abundance as a fixed effect while including pedigree-based kinship as a random effect.

miRNA Target Prediction

A major challenge in understanding the potential functional role of miRNAs is that most are computationally predicted to bind dozens of different mRNAs. To better predict what role

these circulating miRNAs could play in the bone, we used the R package multiMiR (Ru et al. 2014) to query three experimentally validated miRNA target databases – miRecords (F. Xiao et al. 2009), miRTarBase (Hsu et al. 2011), and TarBase (Karagkouni et al. 2018). To be considered a potential target we required experimental validation of miRNA-mRNA interaction via luciferase, qRT-PCR, or Western blot experiments as well as the presence of the target protein in baboon bone (unpublished mass spectrometry data).

Results

miRNAs Associations

After FDR correction, 15 miRNAs were significantly associated with BMD in the DESeq2 analysis of which 5 were associated (likelihood ratio test $p < 0.1$) after correcting for the genetic relatedness of the animals (Table 1). All five of these highly associated miRNAs – miR-197-3p, miR-320a, miR-320b, miR-331-5p, and miR-423-5p – are negatively correlated with BMD (Figure 3) suggesting that increased levels of these miRNAs – which would be expected to decrease mRNA expression – are indicative of a decline in BMD. The heat map in Figure 4 illustrates a high level of heterogeneity in miRNA abundance across animals with varying BMD.

Table 2. miRNAs significantly associated with BMD after FDR correction in DESeq2 analysis. Bolded miRNAs remained significantly associated after correction for pedigree structure.

	β	Mean Expression	p	$LRT\ p$
hsa-miR-423-5p	-0.28	886	2.4×10^{-4}	0.03
hsa-miR-320a	-0.22	2,561	3.9×10^{-3}	0.04
hsa-miR-339-5p	-0.19	1,697	6.2×10^{-3}	0.22
hsa-miR-3173-5p	-0.38	16	3.0×10^{-2}	0.14
hsa-miR-371a-5p	0.6	7	3.0×10^{-2}	0.40
hsa-miR-16-2-3p	-0.44	137	3.3×10^{-2}	0.27
hsa-miR-197-3p	-0.18	577	3.5×10^{-2}	0.09
hsa-miR-21-5p	0.16	30,037	3.5×10^{-2}	0.68
hsa-miR-331-5p	-0.09	808	3.5×10^{-2}	0.07
hsa-miR-374a-3p	0.26	231	3.5×10^{-2}	0.88
hsa-miR-374a-5p	0.19	4,255	3.5×10^{-2}	0.92
hsa-miR-424-3p	-0.16	86	3.5×10^{-2}	0.10
hsa-miR-484	-0.13	2,334	3.5×10^{-2}	0.19
hsa-miR-320b	-0.21	28	3.6×10^{-2}	0.05
hsa-miR-574-3p	-0.16	94	4.1×10^{-2}	0.25

127

128 **Figure 3.** miRNA abundance versus BMD for significantly associated miRNAs.

Figure 4. Heatmap of all miRNAs with nominal $p < 0.05$ in DeSeq2. Columns are color-coded by z-scored BMD tertile with purple indicating the lowest tertile and teal the highest.

Putative miRNA targets

Based on our screening criteria, we identified 60 genes present in bone and experimentally validated as potential targets of our five significant miRNAs (Supplementary Table 1). Analysis in DAVID (Huang, Sherman, and Lempicki 2009) identified several significantly enriched functional annotations within the 60 genes. The top five enriched gene ontology terms representing more than 10% of the genes on the list are myelin sheath (26.4 fold enrichment, FDR = 6.1×10^{-10}), focal adhesion (11.2 fold enrichment, FDR = 2.9×10^{-7}), extracellular matrix (11.1 fold enrichment, FDR = 3.6×10^{-5}), cadherin binding involved in cell-cell adhesion (9.5 fold enrichment, FDR = 9.0×10^{-4}), cell-cell adherens junction (9.0 fold enrichment, FDR = 5.4×10^{-4}). Additionally, these putative targets include seven genes previously linked to osteoporosis in genome wide association studies based on data drawn from the Public Health Genomics and Precision Health Knowledge Base (v7.1) (Yu et al. 2010) (Table 2).

Table 2. miRNA targets linked to osteoporosis in genome wide association studies

GENE NAME	MIRNA
FGB	hsa-miR-197-3p
IGFBP5	hsa-miR-197-3p
SOD1	hsa-miR-197-3p
SOD2	hsa-miR-197-3p
	hsa-miR-331-5p
GAPDH	hsa-miR-320a
	hsa-miR-423-5p
MIF	hsa-miR-320a
MMP9	hsa-miR-320a

Discussion

Most previous research on bone density and miRNAs has focused on patients with osteoporosis or other disorders of bone metabolism. This is reflected in links between osteoporosis or fracture risk in human patients and the miRNAs we have identified, many of which have been suggested as potential clinical biomarkers.

Of the miRNAs we identified, literature linking the miRNA-320 family to bone fragility is the most robust. miR-320b is more common in women with a recent fracture compared to age-matched controls (Weilner et al. 2015). This may be due to its role in osteoblast differentiation. miR-320b over-expression prevents osteoblast differentiation, while inhibition promotes bone matrix mineralization and differentiation via BMP-2 (Laxman et al., 2017). Additionally,

researchers studying low-traumatic fractures in premenopausal, postmenopausal, and idiopathic osteoporosis patients found miR-320a was upregulated in these patients compared to controls (Kocijan et al., 2016). Prior analysis of trabecular bone tissue showed significant dysregulation of miR-320a in patients with osteoporosis, possibly via regulation of osteogenesis by targeting bone-forming genes such as *CTNNB1* (B-catenin) and *RUNX2* (De-Ugarte et al., 2015). These findings led to the inclusion of miR-320a on the commercially available OsteomiR panel (Tamirna).

Similarly, in a study of plasma miRNAs associated with fracture risk, miR-423-5p expression was significantly negatively associated with FRAX score, but not BMD, in osteoporosis patients (Bedene et al. 2016). This finding was reinforced by research on facial bone atrophy where miR-148-3p was shown to promote bone proliferation via prevention of apoptosis in mesenchymal stem cells (Yang et al. 2018). While there is little additional evidence of a role for this miRNA in bone, it has also been linked to regulation of apoptosis in cardiomyocytes (Zhu and Lu 2019), kidney cells (Yuan et al. 2017), retinal pigment epithelial cells (Q. Xiao et al. 2019), and colon cancer cells (Jia et al. 2018).

Circulating miR-331-5p levels have been identified as a potential biomarker for osteoporosis and subsequent bone fracture (Liu et al. 2015), although there is little known about the role of this miRNA in bone. Hints to its function come from studies of vascular smooth muscle cells, where miR-331-5p is induced by BMP2-PPAR γ signaling to regulate cellular proliferation (Calvier et al. 2017). We predict a role for miR-331-5p in regulating *SOD2* expression which directly induces a BMP2 response under hypoxic conditions (Kamiya et al. 2013).

While miR-197-3p has not previously been linked to adult bone metabolism, it was identified in the downregulation of osteogenesis in human amniotic membrane-derived mesenchymal stem cells due to its suppression of SMAD2 in the TGF- β pathway during osteoblast differentiation (Avendaño-Félix et al. 2019). We predict that miR-197-3p may target expression of both *SOD1* and *SOD2*, antioxidative enzymes that respond to vitamin D levels (Lisse 2020) and are thought to play a critical role in the regulation of cellular senescence (Zhang et al. 2017; Jeong and Cho 2015).

Taken together, our BMD-associated miRNAs and their putative targets point towards regulation of extracellular matrix proteins, apoptosis, and cell proliferation – three key components in the maintenance of bone homeostasis. Our findings demonstrate the overlap in

miRNAs associated with bone mineral density and related traits in humans and nonhuman primates, highlighting the utility of this model for understanding aging bone. The association of these miRNAs with variation in bone mineral density within the healthy, non-osteoporotic, range suggests they may be useful in identifying the earliest stages of metabolic shifts in bone. While the impact of epigenetic regulation of gene expression is evident in bone tissue, the use of miRNAs as biomarkers for low BMD and high fracture risk is still relatively new. It is apparent more research is needed to better understand these molecular pathways. Future work should focus on identifying the presence and functional role of these miRNAs in bone tissue to solidify their promise as biomarkers.

Acknowledgements

We thank Lorena Havill for her role in the development of this project. This work was funded by R01 AR064244 to TLB and K01 AG056663 to EEQ. The animals in the SNPRC pedigree were supported by grant P51 OD011133.

- Avendaño-Félix, Mariana, Lizeth Fuentes-Mera, Rosalío Ramos-Payan, Maribel Aguilar-Medina, Vanessa Pérez-Silos, Nidia Moncada-Saucedo, Laurence A. Marchat, et al. 2019. "A Novel OsteomiRs Expression Signature for Osteoblast Differentiation of Human Amniotic Membrane-Derived Mesenchymal Stem Cells." *BioMed Research International* 2019 (March): 1–13. <https://doi.org/10.1155/2019/8987268>.
- Bedene, Ajda, Simona Mencej Bedrač, Lea Ješe, Janja Marc, Peter Vrtačnik, Janez Preželj, Tomaž Kocjan, Tilen Kranjc, and Barbara Ostanek. 2016. "MiR-148a the Epigenetic Regulator of Bone Homeostasis Is Increased in Plasma of Osteoporotic Postmenopausal Women." *Wiener Klinische Wochenschrift* 128: 519–26. <https://doi.org/10.1007/s00508-016-1141-3>.
- Brommage, R. 2001. "Perspectives on Using Nonhuman Primates to Understand the Etiology and Treatment of Postmenopausal Osteoporosis." *Journal of Musculoskeletal & Neuronal Interactions* 1 (4): 307–25.
- Calvier, Laurent, Philippe Chouvarine, Ekaterina Legchenko, Nadine Hoffmann, Jonas Geldner, Paul Borchert, Danny Jonigk, Miklos M. Mozes, and Georg Hansmann. 2017. "PPARγ Links BMP2 and TGFβ1 Pathways in Vascular Smooth Muscle Cells, Regulating Cell Proliferation and Glucose Metabolism." *Cell Metabolism* 25 (5): 1118-1134.e7. <https://doi.org/10.1016/j.cmet.2017.03.011>.
- Cox, Laura a, Anthony G Comuzzie, Lorena M Havill, Genesio M Karere, Kimberly D Spradling, Michael C Mahaney, Peter W Nathanielsz, et al. 2013. "Baboons as a Model to Study Genetics and Epigenetics of Human Disease." *ILAR Journal / National Research Council, Institute of Laboratory Animal Resources* 54 (2): 106–21. <https://doi.org/10.1093/ilar/ilt038>.
- Griffiths-Jones, Sam, Russell J Grocock, Stijn van Dongen, Alex Bateman, and Anton J Enright. 2006. "MiRBase: MicroRNA Sequences, Targets and Gene Nomenclature." *Nucleic Acids Research* 34 (Database issue): D140-4. <https://doi.org/10.1093/nar/gkj112>.
- Griffiths-Jones, Sam, Harpreet Kaur Saini, Stijn van Dongen, and Anton J Enright. 2008. "MiRBase: Tools for MicroRNA Genomics." *Nucleic Acids Research* 36 (Database issue): D154-8. <https://doi.org/10.1093/nar/gkm952>.
- Hansen, Heather L, Todd L Bredbenner, Daniel P Nicolella, Michael C Mahaney, and Lorena M Havill. 2009. "Cross-Sectional Geometry of the Femoral Midshaft in Baboons Is Heritable." *Bone* 45 (5): 892–97. <https://doi.org/10.1016/j.bone.2009.05.028>.

235 Havill, L.M., S.M. Levine, D.E. Newman, and M.C. Mahaney. 2008. "Osteopenia and
236 Osteoporosis in Adult Baboons (*Papio Hamadryas*).
237 146–53. <https://doi.org/10.1111/j.1600-0684.2007.00270.x>.

238 Havill, L.M, JAK Harris, S.M. Levine, and Michael C Mahaney. 2008. "Strong and Significant
239 Genetic Effects on Osteonal Remodeling-Associated Cortical Bone Microstructure in
240 Pedigreed Baboons." *Bone* 42 (Suppl 1): S53-54.

241 Havill, L, Michael C. Mahaney, Stefan a Czerwinski, K D Carey, K Rice, and Jeffrey Rogers.
242 2003. "Bone Mineral Density Reference Standards in Adult Baboons (*Papio Hamadryas*)
243 by Sex and Age." *Bone* 33 (6): 877–88. [https://doi.org/10.1016/S8756-3282\(03\)00231-X](https://doi.org/10.1016/S8756-3282(03)00231-X).

244 Havill, LM, MR Allen, and TL Bredbenner. 2010. "Heritability of Lumbar Trabecular Bone
245 Mechanical Properties in Baboons." *Bone* 46 (3): 835–40.
246 <https://doi.org/10.1016/j.bone.2009.11.002.Heritability>.

247 Hsu, Sheng-Da, Feng-Mao Lin, Wei-Yun Wu, Chao Liang, Wei-Chih Huang, Wen-Ling Chan,
248 Wen-Ting Tsai, et al. 2011. "MiRTarBase: A Database Curates Experimentally Validated
249 MicroRNA-Target Interactions." *Nucleic Acids Research* 39 (Database issue): D163-9.
250 <https://doi.org/10.1093/nar/gkq1107>.

251 Huang, Da Wei, Brad T Sherman, and Richard a Lempicki. 2009. "Bioinformatics Enrichment
252 Tools: Paths toward the Comprehensive Functional Analysis of Large Gene Lists." *Nucleic
253 Acids Research* 37 (1): 1–13. <https://doi.org/10.1093/nar/gkn923>.

254 Jeong, Sin-Gu, and Goang-Won Cho. 2015. "Endogenous ROS Levels Are Increased in
255 Replicative Senescence in Human Bone Marrow Mesenchymal Stromal Cells."
256 *Biochemical and Biophysical Research Communications* 460 (4): 971–76.
257 <https://doi.org/10.1016/j.bbrc.2015.03.136>.

258 Jia, Wenzhuo, Tao Yu, Qi An, Xianglong Cao, and Hongda Pan. 2018. "MicroRNA-423-5p
259 Inhibits Colon Cancer Growth by Promoting Caspase-dependent Apoptosis." *Experimental
260 and Therapeutic Medicine*, June. <https://doi.org/10.3892/etm.2018.6288>.

261 Kamiya, Nobuhiro, Sasha Shafer, Ila Oxendine, Douglas P. Mortlock, Ronald L. Chandler, Leif
262 Oxburgh, and Harry K.W. Kim. 2013. "Acute BMP2 Upregulation Following Induction of
263 Ischemic Osteonecrosis in Immature Femoral Head." *Bone* 53 (1): 239–47.
264 <https://doi.org/10.1016/j.bone.2012.11.023>.

265 Karagkouni, Dimitra, Maria D Paraskevopoulou, Serafeim Chatzopoulos, Ioannis S Vlachos,
266 Spyros Tastsoglou, Ilias Kanellos, Dimitris Papadimitriou, et al. 2018. "DIANA-TarBase v8:
267 A Decade-Long Collection of Experimentally Supported MiRNA–Gene Interactions."
268 *Nucleic Acids Research* 46 (D1): D239–45. <https://doi.org/10.1093/nar/gkx1141>.

269 Karere, Genesio M, Jeremy P Glenn, John L VandeBerg, and Laura A Cox. 2012. "Differential
270 MicroRNA Response to a High-Cholesterol, High-Fat Diet in Livers of Low and High LDL-C
271 Baboons." *BMC Genomics* 13 (July): 320. <https://doi.org/10.1186/1471-2164-13-320>.

272 Lisse, Thomas S. 2020. "Vitamin D Regulation of a SOD1-to-SOD2 Antioxidative Switch to
273 Prevent Bone Cancer." *Applied Sciences* 10 (7): 2554.
274 <https://doi.org/10.3390/app10072554>.

275 Liu, Yanqing, Yueqiu Wang, Nailong Yang, Suning Wu, Yanhua Lv, and Lili Xu. 2015. "In Silico
276 Analysis of the Molecular Mechanism of Postmenopausal Osteoporosis." *Molecular*
277 *Medicine Reports* 12 (5): 6584–90. <https://doi.org/10.3892/mmr.2015.4283>.

278 Love, Michael I, Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold
279 Change and Dispersion for RNA-Seq Data with DESeq2." *Genome Biology* 15 (12): 550.
280 <https://doi.org/10.1186/s13059-014-0550-8>.

281 Mackowiak, Sebastian D. 2011. "Identification of Novel and Known MiRNAs in Deep-
282 Sequencing Data with MiRDeep2." *Current Protocols in Bioinformatics* 12 (12): 1–15.
283 <https://doi.org/10.1002/0471250953.bi1210s36>.

284 R Core Team. 2019. "R: A Language and Environment for Statistical Computing." Vienna,
285 Austria: R Foundation for Statistical Computing.

286 Ru, Yuanbin, Katerina J. Kechris, Boris Tabakoff, Paula Hoffman, Richard A. Radcliffe, Russell
287 Bowler, Spencer Mahaffey, et al. 2014. "The MultiMiR R Package and Database:
288 Integration of MicroRNA–Target Interactions along with Their Disease and Drug
289 Associations." *Nucleic Acids Research* 42 (17): e133–e133.
290 <https://doi.org/10.1093/nar/gku631>.

291 Spradling, Kimberly D, Jeremy P Glenn, Roy Garcia, Robert E Shade, and Laura A Cox. 2013.
292 "The Baboon Kidney Transcriptome: Analysis of Transcript Sequence, Splice Variants, and
293 Abundance." *PloS One* 8 (4): e57563. <https://doi.org/10.1371/journal.pone.0057563>.

294 Therneau, Terry M. 2020. "Coxme: Mixed Effects Cox Models."

- 295 Vagtborg, Harold. 1973. *The Story of the Southwest Research Center: A Private, Nonprofit,*
296 *Scientific Research Adventure.* University of Texas at Austin.
- 297 Wan, Chengyao, Jing Wen, Xiaolin Liang, Qiongni Xie, Wenqi Wu, Meiqing Wu, and Zhenfang
298 Liu. 2021. "Identification of MiR-320 Family Members as Potential Diagnostic and
299 Prognostic Biomarkers in Myelodysplastic Syndromes." *Scientific Reports* 11 (1).
300 <https://doi.org/10.1038/s41598-020-80571-z>.
- 301 Wang, X, J D Mabrey, and C M Agrawal. 1998. "An Interspecies Comparison of Bone Fracture
302 Properties." *Bio-Medical Materials and Engineering* 8 (1): 1–9.
- 303 Weilner, Sylvia, Susanna Skalicky, Benjamin Salzer, Verena Keider, Michael Wagner, Florian
304 Hildner, Christian Gabriel, et al. 2015. "Differentially Circulating MiRNAs after Recent
305 Osteoporotic Fractures Can Influence Osteogenic Differentiation." *Bone* 79 (October): 43–
306 51. <https://doi.org/10.1016/j.bone.2015.05.027>.
- 307 Xiao, Feifei, Zhixiang Zuo, Guoshuai Cai, Shuli Kang, Xiaolian Gao, and Tongbin Li. 2009.
308 "MiRecords: An Integrated Resource for MicroRNA-Target Interactions." *Nucleic Acids*
309 *Research* 37 (Database issue): D105-10. <https://doi.org/10.1093/nar/gkn851>.
- 310 Xiao, Qing, Yinu Zhao, Jia Xu, Wen-Jie Li, Yu Chen, and Hong-Jing Sun. 2019. "NFE2/MiR-
311 423-5p/TFF1 Axis Regulates High Glucose-Induced Apoptosis in Retinal Pigment Epithelial
312 Cells." *BMC Molecular and Cell Biology* 20 (1): 39. [https://doi.org/10.1186/s12860-019-](https://doi.org/10.1186/s12860-019-0223-2)
313 [0223-2](https://doi.org/10.1186/s12860-019-0223-2).
- 314 Yang, Sefei, J. U.N. Guo, L. E.I. Zhou, Helin Xing, Xianli Wang, and Chaofang Dong. 2018.
315 "Mir-148b-3p, Mir-337-5p and Mir-423-5p Expression in Alveolar Ridge Atrophy and Their
316 Roles in the Proliferation and Apoptosis of Ommscs." *Experimental and Therapeutic*
317 *Medicine* 16 (6): 5334–42. <https://doi.org/10.3892/etm.2018.6850>.
- 318 Yu, W., M. Clyne, M. J. Khoury, and M. Gwinn. 2010. "Phenopedia and Genopedia: Disease-
319 Centered and Gene-Centered Views of the Evolving Knowledge of Human Genetic
320 Associations." *Bioinformatics* 26 (1): 145–46. <https://doi.org/10.1093/bioinformatics/btp618>.
- 321 Yuan, Xiao-Peng, Long-Shan Liu, Chuan-Bao Chen, Jian Zhou, Yi-Tao Zheng, Xiao-Ping Wang,
322 Ming Han, and Chang-Xi Wang. 2017. "MicroRNA-423-5p Facilitates
323 Hypoxia/Reoxygenation-Induced Apoptosis in Renal Proximal Tubular Epithelial Cells by
324 Targeting GSTM1 via Endoplasmic Reticulum Stress." *Oncotarget* 8 (47): 82064–77.
325 <https://doi.org/10.18632/oncotarget.18289>.

326 Zhang, Yiqiang, Archana Unnikrishnan, Sathyaseelan S. Deepa, Yuhong Liu, Yan Li, Yuji Ikeno,
 327 Danuta Sosnowska, Holly Van Remmen, and Arlan Richardson. 2017. "A New Role for
 328 Oxidative Stress in Aging: The Accelerated Aging Phenotype in Sod1– Mice Is Correlated
 329 to Increased Cellular Senescence." *Redox Biology* 11 (April): 30–37.
 330 <https://doi.org/10.1016/j.redox.2016.10.014>.

331 Zhu, Xin, and Xiaolan Lu. 2019. "MiR-423-5p Inhibition Alleviates Cardiomyocyte Apoptosis and
 332 Mitochondrial Dysfunction Caused by Hypoxia/Reoxygenation through Activation of the
 333 Wnt/B-catenin Signaling Pathway via Targeting MYBL2." *Journal of Cellular Physiology*
 334 234 (12): 22034–43. <https://doi.org/10.1002/jcp.28766>.

335