Integration of phenomics and transcriptomics data to reveal drivers of inflammatory processes in the skin

Richa Batra^{1,2,3*,†}, Natalie Garzorz-Stark^{1,4,5,†}, Felix Lauffer^{1,†}, Manja Jargosch^{1,6†}, Caroline Pilz¹, Sophie Roenneberg¹, Alexander Schäbitz^{1,4}, Alexander Böhner¹, Peter Seiringer¹, Jenny Thomas⁶, Bentolhoda Fereydouni⁴, Ginte Kutkaite², Michael Menden², Lam C Tsoi⁷, Johann E Gudjonsson⁷, Fabian Theis F^{2,8}, Tilo Biedermann¹, Carsten B Schmidt-Weber⁶, Nikola Müller², ‡, Stefanie Eyerich⁶, ‡, Kilian Eyerich^{1,4,5,*‡}

¹Department of Dermatology and Allergy, Technical University of Munich, Munich, Germany;

²Institute of Computational Biology, Helmholtz Center Munich, Neuherberg, Germany;

³Institute for Computational Biomedicine, Englander Institute for Precision Medicine, Department of Physiology and Biophysics, Weill Cornell Medicine, New York, NY 10021, USA

⁴Division of Dermatology and Venereology, Department of Medicine Solna, and Center for molecular medicine, Karolinska Institutet; Stockholm, Sweden;

⁵Unit of Dermatology, Karolinska University hospital, Stockholm, Sweden;

⁶ZAUM – Center of Allergy and Environment, Helmholtz Center and Technical University of Munich, Munich, Germany;

⁷Department of Dermatology, University of Michigan, Ann Arbor, MI, USA;

⁸Department of Mathematics, Technical University of Munich, Garching, Germany;

†R.B., N.G.Z., F.L. and M.J. joint first authors.

‡N.M., S.E. and K.E. joint last authors.

*To whom correspondence should be addressed: rib4003@med.cornell.edu or kilian.eyerich@ki.se

Key words: chronic inflammatory diseases, psoriasis, atopic dermatitis/eczema, clinical phenotyping, transcriptome, skin, heterogeneity, neutrophil, acanthosis, CEBPB

Funding sources: This work was supported by an ERC grant (IMCIS, 676858) and the German Research Foundation (EY97/3-2).

Conflicts of interest: The authors have declared that no conflict of interest exist in relation to this article.

Abstract

Chronic inflammatory diseases are characterized by complex interactions between genetic predisposition and tissue-specific immune responses. This heterogeneity complicates diagnoses and the ability to exploit omics approaches to improve disease management, develop more effective therapeutics, and apply precision medicine. Using skin inflammation as a model, we developed a method that integrates deep clinical phenotyping information (phenomics) with transcriptome data of lesional and non-lesional skin (564 samples) to identify clinically-relevant gene signatures. It led us to discover so-far unexplored factors, including CCAAT Enhancer-Binding Protein Beta (*CEBPB*) in neutrophil invasion, and Pituitary Tumor-Transforming 2 (*PTTG2*) in the pathogenic epithelial response to inflammation. These factors were validated using genetically-modified human skin equivalents, migration assays, and *in situ* imaging. Thus, by meaningful integration of deep clinical phenotyping and omics data we reveal hidden drivers of clinically-relevant biological processes.

Introduction

Ever since the emergence of high-throughput omics technologies, computational methods have played a pivotal role in biomedicine (1) (2) (3) (4) (5) (6) (7, 8). Although most of the computational algorithms can be applied ubiquitously, specific biomedical questions demand innovation. For example, identifying causative factors in chronic inflammatory processes is challenging as they are clinically heterogeneous and characterized by a complex interplay between genetic predisposition, immune responses against environmental-, self-, or microbiome-derived antigens, and tissue-specific immune response patterns of e.g. the gastrointestinal system, joints, lung, or skin. In addition and contrast to cancer, which is usually characterized by mutation-driven inexorable progression, inflammatory diseases typically show a relapsing-remitting disease course with a broad spectrum of phenotypes (9) ultimately resulting in a pronounced data variety wherein the various distinct signals are not well separated. In line with this, standardized experimental and animal models of human inflammation insufficiently reflect the comprehensive human situation (10-12), and available data sets are insufficiently characterized to dissect disease heterogeneity or identify distinct disease endotypes.

Non-communicable inflammatory skin diseases (ncISD) represent ideal prototype diseases for chronic inflammation to develop a strategy that identifies relevant molecular events within a complex network of overlaying gene expression signatures. ncISD are a group of several hundred diseases including psoriasis and atopic dermatitis (AD). They are defined by physicians using a subjective “best fit” model via investigation of a variety of attributes such as clinical phenotype, tissue and laboratory analysis (13). The transcriptome of ncISD has been widely investigated, greatly enhanced our knowledge(14) and lead to the three umbrella phenotypes which are defined according to the immune response pattern in the skin. Type 1 ncISD, characterized by cytotoxic immune cells in the skin (so-called interface dermatitis) and comprising the diseases lichen planus and cutaneous lupus(15, 16); type 2 ncISD, characterized by impaired epidermal barrier and innate immunity and comprising eczematous diseases(17, 18); and type 3 ncISD defined by Th17 immunity and the presence of neutrophil granulocytes in the skin, e.g. psoriasis and variants(19). However, the subjective nature of diagnosis, together with the heterogeneity and dynamics of ncISD, result in overlapping phenotypes and

potential misdiagnoses. In addition, insufficient clinical metadata hinders discovery of relevant, potentially overlapping, disease mechanisms.

Thus, to achieve a substantial breakthrough in precision medicine, the complex pathogenesis of inflammatory diseases such as ncISD and the resulting disease heterogeneity needs to be acknowledged in research strategies. To this end, we performed deep clinical phenotyping of 287 ncISD patients using 62 attributes, including clinical picture, skin architecture, laboratory parameters, and information on the patient's history, and generated transcriptomic profile of tissue biopsies of paired lesional and non-lesional skin. Furthermore, we developed a novel framework, called **Attribute Gene Expression Regularization (AuGER)**, to integrate phenotypic/phenomic profiles with transcriptomic profiles. This approach enabled us to determine gene signatures of each clinical attribute encompassing known as well as so-far unexplored genes involved in clinically-relevant biological processes. Experimental validation of newly identified factors shed light on the drivers of neutrophil granulocyte migration into tissue and the pathogenic epithelial response to inflammation that ultimately modulates skin tissue architecture.

Results

Deep phenotyping explains substantially more transcriptomic tissue variance than the diagnosis alone

Given the heterogeneity in ncISD patients, we hypothesized that current clinical diagnoses only partially reflects the molecular events occurring in the skin. To test this, we collected lesional skin biopsies from 287 patients covering 13 different diagnoses of ncISD (Fig. 1A, Table S1) and performed RNA-sequencing (RNAseq). Principal component analysis (PCA) of the normalized read count of the RNAseq data showed intermixing of ncISD type 1, type 2, and type 3 diagnoses in the first two major principle components (PC1 and PC2; Fig.1A) (type 1: lichen planus, cutaneous lupus; type 2: atopic dermatitis, dishydrotic eczema, nummular eczema, hyperkeratotic-rhagadiform eczema; type 3: pustular psoriasis, guttate psoriasis, pityriasis rubra pilaris) , which indicates discrepancy between the diagnosis and the transcriptomic profile (at least in some patients). To exclude the possibility that transcriptional heterogeneity is driven by intra-individual differences, we included transcriptomic profiles of autologous non-lesional skin biopsies in the PCA (n = 564 samples in total) (Fig. S1). The status of the skin (lesional or non-lesional) explained the separation of PC1 and in total 32 % of the variance in the transcriptome, with genes related to skin inflammation driving the separation (Fig. S1A, B). Another 9 % of the variance mainly in lesional samples was explained by gender, which was confirmed by the expression of genes located on sex chromosomes (Fig. S1A, B). Thus, we performed unsupervised hierarchical clustering of lesional transcriptome excluding genes encoded on the sex chromosomes. There was still intermixing of patients with different diagnoses in the clusters (Fig. S1C), indicating that gender correction alone is insufficient to improve the concordance of transcriptome and diagnosis.

To dissect transcriptional heterogeneity seen in patients, we performed deep clinical phenotyping of the 287 patients using 62 attributes from the data recorded by dermatologists during diagnosis that reflect clinical picture, histologic architecture, laboratory results, investigation of comorbidities, and personal and family history (Fig. 1B, Fig. S2). This generated 287 unique patient fingerprints (Fig. S3). We analyzed the fingerprints from four patients with the diagnosis psoriasis (n = 2) or AD (n = 2). Three of these clustered closely based on transcriptional profiles, while one AD patient was located far from

the other three patients in the PCA (Fig. 1C). At this single patient level, patients with transcriptionally-similar profiles indeed represented individuals with similar phenomic profiles despite their differing diagnoses (Fig. 1C). To compare the amount of transcriptomic variance explained by attributes derived from deep clinical phenotyping with the clinicians' diagnosis we used a linear mixed effect model. We found that clinicians' diagnoses only explained 7.8 % of the total transcriptional variance in the skin (Figure 1D). In contrast, the 62 attributes of the deep phenotyping together explained 36.7 % of the total transcriptomic variance (Fig. 1D, E). Thus, we concluded that transcriptomic profile of a patient can be better disentangled using multiple, clinically-relevant attributes than the clinician's diagnosis alone.

Unique attribute-gene signatures identified by deconvolution of the transcriptional signal

Usually single attributes are not specific for a distinct disease, for instance, scaling which is considered a hallmark of psoriatic skin, can also occur in other inflammatory skin conditions. However, these overlapping attributes still contribute in shaping the lesional skin transcriptome and thereby the traditional approach of comparing diseases is limited owing to high background noise. To overcome the super-imposed definition of diseases, we evaluated each patient using several clinical attributes. With the assumption that each attribute represents a clinically-relevant biological process, we developed a computational method to identify attribute specific gene signatures. Our approach **Attribute Gene Expression Regularization (AuGER, Fig. 2A)** associates phenomic profiles with transcriptomic profiles using regularized regression. As a first step, we performed attribute reduction using pairwise correlation analysis, which revealed groups of highly correlated attributes (Fig. 2A 1-3, Fig. 2B, Table S4). As expected, attributes that are used to describe a specific ncISD disease type highly correlate with each other: specifically, the ncISD type 1 attributes, "number of dyskeratoses", "interface dermatitis qualitative" and "quantitative", "localization of dyskeratoses"; the ncISD type 2 attributes, "serum crusts", and "spongiosis"; and the ncISD type 3 attributes, "neutrophils qualitative" and "quantitative", and "parakeratosis" (Figure 2B). One representative attribute (= core attribute) per group of highly correlated attributes was selected based on clinical relevance, favouring quantitative

measures over qualitative discretization, resulting in a total of 24 core attributes used for further analysis (Fig. 2C).

Next, we used multivariate regression with elastic net penalty to associate each core attribute with the transcriptome. For each of the 24 core attributes regression analysis was carried out, such that the respective core attribute was used as the dependent variable (output), and the genes of the complete transcriptome data as the predictor variables (input). The remaining 23 core attributes were used as covariates to control for their impact on the dependent variable. For each gene, the regression coefficient was estimated per attribute, and all genes with non-zero regression coefficients were identified (Fig. 2A 4-5). To determine the clinically-relevant associations between individual attributes and genes, genes with non-zero coefficient in lesional and zero coefficient in non-lesional skin were selected (Fig. 2A 6). For robust results, regression analysis of both lesional and non-lesional transcriptome was conducted five times each, using five sets of imputed core attributes. Specific, unique, and shared associations were obtained between a total of 621 genes and 18 of the 24 attributes (Fig. 2A 7, Table S4, Fig. S4). For the first time, AuGER acknowledges disease heterogeneity of ncISD by dissecting diseases into multiple clinically-relevant attributes and assigning gene signatures to these attributes. These gene signatures reveal the molecular basis of complex and biologically-relevant processes, e.g. the presence of neutrophil granulocytes in the skin or pathogenic epithelial thickening as a consequence of inflammation, so-called acanthosis.

The gene expression landscape of ncISD

To understand the interconnections of attributes and its reflection on ncISD types, we created an atlas of skin inflammation (Fig. 2C). This atlas depicts the findings of AuGER as a network, with 18 core attributes and 621 genes as nodes linked by an edge representing a non-zero regression coefficient. On the one hand, this atlas reflects core attributes that represent hallmarks of the three ncISD clusters, type 1 (lichen planus and cutaneous lupus, here identified by the core attribute “interface dermatitis”), type 2 (eczema variants which are typically associated to “dryness” and “eosinophils”), and type 3 (psoriasis variants represented by the core attribute “neutrophils”).

On the other hand, the atlas also illustrates attributes which are not typically associated with one of the three ncISD clusters, but still have an impact on the skin transcriptome. These include scalp or nail involvement, clinical presentation such as scaling, papules, or erythema; or histo-pathological presentation such as acanthosis or orthokeratosis (regular formation of the upper epidermis). These attributes may occur in several diseases and thus increase the unspecific background noise in traditional approaches used to compare different diseases. Finally, external factors influencing the lesional skin transcriptome are also captured. This is represented by numerous genes associated with the core attributes “alcohol consumption” and “smoking” (Fig. 2C).

Notably, AuGER identified only one gene associated with the attribute “pruritus”, namely the T cell recruiting chemokine *CCL18* (20). *CCL18* was shared with the core attribute “eosinophils” and thus connected to type 2 ncISD, namely eczema. This is in line with the fact that most eczema variants are highly pruritic (21). Indeed, levels of *CCL18* were highest in eczema variants (AD and nummular eczema) as compared to psoriasis (Fig. S6A). This was confirmed in an independent data set consisting of skin from healthy volunteers ($n = 38$) and from patients suffering from AD ($n = 27$) or psoriasis ($n = 37$). Here, expression of *CCL18* was again significantly higher in AD than in the healthy controls (log2 fold change: 4.9, $p = 1.4 \times 10^{-13}$) or psoriasis (3.1, $p = 1.4 \times 10^{-7}$) (Fig. S6B). In a third cohort, *CCL18* expression was higher in lichen planus lesional skin ($n = 20$) as compared to healthy controls ($n = 38$). Just like eczema, lichen planus is typically pruritic (22)(Fig. S6B).

Furthermore, all attributes except “mucin” shared at least one link with another attribute in the network, thus constructing a comprehensive gene map of skin inflammation. “Mucin” is a core attribute comprising an enhanced level of extracellular material (23). It is not associated with an inflammation cluster and is usually the result of inflammation of deeper skin layers rather than of the epidermis. Connections of the individual gene signatures of each attribute were also identified at the level of pathway analysis. Some pathways were associated with only one attribute, e.g. the expected cytotoxicity-mediating pathway “IFN-gamma signaling” associated with the ncISD type 1 core attribute “interface dermatitis” (15). Numerous pathways were associated with at least five different attributes. For example, pathways of cell cycle and proliferation, such as “EGF signaling”, “Ras signaling”,

“Oncostatin M signaling”, and the stem cell factor “Wnt signaling”, connected the clinical attributes “elevation” (palpable skin lesions) and “papules”, both of which may be related to exaggerated proliferation of the epidermis (Fig. S5).

Novel insights into neutrophil granulocyte biology in tissue

The comprehensive atlas of the transcriptome landscape of ncISD delivers a plethora of so-far undiscovered connections of clinically-relevant biological processes with gene expression. To understand the pathogenicity of these genes, we explored the genes associated with the attribute “neutrophils”, which was defined as the presence of neutrophil granulocytes in lesional skin as observed by dermatopathology. AuGER identified an association of 35 genes with this attribute (Fig. 3A). While some of these genes such as *CXCL1* or *CXCL8* are well-known drivers of neutrophil migration and/or biology (24), others were previously unknown in this context. As expected, the ncISD type 3 core attribute “neutrophils” was most prominent in psoriasis patients, however, intra- and inter-disease variability was high (Fig. 3B, C). Of the 35 genes majority are associated with inflammation and general processes such as protein binding (Fig. 3D). For biological validation of the attribute-gene association, we focused on five genes *CEBPB*, *HCAR3*, *SHC1*, *HS6ST1* and *SLC23A2* as they were centrally located in the network (Fig. 2C) and are novel in the context of skin diseases. First, we analyzed the effect of stimulation with pro-inflammatory cytokines *IL-17A*, *TNF-α* or *LPS* on gene expression in two potential target cells, primary human keratinocytes and neutrophil-like HL-60 cells. Stimulation induced gene expression only of *CEBPB* consistently in both cell types as compared to unstimulated controls (Fig. 4A).

C/EBP factors, to which *CEBPB* belongs, have been found to regulate cytokine expression in human neutrophils (25). To gain more insight into the function of *CEBPB* in neutrophil biology in particular into the interaction of neutrophils and keratinocytes, we specifically knocked-out *CEBPB* and the unrelated control gene *PTTG2* in keratinocytes using CRISPR/Cas9 technology (controls on knock-out efficiency are given in Fig. S7). Only *CEBPB* knock outs (not *PTTG2*) resulted in significant reduction of central chemokines for neutrophil migration and activation (*CXCL8*, *CXCL1*, *CXCL2*, *CXCL5*, *IL-6*) in the

supernatant of keratinocytes stimulated with *IL-17A* (Fig. 4B). Moreover, *CEBPB* knock-out cells produced significantly higher amounts of the growth factor *GM-CSF* and the chemokines *CCL5* and *CCL22*, indicating that loss of *CEBPB* does not hamper the ability of keratinocytes to promote bone marrow neutrophil maturation, or migration of eosinophils, macrophages or dendritic cells (Fig. 4B). To validate *CEBPB* mediated regulation of neutrophil migration to the skin, we quantified migration of primary human neutrophils towards the supernatant of *CEBPB* or *PTTG2* knock-out keratinocytes, compared to the supernatant of wild type controls. Here, knock-out of *CEBPB* significantly reduced the capacity of keratinocytes to regulate neutrophil migration (Fig. 4C). Furthermore, knock-out of *CEBPB* directly in the neutrophil-like HL-60 cells resulted in diminished migration towards both *CXCL8* and keratinocyte supernatant (Fig. 4D). Taken together, we identify *CEBPB* is a pivotal regulator of neutrophil biology in the skin, acting at both the keratinocyte and the neutrophil level.

Pathogenic epidermal hyperplasia is driven by previously unknown genes

Pathogenic hyperplasia of the epidermis caused by excessive proliferation and disturbed differentiation of keratinocytes, so-called acanthosis, is a process frequently observed in different kinds of skin inflammation. Therefore, acanthosis is a representative example of AuGER's power to select and identify so-far undiscovered drivers of this highly prevalent clinical phenotype. 69 genes were associated with acanthosis (Fig. 2C, Fig. 5A). Acanthosis is neither specific for a ncISD type, nor is it observed in all patients suffering from a ncISD. In line with this, in our cohort, acanthosis as an attribute was variably distributed across the 13 types of ncISD (Fig. 5B) and unsupervised hierarchical clustering of genes associated with acanthosis in all 287 patients did not correlate with distinct diseases (Fig. 5C). The acanthosis gene signature is associated with cell-cell adhesion and activity (Fig. 5D).

For biological validation of the attribute-gene association, we selected *PTTG2*, *SYNE1*, and *CEBPB*. Keratinocytes stimulated with the pro-inflammatory cytokines *IL-17A* and *TNF- α* showed an upregulation of *CEBPB* and a downregulation of *SYNE1* gene expression, while *PTTG2* expression remained unaltered (Fig. 6A). Cellular morphology or the architecture of three-dimensional human skin equivalents remained unaltered in knock-out of *PTTG2*, *SYNE1*, *CEBPB* in primary human

keratinocytes (Fig. 6B, see effectiveness in Fig. S7). However, it is difficult to judge on specific effects in unstimulated models as they only consist of a few cell layers. The cytokine IL-22 is a known inducer of keratinocyte growth and induces acanthosis (26). Next, we investigated the effects of *CEBPB*, *SYNE1* and *PTTG2* knock-out on epidermal thickness as a proxy for acanthosis in response to IL-22. In line with the differences in gene regulation, knock-out of both *PTTG2* and *CEBPB* significantly inhibited the induction of acanthosis by IL-22 compared to the control (noRNP) model, with *CEBPB* preventing it completely. In contrast, knock-out of *SYNE1* markedly increased the thickness of the three-dimensional skin model (Fig. 6B, 6C; Fig. S8). Notably, proliferation of keratinocytes was also reduced by knock-out of *CEBPB* or *PTTG2* but induced by *SYNE1* (Fig. 6C, 6D, 6E). In addition, knock-out of *CEBPB* resulted in altered epithelial differentiation after IL-22 stimulation, namely in clearly diminished expression of the early differentiation marker keratin 10, and keratin 16, which is typically associated with proliferative keratinocytes (27) (Fig. 6E). Hence, *CEBPB* and *PTTG2* are essential for the development of acanthosis, while *SYNE1* is a negative regulator of epidermal thickness.

Discussion

Here, we show that so-far unexplored drivers of key pathogenic events in human inflammatory diseases can be identified from transcriptomics when disease heterogeneity is acknowledged by taking deep clinical phenotyping into account. We investigated non-communicable inflammatory skin diseases (ncISD) as a prototype example of heterogeneous human inflammatory disorders.

We observed that clinician assigned diagnosis explained very little transcriptomic variation. In contrast, metadata with a higher granularity of the observed clinical heterogeneity explains a substantially higher proportion of the transcriptomic variability associated with inflammation. Furthermore, we demonstrate that common hallmarks of the interplay between skin epithelia and immune cells are shared between different patient cohorts independent of the disease background. We and others recently followed this approach for ncISD that share the histological hallmark “interface dermatitis”, which represents cytotoxic activity of lymphocytes at the basal membrane of the epidermis and is frequently observed in type 1 ncISD such as lichen planus and lupus (15, 16). Here, identification of the molecular mechanism, type 1 immune mediated epithelial apoptosis and necroptosis, led to potential novel therapeutic strategies (28). Our current analysis takes into account other dependent clinical attributes and gives an even clearer picture of the genes involved in interface dermatitis.

Even in settings that are heavily researched and where numerous drivers are known, such as neutrophil biology, assigning more granular metadata of clinical phenotyping identifies so-far unexplored, but important mechanistic factors. In particular, our newly developed algorithm AuGER identified *CEBPB* as a master regulator of neutrophil biology in the skin. *C/EBP* proteins are known to be involved in cytokine production such as *CXCL8* and *CCL3* by neutrophils (25, 29). Here, we show that *CEBPB* is also expressed in keratinocytes, and knock-out of *CEBPB* in primary human keratinocytes results in reduced chemokine secretion and reduced neutrophil migration. This newly identified dual mechanism of action to inhibit neutrophil inflammation in tissue renders *CEBPB* a potentially powerful therapeutic target for neutrophilic inflammatory diseases of the skin, such as psoriasis or pyoderma gangrenosum but also for diseases beyond the skin as for example reactive arthritis or neutrophil-rich chronic obstructive lung disease (COPD).

Our approach not only reveals the most significant genes contributing to an attribute, but *vice versa* also identifies the mechanistic significance of single genes for corresponding attributes representing complex mechanistic processes. This was proven for the less restricted attribute “acanthosis”. Acanthosis corresponds to reactive thickening of the epidermis in inflammatory conditions and it can be observed in multiple ncISD but is not part of a disease definition (30). *PTTG2*, a hub gene for acanthosis in our analysis, has not been proposed in the context of acanthosis before. *PTTG2* is involved in cell viability and migration by regulating the expression of vimentin and E-cadherin and overexpression of *PTTG2* was described in psoriatic epidermis (31, 32). We now show that *PTTG2* is directly involved in acanthosis, as inhibiting this single factor significantly impaired acanthosis formation *in vitro*. *SYNE1*, another gene significantly correlated to acanthosis from our network, encodes an important scaffold protein. In genetically modified mice, loss of *SYNE1* leads to thickening of the epidermis (33), an observation we validated in the human system by showing increased thickness of three-dimensional skin equivalents comprised of *SYNE1* knocked-out keratinocytes. Interestingly, *CEBPB*, associated with the attribute “neutrophil” was also associated with “acanthosis” and the third candidate gene we validated for its contribution to acanthosis *in vitro*. In fact, human skin equivalents of keratinocytes lacking *CEBPB* were vital and did not show any morphological alterations, but they did not respond to *IL-22*, an inflammatory stimuli that normally induces acanthosis (26). Hence, given a tolerable safety profile, *CEBPB* could be a target to address diseases in which neutrophils and acanthosis are concomitantly present, in particular in psoriasis, hidradenitis suppurativa, or severe acne.

Our *proof-of-concept* study has limitations. First, phenotyping was partially based on subjective information, such as the patient’s narrative, clinical examination and histological evaluation. Second, though we performed RNA-Sequencing of 287 patient’s tissue samples, the sample size is still limited, and underestimation of real biological effects cannot be ruled out. Higher sample size and/or additional datasets are required to strengthen our approach, AuGER, and its potential to extract markers predictive of medically relevant outcomes, such as response to therapy. AuGER is optimized to identify genes so-far hidden in the background noise, by correcting each attribute signature for

other attributes, thus, gene signatures may seem incomprehensive. Despite the limitations, this study clearly shows that associating granular clinical phenotypes to omics datasets critically determines the quality of outcome of the analysis. Phenotyping methods such as image-based machine-learning as well as adding several molecular information sources such as serum proteome or microbiome (34) will further improve our novel approach in the future.

Taken together, exemplifying ncISD we successfully combined deep clinical phenotyping with transcriptome and established that this novel approach has the potential to define pathogenic events in complex processes such as chronic human inflammation and thus build the basis for precision medicine in the field.

Methods

Data, Code and Material Availability

Available upon request. This study did not generate new unique reagents.

Study cohort and deep clinical phenotyping

Patients with nclSD (n = 287, 13 distinct diagnoses, male n=163, female n=124, age male 54,19±16,69 years, age female 57,92±18,82) from the *Biobank Biederstein* were enrolled into the study. All patients were investigated by dermatology specialists. A washout phase of at least two weeks for topical treatments and five half-life times for systemic treatments was mandatory for study inclusion. The study followed the Declaration of Helsinki and was approved by the local ethics committee (Klinikum Rechts der Isar, 44/16 S). The *Biobank Biederstein* follows data protection rules and is approved by the local ethics committee (Klinikum Rechts der Isar, 5590/12). After written informed consent was obtained from all patients, each patient was deep-phenotyped regarding clinical presentation (distribution and phenotype of lesions), personal and family history (questionnaires), and laboratory parameters (smear test cultures and serum analysis). Furthermore, a non-lesional and a lesional skin biopsy (6 mm each) were obtained under local anesthesia from all patients except for ten patients that did not have non-lesional skin as all skin was affected by inflammation. Biopsy specimens were divided and one part was used for hematoxylin-eosin staining from paraffin-embedded skin. These sections were evaluated by a dermato-pathologist in a blinded manner. A list of all parameters evaluated is given in Fig. S2. Effects of sex were evaluated, reported in the results section and the corresponding figures Fig. 1 and Fig. S1.

Primary human keratinocytes

Primary human epidermal keratinocytes (male n=4, female n=1) were obtained by suction blister as reported previously (10) and cultured in keratinocyte medium (DermaLife Basal Medium supplemented with DermaLife K LifeFactor Kit (Lifeline Cell Technology, LL-0007)) at 37 °C, 5 % CO₂. Second- to third-passage primary human epidermal keratinocytes were used for all experiments.

Primary neutrophils and neutrophil-like cells (HL-60 cells)

Primary human neutrophils were isolated from peripheral blood of healthy donors (male n=1, female n=2). HL-60 cells represent a promyelocytic cell line with phagocytic and chemotactic functions that has been obtained from a 36-year old Caucasian female with acute promyelocytic leukemia.

RNAseq library preparation, sequencing, mapping and quantification

RNA from skin biopsies was isolated using the QIAzol Lysis Reagent (Qiagen) and miRNeasy Mini Kit (Qiagen) according to manufacturer's protocol. RNASeq libraries were generated using the TruSeq Stranded Total RNA Kit (Illumina) according to manufacturer's high sample protocol. Finally, samples were sequenced on an Illumina HiSeq4000 as paired-end with a read length of 2x 150 bp and an average output of 40 Mio reads per sample and end.

Sequence alignment was performed using STAR aligner with human genome reference hg38. RNAseq count data was normalized and then transformed using variance stabilizing transformation from the bioconductor package DESeq2. In total, 564 samples (287 lesional, 277 non-lesional) samples were used for further analysis.

Principal component analysis (PCA)

Principal component analysis was performed using the top 1000 most varying genes across samples. Variance was computed per gene across the samples and 1000 genes with highest variance was used in the principal component analysis. PCA was performed twice, once for all 564 samples and once for 287 lesional samples only.

Variance estimation

To estimate the variance in lesional gene expression explained by differences in diagnosis, gender, age and other clinical attributes, the RNAseq count data was first transformed using voom transformation from R package Limma and then modelled with a linear mixed effect model using bioconductor package variancePartition.

Attribute- Gene Expression Regularization (AuGER)

Step1: Define core attributes - A total of 62 clinical attributes (Fig. S2) from 287 patients/samples were used for this analysis. Missing values ($\leq 20\%$) were imputed using the R package mice. To identify highly correlated clinical attributes, pairwise comparison between attributes was performed using the hetcor function from the R package polycor. This function computes pearson correlation between numeric variables, polyserial correlation between numeric and ordinal variables, and polychoric correlation between ordinal variables. Of the highly correlated attributes, a representative was selected based on clinical relevance for the follow up analysis, leading to a set of 24 core attributes. The core attributes were reimputed generating 5 sets. **Step 2:** Transform the gene expression data – RNASeq count data was split into lesional and non-lesional sets. Both the sets were transformed using variance stabilizing transformation. 10 samples did not have a matching non-lesional sample. These missing samples were imputed with the median value of each gene, of non-lesional samples. **Step 3:** Regularization - Multivariate regression with elastic net penalty was used to identify the association between clinical attributes and gene expression. The regression was performed using R package glmnet, alpha was set to 0.5, lambda was computed using five-fold cross validation, family of regression was selected based on the response variable (binomial for binary, gaussian for numeric and multinomial for categorical variables). Each clinical attribute was used as response variable and gene expression data was used as predictor variables. The attributes other than response were used as covariates to control for the influence on response variable. Clinical attributes age and gender were accounted for in each regression. This step was repeated 5 x 2 times, for 5 imputed clinical attribute sets and the 2 gene expression sets of lesional and non-lesional skin. **Step 4:** Selection of genes –A regression/beta coefficient for each predictor variable was estimated by the regression analysis. All genes with a non-zero coefficient were selected for each attribute from all ten regression sets, however, genes unique to lesional skin were selected for further analysis i.e genes with non-zero coefficient in non-lesional skin were discarded. Next, genes were ranked according to the frequency of their detection in the five sets of lesional skin. We refer to these ranks as strength of association between

an attribute and the gene. **Step 5: Network** – The association of genes and clinical attributes was visualized as bipartite network. Genes and clinical attributes were represented as nodes and edges/links between them indicate associations. The edge/link width indicates the rank of the genes. The network was visualized using cytoscape. Nodes were manually arranged to aid the visualization.

Culture of primary human keratinocytes, primary neutrophils and neutrophil-like cells (HL-60 cells)

Primary human epidermal keratinocytes were cultured in keratinocyte medium (DermaLife Basal Medium supplemented with DermaLife K LifeFactor Kit (Lifeline Cell Technology)) at 37 °C, 5 % CO₂. Second- to third-passage primary human epidermal keratinocytes were used for all experiments. Before stimulation, cells were starved for 5 hours in DermaLife Basal Medium followed by stimulation with human recombinant IL-17A (R&D systems, 50 ng/ml) and TNF-α (R&D systems, 10 ng/ml) for 16 hours (mRNA analysis) or 72 hours (production of supernatant) in keratinocyte medium without hydrocortisone.

Primary human neutrophils were isolated from peripheral blood of healthy donors using LymphoPrep™ (Progen) and 2 % Dextran solution (Sigma Aldrich) and a final erythrocyte lysis. Neutrophils were cultured in RPMI 1640 medium supplemented with 1 % human serum, 0.1 mM NEAA, 2 mM L-Glutamine, 1 mM sodium pyruvate and 100 U/ml penicillin/streptomycin at 37 °C, 5 % CO₂.

HL-60 cells were cultured in HL-60 basal medium (RPMI1640 + HEPES & GlutaMAX (GIBCO) supplemented with 9 % FCS and 100 U/ml penicillin/streptomycin) at 37 °C, 5 % CO₂. To obtain neutrophil-like cells, HL-60 cells were differentiated for 6 days in HL-60 basal medium supplemented with 1.3 % DMSO (PanReac & AppliChem ITW Reagents) as published before (35). For gene expression analysis HL-60 cells were stimulated with 100 ng/ml LPS (InvivoGen) for 6 hours.

Neutrophil migration assay

Migration assay was performed using ChemoTx® Disposable Chemotaxis System (neuroprobe). 3 x 10⁴ neutrophils were added to the top of a 5 µm pore polycarbonate membrane and migrated to

keratinocyte supernatant for two hours. Migrated cells were analysed with an LSRFortessa flow cytometer (BD Biosciences). Migration was performed in triplicates.

CRISPR-Cas9 knock-out

Preparation of RNP complexes: Predesigned crRNA, tracrRNA, tracrRNA-ATT and S.p. HiFi Cas9 Nuclease was ordered from Integrated DNA Technologies in Alt-R® format. CrRNA and tracrRNA was reconstituted to 200 µM with IDTE Buffer. For generation of crRNA-tracrRNA duplex Oligos were mixed at equimolar concentrations and annealed by heating at 95 °C for 5 min following a slow cool-down to room temperature. For each target, a mixture of two crRNA sequences was applied. Finally, for RNP formation 180 pmol crRNA-tracrRNA duplex was mixed with 60 pmol Cas9 protein in Nucleofection Buffer P3 (Lonza) and incubated for at least 10 min at room temperature. crRNA sequences can be found in table S2. **Transfection of RNP complexes:** CRISPR knockout was done by application of RNP complexes with the 4D Nucleofector™ device (4D-Nucleofector™ Core Unit; 4D-Nucleofector™ X Unit, both Lonza) using the P3 Primary Cell 4D-Nucleofector™ X Kit S (Lonza) for keratinocytes or the SF Cell Line 4D-Nucleofector™ X Kit S (Lonza) for HL-60 cells. For transfection, 1.6×10^5 keratinocytes or 1×10^6 HL-60 cells were resuspended in 16 µl Nucleofection Buffer P3 (for keratinocytes) or SF (for HL-60 cells) and mixed with 4 µl RNP complex. The transfection mixture was transferred to Nucleofection™ cuvette strips and electroporated with the impulse DS-138 (for keratinocytes) or EO-115 (for HL-60 cells) in the 4D Nucleofector™ X unit. After nucleofection, cells were rested at room temperature for 10 min. Afterwards, 80 µl prewarmed cell culture media was added and cells were transferred to cell culture plates of choice depending on application.

3D keratinocyte skin model

3D keratinocyte skin models were cultured in polycarbonate inserts (Millipore) placed in a tissue-culture treated 6-Well plate. Inserts was pre-coated with 1 % collagen type I solution (SIGMA) in PBS for 45 min. 0.3×10^6 primary human keratinocytes were seeded directly after transfection in 500 µl keratinocyte medium supplemented with 1.5 mM CaCl₂ (SIGMA) into the insert. 2.5 ml keratinocyte

medium supplemented with 1.5 mM CaCl₂ was added in the surrounding well. The models were cultured at 37 °C and 5 % CO₂. Two days after seeding, airlift was done by aspirating the medium in the insert and the medium in the surrounding 6-well was replaced with 1.8 ml keratinocyte medium supplemented with 1.5 mM CaCl₂ and 50 µg/ml Vitamin C (SIGMA). Every second day the medium was exchanged with 2 ml keratinocyte medium supplemented with 1.5 mM CaCl₂ and 50 µg/ml Vitamin C in the surrounding 6-well. Nine days after airlift, 3D keratinocyte skin models were stimulated with 50 ng/ml IL-22 (R&D) in 2 ml keratinocyte medium supplemented with 1.5 mM CaCl₂ and 50 µg/ml Vitamin C and without hydrocortisone for three days. For histological analysis, inserts were fixed with 4 % Formaldehyde for 24 h at 4 °C and embedded in paraffin. The epidermal thickness of 3D skin models was measured from the top of the corneal layer to the bottom of the basal keratinocyte layer with ImageJ (<https://imagej.nih.gov/ij/download.html>) in four sections with a distance of 100 µm each per sample and the mean was calculated. Next, delta thickness (IL-22 minus unstimulated thickness) was calculated for each knockout group to visualize the acanthosis effect. Finally, this delta thickness was illustrated in relation to the delta thickness from the untransfected noRNP control sample.

Immunohistochemistry

5 µm sections of 3Dskin model samples were air-dried overnight at 37 °C, dewaxed and rehydrated. Staining were performed by an automated BOND system (Leica) according to the manufacturer's instructions: After antigen retrieval in a pH9 EDTA buffer based epitope retrieval solution (Leica), sections were incubated with monoclonal antibodies against Ki67 (rabbit anti-Ki67, Zytomed), and a secondary polymeric alkaline phosphatase (AP)-linked anti-rabbit antibody was applied. The complex was visualized by the substrate chromogen Fast Red. Slides were counterstained with haematoxylin. As a negative control, primary antibodies were omitted. Positive cells were counted in four to nine visual fields per condition.

Immunofluorescence

Paraffin mounted sections were air-dried overnight at 37 °C and dewaxed at 65 °C for 25 minutes and rehydrated by consecutive washes in X-TRA-Solv® (Mediate), isopropanol, ethanol (96 % vol/vol and 70 % vol/vol, respectively) and distilled H₂O. Antigen retrieval was performed in a pressure cooker with boiling citrate buffer (pH6). Hydrogen peroxide (Sigma Aldrich, Germany) blocking was performed in 3 % solution for 15 minutes. Sections were further blocked with 10 % normal goat serum and 10 % normal donkey serum diluted in antibody diluent (Abcam) for one hour and incubated with primary antibody mix (anti-ki67, anti-KT10 antibody & anti-KT16 antibody, all Abcam) for one hour at room temperature and then overnight. For negative controls, antibody diluent only was applied. After overnight incubation, secondary antibodies (AF488 donkey anti-rat, AF647 goat-anti rabbit antibody & AF557 donkey anti-mouse antibody, all Thermo Fisher Scientific) were applied in the dark for one hour at room temperature. Sections were treated with Vector® TrueVIEW® autofluorescence quenching kit (VECTOR Laboratories) and mounted with the included VECTASHIELD® Vibrance™ plus DAPI (VECTOR Laboratories). Stainings were imaged in the blue (DAPI), green (KI67), red (KT10) and far red (KT16) channel of a Leica TCS SP8 confocal laser scanning microscope.

Protein analysis and real time PCR

Protein analysis was performed using Bioplex assays (27-Plex, CXCL-1-, CXCL-2-, CXCL-5-, CCL22- single plex) (Bio-Rad Laboratories) according to the manufacturer's protocol.

For real time PCR, mRNA was isolated using the InviTrap Spin Universal RNA Mini Kit (Strattec) according to manufacturer's protocol. mRNA was transcribed into cDNA with Applied Biosystems High Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific) according to manufacturer's protocol. Using Fast Start Universal SYBRGreen Master Rox (Roche) gene expression was measured on Applied Biosystems ViiA7 Real-Time PCR system (Thermo Fisher Scientific) according to manufacturer's protocol. Primers were ordered from Metabion (www.metabion.com) Sequences can be found in Table S3.

Statistical analysis

Data of epidermal thickness, Ki67 immunohistochemistry, Bioplex, migration and real time PCR assays were visualized using GraphPad Prism 6 software (<https://www.graphpad.com>). The unpaired t-test with Welch's correction was used to test for difference in the means to noRNP sample. The anova test was used to test for differences within the groups for the Ki67 immunohistochemistry assay. Significance level was defined as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***) and $p < 0.0001$ (****).

Acknowledgements:

We thank Helen Pickersgill from Life Science Editors for editing the manuscript and Jana Sanger as well as Kerstin Weber for excellent technical support. In addition, we thank the NGS core facility at the Helmholtz Center Munich for help with RNA sequencing and the Biobank Biederstein for providing patient samples. RB would like to thank her colleagues at Institute of Computational Biology, Helmholtz Center Munich for helpful discussions.

Author contributions:

Conceptualization: R.B., N.G.S, F.L., N.S.M, S.E., K.E; Methodology: R.B., G.K., M.M., N.S.M, M.J.; Validation M.J.; Formal Analysis R.B., L.C.T., J.E.G.; Investigation: M.J., ; Resources: N.G.S, F.L., C.P., S.R., A.S., A.B., P.S, B. F., J.T.; Data Curation: R.B.; Writing –Original Draft: F.L., N.G.S, S.E., K.E.; Writing – Review & Editing Visualization: R.B, C.B.S-W, T.B., F.T., J.E.G.; Supervision: N.S.M., S.E., K.E.; Project Administration: N.S.M., S.E., K.E.; Funding Acquisition: K.E.

References

1. Rahwan I, Cebrian M, Obradovich N, Bongard J, Bonnefon JF, Breazeal C, et al. Machine behaviour. *Nature*. 2019;568(7753):477-86.
2. Esteva A, Kuprel B, Novoa RA, Ko J, Swetter SM, Blau HM, et al. Dermatologist-level classification of skin cancer with deep neural networks. *Nature*. 2017;542(7639):115-8.
3. Dey D, and Lin A. Machine-Learning CT-FFR and Extensive Coronary Calcium: Overcoming the Achilles Heel of Coronary Computed Tomography Angiography. *JACC Cardiovasc Imaging*. 2019.
4. McKinney SM, Sieniek M, Godbole V, Godwin J, Antropova N, Ashrafi H, et al. International evaluation of an AI system for breast cancer screening. *Nature*. 2020;577(7788):89-94.
5. Cristiano S, Leal A, Phallen J, Fiksel J, Adleff V, Bruhm DC, et al. Genome-wide cell-free DNA fragmentation in patients with cancer. *Nature*. 2019;570(7761):385-9.
6. Hayward NK, Wilmott JS, Waddell N, Johansson PA, Field MA, Nones K, et al. Whole-genome landscapes of major melanoma subtypes. *Nature*. 2017;545(7653):175-80.
7. Ott PA, Hu Z, Keskin DB, Shukla SA, Sun J, Bozym DJ, et al. An immunogenic personal neoantigen vaccine for patients with melanoma. *Nature*. 2017;547(7662):217-21.
8. Hilf N, Kuttruff-Coqui S, Frenzel K, Bukur V, Stevanovic S, Gouttefangeas C, et al. Actively personalized vaccination trial for newly diagnosed glioblastoma. *Nature*. 2019;565(7738):240-5.
9. Nathan C, and Ding A. Nonresolving inflammation. *Cell*. 2010;140(6):871-82.
10. Eyerich K, Brown SJ, Perez White BE, Tanaka RJ, Bissonette R, Dhar S, et al. Human and computational models of atopic dermatitis: A review and perspectives by an expert panel of the International Eczema Council. *J Allergy Clin Immunol*. 2019;143(1):36-45.
11. Hawkes JE, Adalsteinsson JA, Gudjonsson JE, and Ward NL. Research Techniques Made Simple: Murine Models of Human Psoriasis. *J Invest Dermatol*. 2018;138(1):e1-e8.
12. Choudhary N, Bhatt LK, and Prabhavalkar KS. Experimental animal models for rheumatoid arthritis. *Immunopharmacol Immunotoxicol*. 2018;40(3):193-200.

13. Eyerich K, and Eyerich S. Immune response patterns in non-communicable inflammatory skin diseases. *J Eur Acad Dermatol Venereol*. 2018;32(5):692-703.
14. Inkeles MS, Scumpia PO, Swindell WR, Lopez D, Teles RM, Graeber TG, et al. Comparison of molecular signatures from multiple skin diseases identifies mechanisms of immunopathogenesis. *J Invest Dermatol*. 2015;135(1):151-9.
15. Lauffer F, Jargosch M, Krause L, Garzorz-Stark N, Franz R, Roenneberg S, et al. Type I Immune Response Induces Keratinocyte Necroptosis and Is Associated with Interface Dermatitis. *J Invest Dermatol*. 2018;138(8):1785-94.
16. Shao S, Tsoi LC, Sarkar MK, Xing X, Xue K, Uppala R, et al. IFN-gamma enhances cell-mediated cytotoxicity against keratinocytes via JAK2/STAT1 in lichen planus. *Science translational medicine*. 2019;11(511).
17. Eyerich K, Eyerich S, and Biedermann T. The Multi-Modal Immune Pathogenesis of Atopic Eczema. *Trends Immunol*. 2015;36(12):788-801.
18. Brunner PM, Leung DYM, and Guttman-Yassky E. Immunologic, microbial, and epithelial interactions in atopic dermatitis. *Ann Allergy Asthma Immunol*. 2018;120(1):34-41.
19. Hawkes JE, Chan TC, and Krueger JG. Psoriasis pathogenesis and the development of novel targeted immune therapies. *J Allergy Clin Immunol*. 2017;140(3):645-53.
20. Gunther C, Bello-Fernandez C, Kopp T, Kund J, Carballido-Perrig N, Hinteregger S, et al. CCL18 is expressed in atopic dermatitis and mediates skin homing of human memory T cells. *J Immunol*. 2005;174(3):1723-8.
21. Weidinger S, Beck LA, Bieber T, Kabashima K, and Irvine AD. Atopic dermatitis. *Nat Rev Dis Primers*. 2018;4(1):1.
22. Steinke S, Zeidler C, Riepe C, Bruland P, Soto-Rey I, Storck M, et al. Humanistic burden of chronic pruritus in patients with inflammatory dermatoses: Results of the European Academy of Dermatology and Venereology Network on Assessment of Severity and Burden of Pruritus (PruNet) cross-sectional trial. *J Am Acad Dermatol*. 2018;79(3):457-63 e5.

23. Fernandez-Flores A, and Saeb-Lima M. Mucin as a diagnostic clue in dermatopathology. *J Cutan Pathol*. 2016;43(11):1005-16.
24. Lauffer F, Jargosch M, Baghin V, Krause L, Kempf W, Absmaier-Kijak M, et al. IL-17C amplifies epithelial inflammation in human psoriasis and atopic eczema. *J Eur Acad Dermatol Venereol*. 2020;34(4):800-9.
25. Cloutier A, Guindi C, Larivee P, Dubois CM, Amrani A, and McDonald PP. Inflammatory cytokine production by human neutrophils involves C/EBP transcription factors. *J Immunol*. 2009;182(1):563-71.
26. Zheng Y, Danilenko DM, Valdez P, Kasman I, Eastham-Anderson J, Wu J, et al. Interleukin-22, a T(H)17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature*. 2007;445(7128):648-51.
27. Moll R, Divo M, and Langbein L. The human keratins: biology and pathology. *Histochem Cell Biol*. 2008;129(6):705-33.
28. Damsky W, Wang A, Olamiju B, Peterson D, Galan A, and King B. Treatment of severe lichen planus with the JAK inhibitor tofacitinib. *J Allergy Clin Immunol*. 2020.
29. Mayer TZ, Simard FA, Cloutier A, Vardhan H, Dubois CM, and McDonald PP. The p38-MSK1 signaling cascade influences cytokine production through CREB and C/EBP factors in human neutrophils. *J Immunol*. 2013;191(8):4299-307.
30. Fernandez-Flores A. Lesions With an Epidermal Hyperplastic Pattern: Morphologic Clues in the Differential Diagnosis. *Am J Dermatopathol*. 2016;38(1):1-16; quiz 7-9.
31. Liu XB, Li F, Li YQ, and Yang F. Pituitary tumor transforming gene PTTG2 induces psoriasis by regulating vimentin and E-cadherin expression. *Int J Clin Exp Pathol*. 2015;8(9):10887-93.
32. Mendez-Vidal C, Gamez-Del Estal MM, Moreno-Mateos MA, Espina-Zambrano AG, Torres B, and Pintor-Toro JA. PTTG2 silencing results in induction of epithelial-to-mesenchymal transition and apoptosis. *Cell Death Dis*. 2013;4:e530.

33. Luke Y, Zaim H, Karakesisoglou I, Jaeger VM, Sellin L, Lu W, et al. Nesprin-2 Giant (NUANCE) maintains nuclear envelope architecture and composition in skin. *J Cell Sci.* 2008;121(11):1887-98.
34. Fyhrquist N, Muirhead G, Prast-Nielsen S, Jeanmougin M, Olah P, Skoog T, et al. Microbe-host interplay in atopic dermatitis and psoriasis. *Nature communications.* 2019;10(1):4703.
35. Rincon E, Rocha-Gregg BL, and Collins SR. A map of gene expression in neutrophil-like cell lines. *BMC Genomics.* 2018;19(1):573.

Figures and figure legends

Figure 1: Transcriptomic heterogeneity of ncISD requires deep clinical phenotyping

A) Principal component analysis (PCA) of gene expression from 287 lesional skin biopsies, spanning 13 different non-communicable inflammatory skin diseases. Each dot represents a patient colored by its diagnosis. PCA reveals three, albeit inconspicuous, subclusters consisting of lichen planus/ lupus (type 1 ncISD) (light and dark orange), nummular eczema/ atopic eczema (type 2 ncISD) (light purple/light blue) and psoriasis (type 3 ncISD) (brown) B) Overview of clinical attributes (n=62) grouped into morphology, comorbidity, histology, patient's history (history), and laboratory parameters (lab), used to obtain a deep clinical phenotype of each patient. C) Exemplary fingerprints of four patients (two atopic dermatitis and two psoriasis patients) attribute composition (radar plots), histological examination (middle) and clinical picture (right) demonstrating that two patients with a similar transcriptional profile also have a similar phenomic profile (radar plot) despite differing clinical diagnosis (patients highlighted in the circle). However, patients with the same diagnosis (e.g. atopic dermatitis as indicated) may have a heterogeneous clinical phenotype combined with transcriptional diversity. D) Transcriptional variance explained estimated using linear mixed effect model. Clinical attributes additively explain higher transcriptional variance (36.7%) compared to the clinician's diagnosis (7.8%) E) Overview of the contribution of each individual clinical attribute to the variance in the transcriptome data set. BMI: body mass index, GPT: Glutamat-Pyruvat-Transaminase, Hb: hemoglobin, IBD: inflammatory bowel disease, RCA: rhinoconjunctivitis allergica.

A) Schematic of the AuGER approach. 1st step: Definition of core attributes. The 62 attributes of all samples $i = \{1 \dots n\}$ were used to identify highly correlated attributes. From the highly correlated set of attributes one representative was selected based on clinical relevance, resulting in a set of 24 core attributes $j = \{1 \dots m\}$ and in the attribute matrix $A_{i,j}$. 2nd step: Gene-attribute association. First, RNASeq count data was split into lesional and non-lesional sets of k genes ($o = \{1 \dots k\}$, gene matrix $G_{i,k}$, respectively). Next, multivariate regression with elastic net penalty was used to identify the association between each core attribute (blue box) and gene expression (here illustrated for lesional skin only, orange box). The $m-1$ attributes other than response were used as covariates to control for the influence on response variable (dashed blue box). Regression or beta coefficients β_{attr} and β_{gene} (yellow box) for each predictor variable were estimated. Then, genes with a non-zero β_{gene} coefficient in lesional skin only were selected for each attribute and the association of genes and clinical attributes was visualized as a bipartite network. B) Exemplary highly correlated attributes belonging to type 1, type 2, or type 3 ncISD clusters with their core attribute name in italics. C) Network visualization of gene sets derived from AuGER. Attributes (green diamonds) and genes (blue ellipses) represent nodes, edge is drawn between an attribute and gene with non-zero regression coefficient, thickness of edges represent strength of an association derived from the frequency of association (see methods). Colors indicate core attributes of the three ncISD clusters

Figure 3: The attribute 'neutrophils'

A) A focus on the clinical attribute 'neutrophils' and its associated genes. Line thickness indicates the strength of gene-attribute association B) Distribution of the attribute 'neutrophils' across the 13 non-

communicable inflammatory skin diseases. Each bar represents one patient, and the height of the bar indicates the number of neutrophils present in skin histology with 0 = no, 1 = low, 2 = moderate, 3 = high numbers of lesional neutrophils (see Fig. S2). C) Hierarchical clustering of neutrophil-associated genes annotated with diagnosis (colors) and presence of neutrophils (grayscale). D) Gene ontology terms associated with the neutrophil gene set. CL: cutaneous lymphoma; CSB: cutaneous side effect of biologics; DE: dishydrotic eczema; GP: guttate psoriasis; Parapso: parapsoriasis; PP: psoriasis pustulosa; PRP: pityriasis rubra pilaris.

Figure 4: *CEBPB* regulates neutrophil granulocyte biology

A) Five genes with a strong association to the attribute 'neutrophils' and a yet unknown function in this field were selected and their relative gene expression was evaluated in primary human keratinocytes (stimulated with IL-17A and TNF- α) and neutrophil-like cells (differentiated HL60 cells) (stimulated with LPS) compared to non-stimulated controls by real time PCR. B) *CEBPB* and *PTTG2* were knocked out in primary human keratinocytes using the RNP-CRISPR technology. Knock-out (RNP) and wild-type (no-RNP) keratinocytes were cultured and stimulated with recombinant IL-17A for 72 hours to induce neutrophil-relevant genes. The supernatant was analyzed for cytokine/chemokine content by multiplex technology and relative changes are given in the graphs as percentage compared to no-RNP controls. C) Frequency of neutrophils migrated towards the supernatant of keratinocytes stimulated with recombinant IL-17A after *CEBPB* or *PTTG2* knock-out compared to IL-17A-stimulated wild type keratinocytes. D) Knockout of *CEBPB* or *PTTG2* in neutrophil-like cells (differentiated HL60

A) A focus on the clinical attribute ‘acanthosis’ and its associated genes. Line thickness indicates the strength of the gene-attribute association. B) Distribution of the attribute ‘acanthosis’ across the 13 non-communicable inflammatory skin diseases. Each bar represents one patient, and the height of the bar indicates the level of acanthosis measured by skin histology, with 0 = no, 1 = low-, 2 = moderate, 3 = strong acanthosis (see also Fig. S2). C) Hierarchical clustering of acanthosis-associated genes annotated with diagnosis (colors) and strength of acanthosis (grayscale). D) Gene ontology terms associated with the acanthosis gene set. CL: cutaneous lymphoma; CSB: cutaneous side effect of biologics; DE: dishydrotic eczema; GP: guttate psoriasis; Parapso: parapsoriasis; PP: psoriasis pustulosa; PRP: pityriasis rubra pilaris.

Figure 6: Validation of acanthosis-related genes

A) Genes with a strong and yet unknown association to the attribute acanthosis were chosen and their expression was evaluated in primary human keratinocytes after *IL-17A* and *TNF- α* stimulation by real time PCR. B) The acanthosis-associated genes *CEBPB*, *SYNE1* and *PTTG2* were knocked out in primary human keratinocytes using CRISPR-Cas9. 3D skin models of knock-out (RNP) and wild-type (no-RNP) were stimulated with recombinant *IL-22* to induce acanthosis or left untreated (control) and stained for the proliferation marker Ki67. Representative staining are shown. Bars represent 40 μ m. C) Relative delta thickness of 3D skin models with RNP-CRISPR knock-out for *CEBPB*, *SYNE1* or *PTTG2* (n = 11). Here, thickness of each *IL-22* stimulated model was normalized to its untreated control followed by calculation of percental changes of RNP vs no-RNP treated models. D) Quantification of Ki67 positive cells in 3D skin models shown in B). E) Representative Keratin 10 (KT10) (red), Keratin 16 (KT16) (yellow), and Ki67 (green) immunofluorescence staining of 3D skin models with a RNP-CRISPR knock out of *CEBPB* with or without (control) *IL-22* stimulation. Nuclei were stained with DAPI (blue). The dotted white line indicates the border between basal keratinocytes and the culture membrane. Results on *SYNE1* and *PTTG2* are given in Fig. S8. Bars represent 50 μ m. KT: keratin

Supplemental material - figures:

Figure S1: Impact of non-lesional skin and gender on transcriptomics-based clustering. Related to Figure 1.

A) Status of the skin (lesional, non-lesional) explains separation of PC1 and in total 32 % of variance in the gene expression data set. The sex (male, female) explains separation of PC2 and 9 % of data set variance. B) Loadings of the principal component analysis depicting the top 10 genes driving PC1 and PC2. C) Hierarchical clustering of gene expression data after correction for gender annotated with diagnosis.

	no.	clinical attribute	data collected	unit	data used in algorithm
age	1	age		years	
	2	arterial hypertension			
	3	arthritis			
	4	asthma			
comorbidity	5	BMI		in kg/m ²	
	6	diabetes mellitus			
	7	hepatitis			
	8	infect associated aggravation			
	9	inflammatory bowel diseases (IBD)			
	10	mucosa			
	11	nail involvement			
	12	rhinoconjunctivitis allergica (RCA)			
	13	renal disease			
	14	scalp involvement			
PGA	15	tonsillitis			
sex	16	PGA			
	17	sex			
histology	18	acanthosis qualitative			
	19	acanthosis quantitative		1-2-3	
	20	depth of lymphocyte infiltration into dermis			
	21	bacteria			
	22	dilated capillaries in papillary dermis			
	23	eosinophils		0-1-2-3	
	24	exocytosis		0-1-2-3	
	25	granulocytes		1-2-3-4-5	
	26	hyperkeratosis quantitative		0-1-2-3	
	27	interface dermatitis qualitative			
	28	interface dermatitis quantitative		0-1-2-3	
	29	keratosis		0-1-2-3	
	30	lymphocytosis		0-1-2-3	
	31	leucocytoclastic vasculitis			
	32	microabscesses			
	33	mucin		0-1-2-3	
	34	neutrophils qualitative			
	35	neutrophils quantitative		0-1-2-3	
	36	orthokeratosis quantitative		0-1-2-3	
	37	parakeratosis qualitative			
	38	parakeratosis quantitative		0-1-2-3	
patient's history	39	suprapapillary epidermis		0-1-2-3	
	40	serum crust			
	41	spongiosis			
	42	dermal localization of lymphocytes			
	43	alcohol consumption		0-1-2-3	
	44	chronic		number of weeks	
lab	45	exanthematic			
	46	family history			
	47	onset		0-1-2-3	
	48	progressive			
	49	pruritus		0-1-2-3	
	50	self limited			
morphology	51	smoker			
	52	GPT			
	53	Hb			
	54	leukocytes		in counts/ μ l	
	55	serum kreatinin		in mg/dl	
	56	dryness			
	57	elevation		0-1-2-3	
	58	erythema			
	59	papules			
	60	pustules			
	61	scaling		0-1-2-3	
	62	vesicles			

dichotomous variables

yes	no
-----	----

ordinal variables

3	2	1	0
---	---	---	---

continuous variables

--	--	--	--

Figure S2: Description of clinical attributes. Related to Figure 1.

In total, 62 clinical attributes were obtained from every patient included in the study. These attributes included sex, age, Physician Global Assessment (PGA), information on existing comorbidities (n = 14), histology (n = 23), patient's history (n = 9), laboratory parameters (lab) (n = 4) and morphology of the skin lesion (n = 7). Data was collected as dichotomous-, ordinal- or continuous variables as indicated. Framed attributes represent the 24 core attributes that were identified and assigned with a gene signature by the AuGER approach.

Figure S3: Fingerprints of 287 nclSD patients. Related to Figure 1

Heatmap illustrating each of the 287 patients of the study with his/her personalized clinical phenotype using all 62 attributes as explained in Figure 1B and Figure S2. Patients have been grouped according to their clinical diagnosis as indicated in the legend.

Figure S4: Overlap between the gene sets derived for each attribute using AuGER. Related to Figure 2.

The identified core attributes are given on the left and the total number of associated genes by the AuGER approach on the right. On the top, the number of genes are given that are either unique to one attribute (left row), shared between two attributes (middle row), or shared between three or four attributes (right rows). For instance, 102 genes were associated with the attribute ‘dryness’. Of these, 95 genes were unique to ‘dryness’ and the remaining 7 genes were shared with other attributes (with 5 genes shared between ‘dryness’ as well as one other attribute (i.e., ‘eosinophils’, ‘family history’, ‘orthokeratosis’, ‘acanthosis’ or ‘interface dermatitis’); one gene being shared with two other attributes (i.e., ‘eosinophils’ and ‘nail’), and one gene being shared with three other attributes (i.e., ‘eosinophils’, ‘smoker’ and ‘nail’). Overall, 95% of all genes were exclusively associated with one attribute, whereas only 34 genes (5 %) were associated with two, three or at most four attributes simultaneously, corroborating the specificity of the approach.

Figure S5: Pathway analysis of each attribute gene signature. Related to Figure 2.

A) Unique pathway annotation for each attribute. B) Pathways annotated to five or more attributes.

Dark grey (0) indicates no annotation, light grey (1) indicates annotation of an attribute.

Figure S6: CCL18 levels in lesional skin are associated with pruritic inflammatory skin diseases. Related to Figure 2.

A) CCL18 levels as detected by RNA sequencing from the main cohort of the manuscript, shown for atopic eczema, nummular eczema, and psoriasis patients. B) CCL18 levels in skin as detected by microarray in a validation cohort of healthy volunteers (n=38) and patients suffering from AE (n=27) and psoriasis (n=37) (left graph) and a RNAseq cohort of healthy volunteers (n=38) and lichen planus (n=20).

Figure S7: Controls to indicate efficient CRISPR-Cas9 knock-out. Related to figure 4 and 6

A) The SURVEYOR assay was used to determine the knock-out efficiency of SYNE1 and PTTG2 in human primary keratinocytes that have been used for 3D skin models. Due to the high GC content of the CEBPB mRNA, a PCR product could not have been established and analysis by the SURVEYOR assay was not possible. B) Knock-out efficiency of CEBPB determined by western blot analysis of unstimulated and IL-17/TNFα stimulated keratinocytes

Figure S8: Immunofluorescence staining of 3D skin models. Related to Figure 6.

Immunofluorescence staining with Keratin 10 (KT10) (red), Keratin 16 (KT16) (yellow), Ki67 (green) and DAPI (Blue) of 3D skin models with *SYNE1* or *PTTG2* CRISPR-Cas9 knock-out. Acanthosis was induced by stimulation with recombinant IL-22. Unstimulated models served as control. Bars represent 50 μ m. Representative Keratin 10 (KT10) (red), Keratin 16 (KT16) (yellow), and Ki67 (green) immunofluorescence staining of 3D skin models with a RNP-CRISPR knock out of *SYNE1* or *PTTG2* with or without (control) IL-22 stimulation. Nuclei were stained with DAPI (blue). The dotted white line indicates the border between basal keratinocytes and the culture membrane. Results on CEBPB are given in Fig. 6 of the main manuscript. Bars represent 50 μ m. KT: keratin

Supplemental material - tables:

Table S1: Patient characteristics

ID	gender	age	diagnosis	BMI	PGA	lesional biopsy	non-lesional biopsy
10178	M	41,96	cutaneous lupus	23	3	x	x
10201	F	17,35	cutaneous lupus	21,7	2	x	x
10213	M	66,03	cutaneous lupus	28,1	3	x	x
10217	F	68,15	cutaneous lupus	25,6	3	x	x
10322	F	52,73	cutaneous lupus	18,2	2	x	
10359	F	69,03	cutaneous lupus	18,6	3	x	x
10381	F	36,01	cutaneous lupus	22,8	3	x	x
10409	F	59,80	cutaneous lupus	20,8	4	x	x
10418	M	26,94	cutaneous lupus	27,8	1	x	x
10421	F	44,47	cutaneous lupus	23,3	4	x	x
10710	F	54,30	cutaneous lupus	24,2	3	x	x
<i>n=11</i>	<i>F: 8; M: 3</i>	<i>48.8 ±17.16</i>		<i>23.1 ±3.26</i>	<i>2.82 ±0.87</i>		
10005	M	44,79	lichen planus	27,7	3	x	x
10009	F	79,30	lichen planus	23,6	3	x	x
10016	F	43,92	lichen planus	18,4	4	x	x
10053	M	51,07	lichen planus		3	x	x
10069	F	65,30	lichen planus	27	4	x	x
10074	M	71,55	lichen planus		2	x	x
10075	F	51,70	lichen planus		2	x	x
10076	F	75,48	lichen planus		2	x	x
10104	M	43,22	lichen planus		2	x	x
10109	M	59,93	lichen planus	32	4	x	x
10113	M	65,17	lichen planus	24,6	3	x	x
10174	F	39,04	lichen planus	19,4	3	x	x
10212	F	74,00	lichen planus	24	2	x	x
10317	M	61,83	lichen planus	32,6	3	x	x
10318	F	54,53	lichen planus	25,2	3	x	x
10340	F	65,22	lichen planus	28,2	4	x	x
10347	F	78,65	lichen planus	25,7	4	x	x
10382	M	63,89	lichen planus	36,8	2	x	x
10422	M	54,78	lichen planus	34	3	x	x
10435	F	33,83	lichen planus	21,1	4	x	x
10458	F	57,63	lichen planus		3	x	x
10492	F	84,65	lichen planus	22,9	2	x	x
10496	M	56,15	lichen planus	30	4	x	x
10532	F	70,61	lichen planus	28,7	2	x	x
10576	F	54,70	lichen planus	32,8	3	x	x
10613	M	28,92	lichen planus	24,2	3	x	x
10622	M	72,18	lichen planus	25,5	3	x	x
10640	F	54,00	lichen planus	26,2	3	x	
10687	F	65,83	lichen planus	25,4	2	x	x

10705	M	43,18	lichen planus	22,2	3	x	x
<i>n=30</i>	<i>F: 17; M: 13</i>	<i>58.83 ±14</i>		<i>26.59 ±4.64</i>	<i>2.93 ±0.74</i>		
10004	F	27,85	AD	19,4	4	x	x
10007	M	27,48	AD	19	4	x	x
10020	F	22,49	AD	20	3	x	x
10025	M	23,61	AD	22,7	4	x	x
10043	M	72,33	AD	19,8	4	x	x
10063	F	27,10	AD	21,97	3	x	x
10085	F	35,09	AD	32	4	x	x
10089	M	62,30	AD		3	x	x
10137	M	29,01	AD	25,8	4	x	x
10146	M	25,89	AD	21	3	x	x
10248	F	53,13	AD	27,8		x	x
10264	M	57,30	AD		3	x	x
10269	M	54,18	AD	21		x	x
10294	M	50,01	AD	24,6	3	x	x
10321	M	79,55	AD		3	x	x
10393	M	39,44	AD	32,6	4	x	x
10401	M	77,67	AD	20,8	3	x	x
10416	M	61,81	AD	30	4	x	x
10442	M	22,75	AD	34,1	3	x	x
10452	M	51,21	AD	20,1	3	x	x
10462	F	66,44	AD	24,7	3	x	x
10655	M	53,69	AD	32,1		x	x
10671	F	23,90	AD	22,1		x	
10682	F	21,50	AD	26,6		x	x
13237	M	56,01	AD	30,1	4	x	x
16083	M	31,45	AD	24,57	4	x	x
24627	F	47,59	AD	34,19	3	x	x
28093	F	26,15	AD	21,33	3	x	x
37685	M	38,74	AD	20,2	3	x	x
48724	M	49,59	AD	21,96	4	x	x
51694	M	55,98	AD	30,75	3	x	x
57856	M	36,77	AD	23,95	3	x	x
71982	M	41,64	AD	29,72	3	x	x
83239	M	33,07	AD	15,56	4	x	x
10464	M	56,69	AD	23,5	3	x	x
10570	F	45,04	AD	31,7		x	x
10367	M	68,47	AD	21,2	4	x	x
10035	F	51,51	AD	45,7	4	x	x
10071	F	79,26	AD	23,5	4	x	x
10091	M	81,55	AD		2	x	x
10147	M	25,31	AD	23	3	x	x
10149	F	87,00	AD	23	4	x	x
10175	M	23,22	AD	27	3	x	x

10320	M	47,50	AD	30,1	3	x	x
10498	M	74,21	AD	26,5	3	x	x
10654	M	56,75	AD	28,7		x	x
10664	F	73,99	AD	23		x	x
<i>n=47</i>	<i>F: 15; M: 32</i>	<i>47.94 ±19.22</i>		<i>25.52 ±5.63</i>	<i>3.38 ±0.54</i>		
10166	F	79,27	DE	26	4	x	x
10223	M	82,09	DE		3	x	x
10410	M	52,49	DE	27,8	3	x	x
<i>n=3</i>	<i>F:1, M: 2</i>	<i>71.28 ±16.33</i>		<i>26.9 ±1.27</i>	<i>2.22 ±0.58</i>		
10028	M	66,52	HRE	21,6	3	x	x
10073	M	21,93	HRE	28,4	4	x	x
10082	M	29,72	HRE	32,7	1	x	x
<i>n=3</i>	<i>F:0, M: 3</i>	<i>39.39 ±23.81</i>		<i>27.57 ±5.6</i>	<i>2.67 ±1.53</i>		
10021	F	66,20	nummular eczema		4	x	x
10084	F	51,72	nummular eczema		2	x	x
10144	F	55,46	nummular eczema	22,3	3	x	x
10150	M	67,60	nummular eczema	23	2	x	x
10165	M	60,35	nummular eczema	18	4	x	x
10220	M	35,70	nummular eczema	44,2	4	x	x
10309	M	78,82	nummular eczema	31,5	2	x	x
10314	F	71,42	nummular eczema		2	x	x
10355	F	87,65	nummular eczema	29,7	3	x	x
10361	F	38,99	nummular eczema	29,8	3	x	x
10387	M	66,60	nummular eczema	45,6	2	x	x
10389	M	49,12	nummular eczema	19,6	3	x	x
10400	M	71,40	nummular eczema	27	3	x	x
10404	F	67,17	nummular eczema	25,9	3	x	x
10406	M	58,18	nummular eczema	30,7	3	x	x
10413	M	33,40	nummular eczema	32	3	x	x
10417	M	74,37	nummular eczema	29,4	3	x	x
10426	F	81,35	nummular eczema	27,1	3	x	x
10443	M	79,37	nummular eczema	27,6	3	x	x
10446	F	79,82	nummular eczema	23,2	3	x	x
10450	F	69,04	nummular eczema	27	3	x	x
10453	F	89,89	nummular eczema	20,5	4	x	x
10455	M	63,95	nummular eczema	42,6	3	x	x
10466	F	29,63	nummular eczema	18,4	2	x	x
10479	F	59,07	nummular eczema	31,1	3	x	x
10481	M	24,57	nummular eczema	20,7	3	x	x
10483	F	77,36	nummular eczema	23,1	2	x	x
10528	M	54,09	nummular eczema	34	3	x	x
10535	M	68,29	nummular eczema	23,8	3	x	x
10539	M	63,14	nummular eczema	29,4	2	x	x

10544	F	48,98	nummular eczema		2	x	x
10558	M	90,76	nummular eczema	22,1		x	x
10573	F	40,55	nummular eczema	29,7	2	x	x
10577	F	83,80	nummular eczema	22,6	3	x	x
10580	M	28,11	nummular eczema	21,7	3	x	x
10581	M	60,12	nummular eczema	27,8	3	x	x
10592	M	35,25	nummular eczema	26,5	3	x	x
10596	F	54,38	nummular eczema		3	x	x
10598	M	52,50	nummular eczema		3	x	x
10599	M	69,40	nummular eczema		3	x	x
10601	M	31,25	nummular eczema	40,23		x	x
10610	F	67,99	nummular eczema	31,6		x	x
10624	M	82,92	nummular eczema	25,3	3	x	x
10634	F	64,78	nummular eczema		3	x	x
10636	M	72,95	nummular eczema		3	x	x
10637	M	63,49	nummular eczema	32,3	3	x	x
10643	M	55,30	nummular eczema			x	x
10644	F	70,49	nummular eczema	25,2		x	x
10659	F	71,22	nummular eczema	26,9		x	x
10670	M	70,32	nummular eczema	25		x	x
10685	M	54,87	nummular eczema		3	x	x
<i>n=51</i>	<i>F: 22, M:29</i>	<i>61.63 ±16.87</i>		<i>27.85 ±6.59</i>	<i>2.86 ±0.55</i>		
10366	M	82,30	cut. lymphoma	24,2	3	x	x
10390	F	85,65	cut. lymphoma		4	x	x
10392	F	92,91	cut. lymphoma	31,2	2	x	x
10402	M	84,29	cut. lymphoma	25,5	1	x	x
10407	M	75,51	cut. lymphoma		4	x	x
10430	M	52,15	cut. lymphoma	27,7	1	x	x
10440	M	70,38	cut. lymphoma	25,8	2	x	x
10467	M	61,97	cut. lymphoma		2	x	x
10504	M	88,62	cut. lymphoma	29,9	2	x	x
10549	F	77,23	cut. lymphoma	25,3	3	x	x
HG16.1069	F	86,53	cut. lymphoma	24,2	1	x	x
HG16.1099	M	73,49	cut. lymphoma	22,7	2	x	x
HG16.1628	M	46,38	cut. lymphoma	24	1	x	x
HG16.537	M	68,15	cut. lymphoma	22	1	x	x
HG16.668	M	63,35	cut. lymphoma	22,1	1	x	x
HG16.711	F	70,83	cut. lymphoma	23,4	2	x	x
HG16.734	F	82,26	cut. lymphoma	29	1	x	x
HG16.975	F	78,57	cut. lymphoma	23	1	x	x
10607	F	43,13	cut. lymphoma	25,4	3	x	x
10658	M	86,45	cut. lymphoma	23,7	4	x	
<i>n=20</i>	<i>F: 8; M: 12</i>	<i>73.51 ±14.14</i>		<i>25.24 ±2.73</i>	<i>2.05 ±1.09</i>		
10283	M	36,69	CSB	23,9	1	x	x

10371	F	39,96	CSB	21,9	2	x	x
10349	M	65,52	CSB	24,4	3	x	x
10632	M	45,19	CSB	29		x	x
10353	F	44,66	CSB		3	x	x
10344	F	58,66	CSB		2	x	x
10346	F	30,58	CSB		3	x	x
10383	M	25,42	CSB	19	3	x	x
10388	F	28,05	CSB	17	2	x	x
10420	M	41,25	CSB	29,3	3	x	x
10447	F	57,53	CSB		2	x	x
10541	F	47,44	CSB	30	3	x	x
<i>n=12</i>	<i>F: 7; M: 5</i>	43.41 <i>±12.53</i>		24.31 <i>±4.88</i>	2.45 <i>±0.69</i>		
10002	F	18,33	psoriasis	26,8	4	x	x
10006	M	42,72	psoriasis	37	4	x	x
10013	M	50,26	psoriasis	33,6	4	x	x
10030	M	38,26	psoriasis	30,6	4	x	x
10033	M	67,58	psoriasis	32,9	4	x	x
10034	F	67,48	psoriasis	29,6	4	x	x
10036	M	52,63	psoriasis	36,7	4	x	x
10046	M	54,78	psoriasis	32,2	3	x	x
10049	M	45,79	psoriasis	22,8	3	x	x
10051	F	42,20	psoriasis	23,2	3	x	x
10052	F	90,62	psoriasis	33,2	3	x	x
10056	M	22,99	psoriasis	22,6	3	x	x
10058	F	70,31	psoriasis	43,7	4	x	x
10060	M	52,93	psoriasis	24	4	x	x
10067	M	42,30	psoriasis	21,9	3	x	x
10072	F	31,69	psoriasis	22	3	x	x
10078	M	51,36	psoriasis	26,1		x	
10079	M	24,72	psoriasis		3	x	x
10083	F	41,79	psoriasis	40,3	4	x	x
10103	M	55,23	psoriasis	31,7	4	x	x
10119	F	57,01	psoriasis	23	4	x	x
10128	F	53,49	psoriasis	30	3	x	x
10133	M	63,70	psoriasis	31,2	3	x	x
10138	F	68,62	psoriasis	21	3	x	x
10153	F	46,87	psoriasis	33	3	x	x
10154	F	55,43	psoriasis	47	4	x	x
10158	M	55,42	psoriasis	24,6	2	x	x
10171	M	56,98	psoriasis	31	2	x	x
10172	F	90,97	psoriasis	26	2	x	x
10195	M	47,40	psoriasis	21,4	3	x	x
10198	M	54,32	psoriasis	29	3	x	x
10199	F	44,12	psoriasis	34	4	x	x
10200	M	38,52	psoriasis	32	4	x	x
10203	M	49,74	psoriasis	26	4	x	x

10206	F	27,58	psoriasis	29	3	x	x
10208	F	65,75	psoriasis	26,7	3	x	x
10210	F	50,70	psoriasis		2	x	x
10227	M	54,64	psoriasis	29,3	3	x	x
10233	M	58,41	psoriasis	23,1	4	x	x
10239	F	88,72	psoriasis	21,9	4	x	x
10242	F	65,87	psoriasis	34,7		x	x
10249	M	72,65	psoriasis	28,7		x	x
10282	M	38,66	psoriasis	26,3	1	x	x
10297	F	66,59	psoriasis	26,4	2	x	x
10298	M	71,42	psoriasis	29,6	3	x	x
10307	M	46,40	psoriasis	32,2	4	x	x
10327	M	44,07	psoriasis	32,7	2	x	x
10343	M	76,58	psoriasis		3	x	x
10354	M	74,73	psoriasis	23,7	4	x	x
10384	M	47,07	psoriasis	39,1	4	x	x
10429	M	60,89	psoriasis		3	x	x
10456	M	33,87	psoriasis	29	3	x	x
10465	M	49,82	psoriasis	23,8	3	x	x
10489	F	84,55	psoriasis		2	x	x
10542	M	57,34	psoriasis			x	
10562	F	42,45	psoriasis	23,2		x	x
10574	M	39,34	psoriasis	20,8		x	x
10578	M	51,88	psoriasis	29,4		x	x
10590	M	61,66	psoriasis	29,9		x	x
10602	M	55,67	psoriasis	21,6		x	x
10609	F	73,35	psoriasis	29,8		x	x
10615	M	66,90	psoriasis	26,7		x	x
10620	M	36,87	psoriasis	29,8		x	x
10630	F	58,20	psoriasis	21,5		x	x
10646	M	58,55	psoriasis	26		x	x
10649	M	53,56	psoriasis	25,4		x	x
10650	F	43,52	psoriasis	29,7		x	x
10656	F	48,49	psoriasis	27,1		x	x
10665	M	25,84	psoriasis	32,4		x	x
10668	F	25,09	psoriasis	26,4		x	x
10691	M	46,75	psoriasis			x	x
10350	F	73,52	psoriasis		3	x	x
10419	F	73,67	psoriasis		3	x	x
10064	F	92,21	psoriasis	31,6	2	x	x
10136	F	67,50	psoriasis	30	3	x	x
10626	M	68,13	psoriasis	22		x	x
10177	M	51,57	psoriasis	34	2	x	x
10369	F	51,61	psoriasis	17,1	3	x	x
10372	M	78,22	psoriasis	35,5	3	x	x
10379	F	68,84	psoriasis	36,3	3	x	x
10482	F	58,82	psoriasis		3	x	x

10631	M	24,64	psoriasis	35,6		x	x
10633	F	37,32	psoriasis	24,8		x	x
10173	F	56,78	psoriasis	36	4	x	x
10211	M	39,41	psoriasis	40,35	4	x	x
10616	M	67,34	psoriasis	27,8	4	x	x
10645	F	59,98	psoriasis	32,9		x	x
10019	M	59,32	psoriasis	31,1	4	x	x
10159	F	53,41	psoriasis	36,3	4	x	x
10386	M	81,76	psoriasis	22,3	2	x	x
10461	F	53,80	psoriasis	48	3	x	x
10235	F	42,82	psoriasis	20,9	1	x	x
<i>n= 92</i>	<i>F: 40; M: 52</i>	54.69 ±16.05		29.25 ±6.14	3.18 ±0.79		
10255	M	46,76	Parapsoriasis		3	x	x
10439	M	67,28	Parapsoriasis	27,2	1	x	x
10449	M	57,99	Parapsoriasis	27,5	1	x	x
<i>n= 3</i>	<i>F: 0; M: 3</i>	57.34 ±10.27		27.35 ±0.21	1.67 ±1.15		
10087	F	32,33	pso guttata	21	2	x	x
10107	F	48,88	pso guttata	21,2	2	x	x
10205	M	46,44	pso guttata	26	3	x	x
10291	M	37,92	pso guttata	25,5	2	x	x
10296	M	34,21	pso guttata		2	x	x
10471	F	34,98	pso guttata	19,3	2	x	x
10276	M	28,77	pso guttata		4	x	x
<i>n=7</i>	<i>F: 3; M: 4</i>	37.65 ±7.41		22.6 ±2.97	2.43 ±0.79		
10086	M	65,12	PRP		3	x	x
10478	M	77,36	PRP	26,1	4	x	x
10524	F	76,73	PRP		3	x	x
10561	F	65,12	PRP	28	3	x	x
10648	M	69,69	PRP	19,7	2,5	x	
10666	M	62,00	PRP	24,8	3	x	x
10690	M	65,26	PRP	25,7	4	x	x
10721	F	46,38	PRP	28,7	3	x	x
<i>n=8</i>	<i>F: 2; M:5</i>	65.96 ±9.72		25.5 ±3.19	3.19 ±0.53		

Table S2: crRNA sequences.

Target	Name	Sequence (5'-3')	PAM
CEBPB	Hs.Cas9.CEBPB.1.AB	GGCCAACTTCTACTACGAGG	GGG
	Hs.Cas9.CEBPB.1.AA	CTCTTCTCCGACGACTACGG	GGG
SYNE1	Hs.Cas9.SYNE1.1.AC	ACGGCACTCAAGTTCCTCGA	AGG
	Hs.Cas9.SYNE1.1.AF	AGATCGTCCACCACCATTGG	AGG

<i>PTTG2</i>	Hs.Cas9.PTTG2.1.AA	TACTCTGATCTACGTTGATA	AGG
	Hs.Cas9.PTTG2.1.AB	CAAGTTTTAACACGACGTTT	TGG

Table S3: Primer sequences.

Gene	Sequence (5'-3')	
<i>CEBPB</i>	Fw	GGGAGCCCGTCGGTAATTTT
	rv	CATGTGCGGTTGGTTTGGAC
<i>SYNE1</i>	Fw	AGGAAAGGAGCATGTCTTCAGT
	rv	GCCACTTTTCGACCTCTTGG
<i>PTTG2</i>	fw	GATGTGCTGAAGCTGGAGTCT
	rv	AAGGCTGATGGAGCATCGTAT
<i>IL19</i>	fw	ATCATCAGGAGCCAAACCCC
	rv	TCTGTTCTGACATTGCCGC
<i>HCAR3</i>	fw	ATGACATAAAGGCAGGCACG
	rv	CTCGCCTAGTGAATGCTCCA
<i>SHC1</i>	fw	GGGAAAGTGGGAAGCGTGAA
	rv	TACACTCACTCCGGCCTCA
<i>HS6ST1</i>	fw	CACGCCCAGGAAGTTCTACTA
	rv	TGCAACGACGTCTTCCACG
<i>SLC23A2</i>	fw	GGAATTACAAAGGTCGGCAGC
	rv	GAACCTCCCGATCATGCCCA

Table S4: Gene lists of each core attribute-gene association.

acanthosis	alcohol	CD44	TUBB8	SPRR2B	TTN	FERMT1	EIF4ENIF1	HS6ST1	EPDR1
ARHGAP26	AK127697	CYP4X1	V.alpha.immuno	DYNLL1.AS1	XIRP1	IDH3A	RHNO1	KCNJ2	ARHGAP31
BC051760	AB209061	ISG15	VAV3.AS1	L3MBTL3	ZNF618	LOC644838	STIL	PRSS22	GPR15
DSC2	LINC00278	MYOF	ZNF560	RUNX2	AFAP1.AS1	NFE2L2	TBX4	S100B	MMP28
EFCAB10	LOC100134368	NEFL	ZNF705A	ZBTB24	AK024936	SNORD109B	AK097242	SMYD3	PAEP
FAM43A	PHPT1	PCDH17	elevation	ZNF575	AK096335	TDP2	AP3M1	SPTSSA	PPTC7
LINC00624	SNCG	ABT1	hCPE.R	AASS	AK096792	BC016972	AX746839	VNN3	ADAMTSL3
PTTG2	TTTY10	AK055856	LINC00624	ARIH2OS	AK097184	BC132944	AZGP1	ZBTB40	AK055811
SYNE1	TXNRD3NB	AK090688	CCSAP	AY343892	AK125001	C21orf90	BC039516	ZNF439	ALDH3A1
TMPRSS11D	UTY	AK124939	GLCC11	C11orf71	BC137484	C5orf48	C3orf79	ZNF883	ANKRD33B
TMPRSS6	GPR22	ANKS6	LDLR	CAMK2G	BTG3	contactin.1	C8orf34	scalp	AX747031
C3orf72	MAPKAPK5	BC040865	UTP14A	CCDC42	C17orf76.AS1	DL491527	CACNA2D3	C1orf226	BC045578
FAM153B	RPS4Y1	BEST3	DSC2	CCKAR	C5orf48	DLL3	CD36	C2orf194	BC142949
GJB6	TMEM191C	C17orf99	LINC00649	CD27	CASP8	GPR27	DLX1	CEACAM1	BOK
KIAA0146	TMSB4Y	PCORF15	VSNL1	CLEC11	CCL5	GTSF1	DLX2	DMBX1	BRE.AS1
PGBD5	AK055570	CA8	ZNF606	DYRK2	CEACAM22P	HLA.H	EF537580	EPHA1.AS1	C2orf91
ZNF841	ANAPC10	CD244	ARNTL	FAIM3	CERKL	LOC285847	FLG	FDFT1	COL6A5
ZNF880	BC042649	CDK2	LOC150568	GCM1	CHRFAM7A	LOC339788	FLJ44838	HDC	CRHR2
ACTN2	BTG3	COL26A1	MIR31HG	IDS	CXCL9	MIR1343	FRMD6.AS2	HMGCS1	DMBX1
BC016035	C10orf35	COL4A3	RALB	LGALS9C	CYP2B7P1	MIRLET7I	GABRB1	HOXA13	EIF4E3
DQ572823	DDX11L2	CRHR1	SOGA2	LOC728637	DAPK1	NFKBIZ	LNK1	HPSE	FOSB
DQ573539	DDX3Y	DLG1.AS1	TARS	PLAC8	DDX28	PGLYRP3	LOC100128822	KCNT2	FST
NRBF2	DMRT2	DQ572285	TTC23L	SSH2	DQ786293	PSMD12	LOC100130264	LINC00637	GGT2
XPNPEP1	GTF2H5	DQ575786	eosinophils	TCRBV6S2A1N	FAM110C	PSORS1C2	MLIP	MSMO1	HABP4
AK127309	GUCY2C	DQ584906	COL6A6	TGM3	FKBP15	SAMD8	OPN1LW	PDHA1	IL15
CEBPB	ID12	DUSP15	AK124272	UGD	FLJ16124	THEG	RP1L1	PRMT2	KLF11
DTNA	KIAA0930	EEF1A2	ACE2	interface derm	GBP5	ZFP92	SGOL2	SLC18A2	MAPK9
GSAP	L3HYPDH	FGF9	ATXN1	C17orf66	HTT.AS1	ZNF137P	SLC30A5	TBX1	MYCN
KCNH1	LOC100652791	HOXA9	CCL18	CXCL11	JPH4	neutrophils	TMCO2	TMEM200A	PCDHGB2
LIFR	ODF3B	HP07349	COL6A5	EHF	LOC151475	CEBPB	ZNF732	TMPRSS11GP	PRRT4
MAT1A	OTOP1	HPR	CYSLTR2	IL22	LRP2	CXCL1	papules	BC040646	RAB43
SLC24A3	PAX9	IFI30	NGFRAP1	LGALS17A	LTF	HS6ST1	AASS	BTBD16	SEMA5B
SOST	ROGDI	IL13RA2	erythema	LOC728989	MARCH4	PLAT	ITGAM	C2orf26	SMARCD1
WNT5A	RPL2B	IL21R.AS1	CHRM4	SLC8A1	MIR1203	SLC23A2	PRKCB	CD1C	TBX6
ZNF160	SEMG2	ISY1.RAB43	HEPHL1	TGM6	NEFL	DTHD1	TBC1D5	CXorf23	TMEM150C
AL389983	SNORA2B	JA611241	PRSS22	TMEM164	NHS	TREM1	C1QB	DL490908	TSPAN10
BC039483	SNORD116.25	LINC00240	PYGL	WNT16	NUP214	HCAR3	PPP2R2A	MYCBP2.AS1	UPB1
CACNA2D1	SNORD75	LINC00622	RBM47	CTLA4	NXF3	IL19	pruritus	STIP1	ZNF155
CEBPD	SPTBN4	LOC101059948	SPRR2B	IGFL2	OR5M11	IL8	CCL18	TCRBV21S1	ZNF256
CNGA3	TM4SF4	LOC145845	TMPRSS4	LOC100130476	PIGX	SHC1	scaling	USP50	ZSCAN12P1
DGCR10	TMCC3	LOC284551	UPP1	TREM2	PKIA	ZNF350	ARHGAP31	AK127609	
DIO2	XIST	LOC348761	LIPG	AK096834	PRSS22	FXYD7	BC007984	ALOX15B	
DNAJB1	ZFY	LOC645895	SPRR2D	CHRM4	SLC26A9	MIR3978	BCL2	ATP1B1	
FYB	ZNF878	MAGEA8.AS1	exocytosis	ERMN	SPIINK2	POU5F2	COL19A1	CD1A	
HLA.DRB5	dryness	MIPEPP3	KDM5B.AS1	F3	TCHHL1	RFESD	GPX3	CGREF1	
IL15	COL6A5	MIR127	GLB1L3	FCRL6	TCR.alpha	SLCO4A1	HSPA12A	DQ576544	
LOC388692	COL6A6	MIR3692	MAP7	HKDC1	TMEM257	THEGL	LAMP2	DQ583161	
LOC401320	CXCR2	MIR4534	PAWR	HLA.DMA	TRAPPC3	AB074166	LCE3A	DQ600342	
MIR425	DOK7	MIR496	ENPP2	MRPS27	mucin	AFAP1L2	MARCKS	G3BP1	
MIR4287	HSD3B1	MIR661	LMO2	SERPING1	IRAK3	AK057897	SNX25	LOC728989	
MYH15	IGFL4	MT3	ARHGAP32	TBX21	TMEM132B	AK091996	SPRR2C	MSTN	
NFX1	LAMP5	MTRNR2L8	CPLX3	TSFM	naïl	BX648501	SPRR2D	NAT10	
PAFAH2	LILRA4	MYO10	EFEMP1	CHIT1	CLEC3A	C3orf20	TGM6	PLIN5	
PCSK5	LIMS3	NCAPD2	GIMAP2	CR627049	COL6A6	CDKAL1	TMC5	PPM1H	
PDE4DIP	OTOP3	NOMO3	GNS	CYP2C9	EPB41L4A	EFCAB10	AASS	PRSS21	
PHLPP2	PLAC8	NR0B1	KCTD4	DL492607	MCTP2	HTJA150	BC038535	RAB44	
PIM1	PLD4	OR6Q1	LGALS9B	DLGAP1.AS5	MID1	KCNJ2	FNIP2	RRP9	
PNMA3	SCUBE2	PCSK1N	LRRIQ4	DTHD1	PON2	KCTD1	IL36A	DNAJA3	
PROKR1	SLC10A3	PDE6G	MILR1	FAM58A	RAB27A	MIR3935	SNORA45	KATNAL1	
PSAPL1	SLC25A43	PHOSPHO1	P2RX5	GCNT1	TMPRSS11A	MIR4524A	TRPM6	KIF13A	
SCML4	VSTM4	PPP2R5D	RCSD1	GTSF1L	LOC729041	P4HB	DAW1	LPCAT3	
SIMC1	ACER2	PSPN	RNASE6	IMPDH2	SFXN3	PCDH9	LRG1	P2RY13	
SLA	CACNB4	RASGEF1C	ZBTB7C	ITGB7	GFRA2	SNX25	PYGO1	SOST	
SLC7A1	COPE	RHBG	family history	LAMC2	LRRC55	ZNF136	RBM47	FKBP4	
SNORD66	HSD3BP4	RSPO2	MGC12916	LCE3A	N4BP1	orthokeratosis	STAT3	HTATSF1P2	
SPATA13	LOC401321	SLC24A3	CDC42SE2	LOC100133920	NOS2	AF086294	ASPHD1	MIR377	
ST3GAL5	OR2T8	SPRR3	PLIN5	LOC646214	SPTLC2	SERPINA12	MIR3935	NSDHL	
THEMIS	NGFRAP1	TBX5.AS1	ZMAT3	NPY4R	AK056253	TMEM231	NES	smoker	
TIMM44	POU2F3	TMCO2	ARL4C	PRR9	CD1E	ZNF367	SYTL5	PTPN9	
TPBG	ANO1	TMX2.CTND1	PRKCB	SFMBT2	COL6A5	ANKRD34C	AX747401	SLC6A6	
UCA1	CCL26	TSHZ2	SLC29A3	SPRR2B	DCHS2	C1orf68	DQ599569	C3orf17	