Mutually opposing activity of PIN7 splicing isoforms is required for auxin-mediated tropic responses in *Arabidopsis thaliana*

Ivan Kashkan^{1,2}, Mónica Hrtyan², Katarzyna Retzer¹, Jana Humpolíčková³, Aswathy Jayasree², Roberta Filepová¹, Zuzana Vondráková¹, Siby Simon¹, Debbie Rombaut^{4,5}, Thomas B. Jacobs^{4,5}, Mikko J. Frilander⁶, Jan Hejátko², Jiří Friml⁷, Jan Petrášek¹, Kamil Růžicka^{1,2,8,*}

¹*Laboratory of Hormonal Regulations in Plants, Institute of Experimental Botany, Czech Academy of Sciences, 16502 Prague, Czech Republic*

²*Functional Genomics and Proteomics of Plants, Central European Institute of Technology and National Centre for Biomolecular Research, Masaryk University, 62500 Brno, Czech Republic*

³*Institute of Organic Chemistry and Biochemistry, Czech Academy of Sciences, 166 10 Prague 6, Czech Republic*

⁴*Department of Plant Biotechnology and Bioinformatics, Ghent University, 9052 Ghent, Belgium*

⁵*VIB Center for Plant Systems Biology, 9052 Ghent, Belgium*

⁶*Institute of Biotechnology, University of Helsinki, 00014 University of Helsinki, Finland*

⁷*Institute of Science and Technology (IST Austria), 3400 Klosterneuburg, Austria*

⁸*Lead Contact*

*Correspondence: kamil.ruzicka@ueb.cas.cz

23 SUMMARY

24 Advanced transcriptome sequencing has uncovered that the majority of eukaryotic genes
 25 undergo alternative splicing (AS). Nonetheless, little effort has been dedicated to investigating
 26 the functional relevance of particular splicing events, even those in the key developmental and
 27 hormonal regulators. Here we reveal, in the plant model *Arabidopsis thaliana*, that the *PIN7*
 28 gene, which encodes a polarly localized transporter for the phytohormone auxin, produces
 29 two evolutionary-conserved transcripts. These isoforms PIN7a and PIN7b, differing in a 4
 30 amino acid motif, are present at nearly equal levels in most cells. Although both variants do
 31 not differ in the subcellular localization and transport auxin with similar capacity, they closely
 32 associate and mutually influence their stability within the plasma membrane. Phenotypic
 33 complementation tests reveal that the functional contribution of PIN7b *per se* is minor but it
 34 markedly attenuates the prominent PIN7a activity, which is required for correct seedling
 35 apical hook formation and auxin-mediated tropic responses. These results establish alternative
 36 splicing of the PIN family as a conserved, functionally-relevant mechanism, unveiling an
 37 additional regulatory level of auxin-mediated plant development.

39 KEYWORDS

40 Auxin, auxin transport, PINs, alternative splicing, plant development, RNA processing, tropic
 41 responses

43 INTRODUCTION

44 Auxin is an essential phytohormone, which plays a role in nearly all aspects of plant
 45 development. To flexibly adapt to rapidly-changing environmental cues, directional auxin
 46 transport represents a highly dynamic means for triggering downstream morphogenetic
 47 processes. PIN-FORMED (PIN) auxin efflux carriers are among the key regulators in this
 48 respect. In the past years, many efforts uncovered multiple mechanisms operating
 49 transcriptionally or post-translationally on the capacity and directionality of PIN-mediated
 50 transport. However, little progress has been made in exploring the contribution of post-
 51 transcriptional regulation (Hrtyan et al., 2015; Adamowski and Friml, 2015).

52 Advances in high throughput sequencing have revealed unexpected complexity within
 53 eukaryotic transcriptomes by alternative splicing (AS). Although the majority of AS
 54 transcripts may be functionally neutral (Tress et al., 2017; Blencowe, 2017; Darracq and
 55 Adams, 2013; Chamala et al., 2015; Reddy et al., 2013), several detailed studies have
 56 highlighted a plausible role for numerous AS events in physiologically relevant contexts,

including those involved in plant developmental and hormonal pathways (Shang et al., 2017; Hrtyan et al., 2015; Staiger and Brown, 2013; Szakonyi and Duque, 2018). Earlier works have described auxin-related defects resulting from the aberrant function of several regulators of AS (Hrtyan et al., 2015; Kalyna et al., 2003; Tsugeki et al., 2015; Casson et al., 2009; Retzer et al., 2014; Bazin et al., 2018). AS changes subcellular localization of the auxin biosynthetic gene *YUCCA 4* (Kriechbaumer et al., 2012) and differential splicing of an exon (Marquez et al., 2015) inside the *AUXIN RESPONSE FACTOR 8* (*ARF8*) results in developmental changes of generative organs (Ghelli et al., 2018). Likewise, a splice variant of *MONOPTEROS* (*MP/ARF5*) acts as a noncanonical regulator of ovule development (Cucinotta et al., 2020). In addition, AS of the Major Facilitator Superfamily transporter *ZIFL1* interferes with auxin transport, influencing the stability of PINs on the plasma membrane (PM) (Remy et al., 2013). These lines of evidence indicate that AS is an important player in auxin-dependent processes. However, no coherent functional model of any auxin-related AS event has been provided so far.

Here, we have investigated the functional significance of the protein isoforms arising through AS of the *Arabidopsis thaliana* *PIN7* gene. *PIN7* is, together with *PIN3* and *PIN4*, a member of the *PIN3* clade of PIN auxin efflux carriers (Bennett et al., 2014), which are required for a broad range of morphogenetic and tropic processes (Adamowski and Friml, 2015). We provide evidence that the evolutionary conserved *PIN7* isoforms mutually influence their dynamics within PM and demonstrate that the coordinated action of both splice variants is required for auxin-mediated differential growth responses.

RESULTS

Arabidopsis PIN7 and *PIN4* produce two evolutionary conserved AS transcripts

Our previous survey (Hrtyan et al., 2015) revealed that several genes involved in auxin-dependent processes undergo AS. Among them, closely related paralogs from the *PIN3* clade of auxin transporters, *PIN4* and *PIN7* (but not *PIN3*), are regulated by the same type of AS (Petrasek et al., 2006; Bennett et al., 2014; Hrtyan et al., 2015). The resulting transcripts, denoted as *a* and *b*, differ in the AS donor site position in the first intron (Figure 1A). The differentially spliced region corresponds to a four amino acid motif inside the large internal hydrophilic loop (Nodzyński et al., 2016; Ganguly et al., 2014) of the integral PM transporter (Figure 1A and 1B). We examined the quantities of individual reads spanning the exon junctions in the respective gene region from the *Arabidopsis* root tip and in several other available transcriptomes from different tissues and organs (Cheng et al., 2017; Klepikova et

al., 2016; Ruzicka et al., 2017) (Figures 1C and Supplemental Table 1). We found that each of the *PIN4* and *PIN7* splice isoforms is expressed abundantly in all tissues, independently of the inspected data set. We also identified a minor *PIN4c* (Hrtyan et al., 2015; Marquez et al., 2012) transcript (but not corresponding *PIN7c*), which comprised around 3-7% of the *PIN4* exon1-exon2 spanning reads (Figures 1A and Supplemental Table 1). Other occasionally observed transcripts (also corresponding to the other exon junctions) were not seen among different RNA-seq data sets. It thereby appears that *PIN7* and *PIN4* are processed into two and three splice isoforms, respectively, and that *PIN7a* and *b* (or *PIN4a* and *b*) are expressed in most of the plant organs at comparable or nearly similar levels.

Functionally relevant AS events are commonly evolutionary conserved (Reddy et al., 2013; Keren et al., 2010), therefore we sought for available validated transcripts to determine whether similar splicing events occur in orthologous *PIN3* clade genes in other dicot species (Bennett et al., 2014; O’Leary et al., 2016). We found examples of such mRNAs besides members of the *Brassicaceae* family, for instance, in *Abrus precatorius* (*Fabaceae*), which documents the conservation of this AS event in rosids, a plant clade that diversified more than 100 million years ago (Li et al., 2019) (Figure 1D). *PIN4c* did not show any deeper evolutionary conservation. Thus, at least some genes of the *PIN3* clade are regulated by the same type of AS, across several plant families, which suggests that these AS events may have a relevant biological function.

***PIN7* splice isoforms are functionally distinct, based on the mutant phenotype rescue tests**

We expressed fluorescently-tagged cDNA versions of the respective *PIN7* and *PIN4* transcripts under their native promoters in *Arabidopsis thaliana*. Their overall expression patterns resembled those of the *PIN7-GFP* and *PIN4-GFP* lines made on the basis of the genomic sequence (Supplemental Figures 1A–1D). Next, we tested their ability to complement the phenotypes associated with the *PIN7* locus. The phototropic hypocotyl bending is among the classical assays for testing the activity of the *PIN3* clade proteins (Friml et al., 2002; Willige et al., 2013). As weak phenotypes of the *pin7-2* loss of function mutants are the result of redundancy with other genes from the *PIN3* clade (Blilou et al., 2005; Willige et al., 2013; Friml et al., 2003), we employed a triple *pin3-3 pin4-101 pin7-102* knock out (*pin347*) as a genetic background, which lacks the phototropic response almost completely (Willige et al., 2013). Here, *PIN7a-GFP* cDNA rescued the phototropic bending, while the *PIN7b-RFP* cDNA did not show any effect, regardless of whether the native (Figure

1E) or a stronger endodermal *SCR* promoter (Rakusová et al., 2011) (Supplemental Figure 1E) was used. The tag choice does not appear to have any effect in these assays, as evidenced by the lines where the fluorophore sequences have been swapped (Figure 1F). Together, these data indicate that the motif changed by AS alters the function of the PIN7 protein in *Arabidopsis*.

The PIN3 clade auxin efflux carriers have been implicated in many other instances of auxin-mediated development (Adamowski and Friml, 2015). We therefore hypothesized whether, in some of them, the role of the particular isoform could be prevalent. These processes include: determining of root protoxylem formation (Bishopp et al., 2011) (Figure 1G), lateral root density (Swarup et al., 2008) (Figure 1H), vertical direction of the root growth (Kleine-Vehn et al., 2010; Friml et al., 2002; Pernisova et al., 2016) (Supplemental Figure 1F), lateral root orthogravitropism (Rosquete et al., 2013) (Supplemental Figure 1G), gravity-induced hypocotyl bending (Rakusová et al., 2011) (Supplemental Figure 1H), number of rosette branches after decapitation (Bennett et al., 2016) (Supplemental Figure 1I) and the overall rosette size (Bennett et al., 2016) (Supplemental Figure 1J). In sum, the *PIN7a-GFP* cDNA usually almost completely rescues the *pin347* phenotypes, while the contribution of *PIN7b-RFP* is smaller or even undetectable.

Fluorescent reporters reveal highly overlapping *PIN7a* and *b* expression

The levels of *PIN7* (and *PIN4*) AS transcripts seem to be at comparable levels in most organs (Figure 1C and Supplemental Table 1). However, this may not describe the actual situation at the resolution of individual cells. To address this, we analyzed the P7A₁G and P7BR fluorescent reporters (Kashkan et al., 2020), which allow for monitoring the activity of the AS of *PIN7* *in planta* and *in situ* (Figure 2A). Indeed, in most cells, including the primary root tip (Figure 2B) or in the hypocotyl during the light bending assay (Figure 2C), we observed generally overlapping expression of both isoforms without any apparent tissue preference. However, there were several instances in the vegetative tissue, where the ratio of reporter signals appears to be uneven. These include early lateral root primordia (Supplemental Figure 1K), the mature pericycle adjacent to the phloem area (Supplemental Figure 1L), the stomatal lineage ground cells of the cotyledons (Supplemental Figure 1M), and the concave side of the opening apical hook (Figure 2D), where the *PIN7b-RFP* signal prevailed over that of *PIN7a-GFP*. In general, these data corroborate the presence of both isoforms in most cells and suggest that they may function in a coordinated manner.

The combined activity of both PIN7a and PIN7b is required for apical hook formation and tropic responses

The generally overlapping expression and also an occasionally exaggerated response of the *pin347* mutants containing the *PIN7a-GFP* transgene (Supplemental Figure 1H) prompted us to carefully record the temporal dynamics of the processes linked with the PIN3 clade function. We were even able to capture subtle phenotypic changes observed in the single *pin7-2* knock out allele (Friml et al., 2003), including those complemented with the *PIN7* cDNAs during hypocotyl phototropism (Supplemental Figure 2A) or apical hook formation (Supplemental Figure 2B), where the time scale measurements of the PIN-mediated morphogenesis have been well established (Zadnikova et al., 2010). Next, we analyzed apical hook development in the *pin4-101 pin7-102 (pin34)* mutants and a newly generated *pin347* line that carried a combination of both *PIN7* cDNA constructs. Similar to the gravi- and phototropic experiments, the *PIN7b-RFP* cDNA generally did not complement the severe *pin347* phenotype. Expression of the *PIN7a-GFP* cDNA in *pin347* indeed led to partial rescue of the phenotypic defects, even surpassing the values observed for *pin34*. Surprisingly, the simultaneous expression of both *PIN7a-GFP* and *PIN7b-RFP* suppressed the dominant effects conferred by the *PIN7a-GFP* cDNA alone and phenocopied the *pin34* mutant (Figure 3A; this effect was not caused by suppressing the expression of PIN7a-GFP by the other transgene, Supplemental Figure 2C). This strongly suggests that both isoforms act in a mutually opposing manner to modulate apical hook development.

Apical hook development is a complex process that involves several bending steps (Zadnikova et al., 2010), and the splicing reporter suggests a slightly different expression pattern of *PIN7a* and *b* isoforms during apical hook development (Figure 2D). The hypocotyl photo- and gravitropic response includes only a single bending (Rakusová et al., 2011, 2016), and the reporter expression appears to be uniform in the respective tissues (Figure 2C). It can thus provide a hint to whether one can account for the antagonistic behavior of both isoforms to differential expression or to another mechanism, likely occurring at the cellular level. Similar to apical hook development (Figure 3A), introducing the *PIN7b-RFP* cDNA had a marginal effect on the *pin347* phenotype, while the expression of *PIN7a-GFP* lead to faster bending than that observed for the *pin34* mutant. Finally, the presence of both *PIN7a-GFP* and *PIN7b-RFP* cDNAs in *pin347* was reminiscent of the *pin34* phenotype in both phototropic and gravity assays (Figures 3B and 3C). We therefore conclude that the shared activity of both PIN7 isoforms, located probably in the same group of cells, is required for proper apical hook formation and auxin-mediated tropic responses.

PIN7a, but not PIN7b, is required for proper formation of auxin maxima

To further investigate the functional impact of AS on PIN7, we validated the formation of the downstream auxin response maxima in apical hooks by crossing the *DR5v2::GFP* transcriptional auxin reporter (Liao et al., 2015) into lines where the phenotypic changes have been temporally monitored (see Figure 3). The reporter expression pattern in the forming apical hooks of *pin34*, *PIN7a-GFP pin347* and *PIN7a-GFP PIN7b-RFP pin347* resembled the situation in the wild type (Zadnikova et al., 2016), whereas the *DR5v2::GFP* distribution in the *PIN7b-RFP pin347* line was indistinguishable from that of *pin347* (Figure 4A). These results underline that the PIN7b isoform alone is unable to establish the morphogenic auxin gradients and that the observed developmental changes occur in line with previously described mechanisms (Friml et al., 2002; Willige et al., 2013; Zadnikova et al., 2010, 2016).

PIN7a and PIN7b transport auxin with comparable rates in tobacco cells

Because PIN7a and PIN7b showed differential ability to generate auxin maxima *in planta*, we asked whether both protein isoforms indeed function as auxin transporters. We expressed the *PIN7a* and *b* cDNA variants tagged with GFP under the control of the β -estradiol-driven promoter in the tobacco BY-2 tissue culture system (Petrasek et al., 2006; Müller et al., 2019). We confirmed that PIN7a (Petrasek et al., 2006), as well as the PIN7b isoform, are functional auxin transporters. To this end, we also analyzed the quantitative aspects of PIN7-mediated transport, using multiple concentrations of β -estradiol. Following induction of both transgenes, we observed a comparable decrease of radioactive-labeled auxin accumulation inside the BY-2 cells for both lines in the β -estradiol dose range used (Figure 4B), consistent with the similar PIN7a and PIN7b protein levels (Supplemental Figures 3A–3C). The time course of the auxin accumulation drop appeared to be similar for both constructs as well (Figure 4B). Hence, *PIN7a* and *PIN7b* code for true auxin exporters, and they seem to transport auxin in tobacco cell cultures at similar rates.

PIN7a and PIN7b differ in protein subcellular dynamics

Polarity and dynamic intracellular trafficking are essential functional attributes of PINs (Adamowski and Friml, 2015). However, the PIN7 isoforms did not prominently differ from each other in terms of polarity or general subcellular localization in the root tip at a given resolution limit (Figure 4C). The anterograde trafficking of proteins towards PM can be

effectively blocked by the fungal toxin brefeldin A (BFA). It leads to internal accumulation of the membrane-bound PINs into characteristic BFA bodies (Geldner et al., 2001; Kleine-Vehn et al., 2010). During time-lapse imaging, we observed that while PIN7a-GFP accumulated readily in these intracellular aggregates, the PIN7b-RFP BFA bodies were less pronounced, co-localizing with PIN7a-GFP incompletely (Figure 4C). To exclude the effects of diverse fluorescent tags, we compared the response of PIN7a-GFP to the PIN7b-GFP cDNA lines. The BFA mediated aggregation of PIN7b-GFP indeed showed a moderate delay, compared with PIN7a-GFP (Figure 4D). It suggests that the PIN7 isoforms differ in the speed of their intracellular trafficking pathways or delivery to the PM, and the choice of the tag does not appear to interfere significantly with the subcellular dynamics of PIN7.

PIN polarity does not seem to strictly require the cytoskeleton (Glanc et al., 2019), but subcellular PIN trafficking has been proposed to be mediated by two distinct pathways (Geldner et al., 2001; Glanc et al., 2019). The first depends on actin filaments (cytochalasin D-sensitive) and occurs in most of the root meristem cells. The second (oryzalin-sensitive) utilizes microtubules and is linked with cytokinesis. Drugs that depolymerize actin filaments (cytochalasin D) and tubulin (oryzalin) (Geldner et al., 2001; Kleine-Vehn et al., 2008) showed only little effect on the intracellular localization of both PIN7 isoforms when applied alone (Supplemental Figures 3D and 3E). Pretreatment with cytochalasin D prevented the formation of the BFA bodies (Geldner et al., 2001) containing both PIN7 isoforms (Supplemental Figure 3F). Yet, when we applied oryzalin prior to the BFA treatment, the BFA compartments containing PIN7a-GFP and PIN7b-RFP associated in only very weakly co-localizing structures (Supplemental Figure 3G). Next, we tested how the cytoskeleton is involved in the trafficking of both PIN7 isoforms from the BFA bodies to the PM by washing out BFA with cytochalasin D or oryzalin (Geldner et al., 2001). In the presence of cytochalasin D, PIN7a-GFP largely persisted inside the BFA bodies, while the PIN7b-RFP signal was almost absent in these aggregates (Supplemental Figure 3H). We generally did not see any difference between both isoforms when BFA was washed out with oryzalin (Supplemental Figure 3I). These data thereby suggest that both PIN7 isoforms use vesicle trafficking pathways assisted by a common cytoskeletal scaffold. These pathways differ in their dynamics and the endomembrane components involved and are consistent with previous findings that individual PINs can utilize multiple PM delivery routes (Kleine-Vehn et al., 2008; Boutté et al., 2006).

PIN7a and PIN7b do not differ in tropic stimuli-mediated polarity change or dimer formation

Auxin transporters of the PIN3 clade change their polar localization on the PM by the reaction to various environmental cues, in particular by switching the light or gravity vectors (Friml et al., 2002; Ding et al., 2011; Rakusová et al., 2011). PIN3 relocation in response to gravity in columella root cells is seen in as little as 2 min, while the relocation of PIN7-GFP requires approximately 30 min to be detected (Kleine-Vehn et al., 2010; Friml et al., 2002; Grones et al., 2018; Pernisova et al., 2016). We examined plants harboring both *PIN7a-GFP* and *PIN7b-RFP* cDNAs under short and long gravitropic stimuli. We did not find any difference in relocation dynamics between both isoforms in these experiments (Supplemental Figures 4A–4B). We also observed no difference in the polarity change between PIN7a-GFP and PIN7b-RFP in hypocotyl gravitropic (Rakusová et al., 2011, 2016) and phototropic (Ding et al., 2011) bending assays (Supplemental Figure 4C; due to limited transparency of hypocotyls, we used lines expressing the cDNAs under control of a stronger *SCR* promoter (Rakusová et al., 2011)). The different subcellular pathways driving both PIN7a-GFP and PIN7b-RFP cargos are thereby not connected with their ability to change polarity in response to tropic stimuli.

AS frequently changes the protein function by altering their ability to multimerize (Kelemen et al., 2013), and PIN proteins have recently been found to associate in higher-order complexes (Teale et al., 2020; Abas et al., 2021). To independently corroborate these findings, we used crude extracts from the *Arabidopsis* seedlings and the BY-2 cells, expressing the respective cDNAs, followed by non-denaturing protein electrophoresis and western blotting (Leitner et al., 2012). Probing with anti-GFP antibody indeed revealed a faint signal at the predicted size of PIN7 dimers in both tissue sources (Figures 4E and 4F; Supplemental Figures 4D and 4E). We further validated these findings with Förster resonance energy transfer by fluorescence lifetime imaging (FRET-FLIM), analyzing the cDNAs expressed stably in *Arabidopsis* seedlings and transiently in the tobacco leaves (Figures 4G and 4H; Supplemental Figures 4F and 4G). Here, we also examined the combinations of the GFP and RFP-tagged *PIN7a* and *PIN7b* cDNAs to determine their pairing preference. Shortening of the GFP fluorescence lifetime was seen regardless of the isoform interaction tested. It therefore hints at the possibility that the PIN7 splice isoforms can mutually associate into dimers (or other higher order complexes), but it seems the strength of their interaction remains comparable.

PIN7a and PIN7b show distinct stability on PM

Numerous studies employed the fluorescence recovery after photobleaching (FRAP) analysis to investigate the dynamic turnover of various proteins, including PINs, on PM (Men et al., 2008; Martinière et al., 2012; Langowski et al., 2016; Laňková et al., 2016). We therefore bleached a region of the PM signal in the root meristem of the *PIN7a-GFP* and *PIN7b-GFP* cDNA lines and measured the FRAP in this area. Notably, PIN7a-GFP showed a slower recovery of fluorescence than PIN7b-GFP (Figure 5A); the use of either GFP or RFP tag did not markedly influence the fluorescence recovery (Supplemental Figure 5A). We also generated plants carrying a *PIN3::PIN3Δ-RFP* cDNA construct, which lacks the GETK motif corresponding to the four amino acids absent in PIN7b (Figures 1B and 5B). The PIN3Δ-RFP signal displayed an incomplete co-localization with the wild type PIN3-GFP variant in the BFA bodies (Supplemental Figure 5B) and faster recovery on the PM, similar to that observed for PIN7a and PIN7b (Figure 5B). It therefore appears that the motif altered by AS of PIN7 is required for the regulation of dynamics of individual isoforms within the PM.

The fluorophore-tagged *PIN7* cDNAs showed a relatively weak signal under natural promoter and biasing the obtained data by extensive bleaching by prolonged confocal imaging (Figure 5A). Therefore, we utilized the *SCR* promoter-driven lines, which displayed higher expression levels, but still lower intensities than the commonly used genomic DNA-derived *PIN7::PIN7-GFP* (Blilou et al., 2005) (Supplemental Figures 5C and 5D). Indeed, these lines allowed for detecting a finer difference in FRAP, as compared to those under the control of the native promoter (Figure 5C). The FRAP experiments (Figure 5A) were conducted initially on the *pin347* lines, which expressed the corresponding transgenes separately. Therefore, we compared the FRAP values of the single PIN7a-GFP and PIN7b-RFP with those carrying these constructs simultaneously in the *pin347* genetic background. Strikingly, the co-expression of PIN7a-GFP decreased the FRAP of PIN7b-RFP and, *vice versa*, the presence of PIN7b-RFP increased the FRAP of PIN7a-GFP (Figure 5C). This shows that the PIN7 isoforms mutually influence each other stability within PM, presumably by the association in a molecular complex.

In addition, both PIN7a and PIN7b did not markedly differ in their capability to transport auxin in the BY-2 cells (Figure 4B and Supplemental Figure 3A). We were unable to detect homologous AS events in plant lineages related to tobacco (see Figure 1D). We tested FRAP of the *Arabidopsis* PIN7a- and PIN7b-GFP on PM in BY-2 cells. Interestingly, the recovery of both isoforms showed a virtually identical course (Figure 5D). Altogether, these observations support the scheme that both isoforms physically interact and mutually

influence each other stability within PM, which leads to a differential impact on the plant phenotype.

DISCUSSION

We provide evidence that the evolutionary conserved AS event increases the diversity among the PIN auxin transporters in *Arabidopsis thaliana*. Focusing on PIN7, we show that both PIN7a and PIN7b isoforms (differing by presence of a 4-amino acid stretch) are required together for the correct apical hook formation and tropic responses. A similar, genetically evidenced cooperative modes of action of splice isoforms has already been proposed during *Arabidopsis* development (Szakonyi and Duque, 2018). The seed dormancy regulator *DELAY OF GERMINATION 1 (DOG1)* is processed into five mRNAs. Only the expression of two or more DOG1 cDNAs under the native promoter rescues the *dog1* phenotype by synergistic stabilization of the protein by its multimerization (Nakabayashi et al., 2015). The transcriptional factor HYH (HY5 HOMOLOG) possesses an isoform, which lacks a domain required for proteasomal degradation, which leads to its increased stability and probably works as a semi-dominant splice variant (Szakonyi and Duque, 2018; Sibout et al., 2006); this can probably also be the case of the AS of *MONOPTEROS* (Cucinotta et al., 2020). Together with a partly similar mechanism described for the temperature-mediated regulation of flowering (Posé et al., 2013; Lee et al., 2013), the example of AS of PIN7 is among rarely described instances where the mutually antagonistic effects of two splice isoforms are observed and can be placed in developmental context.

Based on the current understanding of the PIN-mediated auxin transport (Adamowski and Friml, 2015), we validated the most plausible hypotheses to provide a mechanistic view on the mutual functionality of PIN7a and PIN7b. This AS event neither affects the ability of PIN7 carriers to transport auxin *per se* nor influences their ability to relocate after the tropic stimulus. Although the expression of PIN7b is pronounced at the concave side of the apical hook, we did not see any remarkable expression changes of the P7A₁G/P7BR dual reporter during hypocotyl bending assays (Figure 2C and Supplemental Figure 4C). Instead, our data favor the scenario that PIN7a and PIN7b functionally interact at the protein level. Along with recent findings (Teale et al., 2020; Abas et al., 2021), we confirm with orthogonal experimental tools that PIN7 isoforms closely associate. Moreover, the combined FRAP analysis shows that PIN7a and PIN7b influence each other stability within PM. It has previously been suggested that PINs are complexed inside stable integral PM clusters (Langowski et al., 2016; Feraru et al., 2011; Kleine-Vehn et al., 2011). The concept that

PIN7a, detained in these microdomains, transports auxin, but dynamic PIN7b does not, can also be in accord with the diverse subcellular pathways delivering the PIN7 isoforms to PM. Hence, PIN7b probably binds PIN7a and impedes the polar auxin flow provided by PIN7a in these membrane domains to fine-tune auxin developmental responses (Supplemental Figure 5E).

In general, AS modifies protein properties variably. It can affect protein subcellular localization, ligand binding affinity, enzymatic or transporting activities, protein stability, or the presence of covalent post-translational modifications (Stamm et al., 2005; Kelemen et al., 2013). Different covalent modifications alter the subcellular trafficking of most PINs. Phosphorylation sites on serine, threonine or tyrosine residues of various PINs have been identified; their phosphorylation status also changes PIN-mediated tropic responses (Zwiewka et al., 2019; Barbosa et al., 2018; Rademacher and Offringa, 2012). Also PIN ubiquitination (on lysines) and controlled proteolytic degradation act in auxin-mediated processes (Leitner et al., 2012). However, none of the candidate residues required for these modifications is present in the vicinity of the amino acid motif changed by AS. This region, present inside the PIN long hydrophilic loop, shows low amino acid conservation (as seen particularly on the *PIN4* AS event), and it is perhaps intrinsically disordered. Therefore, one can speculate whether just the length of this motif can be critical for functional interaction among internal structural domains (Buljan et al., 2013). These structural differences may modulate the affinity of PINs to bind the factors required for the entry and presence in the PM microdomains, and thus participate in the PIN subcellular dynamics and directional auxin transport.

EXPERIMENTAL PROCEDURES

Plant material and plant growth conditions

All plant material, except tobacco cell cultures, was in the *Arabidopsis thaliana* (L.) Heynh., Col-0 ecotype. These mutant and transgenic lines were described previously: *PIN3::PIN3-GFP* (Zadnikova et al., 2010), *PIN4::PIN4-GFP*, *PIN7::PIN7-GFP* (Blilou et al., 2005), *pin3-3 pin4-101 pin7-102 (pin347)*, *pin3-3 pin4-101 (pin34)* (Willige et al., 2013), *pin7-2* (Friml et al., 2003).

For the *in vitro* cultivation, the seeds were surface-sterilized for 5 h with chlorine gas, plated on 0.5× Murashige & Skoog medium with 1 % sucrose, and then stratified for 2 d at 4°C in darkness. Unless indicated otherwise, the seedlings were grown on vertically oriented plates for 4-6 days under 16 h : 8 h photoperiod, 22 : 18°C, light : dark.

The following chemicals were used for treatments: brefeldin A (BFA), cytochalasin D (CytD), oryzalin (Ory), β -estradiol, 2,4-dichlorophenoxyacetic acid (2,4-D), all from Sigma (Sigma-Aldrich, St. Louis, MO, USA). Radioactively labeled auxin accumulation assays were performed with [^3H]NAA (naphthalene-1-acetic acid; 20 Ci.mmol $^{-1}$; American Radiolabeled Chemicals, St. Louis, MO, USA).

DNA manipulations and transgenic work

The genomic *PIN7::PIN7-RFP* construct was made by replacing the GFP coding sequence in the original *PIN7::PIN7-GFP* construct (Blilou et al., 2005). For creating the *PIN7(4)::PIN7(4)a-GFP* and *PIN7(4)::PIN7(4)b-RFP* cDNA constructs, the respective cDNAs were cloned into the pDONR221 P5-P2 entry vector (Invitrogen, Life Technologies, Carlsbad, CA, USA) by the Gateway BP reaction (Invitrogen). For placing the fluorophore tag, the *XbaI* restriction site was introduced at the 1350 bp (for *PIN7*) or 1341 bp (for *PIN4*) behind the start codon of the cDNA coding region, in the same amino acid sequence context as in the published *PIN4* and *PIN7-GFP* constructs (Blilou et al., 2005). In parallel, the *PIN7* (or *PIN4*) promoters were inserted into the pH7WG Gateway vector (Department of Plant Systems Biology, Ghent University, Belgium (Karimi et al., 2002)) by the Gibson Assembly kit (New England Biolabs, Ipswich, MA, USA). The tagged *PIN7* and *PIN4* cDNA entry clones were then recombined with the modified pH7WG destination vector by the Gateway LR reaction (Invitrogen). For the cloning of the *PIN7* splicing dual fluorescent reporter, the entry vectors carrying the genomic sequence of *PIN7-GFP* and *PIN7-RFP*, respectively, were modified by inverse PCR. The resulting constructs were then recombined by the Gateway LR reaction with the *PIN7* promoter containing the pH7WG destination vector. The *SCR::PIN7a-GFP* and *SCR::PIN7b-RFP* constructs were obtained by recombination of the *SCR* promoter in pDONR221 P1-P5r and *PIN7* cDNA entry clones with the pH7WG destination vector by the Multisite Gateway LR reaction. *PIN3A-RFP* cDNA was custom synthesized (Gen9, Ginkgo Bioworks, Boston, MA, USA), cloned into pDONR221 P5-P2 vector and together with the *PIN3* promoter pDONR221 P1-P5r construct recombined into the pH7WG vector with Multisite Gateway. The validated binary constructs were transformed into *Arabidopsis* by floral dipping. For making the estradiol-inducible *G10-90::XVE>>PIN7(4)a-GFP*, *PIN7(4)b-GFP*, *PIN7a-RFP* and *PIN7b-RFP* constructs, tagged *PIN7* or *PIN4* cDNA entry clones were recombined with the pMDC7 destination vector (Curtis and Grossniklaus, 2003) by the Gateway LR reaction. Unless stated otherwise, the constructs present in the study were

cloned under their natural *PIN4* or *PIN7* promoter. Primers used in this work are listed in Supplemental Table 1.

For the generation of the stable transgenic lines carrying the cDNA constructs, at least eight independent descendant populations of primary transformants were preselected for the presence of the fluorescent signal. The functional validity of the cDNA constructs was verified in the phototropic bending test, where all candidate lines matched the presented results. Selected lines were used for further phenotypic analysis.

Plant phenotype analysis

The dynamic seedling development was tracked in the custom made dynamic morphogenesis observation chamber equipped with blue and white LED unilateral light sources, infra-red LED back light and a camera for imaging in the infra-red spectra, controlled by the Raspberry Pi3B microcomputer (Raspberry Pi foundation, Cambridge, UK). For the hypocotyl bending assays, the plated seeds were first illuminated for 6 h with white light. The plates were then transferred to the observation chamber for 3-4 d. For the hypocotyl phototropic bending experiments (Friml et al., 2002), the dark-grown seedlings were afterward illuminated for 20 h with unilateral white light and imaged every 20 minutes. For hypocotyl gravitropic bending experiments (Rakusová et al., 2011), the dark-grown seedlings were rotated by 90° clockwise and imaged for 30 h every 60 min. For tracking apical hook development, the seeds were first illuminated for 6 h with white light. They were then transferred to the observation chamber and their development recorded every 4 h for a total 150 - 200 h in infrared spectra. Germination time was set as time 0, when first traces of the main root were observed, individually for each seedling analyzed. Apical hook was determined as an angle between the immobile (non-bending) part of hypocotyl and the distal edge of cotyledons (Zadnikova et al., 2010), using the ImageJ software (Rueden et al., 2017). At least 15 seedlings were analyzed for each line. Each experiment was done at least three times.

For protoxylem defects analysis, 5-d old light-grown seedlings were cleared and analyzed as described previously (Bishopp et al., 2011). For examining lateral root density, 8-d old light-grown seedlings were cleared and observed with a DIC microscope (Dubrovsky et al., 2009). Lateral root density was calculated by dividing the total number of lateral roots and lateral root primordia to the length of the main root as described (Dubrovsky et al., 2009). Vertical Growth Index (VGI), defined as a ratio between main root ordinate and main root length, was quantified on 5-d old seedlings as published previously (Grabov et al., 2005). For

measuring the Gravitropic Set-point Angle (GSA), plates with 14-d old light-grown seedlings were scanned and the angles between the vertical axis and five innermost 0.5 mm parts of lateral root were determined as described previously (Roychoudhry et al., 2017). In all cases, 12-20 seedlings were analyzed for each line. Each experiment was done at least three times.

For decapitation experiments, 4-week short-day grown plants were moved into long-day conditions to induce flowering. After the primary bolt reached 10-15 cm, the plant was decapitated. The number of rosette branches was recorded at 7, 10 and 14-d after decapitation (Waldie and Leyser, 2018; Greb et al., 2003). Rosette size was inspected in 18-d light-grown plants prior to documenting. 10 plants were analyzed for each line, the experiment was done three times.

Microscopy

Bright-field microscopy (differential interference contrast, DIC) was conducted on the Olympus BX61 instrument (Olympus, Shinjuku, Tokyo, Japan) equipped with a DP50 camera (Olympus). Routine confocal microscopy was performed on inverted Zeiss Axio Observer.Z1 containing the standard confocal LSM880 and Airyscan modules with 20x/0.8 DIC M27 air, 40x/1.2 W Kor FCS M27 air and 63x/1.40 Oil DIC M27 objectives (Carl Zeiss AG, Jena, Germany). Gravity-induced polarity change experiments were carried out on Zeiss Axio Observer.Z1 with vertically oriented sample position and the 40x/0.75 glycerol objective.

To observe the light-induced polarity change of PIN7a-GFP and PIN7b-RFP, 4-d dark-grown seedlings were irradiated for 4 h with unilateral white light and then imaged with Zeiss Axio Observer.Z1 LSM880 with a vertically oriented sample position, as described (Willige et al., 2013). For analyzing gravity-induced polarity change, 4-d dark-grown seedlings were reoriented by 90° clockwise and imaged 6 and 24 h after rotation, as described (Rakusová et al., 2011).

For BFA treatments, 5-d light-grown seedlings were transferred to liquid 0.5× MS media containing 50 μM BFA. The membrane/cytosol ratio was determined with ImageJ, it was defined as the mean membrane signal intensity divided by the mean fluorescence in the cytosol. For cytoskeleton depolymerizing drug treatments (Geldner et al., 2001), 5-d light-grown seedlings were transferred to liquid 0.5× MS media supplemented with 20 μM of cytochalasin D or 20 μM oryzalin. The co-treatments were done by the direct addition of BFA. For the BFA removal, prior to the addition of the cytoskeleton depolymerizing compounds, seedlings were twice washed out with fresh media and then transferred to that supplemented with the respective cytoskeleton-depolymerizing drug.

For imaging of the expression of *DR5v2::GFP* in the apical hook, the GFP signal in the 4-days old etiolated seedlings was fixed in 4% paraformaldehyde (PFA, Sigma) overnight. Then, the tissue was hand-sectioned with a razor blade and observed.

Fluorescent recovery after photobleaching (FRAP)

For the FRAP experiments, Zeiss Axio Observer.Z1 equipped with the LSM880 confocal and Airyscan modules and the 40x/1.2 W Kor FCS M27 air objective was used. A rectangular region of interest (ROI) of 40 x 20 pixels was selected on the basal or apical PM of cells inside the vascular cylinder in the primary root meristematic area of 5-d light-growth seedlings. ROI was bleached with the maximum 488 nm laser intensity and fluorescence recovery was documented every 1 or 2 sec for a total 200 sec. Recovery time lapses were analyzed with ImageJ. The Slices Alignment plugin (Tseng et al., 2012) for ImageJ was used for the elimination of cell movement caused by root growth. In parallel, another rectangular ROI (100 x 20 pixels) on the non-bleached cell was selected as a reference. To compensate for the fluorophore bleaching during the recovery period, the data were normalized using equation (Laňková et al., 2016)

$$In = \frac{\left(\frac{It}{It_{ref}}\right) - \left(\frac{Imin}{Imin_{ref}}\right)}{\left(\frac{Imax}{Imax_{ref}}\right) - \left(\frac{Imin}{Imin_{ref}}\right)},$$

where *In* is normalized fluorescence intensity, *It* is the intensity at the specific time point, *Imax* is the intensity after the initial bleaching, *Imin* is the intensity before initial bleaching. *It_{ref}*, *Imax_{ref}* and *Imin_{ref}* represent the same values for the reference ROI. At least 3 membranes in 4-5 root tips were analyzed in each experiment. Single-phase exponential fitting was applied to the normalized FRAP data (SigmaPlot, Systat Software, Chicago, IL, USA) as described (Laňková et al., 2016; Sprague and McNally, 2005). Recovery half-time is defined as a time required for the fluorescence recovery to reach half of the steady-state fluorescence intensity (Sprague and McNally, 2005; Soumpasis, 1983).

FLIM-FRET analysis

For assessing the pairing specificity in *Arabidopsis*, the lines harboring all 4 combinations of *G10-90::XVE* > *PIN7a/b-GFP* (donors) and *PIN7a/b-RFP* (acceptors) were generated. Before imaging, 4-d light-growth seedlings were transferred for two days to the solid ½ MS media supplemented with 5 µM β-estradiol. The transient expression in tobacco leaves was performed as earlier published (Horák et al., 2008), omitting the β-estradiol

induction to keep the transgene expression low (Bashandy et al., 2015). The abaxial epidermis of tobacco leaves was observed 3 days after infiltration.

The procedure of the FRET-FLIM assay was done as previously described (Krejčí et al., 2015). Imaging was performed on inverted Zeiss LSM 780 AxioObserver.Z1 equipped with the M27 Plan-Apochromat 20x air objective, external In-Tune laser (<3 nm width, pulsed, 40 MHz, 1.5 mW) and the GaAsP detectors. For conducting the FLIM-FRET assays, the HPM-100-40 hybrid detector associated with the photon counting module Simple-Tau 150 (compact TCSPC system based on the SPC-150N device) was utilized, using the detector controller card DCC-100 (Becker & Hickl GmbH, Berlin, Germany). The cell membranes containing both donor and acceptor signals were excited with the wavelength 490 nm to generate intensity image and the photons were recorded with the supplied SPCM software. In the image analysis, the pixel per pixel fitting model was used for the two-component system and the one-component system was used for the donor alone (Becker & Hickl SPCImage software package). Multiexponential decay model was used for fitting and FRET efficiencies were calculated using the equation

$$E = 1 - \tau_{DA}/\tau_D,$$

where E is energy transfer efficiency, τ_{DA} donor fluorescent lifetime in the presence of acceptor and τ_D donor fluorescent lifetime in the absence of acceptor. The observation of two lifetimes in the GFP fluorescence spectrum was expected: first, close to 2.6 ns, corresponds to a non-interacting fraction, while the second one, characterized by a shorter lifetime, was considered as associated with interacting molecules.

Tobacco cell lines and auxin accumulation assays

Tobacco cell line BY-2 (*Nicotiana tabacum* L. cv. Bright Yellow 2) was cultivated as described (Müller et al., 2019). BY-2 cells were transformed with the pMDC7 constructs by co-cultivation with *Agrobacterium tumefaciens* strain GV2260 as earlier described (Petrasek et al., 2006; Müller et al., 2019). The transgene expression was maintained by cultivation on media supplemented with 40 $\mu\text{g}.\text{ml}^{-1}$ hygromycin (Roche) and 100 $\mu\text{g}.\text{ml}^{-1}$ cefotaxime (Sigma). Accumulation assays of radioactively labeled auxin were performed as previously published (Petrasek et al., 2006; Delbarre et al., 1996) with cells cultured for 2 days in β -estradiol or DMSO mock treatments. Presented results are from 3 biological replicates for each representative *PIN7a* and *PIN7b* line and were confirmed for each on two other independent genotypes. Independent β -estradiol inductions were done within 3-9 months after

establishing the cell suspension on lines, which showed comparable levels of the expressed PIN7-GFP signal.

Protein extraction and immunoblotting

Protein extraction from the BY-2 tobacco cell lines and the *Arabidopsis* roots was performed as described previously (Leitner et al., 2012), using the modified buffer, which contained only NP-40 as detergent. Otherwise, DTT, SDS and any other detergents were omitted from the buffers. The GFP tagged PIN7 variants were immunoprecipitated with the GFP-Trap Dynabeads (Chromotek, Planegg, Germany) according to the manufacturer's description. The bead elution was done with the glycine buffer (0.2 M glycine, pH 2.5) and the eluate was neutralized with the 1/10 sample volume of 1M Tris-HCl, pH 8.5. The eluted proteins were mixed with a premade 2× Sample Buffer (Biorad, Hercules, CA, USA), omitting β-mercaptoethanol (Sigma) prior to the loading on the gel. To determine the putative dimerization of PIN7-GFP, the total membrane fractions or immunoprecipitated proteins were run on a non-denaturing polyacrylamide gel electrophoresis. Here, the Mini-PROTEAN® TGX™ Stain-Free Precast gels (Biorad) were used, devoid of SDS. The trihalo compounds present in the gel allowed for the visualization of the proteins under UV light to control the loading quality. As a marker, the PageRuler™ Prestained Protein Ladder (10 to 180 kDa, ThermoFisher) was used. For western blotting, the proteins were transferred to the nitrocellulose membrane (GE Healthcare Life Sciences, Chicago, IL, USA), blocked with 3% BSA for 2 h, and incubated overnight at 4°C with monoclonal primary anti-GFP antibody from rabbit (1:2,000) (Invitrogen), and with secondary anti-rabbit antibody (1:30,000) (Invitrogen), as described previously (Leitner et al., 2012).

Statistics and sequence analysis

For the mean comparison of the two groups, Student's *t*-test was applied. Statistical analysis of multiple groups was performed by one-way ANOVA with subsequent Tukey HSD post-hoc test. All statistical tests, including that for equal normality, were performed in R-studio IDE (R-studio, Boston, MA, USA). In the box plots, the whisker length was set as $Q \pm 1.5 \times IQR$, where Q is the corresponding quartile and IQR is the interquartile range. For creating the multiple sequence alignments, the protein sequences were assembled with the Clustal Omega algorithm (Sievers et al., 2011) and graphically outlined by Mega-X, using default ClustalX color code (Kumar et al., 2018).

ACKNOWLEDGMENT

We thank Claus Schwechheimer for the *pin34* and *pin347* seeds, Yuliia Mironova for technical assistance, Ksenia Timofeyenko and Dmitry Konovalov for help with the evolutionary analysis, Konstantin Kutashev and Siarhei Dabravolski for assistance with the FRET-FLIM analysis, Huibin Han for advice with the hypocotyl imaging, Karel Müller for the initial qRT-PCR on the tobacco cell lines, and Jozef Mravec and Lindy Abas for their comments on the manuscript. This work was supported by the Czech Science Foundation (16-26428S and 19-23773S) to I. K., M. H. and K. R., and the Ministry of Education, Youth and Sports of the Czech Republic (MEYS, CZ.02.1.01/0.0/0.0/16_019/0000738) to K. R. The imaging Facilities of the Institute of Experimental Botany and CEITEC are supported by MEYS (LM2018129 - Czech BioImaging and CZ.02.1.01/0.0/0.0/16_013/0001775).

AUTHOR CONTRIBUTIONS

I. K., M. H., K. Re., J. Hu., A. J., R. F., Z. V., D. R., and J. P. conducted experiments. S. S. and J. F. provided unpublished material. I. K., K. Re., J. Hu., J. P., T. B. J., M. J. F., J. He., J. F., J. P., and K. Rů. conceived the research and designed experiments. K. Rů. and I. K. wrote the manuscript. All authors read and commented on the final version of the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

- Abas, L., Kolb, M., Stadlmann, J., Janacek, D.P., Lukic, K., Schwechheimer, C., Sazanov, L.A., Mach, L., Friml, J., and Hammes, U.Z. (2021). Naphthylphthalamic acid associates with and inhibits PIN auxin transporters. PNAS 118.
- Adamowski, M. and Friml, J. (2015). PIN-Dependent Auxin Transport: Action, Regulation, and Evolution. Plant Cell 27: 20–32.
- Barbosa, I.C.R., Hammes, U.Z., and Schwechheimer, C. (2018). Activation and Polarity Control of PIN-FORMED Auxin Transporters by Phosphorylation. Trends in Plant Science 23: 523–538.
- Bashandy, H., Jalkanen, S., and Teeri, T.H. (2015). Within leaf variation is the largest source of variation in agroinfiltration of *Nicotiana benthamiana*. Plant Methods 11.
- Bazin, J., Romero, N., Rigo, R., Charon, C., Blein, T., Ariel, F., and Crespi, M. (2018). Nuclear Speckle RNA Binding Proteins Remodel Alternative Splicing and the Non-coding Arabidopsis Transcriptome to Regulate a Cross-Talk Between Auxin and Immune Responses. Front Plant Sci 9: 1209.

- 630 Bennett, T., Brockington, S.F., Rothfels, C., Graham, S.W., Stevenson, D., Kutchan, T., Rolf,
631 M., Thomas, P., Wong, G.K.-S., Leyser, O., Glover, B.J., and Harrison, C.J. (2014).
632 Paralogous Radiations of PIN Proteins with Multiple Origins of Noncanonical PIN
633 Structure. *Mol Biol Evol* 31: 2042–2060.
- 634 Bennett, T., Hines, G., Rongen, M. van, Waldie, T., Sawchuk, M.G., Scarpella, E., Ljung, K.,
635 and Leyser, O. (2016). Connective Auxin Transport in the Shoot Facilitates
636 Communication between Shoot Apices. *PLOS Biology* 14: e1002446.
- 637 Bishopp, A., Help, H., El-Showk, S., Weijers, D., Scheres, B., Friml, J., Benková, E.,
638 Mähönen, A.P., and Helariutta, Y. (2011). A mutually inhibitory interaction between
639 auxin and cytokinin specifies vascular pattern in roots. *Curr. Biol.* 21: 917–926.
- 640 Blencowe, B.J. (2017). The Relationship between Alternative Splicing and Proteomic
641 Complexity. *Trends in Biochemical Sciences* 42: 407–408.
- 642 Blilou, I., Xu, J., Wildwater, M., Willemsen, V., Paponov, I., Friml, J., Heidstra, R., Aida, M.,
643 Palme, K., and Scheres, B. (2005). The PIN auxin efflux facilitator network controls
644 growth and patterning in Arabidopsis roots. *Nature* 433: 39–44.
- 645 Boutté, Y., Crosnier, M.-T., Carraro, N., Traas, J., and Satiat-Jeunemaitre, B. (2006). The
646 plasma membrane recycling pathway and cell polarity in plants: studies on PIN
647 proteins. *J. Cell. Sci.* 119: 1255–1265.
- 648 Buljan, M., Chalancon, G., Dunker, A.K., Bateman, A., Balaji, S., Fuxreiter, M., and Babu,
649 M.M. (2013). Alternative splicing of intrinsically disordered regions and rewiring of
650 protein interactions. *Current Opinion in Structural Biology* 23: 443–450.
- 651 Casson, S.A., Topping, J.F., and Lindsey, K. (2009). MERISTEM-DEFECTIVE, an RS
652 domain protein, is required for the correct meristem patterning and function in
653 Arabidopsis. *The Plant Journal* 57: 857–869.
- 654 Chamala, S., Feng, G., Chavarro, C., and Barbazuk, W.B. (2015). Genome-wide identification
655 of evolutionarily conserved alternative splicing events in flowering plants. *Front*
656 *Bioeng Biotechnol* 3: 33.
- 657 Cheng, C.-Y., Krishnakumar, V., Chan, A.P., Thibaud Nissen, F., Schobel, S., and Town,
658 C.D. (2017). Araport11: a complete reannotation of the Arabidopsis thaliana reference
659 genome. *The Plant Journal* 89: 789–804.
- 660 Cucinotta, M., Cavalleri, A., Guazzotti, A., Astori, C., Manrique, S., Bombarely, A., Oliveto,
661 S., Biffo, S., Weijers, D., Kater, M.M., and Colombo, L. (2020). Alternative Splicing
662 Generates a MONOPTEROS Isoform Required for Ovule Development. *Current*
663 *Biology*.
- 664 Curtis, M.D. and Grossniklaus, U. (2003). A gateway cloning vector set for high-throughput
665 functional analysis of genes in planta. *Plant Physiol.* 133: 462–469.
- 666 Darracq, A. and Adams, K.L. (2013). Features of evolutionarily conserved alternative splicing
667 events between Brassica and Arabidopsis. *New Phytologist* 199: 252–263.

- 668 Delbarre, A., Muller, P., Imhoff, V., and Guern, J. (1996). Comparison of mechanisms
669 controlling uptake and accumulation of 2,4-dichlorophenoxy acetic acid, naphthalene-
670 1-acetic acid, and indole-3-acetic acid in suspension-cultured tobacco cells. *Planta*
671 198: 532–541.
- 672 Ding, Z., Galván-Ampudia, C.S., Demarsy, E., Łangowski, Ł., Kleine-Vehn, J., Fan, Y.,
673 Morita, M.T., Tasaka, M., Fankhauser, C., Offringa, R., and Friml, J. (2011). Light-
674 mediated polarization of the PIN3 auxin transporter for the phototropic response in
675 *Arabidopsis*. *Nat Cell Biol* 13: 447–452.
- 676 Dubrovsky, J.G., Soukup, A., Napsucialy-Mendivil, S., Jeknic, Z., and Ivanchenko, M.G.
677 (2009). The lateral root initiation index: an integrative measure of primordium
678 formation. *Ann. Bot.* 103: 807–817.
- 679 Feraru, E., Feraru, M.I., Kleine-Vehn, J., Martinière, A., Mouille, G., Vanneste, S.,
680 Vernhettes, S., Runions, J., and Friml, J. (2011). PIN Polarity Maintenance by the Cell
681 Wall in *Arabidopsis*. *Current Biology* 21: 338–343.
- 682 Friml, J., Vieten, A., Sauer, M., Weijers, D., Schwarz, H., Hamann, T., Offringa, R., and
683 Jürgens, G. (2003). Efflux-dependent auxin gradients establish the apical-basal axis of
684 *Arabidopsis*. *Nature* 426: 147–153.
- 685 Friml, J., Wiśniewska, J., Benková, E., Mendgen, K., and Palme, K. (2002). Lateral relocation
686 of auxin efflux regulator PIN3 mediates tropism in *Arabidopsis*. *Nature* 415: 806–809.
- 687 Ganguly, A., Park, M., Kesawat, M.S., and Cho, H.-T. (2014). Functional Analysis of the
688 Hydrophilic Loop in Intracellular Trafficking of *Arabidopsis* PIN-FORMED Proteins.
689 *Plant Cell* 26: 1570–1585.
- 690 Geldner, N., Friml, J., Stierhof, Y.D., Jürgens, G., and Palme, K. (2001). Auxin transport
691 inhibitors block PIN1 cycling and vesicle trafficking. *Nature* 413: 425–428.
- 692 Ghelli, R. et al. (2018). A Newly Identified Flower-Specific Splice Variant of AUXIN
693 RESPONSE FACTOR8 Regulates Stamen Elongation and Endothecium Lignification
694 in *Arabidopsis*. *Plant Cell* 30: 620–637.
- 695 Glanc, M., Fendrych, M., and Friml, J. (2019). PIN2 Polarity Establishment in *Arabidopsis* in
696 the Absence of an Intact Cytoskeleton. *Biomolecules* 9: 222.
- 697 Grabov, A., Ashley, M.K., Rigas, S., Hatzopoulos, P., Dolan, L., and Vicente-Agullo, F.
698 (2005). Morphometric analysis of root shape. *New Phytol.* 165: 641–651.
- 699 Greb, T., Clarenz, O., Schafer, E., Muller, D., Herrero, R., Schmitz, G., and Theres, K.
700 (2003). Molecular analysis of the LATERAL SUPPRESSOR gene in *Arabidopsis*
701 reveals a conserved control mechanism for axillary meristem formation. *Genes Dev.*
702 17: 1175–1187.
- 703 Grones, P., Abas, M., Hajný, J., Jones, A., Waidmann, S., Kleine-Vehn, J., and Friml, J.
704 (2018). PID/WAG-mediated phosphorylation of the *Arabidopsis* PIN3 auxin
705 transporter mediates polarity switches during gravitropism. *Sci Rep* 8: 10279.

706 Horák, J., Grefen, C., Berendzen, K.W., Hahn, A., Stierhof, Y.-D., Stadelhofer, B., Stahl, M.,
707 Koncz, C., and Harter, K. (2008). The Arabidopsis thaliana response regulator ARR22
708 is a putative AHP phospho-histidine phosphatase expressed in the chalaza of
709 developing seeds. *BMC Plant Biology* 8: 77.

710 Hrtyan, M., Šliková, E., Hejátko, J., and Růžička, K. (2015). RNA processing in auxin and
711 cytokinin pathways. *J. Exp. Bot.* 66: 4897–4912.

712 Kalyna, M., Lopato, S., and Barta, A. (2003). Ectopic Expression of atRSZ33 Reveals Its
713 Function in Splicing and Causes Pleiotropic Changes in Development. *Mol. Biol. Cell*
714 14: 3565–3577.

715 Karimi, M., Inzé, D., and Depicker, A. (2002). GATEWAYTM vectors for Agrobacterium-
716 mediated plant transformation. *Trends in Plant Science* 7: 193–195.

717 Kashkan, I., Timofeyenko, K., Kollárová, E., and Růžička, K. (2020). In Vivo Reporters for
718 Visualizing Alternative Splicing of Hormonal Genes. *Plants* 9: 868.

719 Kelemen, O., Convertini, P., Zhang, Z., Wen, Y., Shen, M., Falaleeva, M., and Stamm, S.
720 (2013). Function of alternative splicing. *Gene* 514: 1–30.

721 Keren, H., Lev-Maor, G., and Ast, G. (2010). Alternative splicing and evolution:
722 diversification, exon definition and function. *Nat Rev Genet* 11: 345–355.

723 Kleine-Vehn, J. et al. (2011). Recycling, clustering, and endocytosis jointly maintain PIN
724 auxin carrier polarity at the plasma membrane. *Molecular Systems Biology* 7: 540.

725 Kleine-Vehn, J., Ding, Z., Jones, A.R., Tasaka, M., Morita, M.T., and Friml, J. (2010).
726 Gravity-induced PIN transcytosis for polarization of auxin fluxes in gravity-sensing
727 root cells. *PNAS* 107: 22344–22349.

728 Kleine-Vehn, J., Łangowski, Ł., Wiśniewska, J., Dhonukshe, P., Brewer, P.B., and Friml, J.
729 (2008). Cellular and Molecular Requirements for Polar PIN Targeting and
730 Transcytosis in Plants. *Molecular Plant* 1: 1056–1066.

731 Klepikova, A.V., Kasianov, A.S., Gerasimov, E.S., Logacheva, M.D., and Penin, A.A. (2016).
732 A high resolution map of the Arabidopsis thaliana developmental transcriptome based
733 on RNA-seq profiling. *Plant J.* 88: 1058–1070.

734 Krejčí, J., Stixová, L., Pagáčová, E., Legartová, S., Kozubek, S., Lochmanová, G., Zdráhal,
735 Z., Sehnalová, P., Dabravolski, S., Hejátko, J., and Bártová, E. (2015). Post-
736 Translational Modifications of Histones in Human Sperm: Epigenetics of human
737 sperm. *J. Cell. Biochem.* 116: 2195–2209.

738 Kriechbaumer, V., Wang, P., Hawes, C., and Abell, B.M. (2012). Alternative splicing of the
739 auxin biosynthesis gene YUCCA4 determines its subcellular compartmentation. *Plant*
740 *J.* 70: 292–302.

741 Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular
742 Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and*
743 *Evolution* 35: 1547–1549.

- 744 Langowski, L., Wabnik, K., Li, H., Vanneste, S., Naramoto, S., Tanaka, H., and Friml, J.
745 (2016). Cellular mechanisms for cargo delivery and polarity maintenance at different
746 polar domains in plant cells. *Cell Discov* 2: 16018.
- 747 Laňková, M., Humpolíčková, J., Vosolsobě, S., Cit, Z., Lacey, J., Čovan, M., Čovanová, M.,
748 Hof, M., and Petrášek, J. (2016). Determination of Dynamics of Plant Plasma
749 Membrane Proteins with Fluorescence Recovery and Raster Image Correlation
750 Spectroscopy. *Microscopy and Microanalysis* 22: 290–299.
- 751 Lee, J.H., Ryu, H.-S., Chung, K.S., Posé, D., Kim, S., Schmid, M., and Ahn, J.H. (2013).
752 Regulation of temperature-responsive flowering by MADS-box transcription factor
753 repressors. *Science* 342: 628–632.
- 754 Leitner, J., Petrášek, J., Tomanov, K., Retzer, K., Pařezová, M., Korbei, B., Bachmair, A.,
755 Zažímalová, E., and Luschig, C. (2012). Lysine63-linked ubiquitylation of PIN2
756 auxin carrier protein governs hormonally controlled adaptation of Arabidopsis root
757 growth. *PNAS* 109: 8322–8327.
- 758 Li, H.-T. et al. (2019). Origin of angiosperms and the puzzle of the Jurassic gap. *Nat. Plants* 5:
759 461–470.
- 760 Liao, C.-Y., Smet, W., Brunoud, G., Yoshida, S., Vernoux, T., and Weijers, D. (2015).
761 Reporters for sensitive and quantitative measurement of auxin response. *Nat. Methods*
762 12: 207.
- 763 Marquez, Y., Brown, J.W.S., Simpson, C., Barta, A., and Kalyna, M. (2012). Transcriptome
764 survey reveals increased complexity of the alternative splicing landscape in
765 Arabidopsis. *Genome Res.* 22: 1184–1195.
- 766 Marquez, Y., Höpfner, M., Ayatollahi, Z., Barta, A., and Kalyna, M. (2015). Unmasking
767 alternative splicing inside protein-coding exons defines exons and their role in
768 proteome plasticity. *Genome Res.* 25: 995–1007.
- 769 Martinière, A. et al. (2012). Cell wall constrains lateral diffusion of plant plasma-membrane
770 proteins. *PNAS* 109: 12805–12810.
- 771 Men, S., Boutté, Y., Ikeda, Y., Li, X., Palme, K., Stierhof, Y.-D., Hartmann, M.-A., Moritz,
772 T., and Grebe, M. (2008). Sterol-dependent endocytosis mediates post-cytokinetic
773 acquisition of PIN2 auxin efflux carrier polarity. *Nat. Cell Biol.* 10: 237–244.
- 774 Müller, K., Hošek, P., Laňková, M., Vosolsobě, S., Malínská, K., Čarná, M., Fílová, M.,
775 Dobrev, P.I., Helusová, M., Hoyerová, K., and Petrášek, J. (2019). Transcription of
776 specific auxin efflux and influx carriers drives auxin homeostasis in tobacco cells.
777 *Plant J.* 100: 627–640.
- 778 Nakabayashi, K., Bartsch, M., Ding, J., and Soppe, W.J.J. (2015). Seed Dormancy in
779 Arabidopsis Requires Self-Binding Ability of DOG1 Protein and the Presence of
780 Multiple Isoforms Generated by Alternative Splicing. *PLOS Genetics* 11: e1005737.
- 781 Nodzyński, T., Vanneste, S., Zwiewka, M., Pernisová, M., Hejácí, J., and Friml, J. (2016).
782 Enquiry into the Topology of Plasma Membrane-Localized PIN Auxin Transport
783 Components. *Molecular Plant* 9: 1504–1519.

784 O’Leary, N.A. et al. (2016). Reference sequence (RefSeq) database at NCBI: current status,
785 taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 44: D733-745.

786 Pernisova, M., Prat, T., Grones, P., Harustiakova, D., Matonohova, M., Spichal, L.,
787 Nodzynski, T., Friml, J., and Hejatko, J. (2016). Cytokinins influence root
788 gravitropism via differential regulation of auxin transporter expression and
789 localization in *Arabidopsis*. *New Phytol* 212: 497–509.

790 Petrasek, J. et al. (2006). PIN proteins perform a rate-limiting function in cellular auxin
791 efflux. *Science* 312: 914–918.

792 Posé, D., Verhage, L., Ott, F., Yant, L., Mathieu, J., Angenent, G.C., Immink, R.G.H., and
793 Schmid, M. (2013). Temperature-dependent regulation of flowering by antagonistic
794 FLM variants. *Nature* 503: 414–417.

795 Rademacher, E.H. and Offringa, R. (2012). Evolutionary Adaptations of Plant AGC Kinases:
796 From Light Signaling to Cell Polarity Regulation. *Front. Plant Sci.* 3.

797 Rakusová, H., Abbas, M., Han, H., Song, S., Robert, H.S., and Friml, J. (2016). Termination
798 of Shoot Gravitropic Responses by Auxin Feedback on PIN3 Polarity. *Current*
799 *Biology* 26: 3026–3032.

800 Rakusová, H., Gallego-Bartolomé, J., Vanstraelen, M., Robert, H.S., Alabadí, D., Blázquez,
801 M.A., Benková, E., and Friml, J. (2011). Polarization of PIN3-dependent auxin
802 transport for hypocotyl gravitropic response in *Arabidopsis thaliana*. *Plant J* 67: 817–
803 826.

804 Reddy, A.S.N., Marquez, Y., Kalyna, M., and Barta, A. (2013). Complexity of the alternative
805 splicing landscape in plants. *Plant Cell* 25: 3657–3683.

806 Remy, E., Cabrito, T.R., Baster, P., Batista, R.A., Teixeira, M.C., Friml, J., Sá-Correia, I., and
807 Duque, P. (2013). A major facilitator superfamily transporter plays a dual role in polar
808 auxin transport and drought stress tolerance in *Arabidopsis*. *Plant Cell* 25: 901–926.

809 Retzer, K., Butt, H., Korbei, B., and Luschnig, C. (2014). The far side of auxin signaling:
810 fundamental cellular activities and their contribution to a defined growth response in
811 plants. *Protoplasma* 251: 731–746.

812 Rosquete, M.R., von Wangenheim, D., Marhavý, P., Barbez, E., Stelzer, E.H.K., Benková, E.,
813 Maizel, A., and Kleine-Vehn, J. (2013). An Auxin Transport Mechanism Restricts
814 Positive Orthogravitropism in Lateral Roots. *Current Biology* 23: 817–822.

815 Roychoudhry, S., Kieffer, M., Del Bianco, M., Liao, C.-Y., Weijers, D., and Kepinski, S.
816 (2017). The developmental and environmental regulation of gravitropic setpoint angle
817 in *Arabidopsis* and bean. *Sci Rep* 7: 42664.

818 Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., and
819 Eliceiri, K.W. (2017). ImageJ2: ImageJ for the next generation of scientific image
820 data. *BMC Bioinformatics* 18: 529.

821 Ruzicka, K. et al. (2017). Identification of Factors Required for m6A mRNA Methylation in
822 Arabidopsis Reveals a Role for the Conserved E3 Ubiquitin Ligase HAKAI. *New*
823 *Phytol* 215: 157–172.

824 Shang, X., Cao, Y., and Ma, L. (2017). Alternative Splicing in Plant Genes: A Means of
825 Regulating the Environmental Fitness of Plants. *Int J Mol Sci* 18: 432.

826 Sibout, R., Sukumar, P., Hettiarachchi, C., Holm, M., Muday, G.K., and Hardtke, C.S. (2006).
827 Opposite Root Growth Phenotypes of hy5 versus hy5 hyh Mutants Correlate with
828 Increased Constitutive Auxin Signaling. *PLOS Genetics* 2: e202.

829 Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam,
830 H., Remmert, M., Söding, J., Thompson, J.D., and Higgins, D.G. (2011). Fast,
831 scalable generation of high-quality protein multiple sequence alignments using
832 Clustal Omega. *Molecular Systems Biology* 7.

833 Soumpasis, D.M. (1983). Theoretical analysis of fluorescence photobleaching recovery
834 experiments. *Biophysical Journal* 41: 95–97.

835 Sprague, B.L. and McNally, J.G. (2005). FRAP analysis of binding: proper and fitting. *Trends*
836 *in Cell Biology* 15: 84–91.

837 Staiger, D. and Brown, J.W.S. (2013). Alternative splicing at the intersection of biological
838 timing, development, and stress responses. *Plant Cell* 25: 3640–3656.

839 Stamm, S., Ben-Ari, S., Rafalska, I., Tang, Y., Zhang, Z., Toiber, D., Thanaraj, T.A., and
840 Soreq, H. (2005). Function of alternative splicing. *Gene* 344: 1–20.

841 Swarup, K. et al. (2008). The auxin influx carrier LAX3 promotes lateral root emergence.
842 *Nat. Cell Biol.* 10: 946–954.

843 Szakonyi, D. and Duque, P. (2018). Alternative Splicing as a Regulator of Early Plant
844 Development. *Front. Plant Sci.* 9.

845 Teale, W.D. et al. (2020). Flavonol-mediated stabilization of PIN efflux complexes regulates
846 polar auxin transport. *The EMBO Journal* n/a: e104416.

847 Tress, M.L., Abascal, F., and Valencia, A. (2017). Alternative Splicing May Not Be the Key
848 to Proteome Complexity. *Trends in Biochemical Sciences* 42: 98–110.

849 Tseng, Q., Duchemin-Pelletier, E., Deshiere, A., Balland, M., Guillou, H., Filhol, O., and
850 Théry, M. (2012). Spatial organization of the extracellular matrix regulates cell-cell
851 junction positioning. *Proc. Natl. Acad. Sci. U.S.A.* 109: 1506–1511.

852 Tsugeki, R., Tanaka-Sato, N., Maruyama, N., Terada, S., Kojima, M., Sakakibara, H., and
853 Okada, K. (2015). CLUMSY VEIN, the Arabidopsis DEAH-box Prp16 ortholog, is
854 required for auxin-mediated development. *Plant J.* 81: 183–197.

855 Waldie, T. and Leyser, O. (2018). Cytokinin Targets Auxin Transport to Promote Shoot
856 Branching. *Plant Physiol.* 177: 803–818.

857 Willige, B.C., Ahlers, S., Zourelidou, M., Barbosa, I.C.R., Demarsy, E., Trevisan, M., Davis,
858 P.A., Roelfsema, M.R.G., Hangarter, R., Fankhauser, C., and Schwechheimer, C.
859 (2013). D6PK AGCVIII Kinases Are Required for Auxin Transport and Phototropic
860 Hypocotyl Bending in Arabidopsis. *Plant Cell* 25: 1674–1688.

861 Zadnikova, P. et al. (2010). Role of PIN-mediated auxin efflux in apical hook development of
862 *Arabidopsis thaliana*. *Development* 137: 607–617.

863 Zadnikova, P., Wabnik, K., Abuzeineh, A., Galleml, M., Straeten, D.V.D., Smith, R.S., Inzé,
864 D., Friml, J., Prusinkiewicz, P., and Benková, E. (2016). A Model of Differential
865 Growth-Guided Apical Hook Formation in Plants. *Plant Cell* 28: 2464–2477.

866 Zwiewka, M., Bilanovičová, V., Seifu, Y.W., and Nodzyński, T. (2019). The Nuts and Bolts
867 of PIN Auxin Efflux Carriers. *Front Plant Sci* 10.

868

Figure 1. Alternative splicing (AS) of the *Arabidopsis thaliana* PIN7 and PIN4 genes leads to two evolutionarily conserved functionally different transcripts.

(A) Scheme of coding regions of the PIN3 clade genes in *Arabidopsis thaliana*. The alternative donor splice site at the end of the first exon of PIN7 and PIN4, respectively, but not PIN3, results in two transcripts differing in 12 nucleotides. This sequence (orange) corresponds to the protein motif, located in the long internal hydrophilic loop of the transporter. There is also an additional PIN4c transcript present in publicly available transcriptomes.

(B) Amino acid alignment of the region around the 4-amino acid motif changed by AS in the PIN3 clade proteins (boxed in pink) in *Arabidopsis thaliana*, including their closest PIN paralogs.

(C) Table shows the number of RNA-seq reads spanning the exon1-exon2 junction corresponding to the detected PIN7 transcripts in selected *Arabidopsis thaliana* tissue sources. Their ratio was calculated as a percentage of total reads mapped to this area, as assessed from the genome browser graphic interface.

(D) Protein sequence alignments showing the conservation of AS in the PIN3 clade of auxin transporters among rosids.

(E) and **(F)** Phototropic bending of the etiolated *pin347* seedlings carrying the PIN7a-GFP and PIN7b-RFP **(E)**, and the PIN7a-RFP and PIN7b-GFP constructs **(F)**.

(G) and **(H)** Quantification of the primary root protoxylem defects **(G)** and lateral root primordia **(H)** initiation of the *pin347* seedlings harboring the PIN7a-RFP and PIN7b-GFP constructs.

On **(E)**, **(F)** and **(H)**, the middle line corresponds to median, the box corresponds to the 25% and 75% quantiles, the whiskers are the minima and maxima, dots represent single data points. The asterisks indicate a difference between the respective line and the *pin347* mutant (** $P < 0.01$, *** $P < 0.001$ by two-way ANOVA). For each line in each experiment, at least 15 seedlings were analyzed.

Figure 2. The *PIN7* AS reporter system shows overlapping *PIN7a* and *PIN7b* expression in most tissues.

(A) A scheme of the *PIN7* splicing reporter, which consists of two separate constructs. The *PIN7a*-GFP part (PIN7A₁G) contains two point mutations (marked with dashed arrows) and leads to the frameshift and its subsequent restoration when the *PIN7a* transcript is solely produced. *PIN7b*-RFP reporter (PIN7BR) carries a premature stop codon inside the protruding *PIN7a* region (dashed arrow below).

(B) to (D) Predominantly overlapping *PIN7a*-GFP and *PIN7b*-RFP reporter expression in the root tip **(B)**, in the etiolated hypocotyl following 4 h of unilateral light stimulation **(C)**, and in the area covering apical hook **(D)**.

Yellow arrows: direction of unilateral light, white arrows: differential expression of the reporter. For each tissue, at least 9 images were analyzed. Bars, 50 μ m.

Figure 3. *PIN7a-GFP* and *PIN7b-RFP* cDNAs expressed in the *pin347* knock-out mutant regulate apical hook development and hypocotyl bending in a mutually antagonistic manner.

(A) to (C) A temporal dynamics of etiolated *pin347* seedlings carrying both *PIN7a-GFP* and *PIN7b-RFP* cDNA transgenes during (A) apical hook development, and (B) hypocotyl phototropic and (C) gravitropic bending.

Data are means $\pm$ S. E. For each data point, $n = 15$.

Figure 4. Functional analysis of the PIN7a-GFP and PIN7b-RFP interaction.

(A) Expression of *PIN7a* but not *PIN7b* cDNA gives rise to the auxin reporter *DR5v2::GFP* gradient at the concave side of the apical hooks of the *pin347* seedlings.

(B) Comparable [³H]NAA accumulation kinetics in tobacco BY-2 cells following induction of the *G10-90::XVE>>AtPIN7a-GFP* and *AtPIN7b-GFP* transgenes with decreasing doses of β -estradiol.

(C) The effect of 50 μ M brefeldin A (BFA) on the intracellular distribution of PIN7a-GFP and PIN7b-RFP. Arrows mark the incomplete co-localization of PIN7a and PIN7b in the BFA bodies.

(D) A temporal dynamics of the BFA-mediated aggregation of PIN7a-GFP and PIN7b-GFP inside cells. The values were determined as a ratio of fluorescence intensities between the cytoplasm and PM.

(E) and **(F)** An immunoblot (using anti-GFP antibodies) from the native-PAGE gel showing that both PIN7 isoforms can form multimeric complexes. The gel was loaded with the immunoprecipitation eluate (left) and a crude protein extract (right) from the *Arabidopsis* seedlings harboring the *G10-90::XVE>>PIN7a-GFP* and *PIN7b-GFP* constructs, induced with 5 μ M β -estradiol (**E**), loading control (**F**).

(G) and **(H)** PIN7a and PIN7b can form homo- and heteromers. GFP fluorescence lifetime (τ) determined from the tagged PIN7a and PIN7b proteins and compared to the τ of the PIN7a-GFP alone, presented as a representative image heat map (**G**) quantified on (**H**). Proteins were expressed under the control of the *G10-90::XVE* promoter in the *Arabidopsis* root tips and induced with 5 μ M β -estradiol.

On the box plots, the middle line corresponds to median, the box corresponds to the 25% and 75% quantiles, the whiskers represent the minima and maxima. On (**F**), the asterisks indicate a difference between the respective protein pair and the PIN7a-GFP control (* P < 0.05, *** P < 0.001 by two-way ANOVA). On (**D**), data are means $\pm$ S. E. For each line, on (**A**) and (**B**), $n \geq 9$, on (**C**), (**D**), (**G**) and (**H**), $n \geq 12$. Bars, 50 μ m on (**A**), 20 μ m on (**C**), and 10 μ m on G.

Figure 5. cDNA-encoded PIN7a and PIN7b mutually influence their stability within PM, as assessed by fluorescence recovery after photobleaching (FRAP).

(A) Differential FRAP of *PIN7::PIN7a-GFP* and *PIN7::PIN7b-RFP* on PM. The example images of the bleached region (ROI) are shown on the inset.

(B) The FRAP analysis of PIN3Δ-RFP, which lacks the 4 amino acid motif to mimic the properties of the PIN7b isoform (inset), compared to the control *PIN3-GFP* cDNA line.

(C) The FRAP of *SCR::PIN7a-GFP* and *SCR::PIN7b-RFP* expressed in the *pin347* seedlings. The lines with a single construct are marked as “single”, the lines carrying both transgenes are marked as “cross”.

(D) The FRAP analysis of BY-2 tobacco cell lines expressing *G10-90::XVE>>AtPIN7a-GFP* and *AtPIN7b-GFP*, following induction with 5 μ M β -estradiol.

Data are means $\pm$ S. E. For each data point, at least 20 ROIs were measured. The normalized data points were fit with mono-exponential curves. Bar, 5 μ m on **(A)**.