Organelle genome assembly uncovers the dynamic genome reorganization and cytoplasmic male sterility associated genes in tomato.

Running title: Organelle genomes of CMS tomato

Kosuke Kuwabara¹, Issei Harada¹, Yuma Matsuzawa², Tohru Ariizumi^{1,3*}, and Kenta Shirasawa^{4*}

¹Graduate School of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8577, Japan

²TOKITA Seed Co. LTD., Kazo, Saitama 349-1144, Japan

³Tsukuba Plant Innovation Research Center, Tsukuba, Ibaraki 305-8577, Japan

⁴Kazusa DNA Research Institute, Kisarazu, Chiba 292-0818, Japan

E-mail addresses of all authors

KK: s1921107@s.tsukuba.ac.jp

IH: s2021044@s.tsukuba.ac.jp

YM: ymatsuzawa@tokitaseed.co.jp

TA: ariizumi.toru.ge@u.tsukuba.ac.jp

KS: shirasaw@kazusa.or.jp

***Co-corresponding authors:**

Tohru Ariizumi

Tel. and Fax: +81-29 853 4710

Kenta Shirasawa

Tel.: +81-438 52 3935

Fax: +81-438 52 3934

26 **Abstract**

27 To identify cytoplasmic male sterility (CMS)-associated genes in tomato, we determined the genome
 28 sequences of mitochondria and chloroplasts in three CMS tomato lines derived from independent
 29 asymmetric cell fusions, their nuclear and cytoplasmic donors, and male fertile weedy cultivated tomato
 30 and wild relatives. The structures of the CMS mitochondrial genomes were highly divergent from those
 31 of the nuclear and cytoplasmic donors, and genes of the donors were mixed up in these genomes. On the
 32 other hand, the structures of CMS chloroplast genomes were moderately conserved across the donors, but
 33 CMS chloroplast genes were unexpectedly likely derived from the nuclear donors. Comparative analysis
 34 of the structures and contents of organelle genes and transcriptome analysis identified three genes that
 35 were uniquely present in the CMS lines, but not in the donor or fertile lines. RNA sequencing analysis
 36 indicated that these three genes transcriptionally expressed in anther, two of which were also expressed in
 37 pollen. They could be potential candidates for CMS-associated genes. This study suggests that organelle
 38 reorganization mechanisms after cell fusion events differ between mitochondria and chloroplasts, and
 39 provides insight into the development of new F1 hybrid breeding programs employing the CMS system in
 40 tomato.

41

42 **Keywords:** Tomato, cytoplasmic male sterility, organelle genomes, mitochondria, RNA-Seq

43

44 Introduction

45 Cytoplasmic male sterility (CMS) is broadly found in the kingdom of Plantae¹. CMS plants cannot
46 produce seeds by self-pollination due to a lack of male fertility; therefore, pollen from other plants is
47 always required for these plants to produce seeds. CMS is caused by the incompatibility of interactions of
48 genetic information between nuclei and organelles, especially mitochondria¹. The genes in nuclei and
49 organelles are called *restore of fertility* (*RF*) genes and CMS-associated genes, respectively. Therefore,
50 CMS plants have been used as materials for studies of interactions between nuclear and cytoplasmic
51 genes. Moreover, CMS is used in breeding programs to produce F1 hybrid seeds¹, in which cytoplasmic
52 and pollen donors are employed as maternal and paternal parents, respectively.

53 CMS plants can be artificially generated by recurrent backcrossing or transgenic approaches^{2,3},
54 which leads to incompatibility between nuclei and organelles. A tomato CMS line, called CMS-pennellii,
55 which possesses nuclei and cytoplasm from *Solanum pennellii* and *Solanum peruvianum*, respectively,
56 has been developed by recurrent backcrossing². A gene knockdown strategy is also used to develop CMS
57 tomato lines, for which expression of a nuclear gene that regulates mitochondrial substoichiometric
58 shifting has been suppressed³. In addition, other types of CMS tomato lines have been generated via
59 asymmetric cell fusion between cultivated tomato lines, namely, *Solanum lycopersicum* as the nuclear
60 donor and a wild potato relative, *Solanum acaule*, as the cytoplasmic donor⁴. Among CMS lines, MSA1
61 has been well-studied to reveal nucleus-organelle incompatibility⁴. A physical map of the mitochondrial
62 genome of MSA1 indicates that this asymmetric cell fusion hybrid has a complex mitochondrial genome
63 structure consisting of the parental genomes⁵. Transcripts of an open reading frame (ORF), *orf206*, of the
64 hybrid mitochondrial genome are heterogeneously edited⁶. However, no candidates of CMS-associated
65 genes have been identified in tomato.

66 Although CMS-associated gene sequences are not conserved across plant species, they have
67 common features⁷. Most CMS-associated gene candidates usually possess transmembrane regions and
68 chimeric structures, so-called fusion genes, of genes involved in respiration. Based on this information,
69 CMS-associated genes have been identified in *Oryza sativa*^{8,9}, *Helianthus annuus*¹⁰, and *Gossypium*
70 *hirsutum*¹¹. RNA-sequencing (RNA-Seq) based on next-generation sequencing technology has been
71 employed to select candidates uniquely expressed in CMS lines of *Brassica juncea*¹². Further functional
72 studies are required to confirm that these candidates are involved in CMS. Introduction of *RF* genes into
73 CMS lines would be a useful approach because CMS-associated genes can be downregulated in the
74 presence of *RF* genes⁷. Another approach is to introduce CMS-associated genes into fertile lines to induce
75 sterility¹³. More recently, it has become possible to alter or edit gene sequences of mitochondrial genomes
76 with TALEN technology¹⁴. This technology has been used to disrupt CMS-associated genes in
77 mitochondrial genomes and thereby generate *Arabidopsis thaliana*, *Oryza sativa*, and *Brassica napus*
78 with CMS^{14,15}.

79 In parallel with MSA1, as shown in Figure 1, two asymmetric cell fusions were developed

between cultivated tomato lines *S. lycopersicum* ('O' and 'P') as nuclear donors and a wild potato relative, *S. acaule*, as the cytoplasmic donor¹⁶. The nuclear genome backgrounds of the three cell fusion lines including MSA1 were replaced with the genomes of cultivated tomato lines by a repeated backcrossing strategy. The resultant CMS lines are designated 'CMS[MSA1]', 'CMS[O]', and 'CMS[P]'. Therefore, it may be possible to identify CMS-associated genes by comparative analysis of the genomes and transcriptomes of the CMS lines and their nuclear donors. In this study, we determined the sequences of the organelle genomes of the CMS lines and their donors. Subsequently, the genome sequences and gene expression patterns were compared to identify CMS-associated gene candidates. Furthermore, the results of this analysis may provide insights into the cytoplasmic genome features of asymmetric cell fusions.

Results

De novo assembly of chloroplast and mitochondrial genomes

A total of 10.5 Gb reads per sample were obtained from three CMS tomato lines ('CMS[MSA1]', 'CMS[O]', and 'CMS[P]'), three nuclear donors ('Sekai-ichi', 'O', and 'P'), and one cytoplasmic donor (*S. acaule*). Of them, 374 Mb (3.6%) and 566 Mb (5.4%) of reads per sample were aligned on publicly available sequences of mitochondrial and chloroplast genomes, respectively. The reads mapped on the two sets of reference sequences were separately assembled into contig sequences.

Mitochondrial genome sequences were constructed with reads mapped on the mitochondrial reference sequences (Table 1). The mitochondrial genomes of the nuclear donors 'Sekai-ichi', 'O', and 'P' were all constructed from only contigs with assembly sizes of 562.6 kb ($n = 2$, n represents contig numbers), 536.9 kb ($n = 2$), and 553.3 kb ($n = 2$), respectively. In *S. acaule*, 728.4 kb contigs ($n = 7$) for the mitochondrial genome were established. The assembly sizes were longer in the CMS lines than in the nuclear and cytoplasmic donors, specifically, they were 995.2 kb ($n = 7$) in 'CMS[MSA1]', 968.4 kb ($n = 7$) in 'CMS[O]', and 829.3 kb ($n = 5$) in 'CMS[P]'. For chloroplast genomes, total sequence lengths of 389.2 kb ($n = 2$), 349.5 kb ($n = 2$), and 346.9 kb ($n = 2$) were constructed for 'Sekai-ichi', 'O', and 'P', respectively (Table 1). There were two contig sequences in each of the three nuclear donors. The assembly sizes were shorter in 'CMS[MSA1]' (296.6 kb, $n = 1$) and 'CMS[O]' (307.1 kb, $n = 1$) than in the nuclear donors, but longer in 'CMS[P]' (454.1 kb, $n = 3$).

Comparative genome analysis revealed that the mitochondrial genomes of the CMS lines consisted of highly fragmented, repeated, and duplicated sequences derived from both donors throughout the genome (Figure 2). On the other hand, the structures of the chloroplast genomes of the CMS lines were moderately conserved across the nuclear and cytoplasmic donors (Figure 2).

In parallel, we determined the mitochondrial and chloroplast genome sequences of *Solanum pimpinellifolium* LA1670 and *S. lycopersicum* var. *cerasiforme* LA1673 (Table 1). Sequence reads were obtained from a public DNA database and processed as described above. Assembly sizes of the mitochondrial and chloroplast genomes were 620.6 kb ($n = 3$) and 299.4 kb ($n = 1$) for *S. pimpinellifolium*

LA1670, respectively, and 569.9 kb ($n = 2$) and 337.7 kb ($n = 2$) for *S. lycopersicum* var. *cerasiforme* LA1673, respectively.

118

Gene prediction from the organelle genomes

ORFs encoding ≥ 25 amino acids were extracted from the assembled sequences to predict potential genes. The number of potential genes predicted from the chloroplast genome assemblies ranged from 5,130 (*S. acaule*) to 8,165 (‘CMS[P]’) and the number of potential genes predicted from the mitochondrial sequences ranged from 10,326 (‘O’) to 19,170 (‘CMS[MSA1]’) (Table 1).

The ORFs were clustered to identify genes unique to and shared among the CMS lines, nuclear donors, and cytoplasmic donor (Figure 3). The ORFs in the CMS mitochondrial genomes consisted of four types of genes, namely, those unique to the CMS lines (Type 1: 9.4–11.9%), those shared with the nuclear donors only (Type 2: 14.1–17.0%), those shared with the cytoplasmic donor only (Type 3: 8.9–13.2%), and those shared with both the nuclear and cytoplasmic donors (Type 4: 61.8–64.1%). By contrast, the ORFs in the CMS chloroplast genomes mostly consisted of three types of genes, namely, those unique to the CMS lines (Type 1: 1.2–5.9%), those shared with the nuclear donors only (Type 2: 31.2–33.1%), and those shared with both the nuclear and cytoplasmic donors (Type 4: 62.9–65.7%). Few genes shared with the cytoplasmic donor only were found (Type 3: up to 0.1%).

The genome positions of the genes differed according to the gene type and organelle (Figure 4). Type 1 genes in mitochondria were distributed across the genome with some gaps. The positions of Type 2 genes were basically the same as those of Type 1 genes, while Type 3 genes were located in the gaps between Type 1 genes. Type 4 genes were also located in the gaps and at the ends of contig sequences. On the other hand, in chloroplast genomes, the positions of Type 1 and 2 genes overlapped and Type 4 genes were located at the ends of contigs.

139

Screening of CMS-associated gene candidates

To identify candidates of CMS-associated genes in the mitochondrial genomes, we set the following four criteria: 1) amino acid length ≥ 70 , 2) absent from male fertile lines, 3) present in all three CMS lines, and 4) expressed in anthers of the CMS lines. Among the predicted genes in the ‘CMS[P]’, ‘CMS[MSA1]’, and ‘CMS[O]’ mitochondrial genomes, 831, 1,025, and 969 genes encoded ≥ 70 amino acids, respectively. The gene sequences from the CMS lines were compared with the mitochondrial genomes of the nuclear donors (‘Sekai-ichi’, ‘P’, and ‘O’) and *S. pimpinellifolium* LA1670, *S. lycopersicum* var. *cerasiforme* LA1673), *S. pennellii*, and *Nicotiana tabacum*. In total, 183, 272, and 140 genes were selected because they were absent from the nuclear donors and Solanaceae relatives, all of which possess male fertility. Furthermore, we selected 36, 41, and 33 genes commonly present in the CMS lines. The copy numbers of the genes varied. Finally, RNA-Seq reads were mapped on the mitochondrial genomes of the CMS lines. This analysis limited the number of CMS-associated gene candidates to four, including

two identical sequences. The three genes were named *orf137* (two copies in the genome of each CMS line: CMS-PMt002g07240 and CMS-PMt005g13392), *orf193* (one copy: CMS-PMt002g06465), and *orf265* (one copy: CMS-PMt010g15739).

De novo transcriptome assembly was performed in parallel. RNA-Seq data were obtained from the anthers of ‘P’ and ‘CMS[P]’, and assembled into 62 and 43 transcript sequences, respectively, of which 37 ‘P’ and 18 ‘CMS[P]’ transcripts were predicted to have transmembrane domains. Of these sequences, eight were uniquely detected in ‘CMS[P]’. Two genes (STRG.32.1.p1 and STRG.39.1.p1) were identical to *orf137* and *orf265*.

Because two genes were commonly identified in both analyses, a total of nine genes were finally selected as candidates of CMS-associated genes (Table 2). Sequence similarity searches with the mitochondrial and chloroplast genomes indicated that two copies of the STRG.32.1.p1 (*orf137*) sequence (CMS-PMt002g07240 and CMS-PMt005g13392) were present in the mitochondrial genomes of the three CMS lines. A single copy sequence of *orf193* (CMS-PMt002g06465) and a single copy sequence of STRG.39.1.p1 (*orf265*, CMS-PMt010g15739) were found in the mitochondrial genomes of the three CMS lines in addition to that of *S. acaule*. The presence of the three genes in the CMS lines was validated by a PCR assay with the three CMS lines and six fertile lines. The remaining six genes were found in both the CMS and fertile lines. We selected three genes, *orf137*, *orf193*, and *orf265*, as highly potential candidates for CMS-associated genes due to their presence specifically in the CMS mitochondrial genomes and their expression in anthers.

Sequence similarity analysis of the candidate genes

The sequence similarity of the candidate genes including their flanking genome regions in the mitochondrial genome of ‘CMS[P]’ was investigated. A 3,045 bp genome sequence around *orf193* showed high sequence similarity to a 4,682 bp region of the tomato chloroplast genome sequences. The 3,045 bp sequence was split into three sequences containing 1,590, 488, and 1,007 bp (Figure 5A) with highly conserved boundary sequences (Figure 5B). In the 1,590 bp chloroplast genome sequence, a gene encoding *cytochrome f* was encoded; however, the corresponding sequence in the mitochondrial genome had a single base insertion causing a frame-shift mutation (Figure 5C). This mutation broke the ORF of the *cytochrome f* gene and generated two small ORFs, *orf116* and *orf193*.

A portion of *orf265* and its upstream sequences (177 bp in total) showed high similarity to the *ATP synthase subunit 8 (atp8)* gene encoded in the tomato mitochondrial genome (Figure 5D). The remaining sequences of *orf265* lacked similarity to reported sequences. *orf265* was located upstream of the *nad3* and *rps12* genes in the mitochondrial genome. No sequence similarity was observed for *orf137* and the flanking sequence.

Expression analysis of the candidate genes

188 The expression patterns of the candidate genes, *orf137*, *orf193*, and *orf265*, were investigated by RT-PCR.
 189 First, we validated the results of the transcriptome analysis by detecting expression of the three genes in
 190 anthers of ‘CMS[P]’ and ‘CMS[MSA1]’ (Figure 6A). *orf265* was tandemly arrayed with *nad3* and
 191 *rps12*; therefore, we assumed that these three genes were co-transcribed as an operon. As expected,
 192 transcripts spanning the three genes were also detected (Figure 6A). Next, we analyzed gene expression
 193 in leaves, stems, roots, ovaries, and pollen in addition to anthers of Dwarf ‘CMS[P]’ which was a BC3
 194 generation of ‘CMS[P]’ backcrossed with a tomato dwarf cultivar ‘Micro-Tom’. Expression of *orf137*
 195 and *orf265* was detected in all tested tissues, while that of *orf193* was observed in leaves, stems, roots,
 196 ovaries, and anthers (Figure 6B).

197

198 Discussion

199 We determined the mitochondrial and chloroplast genome sequences of CMS lines derived from
 200 asymmetric cell fusions and those of their nuclear and cytoplasmic donors (Table 1). Comparative
 201 analysis of the structures unexpectedly revealed that the cytoplasmic genome structures of the fusions
 202 were rearranged and divergent from those of the cytoplasmic donor (*S. acaule*) and nuclear donors (*S.*
 203 *lycopersicum*) (Figure 2). CMS-PMt003g09846 and CMS-PMt003g11185 were encoded in both the
 204 mitochondrial and chloroplast genomes (Table 2), suggesting that mitochondria and chloroplasts from the
 205 two donors were fused with each other and reorganized even though the cytoplasm of the nuclear donors
 206 was chemically inactivated to generate asymmetric cell fusions. Interestingly, the mitochondrial genomes
 207 of the CMS lines were larger than those of the donors, while the size of chloroplast genomes among the
 208 CMS lines were equivalent (Table 1). In addition, the structures of the mitochondrial genomes were
 209 divergent, while those of the chloroplast genomes were rather conserved (Figure 2). Gene clustering
 210 analysis suggested that both the cytoplasmic and nuclear donors contributed to form mitochondria in the
 211 CMS lines (Figure 3). Furthermore, the structures of the CMS mitochondrial genomes contained patches
 212 of the two genomes of the donors (Figure 4). These results suggest that the mitochondrial genomes of
 213 both donors were highly fragmented at the time of asymmetric cell fusion and reorganized to form a new
 214 mitochondrial genome⁵. This is completely different from our expectation that genomes of the
 215 cytoplasmic donors should be present in CMS lines derived from asymmetric cell fusions. More
 216 interestingly, chloroplasts of the CMS lines consisted only of genes from nuclear donors, not from the
 217 cytoplasmic donor. This unexpected finding has been frequently made in tomato¹⁷, tobacco¹⁸, and
 218 *Brassica*¹⁹. We speculate that interactions of genetic information between nuclei and organelles might be
 219 strict with chloroplasts rather than with mitochondria. The genome and/or organelle reorganization
 220 mechanisms after cell fusions might differ between mitochondria and chloroplasts.

221 Based on genome and transcriptome analyses, nine genes encoded in the mitochondrial genome of
 222 ‘CMS[P]’ were selected as candidate CMS-associated genes (Table 2). Among them, three genes
 223 (*orf193*, STRG.32.1.p1 = *orf137* and STRG.39.1.p1 = *orf265*) were uniquely present in the genomes of

the CMS lines and expressed in their anthers (Table 2 and Figure 6). STRG.32.1.p1 (*orf137*) showed sequence similarity with the CMS-associated protein encoding *cytochrome c subunit 1* (Figure 5). STRG.39.1.p1 (*orf265*) was similar to *ATP synthase subunit 8* at the N-terminus, but lacked similarity in the remaining regions (Figure 5). CMS-associated genes are generally involved in cellular respiration producing energy to generate pollen²⁰, and this is true of both these genes. In many cases, fusion genes have been reported to be CMS-associated genes, e.g., *orf307* in *Oryza sativa*²¹ and *orf72* in *Brassica oleracea*²², and to produce cytotoxic proteins, which lead to male sterility. Knockout mutagenesis with mitoTALENs^{14,15} targeting the candidate genes would be useful to identify CMS-associated genes and to generate CMS lines from normal tomato cultivars.

CMS lines are powerful tools to produce F1 hybrid seeds in breeding programs¹. However, in cereals and fruits including tomato, the *RF* genes are essential for F1 plants to set seeds and bear fruits. Restorer genes for CMS lines have been identified in wild tomato relatives, e.g., *S. pimpinellifolium* LA1670 and *S. lycopersicum* var. *cerasiforme* LA1673¹⁶. Recently, we published the genome sequence data of these two wild relatives²³. We expect *RF* genes for CMS lines to be discovered soon based on this information, although no candidate genes or genetic loci have been reported. Once CMS-associated genes and restorer genes are identified, tomato F1 hybrid seeds can be produced by employing insect pollinators instead of the currently used hand-pollination systems. We propose that CMS-based F1 hybrid breeding programs with insect pollinators can be implemented in tomato breeding programs to reduce the costs of F1 seed production in the future.

243

244 **Materials and methods**

245 *Plant materials*

246 Three tomato CMS lines (‘CMS[MSA1]’, ‘CMS[O]’, and ‘CMS[P]’), three cultivated tomato lines (*S.*
247 *lycopersicum* ‘Sekai-ichi’, ‘O’, and ‘P’), and one potato wild relative (*S. acaule*) were used (Figure 1).

248 ‘CMS[MSA1]’ was developed by repeated backcrossing using ‘O’ as a recurrent parent and a
249 male-sterile tomato, MSA1, as a cytoplasmic donor. MSA1 is an asymmetric cell fusion between the
250 tomato cultivar Sekai-ichi (as the nuclear donor) and the potato wild relative *S. acaule* (as the cytoplasmic
251 donor)⁴. ‘CMS[O]’ was a progeny in repeated backcrossing using ‘O’ as the paternal parent and an
252 asymmetric cell fusion between ‘O’ (as the nuclear donor) and *S. acaule* (as the cytoplasmic donor).

253 ‘CMS[P]’ was also a progeny in backcrossing using ‘P’ as the paternal parent and an asymmetric cell
254 fusion between ‘P’ (as the nuclear donor) and *S. acaule* (as the cytoplasmic donor). Dwarf ‘CMS[P]’ was
255 developed from ‘CMS[P]’ by backcrossing with *S. lycopersicum* ‘Micro-Tom’ (TOMJPF0001), which
256 is a miniature dwarf cultivar²⁴. The putative nuclear and cytoplasmic genomes of the materials are shown
257 in Figure 1.

258

259 *Genome sequence analysis*

260 Total genomic DNA was extracted from young leaves of the six tomato lines (‘CMS[MSA1]’,
261 ‘CMS[O]’, ‘CMS[P]’, ‘Sekai-ichi’, ‘O’, and ‘P’) and *S. acaule* with a Maxwell 16 Instrument and
262 Maxwell 16 Tissue DNA Purification Kits (Promega, Madison, WI, USA). SMRT sequence libraries were
263 constructed with an SMRTbell Express Template Prep Kit (PacBio, Menlo Park, CA, USA) and used for
264 sequencing on a PacBio Sequel system (PacBio). Genome sequence data for *S. pimpinellifolium* LA1670
265 and *S. lycopersicum* var. *cerasiforme* LA1673 were obtained from a public DNA database (DRA
266 accession numbers DRX231405 and DRX231409)²³.

267

268 *Genome assembly and gene prediction*

269 Sequence reads were mapped on reference genome sequences for mitochondria (GenBank accession
270 numbers MF034192, MF034193, NC_035964, and MF98995–MF989957) or chloroplasts (NC_007898)
271 with Organelle_PBA²⁵. Reads mapped on the reference sequences were assembled into contig sequences
272 with Canu²⁶. Potential sequence errors in the contig sequences were corrected twice with the sequence
273 reads by Arrow (PacBio). The corrected contig sequences were aligned back to the reference sequences
274 with Nucmer²⁷ to select highly confident organelle genomes. ORFs (≥ 75 bases) in the organelle genomes
275 were selected as potential genes with ORFfinder (<https://www.ncbi.nlm.nih.gov/orffinder>). The ORF
276 sequences were clustered with CD-HIT²⁸. Transmembrane domains in the gene sequences were predicted
277 by TMHMM²⁹. Sequence similarity searches with the mitochondrial genomes of *S. pennellii*
278 (NC_035964) and *N. tabacum* (NC_006581) were performed by BLAST³⁰ with a threshold E-value of
279 1e-50.

280

281 *RNA expression analysis*

282 Total RNA was extracted from the anthers of P and ‘CMS[P]’ with an RNeasy Plant Mini Kit (QIAGEN,
283 Hilden, Germany). RNA was treated with RNase-free DNase (QIAGEN) and used for sequence library
284 preparation with a TruSeq Stranded mRNA Library Prep Kit (Illumina, San Diego, CA, USA). The
285 resultant libraries were sequenced on NextSeq500 (Illumina) in paired-end, 151 bp mode. After trimming
286 adaptors and low-quality reads by Trim_galore (<https://github.com/FelixKrueger/TrimGalore>) with option
287 -q 30 --length 100 followed by fastp³¹ with option -l 100, transcriptomes were *de novo* assembled by the
288 HiSat2-Stringtie pipeline³² and putative ORFs were searched for annotation by BLASTP³⁰ against the
289 SWISS-PROT database³³.

290 To validate the RNA-Seq results, RT-PCR was performed. In total, 800 ng of total RNA isolated
291 from anthers of the CMS lines or seedlings of *S. acaule* was converted into cDNA with ReverTra Ace
292 (TOYOBO, Osaka, Japan) using a random primer (TAKARA BIO, Kusatsu, Japan). cDNA diluted
293 10-fold with water was used as a template for PCR. The PCR mixture (10 μ L) contained 0.5 μ L cDNA,
294 0.3 μ M primers (Table 3), 2 $\times$ PCR buffer (TOYOBO), 400 μ M dNTPs, and 1 U DNA polymerase (KOD
295 FX Neo, TOYOBO). The thermal cycling conditions were as follows: initial denaturation at 94°C for 3

min; 35 cycles of denaturation at 98°C for 15 s, annealing at 60°C for 30 s, and extension at 68°C for 60 s; and a final extension at 68°C for 3 min. PCR products were separated by electrophoresis in a 1% agarose gel with TAE buffer. Gels were stained with Midori Green Advance (NIPPON Genetics, Tokyo, Japan) to detect DNA bands under ultraviolet illumination.

Data availability

The DDBJ accession numbers of the assembled sequences are LC613090-LC613141. Genome information is available at KaTomicsDB (<http://www.kazusa.or.jp/tomato>).

Acknowledgments

We are grateful to Prof. Kazuo Watanabe (University of Tsukuba, Japan) for providing the DNA material of *S. acaule*. This work was supported by the Kazusa DNA Research Institute Foundation and the Project of the NARO Bio-oriented Technology Research Advancement Institution (Research Program on Development of Innovative Technology, Grant number: 30010A). Seeds of Micro-Tom (TOMJPF0001) was provided from National BioResource Project Tomato (NBRP tomato).

Conflicts of interest

YM is an employee of TOKITA Seed Co. LTD. All other authors declare no competing interests.

Contributions

TA and KS conceived and coordinated the project. YM established the plant materials. KK, IH, and KS collected the data. KK, IH, TA, and KS analyzed and interpreted the data. KK and KS wrote the manuscript with contributions from TA. All authors read and approved the final manuscript.

References

1. Bohra, A., Jha, U.C., Adhimoolam, P., Bisht, D. & Singh, N.P. Cytoplasmic male sterility (CMS) in hybrid breeding in field crops. *Plant Cell Rep* **35**, 967-93 <http://dx.doi.org/10.1007/s00299-016-1949-3> (2016).
2. Petrova, M. *et al.* Characterisation of a cytoplasmic male-sterile hybrid line between *Lycopersicon peruvianum* Mill. × *Lycopersicon pennellii* Corr. and its crosses with cultivated tomato. *Theoretical and Applied Genetics* **98**, 825-830 <http://dx.doi.org/10.1007/s001220051139> (1999).
3. Sandhu, A.P., Abdelnoor, R.V. & Mackenzie, S.A. Transgenic induction of mitochondrial rearrangements for cytoplasmic male sterility in crop plants. *Proc Natl Acad Sci U S A* **104**, 1766-70 <http://dx.doi.org/10.1073/pnas.0609344104> (2007).
4. Melchers, G., Mohri, Y., Watanabe, K., Wakabayashi, S. & Harada, K. One-step generation of

- 332 cytoplasmic male sterility by fusion of mitochondrial-inactivated tomato protoplasts with
333 nuclear-inactivated *Solanum* protoplasts. *Proc Natl Acad Sci U S A* **89**, 6832-6
334 <http://dx.doi.org/10.1073/pnas.89.15.6832> (1992).
- 335 5. Shikanai, T., Kaneko, H., Nakata, S., Harada, K. & Watanabe, K. Mitochondrial genome
336 structure of a cytoplasmic hybrid between tomato and wild potato. *Plant Cell Rep* **17**, 832-836
337 <http://dx.doi.org/10.1007/s002990050493> (1998).
- 338 6. Shikanai, T., Nakata, S., Harada, K. & Watanabe, K. Analysis of the heterogeneous transcripts of
339 the highly edited orf206 in tomato mitochondria. *Plant Cell Physiol* **37**, 692-6
340 <http://dx.doi.org/10.1093/oxfordjournals.pcp.a029000> (1996).
- 341 7. Chen, L. & Liu, Y.G. Male sterility and fertility restoration in crops. *Annu Rev Plant Biol* **65**,
342 579-606 <http://dx.doi.org/10.1146/annurev-arplant-050213-040119> (2014).
- 343 8. Igarashi, K., Kazama, T., Motomura, K. & Toriyama, K. Whole genomic sequencing of RT98
344 mitochondria derived from *Oryza rufipogon* and northern blot analysis to uncover a cytoplasmic
345 male sterility-associated gene. *Plant Cell Physiol* **54**, 237-43
346 <http://dx.doi.org/10.1093/pcp/pcs177> (2013).
- 347 9. Okazaki, M., Kazama, T., Murata, H., Motomura, K. & Toriyama, K. Whole mitochondrial
348 genome sequencing and transcriptional analysis to uncover an RT102-type cytoplasmic male
349 sterility-associated candidate Gene Derived from *Oryza rufipogon*. *Plant Cell Physiol* **54**,
350 1560-8 <http://dx.doi.org/10.1093/pcp/pct102> (2013).
- 351 10. Makarenko, M.S. *et al.* Characterization of the mitochondrial genome of the MAX1 type of
352 cytoplasmic male-sterile sunflower. *BMC Plant Biol* **19**, 51
353 <http://dx.doi.org/10.1186/s12870-019-1637-x> (2019).
- 354 11. Li, S. *et al.* The comparison of four mitochondrial genomes reveals cytoplasmic male sterility
355 candidate genes in cotton. *BMC Genomics* **19**, 775 <http://dx.doi.org/10.1186/s12864-018-5122-y>
356 (2018).
- 357 12. Wu, Z. *et al.* Mitochondrial genome and transcriptome analysis of five alloplasmic male-sterile
358 lines in *Brassica juncea*. *BMC Genomics* **20**, 348 <http://dx.doi.org/10.1186/s12864-019-5721-2>
359 (2019).
- 360 13. Yang, J., Liu, X., Yang, X. & Zhang, M. Mitochondrially-targeted expression of a cytoplasmic
361 male sterility-associated orf220 gene causes male sterility in *Brassica juncea*. *BMC Plant Biol* **10**,
362 231 <http://dx.doi.org/10.1186/1471-2229-10-231> (2010).
- 363 14. Kazama, T. *et al.* Curing cytoplasmic male sterility via TALEN-mediated mitochondrial genome
364 editing. *Nat Plants* **5**, 722-730 <http://dx.doi.org/10.1038/s41477-019-0459-z> (2019).
- 365 15. Arimura, S.I. *et al.* Targeted gene disruption of ATP synthases 6-1 and 6-2 in the mitochondrial
366 genome of *Arabidopsis thaliana* by mitoTALENs. *Plant J* **104**, 1459-1471
367 <http://dx.doi.org/10.1111/tpj.15041> (2020).

- 368 16. Harada, K., Kondo, K. & Watabe, K. Creation of fl tomato plant recovered in fertility. (1995).
- 369 17. Bonnema, A.B., Melzer, J.M. & O'Connell, M.A. Tomato cybrids with mitochondrial DNA from
370 *Lycopersicon pennelli*. *Theor Appl Genet* **81**, 339-48 <http://dx.doi.org/10.1007/BF00228674>
371 (1991).
- 372 18. Sidorov, V.A., Menczel, L., Nagy, F. & Maliga, P. Chloroplast transfer in *Nicotiana* based on
373 metabolic complementation between irradiated and iodoacetate treated protoplasts. *Planta* **152**,
374 341-5 <http://dx.doi.org/10.1007/BF00388259> (1981).
- 375 19. Morgan, A. & Maliga, P. Rapid chloroplast segregation and recombination of mitochondrial
376 DNA in *Brassica* cybrids. *Mol Gen Genet* **209**, 240-6 <http://dx.doi.org/10.1007/BF00329649>
377 (1987).
- 378 20. Touzet, P. & Meyer, E.H. Cytoplasmic male sterility and mitochondrial metabolism in plants.
379 *Mitochondrion* **19 Pt B**, 166-71 <http://dx.doi.org/10.1016/j.mito.2014.04.009> (2014).
- 380 21. Fujii, S., Kazama, T., Yamada, M. & Toriyama, K. Discovery of global genomic re-organization
381 based on comparison of two newly sequenced rice mitochondrial genomes with cytoplasmic
382 male sterility-related genes. *BMC Genomics* **11**, 209 <http://dx.doi.org/10.1186/1471-2164-11-209>
383 (2010).
- 384 22. Shinada, T., Kikuchi, Y., Fujimoto, R. & Kishitani, S. An alloplasmic male-sterile line of
385 *Brassica oleracea* harboring the mitochondria from *Diplotaxis muralis* expresses a novel
386 chimeric open reading frame, orf72. *Plant Cell Physiol* **47**, 549-53
387 <http://dx.doi.org/10.1093/pcp/pcj014> (2006).
- 388 23. Takei, H. *et al.* De novo genome assembly of two tomato ancestors, *solanum pimpinellifolium*
389 and *S. lycopersicum* var. *cerasiforme*, by long-read sequencing. *DNA Res*
390 <http://dx.doi.org/10.1093/dnares/dsaa029> (2021).
- 391 24. Scott, J.W. & Harbaugh, B.K. Micro-tom: A Miniature Dwarf Tomato (Agricultural Experiment
392 Station, Institute of Food and Agricultural Sciences, University of Florida, 1989).
- 393 25. Soorni, A., Haak, D., Zaitlin, D. & Bombarely, A. Organelle_PBA, a pipeline for assembling
394 chloroplast and mitochondrial genomes from PacBio DNA sequencing data. *BMC Genomics* **18**,
395 49 <http://dx.doi.org/10.1186/s12864-016-3412-9> (2017).
- 396 26. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting
397 and repeat separation. *Genome Res* **27**, 722-736 <http://dx.doi.org/10.1101/gr.215087.116> (2017).
- 398 27. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome Biol* **5**, R12
399 <http://dx.doi.org/10.1186/gb-2004-5-2-r12> (2004).
- 400 28. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-generation
401 sequencing data. *Bioinformatics* **28**, 3150-2 <http://dx.doi.org/10.1093/bioinformatics/bts565>
402 (2012).
- 403 29. Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E.L. Predicting transmembrane protein

404 topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**,
405 567-80 <http://dx.doi.org/10.1006/jmbi.2000.4315> (2001).

406 30. Altschul, S.F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein database
407 search programs. *Nucleic Acids Res* **25**, 3389-402 <http://dx.doi.org/10.1093/nar/25.17.3389>
408 (1997).

409 31. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
410 *Bioinformatics* **34**, i884-i890 <http://dx.doi.org/10.1093/bioinformatics/bty560> (2018).

411 32. Pertea, M., Kim, D., Pertea, G.M., Leek, J.T. & Salzberg, S.L. Transcript-level expression
412 analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc* **11**, 1650-67
413 <http://dx.doi.org/10.1038/nprot.2016.095> (2016).

414 33. Bairoch, A. & Apweiler, R. The SWISS-PROT protein sequence database and its supplement
415 TrEMBL in 2000. *Nucleic Acids Res* **28**, 45-8 <http://dx.doi.org/10.1093/nar/28.1.45> (2000).

416

417

418 **Figure legends**

419 **Figure 1** Pedigree of the CMS tomato lines.

420 Squares and circles indicate cytoplasm and nuclei, respectively. Arrows with dashed lines and doubled
421 lines indicate cell fusions and crossings, respectively.

422 **Figure 2** Comparative maps of the organelle genomes of the CMS tomato lines.

423 Mitochondrial (A) and chloroplast (B) genomes of the three CMS lines, nuclear donors, and cytoplasmic
424 donor. Dots indicate sequence similarity between the genome sequences.

425 **Figure 3** Organelle genes in the CMS lines, nuclear donors, and cytoplasmic donor.

426 Numbers of genes unique to the CMS lines, nuclear donors, and cytoplasmic donor are indicated in bold,
427 standard, and italic fonts, respectively. Percentages of genes are shown in parentheses.

428 **Figure 4** Distributions of CMS tomato genes across the organelle genomes.

429 Dots indicate gene positions on contig sequences of the organelle genomes. Genes are grouped into the
430 following four types: Type 1, genes unique to the CMS lines; Type 2, genes shared with the nuclear
431 donors; Type 3, genes shared with the cytoplasmic donor; and Type 4, genes shared with both the nuclear
432 and cytoplasmic donors.

433 **Figure 5** Structures of mitochondrial genes in ‘CMS[P]’.

434 A. Genome structure of the *orf193* region. Homologous sequences between the two genomes are
435 indicated by gray boxes. Highly conserved sequences at the borders are shown in red and blue. B.
436 Sequence alignments of the borders. C. Details of the genome structure of the *orf193* region. A single
437 nucleotide insertion causing a frame-shift mutation is indicated with a red arrow. D. Genome structure of
438 the *orf265* region.

439 **Figure 6** RT-PCR analysis of the CMS-associated gene candidates.

440 Gene expression patterns in anthers of two CMS lines (A) and in seven samples of Dwarf ‘CMS[P]’ (B).
441 *cox2* is a positive control.

442

Table 1 Assembly data of the organelle genomes.

Organelle	Male sterile			Male fertile					
	CMS lines			Nuclear donors			Cytoplasmic donor	Tomato wild relative and weedy tomato	
	CMS[MSA1]	CMS[O]	CMS[P]	Sekai-ichi	O	P	<i>S. acaule</i>	LA1670	LA1673
<i>Mitochondrion</i>									
Number of sequences	7	7	5	2	2	2	7	3	2
Total length (bp)	995,217	968,425	829,310	562,630	536,932	553,289	728,387	620,567	569,852
Number of genes	19,170	18,623	15,912	10,782	10,326	10,653	13,898	11,920	10,965
<i>Chloroplast</i>									
Number of sequences	1	1	3	2	2	2	1	1	2
Total length (bp)	296,583	307,105	454,083	389,209	349,506	346,936	284,040	299,393	337,655
Number of genes	5,279	5,456	8,165	6,971	6,267	6,192	5,130	5,346	5,995

Table 2 Copy numbers of CMS-associated gene candidates in the organelle genomes.

Gene ID of CMS[PF]	Candidates from genome analysis	Candidates from transcriptome analysis	Copy number in mitochondrial genome							Copy number in chloroplast genome						
			CMS[MSAI],	CMS[O]	CMS[P]	Sekai-ichi	O	P	<i>S. acule</i>	CMS[MSAI],	CMS[O]	CMS[P]	Sekai-ichi	O	P	<i>S. acule</i>
CMS-PMt002g06465	<i>orf193</i>		1	1	1	0	0	0	1	0	0	0	0	0	0	0
CMS-PMt002g07240 and CMS-PMt005g13392	<i>orf137</i>	STRG.32.1.p1	2	2	2	0	0	0	0	0	0	0	0	0	0	0
CMS-PMt002g07993, CMS-PMt003g11130, and CMS-PMt004g12510		STRG.22.1.p1	1	1	3	1	1	1	1	0	0	0	0	0	0	0
CMS-PMt003g09515		STRG.5.1.p1	0	0	1	0	0	0	0	0	0	0	0	0	0	0
CMS-PMt003g09846		STRG.8.1.p1	0	0	1	0	0	0	0	1	1	3	2	1	2	1
CMS-PMt003g11185		STRG.18.1.p1	0	0	1	0	0	0	0	2	2	3	1	2	2	2
CMS-PMt010g15327 and CMS-PMt010g15548		STRG.31.1.p1	2	2	2	0	1	1	1	0	0	0	0	0	0	0
CMS-PMt010g15739	<i>orf265</i>	STRG.39.1.p1	1	1	1	0	0	0	1	0	0	0	0	0	0	0
CMS-PMt010g15740		STRG.39.1.p3	1	1	1	1	1	1	1	0	0	0	0	0	0	0

Table 3 Oligonucleotide sequences of PCR primers

Target gene	Forward primer (5' - 3')	Reverse primer (5' - 3')
<i>orf137</i>	CGATTGAGAAAGCGGCAGGC	GTTATTTTCGCTGCAACGGCG
<i>orf193</i>	GGGGAATCGGCCTTCTTTAGTC	GGGGAGGGTTTAATAAAGGAGCTG
<i>orf265</i>	CGGAGTGAAGCTGTATTGAGGG	GAGGAGAGGAACGAAGAACGAAAC
<i>orf265-rps12</i>	CGGAGTGAAGCTGTATTGAGGG	GATCCGGAATTCCCAGCAAATCC
<i>cox2</i>	CCCGCAAAGGATTGTTTCATGG	CGTATAGGGCTCTTTGCTGGTAG

A

Mitochondrion

Sekai-ichi

S. acaule

O

S. acaule

P

S. acaule

B

Chloroplast

Sekai-ichi

S. acaule

O

S. acaule

P

S. acaule

CMS[MSA1]

CMS[MSA1]

CMS[O]

CMS[O]

CMS[P]

CMS[P]

Mitochondrion

Chloroplast

CMS[MSA1] Mt genome

CMS[MSA1] Cp genome

CMS[O] Mt genome

CMS[O] Cp genome

CMS[P] Mt genome

CMS[P] Cp genome

A

B

C

D

A

'CMS[P]' anther

'CMS[MSA1]' anther

*orf137**orf193**orf265**orf265 - rps12**cox2***B**

'Dwarf CMS[P]'

Pollen (Dried)

Pollen (Incubated)

Anther

Leaf

Stem

Root

Ovary