TRAIT2D: a Software for Quantitative Analysis of Single Particle Diffusion Data

Francesco Reina^{1,2,✉}, John M. A. Wigg², Mariia Dmitrieva³, Joël Lefebvre⁴, Jens Rittscher³, and Christian Eggeling^{1,2,5}

¹Leibniz-Institut für Photonische Technologien e.V., Jena, Germany

²Institute of Applied Optics and Biophysics, Friedrich Schiller Universität, Jena, Germany

³Department of Engineering Science, University of Oxford, Oxford, UK

⁴Département d'informatique, Université du Québec à Montréal, Québec, Canada

⁵MRC Human Immunology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, UK

Summary: Single Particle Tracking (SPT) is one of the most widespread techniques to evaluate particle mobility in a variety of situations, such as in cellular and model membrane dynamics. The proposed TRAIT2D Python library is developed to provide object tracking, trajectory analysis and produce simulated datasets with graphical user interface. The tool allows advanced users to customise the analysis to their requirements.

Availability and implementation: the software has been coded in Python, and can be accessed from: <https://github.com/Eggeling-Lab-Microscope-Software/TRAIT2D>, or the pypi and conda-forge repositories.

A comprehensive user guide is provided at <https://eggeling-lab-microscope-software.github.io/TRAIT2D/>.

Data Analysis | Object Tracking | Single Particle Tracking | Microscopy | Release Software

Correspondence: francesco.reina@uni-jena.de

Introduction

Single Particle Tracking (SPT) is a method to quantify particle dynamics in a sample, such as lipids diffusing on biological membranes (1). This is indifferent to the microscopy technique adopted for detection, and in principle only requires the sample to be sparse enough so that the target particles be identifiable singularly.

Among the earliest example of SPT experiments, we can mention the testing of Einstein's theory of Brownian motion (2). The introduction of electronic camera detection and innovative microscopy systems have made it possible to collect SPT data with framerates up to 50kHz (3–6).

While seemingly straightforward, the analysis of SPT data is not immune from potential pitfalls. Conventionally, trajectory analysis in diffusive systems is performed following Einstein's description of Brownian motion (7) and similar models (8, 9). This theoretical derivation, however, does not take into account the localization error (10) and motion blurring (11). Overlooking these sources of error could lead to overestimate diffusion coefficients (12), or to erroneously detected subdiffusion (13).

Several toolboxes and algorithms have been produced to fulfill the need to identify and track single particles (14, 15, etc), and analyse their motion employing conventional Mean Squared Displacement analysis (16), Deep Learning (17, 18) and other methods (19, 20). However, while in many cases the source code is freely available, they are not platform-agnostic and fully open-source. On the other hand, tools

available on free platforms, such as FIJI or ICY, are not easily customisable.

With the TRacking AnlysIs Toolkit, TRAIT2D, we aim to provide a fully customisable tool for particle tracking, simulation, and analysis. In TRAIT2D, intuitive graphical user interfaces facilitate users with little to no coding experience to develop their own analysis pipelines. A simple 2D tracking algorithm is provided to extract trajectories from SPT data. The source code allows more advanced users to customize data analysis. Furthermore, a track simulator is included as a validation tool of individual analysis pipelines. This allows the user to generate movies of diffusing particles at variable levels of signal-to-noise ratio, with a user-specified point spread function.

Implementation

TRAIT2D is a cross-platform Python software package with graphical user interfaces (GUIs) to support SPT experiments. For simplicity, the software can be divided into three main sections: the tracker, the simulator and the data analyser, which can be used jointly (Figure 1). The details presented here are expanded in the Supplementary Notes.

The tracker extracts particle trajectories from a temporal image sequence. It allows a user to load and visualise TIFF timelapse stack. Once the image sequence is loaded, the parameters of the tracking can be set for detection, sub-pixel localisation and linking. The detection is implemented with Spot Enhancing Filter (SEF) (21), while sub-pixel localisation is achieved using the radial symmetry center approach (22). Particle trajectory linking takes into account spatial and temporal distances between detections and exploits the Hungarian algorithm (23) for data association.

The simulator generates tracks in a virtual sandbox of arbitrary dimensions based on user-adjustable parameters, such as particle diffusion coefficient and time interval between localization. It is possible to simulate particles diffusing in simple Brownian diffusion and hop diffusion modes (24). Once the particle track has been generated, it is also possible to generate a complete movie by performing the convolution of the track with a point spread function, using varying levels of Signal-to-Noise ratio and pixelation. The last phenomenon has been implemented to model camera detectors usually employed for SPT. The purpose of this module is to provide

Fig. 1. Workflow of the TRAIT2D software package. There are three possible avenues for the user to analyse their data: the user can import their timelapses, as tiff stacks, into the Tracker GUI and extract particle trajectories (*right*), pre-determined tracks can be imported in the Analyser GUI from CSV files, with the importer function present in the Analyser GUI *center*, or control datasets can be simulated in the Simulator GUI, and analysed like experimental trajectories. Finally, the Analyser GUI will perform the analysis by following the Mean Squared Displacement (MSD) or the Apparent Diffusion Coefficient (ADC) pipelines (Supplementary Note 4).

computational controls and validation data sets for SPT detection and tracking algorithms.

Finally, the analyser allows the user to quantitatively evaluate tracks extracted from SPT movies following a Mean Squared Displacement (MSD) and Apparent Diffusion Coefficient (ADC) - based approach. Data analysis algorithms are accessible via the code or from a readily compilable GUI. Data can be obtained through the TRAIT2D-tracker, simulated by the simulator GUI or library, or imported from an externally generated comma-separated value (csv) file (Figure 1). The data analysis pipeline hereby employed is described in the Supplementary Materials (11, 12). The presence of a GUI for model fitting makes this procedure especially accessible and intuitive. The user can choose to fix or to put constraints on the fitting variables, as well as set the interval of data to be considered for the operation. It is also possible for the user to adopt alternative motion models which are not pre-compiled in the system (see documentation). The source code, together with exemplary Jupyter notebooks for easier understanding, can be retrieved at <https://github.com/Eggeling-Lab-Microscope-Software/TRAIT2D>. A comprehensive documentation can be found at <https://eggeling-lab-microscope-software.github.io/TRAIT2D/>.

Conclusions

TRAIT2D is a Python-based, open source and easily deployable toolkit for the analysis and simulation of bidimensional SPT data. The combination of particle tracking, simulation, and trajectory analysis in the same package provides a num-

ber of benefits for the user. Another feature of this software are its user-friendliness for inexperienced users, owing to the presence of a clear graphical user interface, while retaining the potential for customisation for more advanced users.

Acknowledgements

FR and CE would like to acknowledge Dr. B. Christoffer Lagerholm for his invaluable scientific advice. The authors would like to thank Dr. J. Keller-Findeisen for allowing the translation of the original algorithm for compartmentalised diffusion simulation.

Funding

FR, CE and JMAW were supported by Wolfson Foundation, the Medical Research Council (MRC, grant number MC_UU_12010/unit programmes G0902418 and MC_UU_12025), MRC/BBSRC/EPSC (grant number MR/K01577X/1), the Wellcome Trust (grant ref. 104924/14/Z/14), the Deutsche Forschungsgemeinschaft (Research unit 1905 “Structure and function of the peroxisomal translocon”, Jena Excellence Cluster “Balance of the Microverse”, Collaborative Research Center 1278 “Polytarget”), Oxford-internal funds (John Fell Fund and EPA Cephalosporin Fund) and Wellcome Institutional Strategic Support Fund (ISSF). JL was supported by a FRQNT postdoctoral fellowship (257844). MD and JR were funded by a Wellcome Collaborative award (203285).

Bibliography

1. G. J. Schütz, H. Schindler, and Th Schmidt. Single-molecule microscopy on model membranes reveals anomalous diffusion. *Biophysical Journal*, 73(2):1073–1080, 1997. ISSN 00063495. doi: 10.1016/S0006-3495(97)78139-6.
2. Jean Perrin. *Brownian Movement and Molecular Reality*. Taylor & Francis, London, 1910.
3. Akihiro Kusumi, Chieko Nakada, Ken Ritchie, Kotonori Murase, Kenichi Suzuki, Hideji Murakoshi, Rinsai S. Kasai, Junko Kondo, and Takahiro Fujiwara. Paradigm Shift of the Plasma Membrane Concept from the Two-Dimensional Continuum Fluid to the Partitioned Fluid: High-Speed Single-Molecule Tracking of Membrane Molecules. *Annual Review of Biophysics and Biomolecular Structure*, 34(1):351–378, 2005. ISSN 1056-8700. doi: 10.1146/annurev.biophys.34.040204.144637.
4. Katelyn M Spillane, Jaime Ortega-Arroyo, Gabrielle de Wit, Christian Eggeling, Helge Ewers, Mark I Wallace, and Philipp Kukura. High-Speed Single-Particle Tracking of GM1 in Model Membranes Reveals Anomalous Diffusion due to Interleaflet Coupling and Molecular Pinning. *Nano letters*, 14(9):5390–7, 2014. ISSN 1530-6992. doi: 10.1021/nl502536u.
5. Francisco Balzarotti, Yvan Eilers, Klaus C Gwosch, Arvid H Gynnä, Volker Westphal, Fernando D Stefani, Johan Elf, and Stefan W Hell. Nanometer resolution imaging and tracking of fluorescent molecules with minimal photon fluxes. *Science*, 355(6325):606–612, feb 2017. ISSN 0036-8075. doi: 10.1126/science.aak9913.
6. Richard W Taylor and Vahid Sandoghdar. *Interferometric Scattering (iSCAT) Microscopy and Related Techniques*, pages 25–65. Springer International Publishing, Cham, 2019. ISBN 978-3-030-21722-8. doi: 10.1007/978-3-030-21722-8_2.
7. Albert Einstein. On the Motion of Small Particles Suspended in a Stationary Liquid, as Required by the Molecular Kinetic Theory of Heat. *Annalen der Physik*, 322:549–560, 1905. ISSN 1521-3889. doi: 10.1002/andp.19053220806.
8. M. J. Saxton. Single-particle tracking: models of directed transport. *Biophysical Journal*, 67(5):2110–2119, 1994. ISSN 00063495. doi: 10.1016/S0006-3495(94)80694-0.
9. Stefan Wieser, Manuel Moertelmaier, Elke Fuertbauer, Hannes Stockinger, and Gerhard J. Schütz. (Un)confined diffusion of CD59 in the plasma membrane determined by high-resolution single molecule microscopy. *Biophysical Journal*, 92(10):3719–3728, 2007. ISSN 00063495. doi: 10.1529/biophysj.106.095398.
10. Xavier Michalet. Mean square displacement analysis of single-particle trajectories with localization error: Brownian motion in an isotropic medium. *Physical Review E*, 82(4):041914, oct 2010. ISSN 1539-3755. doi: 10.1103/PhysRevE.82.041914.
11. Mark Goulian and Sanford M. Simon. Tracking Single Proteins within Cells. *Biophysical Journal*, 79(4):2188–2198, oct 2000. ISSN 0006-3495. doi: 10.1016/S0006-3495(00)76467-8.
12. B. Christoffer Lagerholm, Débora M. Andrade, Mathias P. Clausen, and Christian Eggeling. Convergence of lateral dynamic measurements in the plasma membrane of live cells from single particle tracking and STED-FCS. *Journal of Physics D: Applied Physics*, 50(6): 063001, 2017. ISSN 0022-3727. doi: 10.1088/1361-6463/aa519e.

13. Douglas S. Martin, Martin B. Forstner, and Josef A. Käs. Apparent subdiffusion inherent to single particle tracking. *Biophysical Journal*, 83(4):2109–2117, 2002. ISSN 00063495. doi: 10.1016/S0006-3495(02)73971-4.
14. Jean Yves Tinevez, Nick Perry, Johannes Schindelin, Genevieve M. Hoopes, Gregory D. Reynolds, Emmanuel Laplantine, Sebastian Y. Bednarek, Spencer L. Shorte, and Kevin W. Eliceiri. TrackMate: An open and extensible platform for single-particle tracking. *Methods*, 115:80–90, 2017. ISSN 10959130. doi: 10.1016/j.ymeth.2016.09.016.
15. Simon Christoph Stein and Jan Thiart. TrackNTTrace: A simple and extendable open-source framework for developing single-molecule localization and tracking algorithms. *Scientific Reports*, 6(November):37947, dec 2016. ISSN 2045-2322. doi: 10.1038/srep37947.
16. Frederik W. Lund, Maria Louise V. Jensen, Tanja Christensen, Gitte K. Nielsen, Christian W. Heegaard, and Daniel Wüstner. SpatTrack: An imaging toolbox for analysis of vesicle motility and distribution in living cells. *Traffic*, 15(12):1406–1429, 2014. ISSN 16000854. doi: 10.1111/tra.12228.
17. Naor Granik, Lucien E. Weiss, Elias Nehme, Maayan Levin, Michael Chein, Eran Perlson, Yael Roichman, and Yoav Shechtman. Single-particle diffusion characterization by deep learning. *Biophysical Journal*, 117(2):185 – 192, 2019. ISSN 0006-3495. doi: <https://doi.org/10.1016/j.bpj.2019.06.015>.
18. Marloes Arts, Ihor Smal, Maarten W. Paul, Claire Wyman, and Erik Meijering. Particle mobility analysis using deep learning and the moment scaling spectrum. *Scientific Reports*, 9, 2019.
19. Samudrajit Thapa, Michael A. Lomholt, Jens Krog, Andrey G. Cherstvy, and Ralf Metzler. Bayesian analysis of single-particle tracking data using the nested-sampling algorithm: maximum-likelihood model selection applied to stochastic-diffusivity data. *Physical Chemistry Chemical Physics*, 20(46):29018–29037, 2018. ISSN 1463-9076. doi: 10.1039/C8CP04043E.
20. Anders S. Hansen, Maxime Woring, Jonathan B. Grimm, Luke D. Lavis, Robert Tjian, and Xavier Darzacq. Robust model-based analysis of single-particle tracking experiments with spot-on. *eLife*, 7:1–33, 2018. ISSN 2050084X. doi: 10.7554/eLife.33125.
21. I Smal, M Loog, W Niessen, and E Meijering. Quantitative comparison of spot detection methods in fluorescence microscopy. *IEEE Trans. on Medical Imaging*, 29(2):282–301, 2010.
22. Parthasarathy R. Rapid, accurate particle tracking by calculation of radial symmetry centers. *Nature methods*, (9(7)):724–726, 2012. doi: 10.1038/nmeth.2071.
23. H. W. Kuhn. The hungarian method for the assignment problem. *Naval Research Logistics Quarterly*, 2(1-2):83–97, 1955.
24. Débora M. Andrade, Mathias P. Clausen, Jan Keller, Veronika Mueller, Congying Wu, James E. Bear, Stefan W. Hell, B. Christoffer Lagerholm, and Christian Eggeling. Cortical actin networks induce spatio-temporal confinement of phospholipids in the plasma membrane – a minimally invasive investigation by STED-FCS. *Scientific Reports*, 5(May):11454, 2015. ISSN 2045-2322. doi: 10.1038/srep11454.