Title

Localised expression of *OsIAA29* suggests a key role for auxin in regulating development of the dorsal aleurone of early rice grains.

Authors

Maftroz A. Basunia^{a,b}, Heather M. Nonhebel^a, David Backhouse^c and Mary McMillan^a.

Affiliations

^a School of Science and Technology, University of New England, Armidale, New South Wales 2351, Australia

^b Present address, Department of Biochemistry and Molecular Biology, Jahangirnagar University, Dhaka 1342, Bangladesh

^c School of Environmental and Rural Science, University of New England, Armidale, New South Wales 2351, Australia

Correspondence

H. M. Nonhebel; Email, hnonheb2@une.edu.au; Phone, +61 267732083

ORCID IDs

MAB 0000-0002-3433-2615, HMN 0000-0001-9055-3095, DB 0000-0003-0663-6002, MM 0000-0002-2336-3985

Main Conclusion

Non-canonical AUX/IAA protein, *OsIAA29*, and ZmMPR-1 homologues, *OsMRPLs*, are part of an auxin-related signalling cascade operating in the dorsal aleurone during early rice grain development.

Abstract

Endosperm of rice and other cereals accumulates high concentrations of the predominant *in planta* auxin, indole-3-acetic acid (IAA) during early grain development. However, IAA signalling and function during endosperm development are poorly understood. Here, we report that *OsYUC12* (an auxin biosynthesis gene) and *OsIAA29* (encoding a non-canonical AUX/IAA) are both expressed exclusively in grains, reaching a maximum 5 to 6 days after pollination. *OsYUC12* expression is localized in the aleurone, sub-aleurone and embryo, whereas *OsIAA29* expression is restricted to a narrow strip in the dorsal aleurone, directly under the vascular bundle. Although rice has been reported to lack endosperm transfer cells (ETCs), this region of the aleurone is enriched with sugar transporters and is likely to play a key role in apoplastic nutrient transfer, analogous to ETCs in other cereals. *OsIAA29* has orthologues only in grass species; expression of which is also specific to early grain development. *OsYUC12* and *OsIAA29* are temporally co-expressed with two genes (*ALI* and *OsPR602*) previously linked to the development of dorsal aleurone or ETCs. Also up regulated at the

same time are a cluster of MYB-related genes (designated *OsMRPLs*) homologous to *ZmMRP-1*, which regulates maize ETC development. Wheat homologues of *ZmMRP-1* are also expressed in ETCs. Although previous work has suggested that other cereals do not have orthologues of *ZmMRP-1*, our work suggests *OsIAA29* and *OsMRPLs* and their homologues in other grasses are part of an auxin-regulated, conserved signalling network involved in the differentiation of cells with ETC-like function in developing cereal grains.

Key words

Dorsal aleurone, auxin, endosperm, *ZmMRP-1*, *OsYUC12*, *OsIAA29*

Introduction

The yield and quality of cereal grains is dependent on the coordinated regulation of endosperm development, uptake of photosynthate and production of storage molecules, and is highly susceptible to adverse environmental influences (Yu et al. 2015). In addition, the molecular and physiological events taking place during early endosperm development determine to a large extent the final grain size and weight in rice and other cereals (Mizutani et al. 2010; Fahy et al. 2018). There is a large body of literature reporting the involvement of plant hormones in the regulation of early endosperm development as well as its response to environment (reviewed by Basunia and Nonhebel 2019). However, little is known of the detailed role of hormonal signalling and how this influences key processes of endosperm cellularisation, differentiation of cells responsible for nutrient uptake or the expression of starch and storage protein synthesis genes.

Previous work in our laboratory has shown that a large increase in the auxin, indole-3-acetic acid (IAA), occurs during endosperm cellularisation, aleurone development and the initiation of starch production in rice grains, driven by strong up-regulation of key auxin biosynthesis genes, *OsTAR1*, *OsYUC9*, *OsYUC11* and *OsYUC12* (Abu-Zaitoon et al. 2012; Russell French et al. 2014; Nonhebel and Griffin 2020). This work identified small differences in the expression profiles of *OsYUC9*, *OsYUC11* and *OsYUC12*, suggesting that some sub-functionalisation may occur. *OsYUC12* was found to be exclusively expressed in endosperm for a short period between approximately four and seven days after pollination (DAP). The up- and down-regulation of *OsYUC12* also appeared to coincide with that of *OsIAA29*, encoding an atypical AUX/IAA protein that may play a role as a transcriptional co-regulator in auxin signalling.

A co-expression analysis using *OsYUC12* and *OsIAA29* as bait genes, and online accessed microarray data from several experiments, revealed a small group of genes with the same expression profile restricted to the endosperm from approximately 3 to 7 DAP (Nonhebel and Griffin 2020). This included genes that are exclusively expressed in the dorsal aleurone of rice, *i.e.* *OsPR602*, *OsPR9a*, *ALI* and *OsNF-YB1* (at the early stage of expression) (Li et al. 2008; Kuwano et al. 2011; Xu et al. 2016). Of these, *OsNF-YB1* is reported to play a key role in endosperm development; plants in which the gene was down-regulated had small grains with chalky endosperm (Xu et al. 2016). The other dorsal aleurone-specific genes, *OsPR602*, *OsPR9a* and *ALI* have been reported to have promoters that direct expression to endosperm transfer cells (Li et al. 2008) or contained *cis*-elements similar to those interacting with

ZmMRP-1 (Kuwano et al. 2011), an atypical MYB transcription factor that regulates development of endosperm transfer cells (ETCs) in maize (*Zea mays*) (Gómez et al. 2009). The co-expressed gene group also contained a previously un-reported cluster of MYB-related transcription factor-like genes from rice that are the closest rice homologues to *ZmMRP-1*. These data suggested a possible auxin-signalling network associated with development of cells responsible for uptake of nutrients into the developing endosperm.

Investigations of the role of IAA in rice grain development have been limited by a lack of mutants. More extensive work has been carried out on its importance for the development of maize endosperm, where mutants, *defective endosperm18 (de18)* and *defective kernel18 (dek18)* have reduced expression of IAA biosynthesis genes, reduced IAA content as well as an aberrant basal endosperm transfer layer (BETL) and small, shrivelled grains (Bernardi et al. 2012, 2016). In maize grains, IAA accumulates specifically in the BETL and aleurone (Forestan et al. 2010). Furthermore, treatment of developing maize grains with the auxin transport inhibitor, N-1-naphthylphthalamic acid (NPA), resulted in the formation of a multi-layered aleurone instead of a single layer, suggesting that an IAA maximum at the endosperm periphery acts as a signal for aleurone differentiation. The most recent work by Bernardi et al. 2019 investigated the transcriptome of *de18* maize grains. Their results suggest that *ZmMRP-1* as well as genes that are regulated by this transcription factor are down-regulated in the auxin deficient mutant. Thus, IAA may regulate BETL development in maize by controlling expression of *ZmMRP-1*.

We have previously reported evidence for conservation of auxin signalling networks operating during grain development of different cereals (Russell French et al. 2014). Rice does not have a well-defined ETC layer (Hands et al. 2012). However, the dorsal aleurone is likely to play a similar role in apoplastic nutrient transfer to the developing endosperm given its proximity to the vascular trace and enrichment with sugar transporters (Bai et al. 2016; Xu et al. 2016). Based on our previous observations in rice as well as the information from maize, we suggest that IAA may regulate the development of the dorsal aleurone in rice. We therefore hypothesise that expression of *OsYUC12* and *OsIAA29* may be localised to these cells. We tested this hypothesis by investigating, via *in situ* hybridisation, the localisation of *OsYUC12* and *OsIAA29* and comparing this with the previously studied and dorsal aleurone-specific *OsPR602*. To identify precisely the timing of maximum expression of *OsYUC12* and *OsIAA29* as well as test their temporal co-expression with *OsPR602*, *OsPR9a*, *ALI* and three rice homologues of *ZmMRP-1*, here designated as rice *MRP-1-like* or *OsMRPL*. (*OsMRPL1*,

OsMRPL3 and *OsMRPL4*), we carried out a quantitative expression study from grain samples harvested at daily intervals from 1 to 10 DAP.

The existence of a conserved signalling network within cereals, by which auxin regulates the development of ETCs or cells with ETC-like properties, requires the presence in other cereals of orthologues of key proteins with the same expression profile. As Hands et al. (2012) have previously reported the absence of any orthologues of ZmMRP-1 in other cereals, we investigated the phylogeny and protein structures of ZmMRP-1 and its closest homologues in maize, rice, wheat (*Triticum aestivum*) and *Brachypodium distachyon*. Expression of the wheat *MRP-like* genes was investigated for comparison, using the Wheat Expression Browser (Borrill et al. 2016; Ramírez-González et al. 2018) which has a large number of samples including from dissected grains. A similar phylogenetic and *in silico* expression analysis was carried out on putative cereal orthologues of OsIAA29. Finally, we investigated whether OsIAA29-like proteins are restricted to cereals or whether they occur in dicots and non-grass monocots.

Materials and Methods

Plant material and growing conditions

Rice plants (*Oryza sativa* ssp. *japonica* cv. Reiziq) were grown in a greenhouse at the University of New England under natural light with 30° C/18°C day/night temperatures. Seven rice seeds were sown directly into flooded cylindrical plastic pots (50 cm × 15 cm) filled with cracking clay soil (vertisol). When the seedlings reached the 2-3 leaf growth stage, they were thinned to three plants per pot. Plants were watered daily and fertilised fortnightly with the commercial fertiliser Aquasol® (2.0 gm/L) until panicle initiation. Panicles in which approximately half of the spikelets reached anthesis were tagged in the afternoon. The date of tagging was recorded as the day of pollination; the following day was designated as 1 day after pollination (DAP) and so on. Tagged panicles were harvested daily in the afternoon from 1 to 10 DAP. Only superior caryopses were collected, weighed, frozen immediately in liquid nitrogen and stored at -80°C until further use.

RNA extraction, reverse transcription and quantitative real-time PCR

Total RNA was extracted from 80-100 mg grain samples using Bioline® ISOLATE II RNA Plant Kit (Meridian Bioscience). RNA concentration and purity were measured by NanoDrop™ 8000 Spectrophotometer (Thermo Fisher Scientific). RNA quality was checked by the presence of two clear bands of 18S and 28S rRNAs following agarose gel

electrophoresis (Nolan et al. 2006). Only high-quality RNA with A260/A280 ratio in the range of 1.8-2.0 was used in downstream applications.

Transcript sequences of *OsYUC12*, *OsIAA29*, *AL1*, *OsPR602*, *OsPR9a*, *OsMRPL1*, *OsMRPL3* and *OsMRPL4* were downloaded from PHYTOZOME 12.0 (Goodstein et al. 2012). Primer pairs were designed using Primer3 software (Koressaar et al. 2018) (Supplementary Table S1). Either the left or the right primer of each gene was designed to span an exon-exon boundary in order to avoid amplification of any residual genomic DNA contaminant. Amplification of a single product of the expected size by a primer pair was first confirmed by RT-PCR using Qiagen® One-Step RT-PCR kit (Qiagen) followed by agarose gel analysis of the amplified products. The gene for rice ubiquitin-conjugating enzyme E2 (*OsUBC*; *LOC_Os02g42314*) was used as the reference gene (Li et al. 2010).

Quantitative real-time RT-PCR was done in two steps. Bioline® SensiFAST™ cDNA Synthesis Kit (Meridian Bioscience) was used to synthesize cDNA from 1.0 µg of total RNA template per reaction according to manufacturer's instructions. A no-RT control that contained all reaction components except the reverse transcriptase was also included. Bioline® SensiFAST™ SYBR® No-ROX Kit (Meridian Bioscience) was used for the quantitative PCR. Each well of a 96-well plate contained 10 ng of cDNA per 20 µl of final reaction volume. All other reagents were added as per the manufacturer's instructions. A no-template control was included as negative control. Three biological replicates and two technical replicates were included for each primer set. Reactions were carried out in CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad Laboratories). The amplification program used was as follows: 95°C for 2 min and 40 cycles of 95°C for 5 s, 60°C for 10 s and 72°C for 5 s. Melt curve analysis confirmed the amplification of a single uniform product by each primer pair. This was further confirmed by an agarose gel electrophoresis. Primers with poor amplification efficiency or melt curve analysis implying the presence of more than one product were replaced with new primer sets and the experiment repeated. The manufacturer's software was used to calculate the expression of the targeted genes relative to the expression of the reference gene. Data from the two technical replicates were first averaged, then the mean and standard error of the three biological replicates from the same developmental stage were calculated.

***In situ* mRNA hybridisation**

In situ mRNA hybridisation was carried out using the protocol of Drews 1998 with some modifications. Immature rice grains collected at 5, 6 and 7 DAP were trimmed at both ends, and their palea and lemma were carefully removed. Trimmed grains were immediately fixed in freshly prepared formalin-acetic acid-alcohol (FAA; 3.7% formaldehyde, 5% acetic acid and 50% ethanol) fixative first under gentle vacuum on ice for 15 min and then overnight at 4°C. The fixed grains were dehydrated by a graded ethanol series and xylene before infiltrating them with paraffin wax (Paraplast Plus) in an automated tissue processor (TP1020, Leica Biosystems). The grains were embedded in paraffin wax on an embedding centre (EG1150, Leica Biosystems). Paraffin sections (8.0 µm thick) were cut using a rotary microtome (RM2235, Leica Biosystems) and transferred to glass slides coated with poly-L-lysine (Sigma-Aldrich). The slides were air-dried overnight and stored at 4°C until further use.

Purified cDNAs from *OsYUC12*, *OsIAA29* and *PR602* templates were cloned into pGEM®-T vector (Promega) following manufacturer's instructions (refer to Supplementary Table S2 for primer pairs used to amplify the templates). Gene inserts were amplified from the plasmid by T7 and SP6 primers which annealed to T7 and SP6 promoters flanking the inserts. Purified DNA amplicons with flanking T7 and SP6 promoters were used for *in vitro* synthesis of digoxigenin (DIG)-UTP-labelled single-stranded RNA sense and anti-sense probes by using T7 and SP6 polymerases from a DIG RNA Labelling Kit (Roche). The non-complementary sense probe was used as negative control for each gene. The probes were hydrolysed with 200 mM carbonate buffer at 60°C for 80-90 min to generate 150-200 bp fragments.

Selected paraffin sections were de-waxed with HistoClear (Sigma-Aldrich), and rehydrated with a graded ethanol series and PBS. They were treated with 1.0 µg/mL proteinase K solution for 30 min at 37°C. The sections were dehydrated with a graded ethanol series and probes were applied (320 ng probe in 100 µl hybridisation buffer per slide). Hybridisation was carried out overnight in a humidified box at 55°C. After a series of washes, immunodetection of the DIG-labelled probes was carried out by using an anti-DIG antibody coupled with alkaline phosphatase and a ready-to-use 5-bromo-4-chloro-3-indolyl phosphate (BCIP)/nitro-blue tetrazolium (NBT) solution (Sigma-Aldrich) as the chromogenic substrate for the enzyme. The slides were mounted with an aqueous medium made of glycerol and TE buffer (50% v/v). Images of the sections were acquired by a high-definition slide scanner (NanoZoomer 2.0-RS, Hamamatsu Photonics).

Phylogenetic analysis and database mining

BLASTP search (Altschul et al. 1997) was conducted using *OsIAA29* and *ZmMRP-1* peptides as queries against proteomes of selected monocots and eudicots on PHYTOZOME 12.0 (Goodstein et al. 2012) and Ensembl Plants 47 (Kersey et al. 2016). Members of the CCA1-like subgroup of MYB-related transcription factors were selected based on information in Du et al. (2013). Molecular Evolutionary Genetics Analysis (MEGA) version X (Kumar et al. 2018) was used for the phylogenetic analysis. The sequences were aligned by MUSCLE (Edgar, 2004). Phylogenetic trees were constructed by the Maximum Likelihood method based on JTT matrix-based model (Jones et al. 1992). The reliability of each node in the tree was determined by the bootstrap test from 500 replicates (Felsenstein 1985). *OsIAA29* orthologues were compared with homologues from bamboo and non-grass monocots by multiple sequence alignment using Clustal Omega (<https://www.ebi.ac.uk>). RNA-seq database of Rice Genome Annotation Project (Kawahara et al. 2013) was used to analyse expression profiles of rice genes. Expression data of *OsIAA29* and *OsMRPL* orthologues in wheat and barley (*Hordeum vulgare*) were retrieved from online RNA-seq databases, Wheat Expression Browser <http://www.wheat-expression.com> (Borrill et al. 2016; Ramírez-González et al. 2018) and BaRTv1.0 <https://bio.tools/BaRTv1.0> (Rapazote-Flores et al. 2019), respectively. Co-expression and *cis*-regulatory element (CRE) analyses were conducted as described previously by Nonhebel and Griffin (2020).

Results

Gene expression during the first 10 days after pollination

The expression of *OsIAA29* and *OsYUC12* was compared with that of *OsPR602*, *OsPR9a*, *ALI* and three *OsMRPLs* (*OsMRPL1*, *OsMRPL3* and *OsMRPL4*) in immature grains from 1 to 10 DAP by qRT-PCR. All genes showed a somewhat similar expression profile (Fig. 1). In particular, all genes except *OsMRPL3* were not expressed until 2 to 3 DAP and expression was maximal at 5 to 6 DAP, before declining to low levels by 9 DAP. *OsMRPL3* followed a similar pattern of up-regulation followed by down-regulation, but this was earlier, with expression detected at 1 DAP and maximal expression at 4 to 5 DAP. *OsMRPL4* had the most restricted expression, with very little transcript detected apart from 5- and 6-DAP samples. On the other hand, *ALI* still continued to be active at 10 DAP. As gene expression was only examined in developing grain samples, global expression of genes in other parts of the plant was investigated using RNA-seq datasets available via the Rice Genome Annotation Project. As shown in Fig. 2, expression was mostly restricted to early grain development with

no gene activity found in vegetative tissues, floral tissues, or more mature grains at 25 DAP. However, *OsMRPL3*, *OsMRPL4* and *OsMRPL5* did show slight expression in anthers.

Localisation of *OsPR602*, *OsIAA29* and *OsYUC12* expression in early grains

Spatial expression of *OsPR602*, *OsIAA29* and *OsYUC12* was examined in immature rice grains by *in situ* mRNA hybridisation. The results reported are from sections of grains collected at 7 DAP, just past the peak of expression, as the larger grains were easier to section and gene expression was still detectable. The sense probes of the genes tested did not give any hybridisation signal, validating the experimentation. Signal from the anti-sense probe of *OsPR602* confirmed its spatial expression in the dorsal aleurone (Fig. 3 b-c and f-g). The spatial expression of *OsIAA29* was similar to that of *OsPR602*, with its hybridisation signal restricted exclusively to the dorsal aleurone (Fig. 4 b-c and f-g). The expression of *OsIAA29* was confined to a narrow strip directly under the major vascular bundle (Fig. 4 f-g). We did not detect any signal for *OsIAA29* expression in the ventral aleurone, starchy endosperm, embryo (Fig. 4d), pericarp and vascular bundles. Spatial expression of *OsYUC12*, on the other hand, followed a much broader pattern than that of *OsPR602* and *OsIAA29*. As shown by both longitudinal and transverse sections, its hybridisation signal was detected in the aleurone and sub-aleurone layers; the signal was distributed both in the dorsal and ventral sides of aleurone (Fig. 5 b-c and f-g). We also detected the hybridisation signal in the embryo (Fig. 5d). However, there was no signal in the starchy endosperm, pericarp and vascular bundles.

Phylogenetic analyses of *OsIAA29* and *OsMRPLs* and expression of orthologues in other cereals

As little is known about the non-canonical AUX/IAA protein *OsIAA29* or any of its orthologues, we carried out a comprehensive search for, and phylogenetic analysis of similar proteins to explore whether this protein was restricted to cereals. A BLASTP search on PHYTOZOME 12.0 (Goodstein et al. 2012) using *OsIAA29* as query against proteomes of selected species found proteins with high homology in both cereal and non-cereal grass species, excepting sequences from moso bamboo (*Phyllostachys edulis* J.Houz.) which showed low peptide homology. Amino acid identity was also low for sequences from non-grass monocots and eudicots. The phylogenetic tree comparing all AUX/IAA proteins from rice with closest homologues from other monocots and eudicots (Fig. 6) confirmed that *OsIAA29* had putative orthologues in all grass species tested except moso bamboo. No putative orthologues were found in eudicots. The multiple sequence alignment shown in

Suppl. Fig. S1 showed that, like OsIAA29, putative orthologues lacked the N-terminal domains I and II of canonical AUX/IAA proteins and had an acidic C-terminal extension. On the other hand, closest homologous peptides from moso bamboo, banana and the sea grass *Zostera marina* showed typical AUX/IAA domain structure. We explored the expression of barley and wheat orthologues of *OsIAA29* via publicly available RNA-seq databases. The barley orthologue is expressed exclusively in immature grains at 5 DAP, with no expression recorded in embryo at 4 DAP or other tissues (Suppl. Fig. S2A). The expression of wheat orthologues is also restricted to early grain/endosperm around 10 DAP (Suppl. Fig. S2B).

Apart from our previous report of *OsMPRL* genes, no orthologues of ZmMRP-1 have been described. Fig. 7 shows a phylogenetic analysis of all members of the Circadian Clock Associated 1-like (CCA1-like) sub-group of MYB-related transcription factors from maize, rice and *B. distachyon* as well as wheat proteins homologous to ZmMRP-1. The phylogram showed a well-supported clade (bootstrap value of 85) containing ZmMRP-1 and five closely related maize proteins, as well as OsMRPL1-5, four proteins from *B. distachyon* and 21 wheat homologues. As MYB-related transcription factors are highly diverse, the reliability of these relationships was tested by using other methods of phylogenetic analysis: Neighbour Joining (Saitou and Nei 1987) and Minimum Evolution (Rzhetsky and Nei 1992). All analyses reliably grouped the OsMRPLs with ZmMRP-1. All species investigated had multiple paralogues, existing as tandem repeats indicating a high level of recent, lineage-specific gene expansion. OsMRPL1 and OsMRPL2 as well as OsMRPL4 and OsMRPL5 were pairs of tandem duplicates. Although the tree suggests that OsMRPLs as well as proteins from wheat and *B. distachyon* are potentially orthologous to ZmMRP-1, the amino acid identity between OsMRPLs and ZmMRP-1 is very low, ranging from 30 to 38%. However, the multiple sequence alignment (ClustalO) shown in Fig 8 demonstrates homology between ZmMRP-1 and putative orthologues in rice, wheat and *B. distachyon* in the N-terminal domain as well as the MYB domain. It is also noteworthy that ZmMRP-1 itself is missing part of this N-terminal domain compared with other homologues in maize as shown by the inclusion of the sequence encoded by GRMZM2G121111_T01. Although all *OsMRPL* genes appeared to be expressed, a detailed investigation of the peptide sequences (Suppl. Fig. S3) indicated that OsMRPL5 lacked a section of the N-terminal domain found in OsMRPL4 and other proteins. It also had a large insert within the MYB domain. In addition, a comparison of tandem duplicates OsMRPL1 and OsMRPL2 indicated a number of short indels between them.

To investigate possible orthologous expression of MRPL proteins within the grains of another cereal, we used the Wheat Expression Browser as we were unsuccessful in obtaining *in situ* hybridisation results for the *OsMRPLs*. The results shown in Fig. 9 demonstrated expression of 11 out of the 21 wheat *MRPL* genes. These were active exclusively in developing grains with significant expression seen in whole grain samples only at 10 DAP, coinciding with expression of wheat co-orthologues of *OsIAA29*. Although expression in whole grains was very low at 20 DAP, in manually dissected samples, high relative expression was seen exclusively in the endosperm transfer cell (ETC) layer, indicating highly localised up-regulation of these genes. Supplementary Fig. S4 compares expression of *ZmMRP-1* and other full-length members of this clade in maize with that of *ZmIAA38*, the putative orthologue of *OsIAA29*. As with the other cereals there is a strong co-expression seen with maximal expression at 12 DAP.

Promoter analysis of genes co-expressed with *OsIAA29*, *OsPR602* and *OsMRPLs*

The existence of putative orthologues of ZmMRP-1 in rice and other cereals with similar expression profiles suggested conservation of function of these proteins. ZmMRP-1 interacts with a 12-bp *cis*-acting regulatory element (CRE) consisting of two tandem repeats of TATCTC (TC-box) in the promoter region of its target genes (Barrero et al. 2006). With a view to examining the putative enrichment of this CRE in promoter regions of rice genes, we first determined the genes co-expressed with *OsIAA29*, *OsMRPL1* and *OsPR602*, and ranked them as described previously by Nonhebel and Griffin 2020. We then searched for statistically significant enrichment of the CRE in 1000-bp promoter regions of top 300 co-expressed genes, using the RiceFREND platform (Sato et al. 2013). The 12-bp CRE was not found. However, a single TATCTC element was significantly enriched on both promoter strands of the top 100 co-expressed genes (Table 1). Enrichment was strongly associated with co-expression rank.

Discussion

In this study, we aimed to investigate IAA biosynthesis and signalling during crucial early stages of endosperm development in rice by exploring in detail the temporal and spatial expression of putative IAA biosynthesis and signalling genes, viz. *OsYUC12* and *OsIAA29*, respectively. Spatial expression was compared with that of *OsPR602*, a gene that has previously been reported as expressed exclusively in the dorsal aleurone of developing rice grains. The precise timing of up-regulation was also compared with that of a group of genes previously suggested (Nonhebel and Griffin 2020) to be co-expressed with *OsIAA29* and

OsYUC12, i.e. *OsPR9a*, *ALI* and three homologues of *ZmMRP-1* that we have designated as *OsMRPLs*.

Our results confirmed that all genes were co-expressed, except for *OsMRPL3* which was up-regulated a little earlier. Expression was grain-specific and short-lived, with genes most active between 3 and 7 DAP. Their up-regulation thus coincides with initiation of coenocyte cellularisation, its completion and differentiation of the early aleurone/starchy endosperm, which take place at 3 DAP, 5 DAP and 7 DAP, respectively (Wu et al. 2016). This supports their possible involvement in an auxin-related signalling pathway during a key stage of transcriptional reprogramming associated with coenocyte cellularisation and differentiation of the early aleurone and starchy endosperm.

Several genes confirmed as temporally co-expressed with *OsIAA29* have localised expression within the dorsal aleurone in rice (Li et al. 2008; Kuwano et al. 2011). Here, we have also demonstrated dorsal aleurone-specific expression of *OsIAA29*. This part of rice aleurone lying adjacent to the major vascular bundles is enriched with sugar transporters and may play an important role in apoplastic nutrient transfer into developing grains (Bai et al. 2016; Xu et al. 2016). Unlike maize, wheat and barley, rice aleurone lacks a well-defined ETC layer (Hands et al. 2012). However, a number of observations point to an ETC-like identity and activity of the dorsal aleurone. Dorsal aleurone-specific *OsPR602* has orthologues in barley (*END1*) and wheat (*TaPR60*), which are also expressed only in ETCs (Doan et al. 1996; Li et al. 2008; Kovalchuk et al. 2009). A *GUS*-reporter gene coupled with *OsPR602* promoter was expressed exclusively in transgenic barley ETCs (Li et al. 2008). Spatio-temporal expression pattern of *OsIAA29* is almost identical to that of *OsPR602*. This suggests a role for the atypical AUX/IAA protein encoded by this gene in signalling during differentiation of dorsal aleurone cells that regulate apoplastic nutrient transfer from maternal pericarp into developing endosperm.

The temporal co-expression of rice homologues of *ZmMRP-1* provides further evidence for a signalling node regulating development of cells with an ETC-like function. Hands et al (2012) have previously reported that no orthologues of *ZmMRP-1* could be identified in temperate cereals or in rice. They further observed that expression of the closest homologue in *B. distachyon* (Bradi1g72300) could not be detected in developing grain tissue by reverse transcriptase PCR. The very high sequence diversity seen between MYB-related transcription factors makes it very difficult to predict the existence of orthologues of *ZmMRP-1*. However, our phylogenetic and multiple sequence analyses of homologues of CCA1-like sub-group of

MYB-related transcription factors does indicate the existence of a well-supported clade of similar proteins in rice, wheat and *B. distachyon*. Our expression analysis via a combination of reverse transcriptase PCR and RNA-seq data also shows that activity of rice and wheat homologues is restricted to a short period during early grain development and furthermore that expression in wheat is highest or possibly restricted to ETCs. The short-lived nature of expression may have led to lack of detection of Bradi1g72300 by Hands et al. (2012). It is also of note that all *MRPLs* appear to have undergone several episodes of gene expansion, with evidence of recent tandem duplication as well as earlier duplication events. Not all genes in each species are expressed; this could also explain the lack of detectable expression of Bradi1g72300. ZmMRP-1 transactivates BETL-specific genes by interacting with a TC-box motif consisting of two 6-bp tandem repeats (TATCTCTATCTC) in their promoter (Barrero et al. 2006). Publications on *AL1* (Kuwano et al. 2011), *OsPR602* and *OsPR9a* (Li et al. 2008) have already suggested that regulation of gene expression within the dorsal aleurone of rice may be regulated in a conserved manner in cereals. Supporting this we found significant enrichment of the TC-box like motif, TATCTC (one of the tandem repeats found in maize) within the promoter regions of genes co-expressed with *OsIAA29*, *OsPR602* and *OsMRPL1*. Future research focusing on the question of whether OsMRPLs bind to this *cis*-element and regulate gene expression should shed more light on their functions during early endosperm development.

In maize, a study of gene expression in the auxin-deficient *de18* mutant by Bernardi et al. (2019) has provided evidence for the regulation of *ZmMRP-1* and other members of the gene cluster by IAA. The temporal co-expression of *OsMRPLs* with *OsYUC12* and *OsIAA29* suggests that this regulation may be conserved across different cereals. We have previously suggested that three rice *YUCCA* genes predominantly expressed in grains but with differing patterns of temporal expression point to sub-functionalisation and highly localised IAA production at different stages of endosperm development (Basunia and Nonhebel 2019). We hypothesised that the IAA fraction contributed by *OsYUC12* may regulate the molecular events occurring during early stages of endosperm development such as coenocyte cellularisation and early aleurone differentiation. Further support for this view came from its spatial expression pattern. *In situ* hybridisation signal from *OsYUC12* transcripts was detected in the aleurone and sub-aleurone at 7 DAP (Fig. 5), suggesting localised biosynthesis of IAA in early aleurone. In separate work from our laboratory (Kabir et al. 2020), we have also recently reported highest expression of wheat auxin biosynthesis genes

TaYUC9-B1 and *TaYUC9D1* in aleurone and transfer cells although other genes *TaYUC10-A* and *TaYUC10-D* were most highly expression in the starchy endosperm. Forestan et al. 2010 reported strong IAA signal in the aleurone and BETL of maize kernels; *ZmYUC1* expression occurs predominantly in maize aleurone (Zhan et al. 2015; Doll et al. 2017). All results indicate the production of IAA within the aleurone layer of developing cereals grains. Furthermore, *OsYUC12* expression appears to be regulated by two MADS-box transcription factors, MADS78 and MADS79 (Paul et al. 2020). Their expression peaks at 2 DAP and becomes suppressed by 4 DAP. According to the model of Paul et al., they inhibit *OsYUC12* transcription, allowing thereby the proliferation of the coenocyte nuclei. *OsYUC12* transcription is triggered when *MADS78* and *MADS79* expression is suppressed, allowing biosynthesis of IAA which mediates the transition and progression from the coenocytic to the cellularised endosperm. A critical level of endogenous IAA is also required for the cellularisation to proceed in *Arabidopsis* endosperm (Batista et al. 2019).

Although the *in situ* results for *OsYUC12* and *OsIAA29* suggest the production and function of IAA in the dorsal aleurone, the role of *OsIAA29* within an auxin signalling pathway is unclear as this protein and its orthologues in other cereals lacks the N-terminal region required for interaction with the TIR1-like auxin receptor. IAA33, a non-canonical AUX/IAA from *Arabidopsis* lacking the degron and EAR domains like *OsIAA29*, negatively regulates auxin signalling in root tip to maintain root distal stem cell identity (Lv et al. 2020). IAA33 competes with the canonical IAA5 for binding to ARF10 and ARF16. *OsIAA29* may function in a similar way by competing with a canonical *OsAUX/IAA* protein for binding to one or more, as yet unidentified, *OsARFs*. *OsIAA29* has orthologues only in cereal and non-cereal grass species. Surprisingly, given the close phylogenetic relationship between the sub-families Oryzoideae and Bambusoideae, no orthologue was detected in moso bamboo. However, over the course of evolution, bamboo species have acquired enormous morphological variation, including different caryopses (Ruiz-Sanchez and Sosa 2015). The ancestral gene for *OsIAA29* was probably lost in moso bamboo. Published reports on the expression and function of *OsIAA29* orthologues are scarce. The sorghum orthologue is expressed in immature grains at 4 DAP (Nonhebel and Griffin 2020). Analysis of RNA-seq databases indicate early grain- or endosperm-specific expression of the barley, wheat and maize orthologues (Figs. S2 and S4). This gene thus appears to have evolved independently in the grass family (Poaceae) and acquired a novel and grain-specific function during early endosperm development.

In conclusion, we have confirmed the temporal co-expression of *OsYUC12* and *OsIAA29* with genes that have an association with ETC-like cells, *ALI*, *OsPR602*, *OsPR9a*, *OsMRPL1* and *OsMRPL4*. In addition, we have shown that expression of *OsIAA29* is restricted to a narrow strip in the dorsal aleurone very similar to that reported for *OsPR602* and *OsPR9a*. We suggest this gene may act with OsMRPLs to regulate cell differentiation in this region. Further work is required to confirm the localisation of OsMRPLs as well as details of the signalling pathway. Although *OsYUC12* expression appeared to occur over a broader region of the aleurone to *OsIAA29*, our data suggest the production of auxin within the aleurone as has been observed in maize. Future studies focusing on protein-protein interactions of OsIAA29 with OsAUX/IAAs and OsARFs, and targeted inactivation of these genes will provide more information on auxin signalling during early endosperm development.

References

- Abu-Zaitoon YM, Bennett K, Normanly J, Nonhebel HM (2012) A large increase in IAA during development of rice grains correlates with the expression of tryptophan aminotransferase *OsTAR1* and a grain-specific *YUCCA*. *Physiol Plant* 146:487–499.
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402.
- Bai AN, Lu XD, Li DQ, Liu JX, Liu CM (2016) NF-YB1-regulated expression of sucrose transporters in aleurone facilitates sugar loading to rice endosperm. *Cell Res* 26:384–388.
- Barrero C, Muñiz LM, Gómez E, Hueros G, Royo J (2006) Molecular dissection of the interaction between the transcriptional activator ZmMRP-1 and the promoter of *BETL-1*. *Plant Mol Biol* 62:655–668.
- Basunia MA, Nonhebel HM (2019) Hormonal regulation of cereal endosperm development with a focus on rice (*Oryza sativa*). *Funct Plant Biol* 46:493–506.
- Batista RA, Figueiredo DD, Santos-González J, Köhler C (2019) Auxin regulates endosperm cellularisation in Arabidopsis. *Genes Dev* 33:466–476.
- Bernardi J, Lanubile A, Li QB, Kumar D, Kladnik A, Cook SD, Ross JJ, Marocco A, Chourey PS (2012) Impaired auxin biosynthesis in the *defective endosperm18* mutant is due to mutational loss of expression in the *ZmYuc1* gene encoding endosperm-specific YUCCA1 protein in maize. *Plant Physiol* 160:1318–1328.
- Bernardi J, Li QB, Gao Y, Zhao Y, Battaglia R, Marocco A, Chourey PS (2016) The auxin-deficient *defective kernel18* (*dek18*) mutation alters the expression of seed-specific biosynthetic genes in maize. *J Plant Growth Regul* 35:770–777.

- Bernardi J, Battaglia R, Bagnaresi P, Lucini L, Marocco A (2019) Transcriptomic and metabolomic analysis of *ZmYUC1* mutant reveals the role of auxin during early endosperm formation in maize. *Plant Sci* 281:133-145.
- Borrill P, Ramirez-Gonzalez R, Uauy C (2016). expVIP: a customizable RNA-seq data analysis and visualization platform. *Plant Physiol* 170:2172-2186.
- Doan DN, Linnestad C, Olsen OA (1996) Isolation of molecular markers from the barley endosperm coenocyte and the surrounding nucellus cell layers. *Plant Mol Biol* 31:877-886.
- Doll NM, Depège-Fargeix N, Rogowsky PM, Widiez T (2017) Signaling in early maize kernel development. *Mol Plant* 10:375–388.
- Drews GN (1998) *In situ* hybridisation. In: Martinez-Zapater JM, Salinas J (eds) *Arabidopsis Protocols, Methods in Molecular Biology™*, Volume 82, Humana Press, New York, pp. 353-371.
- Du H, Wang YB, Xie Y, Liang Z, Jiang SJ, Zhang SS, Huang YB, Tang YX (2013) Genome-wide identification and evolutionary and expression analyses of MYB-related genes in land plants. *DNA Res* 20:437-448.
- Edgar RC (2004) Muscle: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797.
- Fahy B, Siddiqui H, David LC, Powers SJ, Borrill P, Uauy C, Smith, AM. (2018). Final grain weight is not limited by the activity of key starch-synthesising enzymes during grain filling in wheat. *J Exp Bot* 69: 5461-5475.
- Felsenstein J (1985) Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39:783–791.
- Forestan C, Meda S, Varotto S (2010) ZmPIN1-mediated auxin transport is related to cellular differentiation during maize embryogenesis and endosperm development. *Plant Physiol* 152:1373–1390.
- Gómez E, Royo J, Muñiz LM, Sellam O, Paul W, Gerentes D, Barrero C, López M, Perez P, Hueros G (2009) The maize transcription factor Myb-Related Protein-1 is a key regulator of the differentiation of transfer cells. *Plant Cell* 21:2022-2035.
- Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, Rokhsar DS (2012) Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res* 40:D1178–D1186.
- Hands P, Kourmpetli S, Sharples D, Harris RG, Drea S (2012) Analysis of grain characters in temperate grasses reveals distinctive patterns of endosperm organization associated with grain shape. *J Exp Bot* 63:6253–6266.
- Jones DT, Taylor WR, Thornton JM (1992) The rapid generation of mutation data matrices from protein sequences. *Bioinformatics* 8:275-282.
- Kawahara Y, de la Bastide M, Hamilton JP, Kanamori H, McCombie WR, Ouyang S, Schwartz DC, Tanaka T, Wu J, Zhou S, Childs KL, Davidson RM, Lin H, Quesada-Ocampo L, Vaillancourt B, Sakai H, Lee SS, Kim J, Numa H, Itoh T, Buell CR, Matsumoto T (2013) Improvement of the *Oryza sativa* Nipponbare reference genome using next generation sequence and optical map data. *Rice* 6:4.

- Kabir MR, Backhouse D, Winter G, Nonhebel HM (2020) Expression of TaTAR2.3-1B, TaYUC9-1 and TaYUC10 correlates with auxin and starch content of developing wheat grains. *bioRxiv*. <https://doi.org/10.1101/2020.10.12.336560>
- Kersey PJ, Allen JE, Armean I, Boddu S, Bolt BJ, Carvalho-Silva D, Christensen M, Davis P, Falin LJ, Grabmueller C, Humphrey J (2016) Ensembl Genomes 2016: more genomes, more complexity. *Nucleic Acids Res* 44:D574-D580.
- Koressaar T, Lepamets M, Kaplinski L, Raime K, Andreson R, Remm M (2018) Primer3_masker: integrating masking of template sequence with primer design software. *Bioinformatics* 34:1937-1938.
- Kovalchuk N, Smith J, Pallotta M, Singh R, Ismagul A, Eliby S, Bazanova N, Milligan AS, Hrmova M, Langridge P, Lopato S (2009) Characterization of the wheat endosperm transfer cell-specific protein TaPR60. *Plant Mol Biol* 71:81–98.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35:1547-1549.
- Kuwano M, Masumura T, Yoshida KT (2011) A novel endosperm transfer cell-containing region-specific gene and its promoter in rice. *Plant Mol Biol* 76:47–56.
- Li M, Singh R, Bazanova N, Milligan AS, Shirley N, Langridge P, Lopato S (2008) Spatial and temporal expression of endosperm transfer cell-specific promoters in transgenic rice and barley. *Plant Biotech J* 6:465–476.
- Li QF, Sun SSM, Yuan DY, Yu HX, Gu MH, Liu QQ (2010) Validation of candidate reference genes for the accurate normalization of real-time quantitative RT-PCR data in rice during seed development. *Plant Mol Biol Rep* 28:49–57.
- Lv B, Yu Q, Liu J, Wen X, Yan Z, Hu K, Li H, Kong X, Li C, Tian H, De Smet I, Zhang XS, Ding Z (2020) Non-canonical AUX/IAA protein IAA33 competes with canonical AUX/IAA repressor IAA5 to negatively regulate auxin signaling. *EMBO J* 39:e101515.
- Mizutani M, Naganuma T, Tsutsumi K, Saitoh Y (2010) The syncytium specific expression of the *Oryza*;KRP3 CDK inhibitor: implication of its involvement in the cell cycle control in the rice (*Oryza sativa* L.) syncytial endosperm. *J Exp Bot* 61:791–798.
- Nolan T, Hands RE, Bustin SA (2006) Quantification of mRNA using real-time RT-PCR. *Nat Protoc* 1:1559–1582.
- Nonhebel HM, Griffin K (2020) Production and roles of IAA and ABA during development of superior and inferior rice grains. *Funct Plant Biol* 47:716-726.
- Paul P, Dhatt BK, Miller M, Folsom JJ, Wang Z, Krassovskaya I, Liu K, Sandhu J, Yu H, Zhang C, Obata T, Staswick P, Walia H (2020) *MADS78* and *MADS79* are essential regulators of early seed development in rice. *Plant Physiol* 182:933-948.
- Ramírez-González R, Borrill P, Lang D, Harrington SA, Brinton J, Venturini L et al. (2018). The transcriptional landscape of hexaploid wheat across tissues and cultivars. *Science* 361:eaar6089.
- Rapazote-Flores P, Bayer M, Milne L, Mayer CD, Fuller J, Guo W, Hedley PE, Morris J, Halpin C, Kam J, McKim SM, Zwirek M, Casao MC, Barakate A, Schreiber M, Stephen G, Zhang R, Brown JWS, Waugh R, Simpson CG (2019) BaRTv1.0: an

- improved barley reference transcript dataset to determine accurate changes in the barley transcriptome using RNA-seq. *BMC Genomics* 20:968.
- Ruiz-Sanchez E, Sosa V (2015) Origin and evolution of fleshy fruit in woody bamboos. *Mol Phylogenet Evol* 91:123-134.
- Russell French S, Abu-Zaitoon Y, Uddin MM, Bennett K, Nonhebel HM (2014) Auxin and cell wall invertase related signalling during rice grain development. *Plants* 3:95–112.
- Rzhetsky A and Nei M (1992) A simple method for estimating and testing minimum evolution trees. *Mol Biol Evol* 9:945-967.
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425.
- Sato Y, Namiki N, Takehisa H, Kamatsuki K, Minami H, Ikawa H, Ohyanagi H, Sugimoto K, Itoh JI, Antonio BA, Nagamura Y (2013) RiceFRIEND: a platform for retrieving co-expressed gene networks in rice. *Nucleic Acids Res* 41:D1214-D1221.
- Stelpflug, SC, Sekhon RS, Vaillancourt B, Hirsch CN, Buell CR, de Leon N, Kaeppler SM (2016) An expanded maize gene expression atlas based on RNA sequencing and its use to explore root development. *Plant Genome* 9:1.
- Wu X, Liu J, Li D, Liu CM (2016) Rice caryopsis development II: Dynamic changes in the endosperm. *J Integr Plant Biol* 58:786–798.
- Xu JJ, Zhang XF, Xue HW (2016) Rice aleurone layer specific OsNF-YB1 regulates grain filling and endosperm development by interacting with an ERF transcription factor. *J Exp Bot* 67:6399–6411.
- Yu SM, Lo SF, Ho TH (2015) Source–sink communication: regulated by hormone, nutrient, and stress cross-signaling. *Trends Plant Sci* 20:844-857.
- Zhan J, Thakare D, Ma C, Lloyd A, Nixon NM, Arakaki AM, Burnett WJ, Logan KO, Wang D, Wang X, Drews GN, Yadegari R (2015) RNA sequencing of laser-capture microdissected compartments of the maize kernel identifies regulatory modules associated with endosperm cell differentiation. *Plant Cell* 27:513–531.

Tables

Table 1. Enrichment of TC-box like TATCTC element in 1000-bp promoter regions of genes co-expressed with *OsIAA29*, *OsPR602* and *OsMRPL1*. Statistical significance was calculated using Chi-squared tests by comparison to the frequency of this element in all genes (Sato et al. 2013). Its significant enrichment in promoter strands of top 100 co-expressed genes is shown in bold.

Gene sample	TATCTC	
	sense strand	antisense strand
Top 100 co-expressed genes (RiceFRIEND)	38% (P= 0.009)	38% (P= 0.006)
Co-expressed genes ranked 101-200 (RiceFRIEND)	35% (P= 0.03)	27% (P= 0.44)
Co-expressed genes ranked 201-300 (RiceFRIEND)	22% (P= 0.78)	28% (P= 0.32)
All genes (RiceFRIEND)	24%	23%

Figures

Fig. 1. Expression of *OsYUC12*, *OsIAA29*, *OsPR602*, *OsPR9a*, *AL1*, *OsMRPL1*, *OsMRPL3* and *OsMRPL4* in whole rice grains from 1 to 10 days after pollination (DAP). Expression of the genes was calculated relative to the expression level of the reference gene *OsUBC* (*LOC_Os02g42314*), using the software provided by the manufacturer of the CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad Laboratories). Results shown are the means $\pm$ the standard errors of the mean (SEMs) of three biological replicates.

Fig 2. Heat map showing expression of OsYUC12, OsIAA29 and OsMRPL1-5 in different vegetative and reproductive tissues of rice. The expression profiles were retrieved from RNA-seq datasets available on Rice Genome Annotation Project (Kawahara et al. 2013). FPKM = Fragments Per Kilobase of transcript per Million mapped reads.

Fig. 3 (a-g). *In situ* hybridisation of *OsPR602* transcripts in immature rice grains. Hybridisation was done using longitudinal (a-d) and transverse sections (e-g) of immature rice grains at 7 DAP. Images show the absence of *in situ* hybridisation signal from the sense, (a and e) probes and its presence from the anti-sense (b-d and f-g) probes. Image c and g are magnified images of marked areas from b and f, respectively. Image d is derived from the same section as b and shows the absence of hybridisation signal in the embryo. Sense probes (a and e) were used as negative control in the experimentation. Note the strong hybridisation signal in the dorsal aleurone (b-c and f-g). The dorsal side in longitudinal grain sections can be determined by the position of the major vascular bundles running in the pericarp along the dorsal side or by the position of the embryo located always on the ventral side. In case of transverse sections, the dorsal side is determined by the position of the major vascular bundles. Scale bars: (a and b) 1000 μ m; (c and d) 250 μ m; (e and f) 500 μ m; (g) 100 μ m. Abbreviations used for annotation: al = aleurone; em = embryo; pc = pericarp; se = starchy endosperm; vb = vascular bundle.

Fig. 4 (a-g). *In situ* hybridisation of *OsIAA29* transcripts in immature rice grains. Hybridisation was done using longitudinal (a-d) and transverse sections (e-g) of immature rice grains at 7 DAP. Images show the absence of *in situ* hybridisation signal from the sense (a and e) probes and its presence from the anti-sense (b-d and f-g) probes. Image c and g are magnified images of marked areas from b and f, respectively. Image d is derived from the same grain as b and shows the absence of hybridisation signal in the embryo. Sense probes (a and e) were used as negative control in the experimentation. Note the strong hybridisation signal in the dorsal aleurone (b-c and f-g). The dorsal side in longitudinal grain sections can be determined by the position of the major vascular bundles running in the pericarp along the dorsal side or by the position of the embryo located always on the ventral side. In case of transverse sections, the dorsal side is determined by the position of the major vascular bundles. Scale bars: (a and b) 1000 μ m; (c) 250 μ m; (d) 100 μ m; (e and f) 500 μ m; (g) 100 μ m. Abbreviations used for annotation: al = aleurone; em = embryo; pc = pericarp; se = starchy endosperm; vb = vascular bundle.

Fig. 5 (a-g). *In situ* hybridisation of *OsYUC12* transcripts in immature rice grains. Hybridisation was done using longitudinal (a-d) and transverse sections (e-g) of immature rice grains at 7 DAP. Images show the absence of *in situ* hybridisation signal from the sense (a and e) probes and its presence from the anti-sense (b-d and f-g) probes. Image c and g are magnified images of marked areas from b and f, respectively. Image d is derived from the same grain as b and shows the presence of hybridisation signal in the embryo. Sense probes (a and e) were used as negative control in the experimentation. Note the strong hybridisation signal in the aleurone, sub-aleurone and embryo (b-d and f-g). The dorsal side in longitudinal grain sections can be determined by the position of the major vascular bundles running in the pericarp along the dorsal side or by the position of the embryo located always on the ventral side. In case of transverse sections, the dorsal side is determined by the position of the major vascular bundles. Scale bars: (a and b) 1000 μ m; (c and d) 250 μ m; (e and f) 500 μ m; (g) 250 μ m. Abbreviations used for annotation: al = aleurone; em = embryo; pc = pericarp; se = starchy endosperm; vb = vascular bundle.

Fig. 6. Phylogenetic tree showing relationship of rice AUX/IAA proteins including OsIAA29 with closest homologous peptides from other grasses, non-grass monocots and eudicots. The phylogram was generated by MEGA 10.0 (Kumar et al. 2018), using the Maximum Likelihood method (Jones et al. 1992). Multiple sequence alignments were done using MUSCLE (Edgar 2004). Bootstrap confidence levels were obtained from 500 replicates (Felsenstein, 1985). Monocots included were: *Oryza sativa* (rice), *Phyllostachys edulis* (moso bamboo), *Brachypodium distachyon*, *Hordeum vulgare* (barley), *Triticum aestivum* (wheat), *Zea mays* (maize), *Sorghum bicolor* (sorghum), *Setaria italica* (millet), *Panicum virgatum* (switchgrass), *Miscanthus sinensis*, *Musa acuminata* (banana), *Ananas comosus* (pineapple), *Dioscorea alata* (purple yam), *Zostera marina* (a sea grass), *Spirodela polyrrhiza* (a duckweed) and *Asparagus officinalis* (asparagus). Eudicots included were: *Arabidopsis thaliana*, *Medicago truncatula*, *Phaseolus vulgaris* (common bean), *Solanum lycopersicum* (tomato), *Vitis vinifera* (grape) and *Manihot esculenta* (cassava). The tree is rooted with a sequence from the moss *Physcomitrella patens*. Putative orthologues of OsIAA29 are shown in bold. Scale bar = 0.50 denotes amino acid substitutions per site.

Fig. 7. Phylogenetic tree showing the relationships between CCA1-like MYB-related proteins from maize, rice, *Brachypodium distachyon*, and wheat. Due to the large number of wheat proteins, only those in the ZmMRP-1 clade are shown. The consensus phylogram was generated by MEGA 10.0 (Kumar et al. 2018), using the Maximum Likelihood method (Jones et al. 1992). Multiple sequence alignments were done using MUSCLE (Edgar 2004). Bootstrap confidence levels were obtained from 500 replicates (Felsenstein, 1985). The cut-off value for the consensus tree was set at 50. The clade containing ZmMRP-1 is shown in red.

Fig. 8 Multiple sequence alignment comparing ZmMRP-1 and one other MRP-1 like protein from maize encoded by GRMZM2G121111_T01 with putative orthologues from rice (OsMRPL1), wheat (TraesCS7D02G135500.1), *B. distachyon* (Bradi1g48002). The alignment was carried out in CLUSTAL Omega accessed via <https://www.ebi.ac.uk> and displayed using Boxshade (https://embnet.vital-it.ch/software/BOX_form.html). Shaded symbols indicate amino acids conserved in at least 50% of sequences.

Fig. 9. Heat map showing expression of putative wheat orthologues of ZmMRP-1 in different vegetative and reproductive tissues. RNA-seq data is derived from two studies, Wheat development time course (Choulet et al. 2014) and Grain development time course (Pfeifer et al. 2014), both studies using the Chinese Spring variety and accessed via the Wheat Expression Browser (Borrill et al. 2016; Ramírez-González et al. 2018). Gene expression is expressed as log2 TPM (Transcripts Per Million).

Supplementary material

Suppl. Table S1: Primer pairs used for qRT-PCR. F=forward primers; R=reverse primers.

MSU7 Locus ID	Gene name	Primer sequence (5'-3')	Product size (bp)
<i>LOC_Os02g17230</i>	<i>OsYUC12</i>	F: GAGTTATCGACGTGCTCGAA R: TGCTTTCACCATTCTTTAGCC	147
<i>LOC_Os11g11430</i>	<i>OsIAA29</i>	F: CAACTATTCCATGTCACCATTTG R: ATGGGACATCACCAAGGAAG	131
<i>LOC_Os01g28474</i>	<i>AL1</i>	F: GCATGCCAAGCCTCATAAGT R: GCTGAGGTGAGTTACTTGTACG	100
<i>LOC_Os06g07740</i>	<i>OsMRPL1</i>	F: CGATGTGCTTTGGACCGATT R: TATCGGCGTACATGGAGACC	137
<i>LOC_Os06g07640</i>	<i>OsMRPL3</i>	F: ACAGAGGAAGAACACAGGCT R: AAGAACTTCTGCGCGTGG	131
<i>LOC_Os06g07650</i>	<i>OsMRPL4</i>	F: GTGGTGCCGGTGGTGATG R: CCTGTGCTCTTCAGTGGTC	117
<i>LOC_Os01g58660</i>	<i>OsPR602</i>	F: TGCATGGAGAAGGTCGTGTA R: AGCCATCATCTTCATGCTTGA	130
<i>LOC_Os02g39625</i>	<i>OsPR9a</i>	F: TATGCAGCCGATGAAGGGA R: ACGGAGCATTGTCAGGGTAA	100
<i>LOC_Os02g42314</i>	<i>OsUBC</i>	F: CTGCGAGCTGAAACACTTTG R: TTCTCGCTGCACCTCCTTAT	105

Suppl. Table S2: Primer pairs used for probe preparation. F=forward primers; R=reverse primers.

MSU7 Locus ID	Gene name	Primer sequence (5'-3')	Product size (bp)
<i>LOC_Os02g17230</i>	<i>OsYUC12</i>	F: TGGAGGAGGTGGTTGTTCTC R: ATGCTTTCACCATTCTTTAGCC	988
<i>LOC_Os11g11430</i>	<i>OsIAA29</i>	F: GCCTATTGTGGTCAGCTTTCA R: CCCCTGGTCAGGAGTAACAA	817
<i>LOC_Os01g58660</i>	<i>OsPR602</i>	F: CACCACAACAATATAGCCACTGT R: GGCCACAATTCCCCACAAAT	527

Suppl. Fig. S1. Multiple sequence alignment of OsIAA29, with putative orthologues from other cereals (Pavir.Ha00742 *Panicum virgatum*, 8G092100 *Setaria italica*, ZmGRMZM2G148188_T01 maize, Sobic.005G098300 sorghum, Misin10G079700 *Miscanthus sinensis*, Bradi4g22050 *Brachypodium distachyon*, HORVU4Hr1G016110 barley, TraesCS4B02G108700 etc wheat) and the closest homologous proteins from moso bamboo (PH01000627G0150), banana (GSMUA_Achr6T29100_001) and *Zostera marina* Zosma287g00210.1, as well as canonical AUX/IAA protein OsIAA2. The alignment was carried out in CLUSTAL Omega accessed via <https://www.ebi.ac.uk> and displayed using Boxshade (https://embnet.vital-it.ch/software/BOX_form.html). Shaded symbols indicate amino acids conserved in at least 50% of sequences.

A.

B.

Suppl. Fig. S2. Expression of barley and wheat orthologues of OsIAA29. (A) RNA-seq data for expression of the barley orthologue (*HORVU4Hr1G016110*) retrieved from BaRTv1.0 (Rapazote-Flores et al. 2019). FPKM = Fragments Per Kilobase of transcript per Million mapped reads. (B) Heat map showing expression of the three wheat orthologues of OsIAA29 in different vegetative and reproductive tissues. RNA-seq data is derived from two studies, Wheat development time course (Choulet et al. 2014) and Grain development time course (Pfeifer et al. 2014), both studies using the Chinese Spring variety and accessed via the Wheat Expression Browser (Borrill et al. 2016; Ramírez-González et al. 2018). Gene expression is expressed as log₂ TPM (Transcripts Per Million).

CLUSTAL O(1.2.4) multiple sequence alignment

```

LOC_Os06g07640      -----MGSIVIEGWTASEIEEARSLLTSPNNG---GEGGDG   33
LOC_Os06g07650      -----MEWTAAEMDEARSIIARLTNAYDSGTLVAG   30
LOC_Os06g07660      -----                                0
LOC_Os06g07700      -----MEWTAELAEARSVIARVSDAYNSGVGS-S   29
LOC_Os06g07740      MLLSHCFVAFALSALLAGLALAMD DATFGMEWTAELGEARSVIARVSNAYDSGAGS-S   59

LOC_Os06g07640      EGNKQKHCIGHIVMELHEWF PWKTIGQVIGLYMKLNAGKPMVMHSLNKSDANN SI---GEV   90
LOC_Os06g07650      AGNGDTRHDRVRELQAWLPWRTMDQLIGLYIELMAEEPAAA---QPQY---FDAGAVV   83
LOC_Os06g07660      -----MAEEPAAA---QPQY---FDAGAVV   19
LOC_Os06g07700      SSACDTKHDRIMRELQARFSPRTMVQVIDLYVNLTVETAA-----QPQDAGSAGDAAAVV   84
LOC_Os06g07740      NSAGDTKHDRIMRELQARFSPRTMVQVIDLYLNLTAETAQAQAGAAQPQDAGGAGDA-AVV   118
                                .      :      .      .      *

LOC_Os06g07640      DHVSALANGN-----PVRL---EEHR---PMLNNVGLVFDYPL-----EEMEMEN   129
LOC_Os06g07650      DPTFDFFNDHNNFLGMPPPPVQQAADDHNMNNVVADAGMNYYYGGGGAGGAMVFGGAPMGE   143
LOC_Os06g07660      DPTFDFFIDHNNFLGMPPPPVQQAADDHAMNNVVADASMNYFYGGGA---MVFGGAPMGE   76
LOC_Os06g07700      HPTFA-----GGMPVVN---NN-----D-GMVHGGGAAMEV   110
LOC_Os06g07740      HPTFGLANDN---FGMPVAN---NND-----DGVDA-GMVFGGAPMEE   154
                                .      .      *      :

LOC_Os06g07640      QTDQ-EPKMVVEEVVQPEGLVIKEKEAGVSKIHTNSQHVTPSIKRRVIWTEEEHR----   184
LOC_Os06g07650      TVEQAAPPVP----V---VPVVMNRDDDEVNNQGGGRHRAAPTNTTRRFWTTEEHR----   192
LOC_Os06g07660      TVEQAAPPVPAPVAV---APVVMNRDDDEVNNQGGGRHRAAPTNTTRRFWTTEEHRATLI   133
LOC_Os06g07700      GA-----VAVNGGDGEVVN-----PDNADDVLWTDYEHR-----   140
LOC_Os06g07740      GA-----VAVNGGDGEVVN-----PDNADDVLWTDYEHR-----   184
                                .      : : :      :      *      :      .      : **      ***

LOC_Os06g07640      -----LFMVGLRVFGRGDWKNISKHLVTTRTAA   212
LOC_Os06g07650      -----QFLRGLRVYGRGEWKSISMNFVRSKTPV   220
LOC_Os06g07660      YSSYDFDDPLITNLLITMTMRKKNMYFLCFTMQFLRGLRVYGRGEWKSISMNFVRSKTPV   193
LOC_Os06g07700      -----LFLTGMRVYGRGDWRNI SRYFVRSKTPE   168
LOC_Os06g07740      -----LFLTGMRVYGRGDWRNIARYFVGSKTPE   212
                                * :      * : ** : *** : * : : *

LOC_Os06g07640      QVSSHAQKFFLKMEARGEAVPPPAKR-----RRRRITGDQQA---AAEHAAA-----   257
LOC_Os06g07650      QVSSHAQKYFRRVESAAADKQRY SINDVGL-----NDDTAAMDGTNSYSNNNF   268
LOC_Os06g07660      QVSSHAQKYFRRVESAAADKQRY SINDVGL-----NDDTAAMDGTNSYSNNNF   241
LOC_Os06g07700      QISMADNYFHMMEIAAAAMEADGGDDDDGHHEI-----NNNNNNL   208
LOC_Os06g07740      QVSMYADNYFHMMEIAAAAMEADGGDDDDHHENNNNNLGGGQLHAVVGAVEHHENYNNNNL   272
                                * : *      : * : : *      : *

LOC_Os06g07640      -----LRRRMPV-----PPPPFNPFLPSLVAPVMH   283
LOC_Os06g07650      GGWQSLAFAGGHLEFPVSGGGAARQVIAPASS-----AAAMNSAAQFWAPMLFNPQIQ   321
LOC_Os06g07660      GGWQSLAFAGGHLEFPVSGGGAARQVIAPASS-----AAAMNSAAQFWAPMLFNPQIQ   294
LOC_Os06g07700      GGGQLHAVVGA----VGHGPGAGH IAPATPSNNNTAAAVNNNNVDTPFVWPLLYNPEIE   263
LOC_Os06g07740      GGGQLNAGLGA----VGHGPGAGH IAPATSSNNNVAAAAANNVDAFPWVPLLYNLEIE   327
                                *      .      *      *      *      : .

LOC_Os06g07640      RLLPPGSQAAGAAASGSGGQASLPQMPWINGANGMGR*-----   321
LOC_Os06g07650      QQFMQM-----QAQTQQA WNDQHMMMAAAP---MEGATDTNFEPAGAVNY   363
LOC_Os06g07660      QQFMQM-----QAQTQQA WNDQHMMMAAAP---MEGATDTNFEPAGAVNY   336
LOC_Os06g07700      QRMMEM-----QAQSQKAWDDQ QMKMAEAATPKEEGAADK*-----   298
LOC_Os06g07740      QRMMEM-----QAQSQKAWDDQ QMKMAEAATDPKEGAADK*-----   362
                                :      :      :      *      *      :      * .

LOC_Os06g07640      -----   321
LOC_Os06g07650      YYYQQQEEEEEGGAYGVPADQWMMNQNNNMC*   394
LOC_Os06g07660      YYYQQQEEEEEGGAYGVPADQWMMNQNNNMF*   367
LOC_Os06g07700      -----   298
LOC_Os06g07740      -----   362

```

Suppl. Fig. S3 Multiple sequence alignment comparing OsMRPL proteins The alignment was carried out in CLUSTAL Omega accessed via <https://www.ebi.ac.uk>.

Suppl. Fig. S4. Comparing expression of ZmMRP-1 and related genes with that of ZmIAA38, the maize orthologue of OsIAA29 using RNA-seq data from Stelpflug, et al. (2016). As no expression of any genes was seen in any of the root, leaf, stem or internode samples, a single sample of each only was included in this figure.