Snowball Earths, population bottlenecks, and the evolution of marine photosynthetic bacteria

Hao Zhang^{1^}, Ying Sun^{1^}, Qinglu Zeng², Sean A. Crowe³, Haiwei Luo^{1*}

¹Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

²Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong SAR

³Department of Earth Sciences, School of Biological Sciences, and Swire Institute for Marine Science (SWIMS), University of Hong Kong, Pokfulam Road, Hong Kong SAR

[^]These authors contributed equally to this study.

***Corresponding author:**

Haiwei Luo
The Chinese University of Hong Kong
Shatin, Hong Kong SAR
Phone: (+852) 39436121
E-mail: hluo2006@gmail.com

Running Title: *Prochlorococcus* and Snowball Earths

Keywords: *Prochlorococcus*, Neoproterozoic Snowball Earth, genome reduction, molecular dating.

Abstract

Prochlorococcus are the most abundant photosynthetic organisms in the modern ocean. A massive DNA loss event occurred in their early evolutionary history, leading to highly reduced genomes in nearly all lineages, as well as enhanced efficiency in both nutrient uptake and light absorption. The environmental landscape that shaped this ancient genome reduction, however, remained unknown. Through careful molecular clock analyses, we established that this *Prochlorococcus* genome reduction occurred during the Neoproterozoic Snowball Earth climate catastrophe. The lethally low temperature and exceedingly dim light during the Snowball Earth event would have inhibited *Prochlorococcus* growth and proliferation and caused severe population bottlenecks. These bottlenecks are recorded as an excess of deleterious mutations that accumulated across genomic regions in the descendant lineages. *Prochlorococcus* adaptation to extreme environmental conditions during Snowball Earth intervals can be inferred by tracing the evolutionary paths of genes that encode key metabolic potential. This metabolic potential includes modified lipopolysaccharide structure, strengthened peptidoglycan biosynthesis, the replacement of a sophisticated circadian clock with an hourglass-like mechanism that resets daily for dim light adaption, and the adoption of ammonia diffusion as an efficient membrane transporter-independent mode of nitrogen acquisition. In this way, the Neoproterozoic Snowball Earth event altered the physiological characters of *Prochlorococcus*, shaping their ecologically vital role as the most abundant primary producers in the modern oceans.

Introduction

Prochlorococcus are the smallest and most abundant photosynthetic organisms on Earth¹. They are prevalent throughout the photic zone of the oligotrophic oceans between 40°N and 40°S¹, where they account for more than 40% of the biomass and contribute almost half of the net primary production². *Prochlorococcus* have diversified into two major phylogenetic groups with distinct ecology (ecotypes), with the high-light (HL) adapted monophyletic group imbedded in the low-light (LL) adapted paraphyletic group³. The distinct ecotypes of *Prochlorococcus* evolved different pigments, light-harvesting systems, and the phycobiliproteins, which allowed for efficient light absorption in the water column⁴, and thus increased growth rates and primary production⁵.

Prochlorococcus genomes have been shaped by stepwise streamlining, including a major genome reduction in their early evolution and a few minor modifications that followed^{6,7}. Modern marine *Prochlorococcus* lineages, in particular those with small genomes, show very low ratios of nonsynonymous (d_N) to synonymous (d_S) nucleotide substitution rates, suggesting that natural selection is a highly efficient throttle on the accumulation of deleterious mutations (i.e., nonsynonymous mutations) in *Prochlorococcus*^{6,8}. On long time scales, however, nucleotide substitutions at synonymous sites become saturated, invalidating the use of d_N/d_S to infer selection efficiency in deep time⁹. Thus, an alternative approach focuses instead on different types of nonsynonymous substitutions leading to radical versus conservative changes in amino acid sequences, with the former more likely to be deleterious^{10,11}. Excess radical mutations accumulate from random fixation of deleterious mutations by genetic drift (i.e., reduced efficiency of selection). Using this approach reveals that the major genome

reduction in *Prochlorococcus* took place under reduced selection efficiency⁹ and implies that the ancient population went through severe bottlenecks as the likely result of environmental catastrophe.

The environmental context underlying *Prochlorococcus* genome reduction remains unknown, however, and precise molecular dating is needed to link this important evolutionary event to its possible environmental drivers. Implementing comprehensive molecular clock analyses, we now link the early major genome reduction event of *Prochlorococcus* to the Neoproterozoic Snowball Earth events. These catastrophic disruptions to the Earth system would likely have challenged warm-water-loving photosynthetic *Prochlorococcus*, with strong potential to cause the population bottlenecks inferred from the genome sequences described above. *Prochlorococcus* survived this catastrophe through likely gains and losses of key metabolic functions reconstructed from the same genome sequences, which have far-reaching impact on their success in today's oceans. *Prochlorococcus* are thus vital “guardians of metabolism”¹², shepherding genes critical to the functioning of the biosphere across environmental catastrophes, including global glaciations.

Results and Discussion

Prochlorococcus experienced a massive gene loss event on the ancestral branch leading to the last common ancestor (LCA) of clades HL, LLI and LLII/III^{6,7,13}. This is confirmed by our analysis, which reconstructed 366 and 107 gene family losses and gains on this branch, respectively (Fig. 1A & S1), among which 163 and 84 encode unknown proteins. On the same ancestral branch, it was shown that d_R/d_C is significantly elevated compared to the sister ancestral branch leading to the LCA of the LLIV⁹, which was

validated here (both sign test and paired t -test, $p < 0.001$; Fig. 1B). These results confirm that the major genome reduction event occurring on this branch was likely driven by genetic drift as a result of one or recurrent population bottlenecks⁹. Given the global distribution and abundance of *Prochlorococcus*, and cyanobacteria more generally, such a bottleneck would likely require a global-scale event, like an environmental or climate catastrophe (e.g. meteorite impact, large igneous province emplacement, or glaciation).

To establish the environmental context for the large, ancient genome reduction, we estimated the timeline of *Prochlorococcus* evolution by implementing molecular clock analyses based on essential calibrations available in the cyanobacterial lineage. We recognize that the use of calibration sets adapted from previous studies (under calibrations C1-C6 in Table S1 with related references included there) results in up to ~320 Ma disparity (Fig. S2A) in the estimated time for the LCA of *Prochlorococcus* HL, LLI and LLII/III clades that emerged with the major genome reduction. We note that the calibrations in previous studies were not properly used. For example, the akinete fossil identified to 2,100 Mya was used as either the maximum bound or the minimum bound to calibrate the crown group of Nostocales^{14,15}. However, given the fact that apomorphic character must evolve earlier than the divergence of crown group, morphological fossils can only serve as the minimum bounds on total groups of assigned lineages¹⁶ (see Section 2.3 in Supplemental Methods for details). Thus, in the present study, we modified the calibration sets by constraining the lower bounds of the Nostocales (and the Pleurocapsales) total groups with morphological fossils and by leaving their upper bounds open (C9-C14; Table S1). Intriguingly, the variation is reduced to less than 10 Ma when these modified calibration sets are used (Fig. S2A).

Recent identification of non-oxygenic Cyanobacteria lineages such as Melainabacteria and Sericytochromatia as sister groups of oxygenic Cyanobacteria^{17,18} provides an alternative way to constrain the evolution of oxygenic Cyanobacteria. Specifically, given that oxygenic photosynthesis evolved at the stem lineage of oxygenic Cyanobacteria, we constrained the minimum age of total Cyanobacteria group at 3.0 Ga, which is supported by geochemical evidence as the time when atmospheric oxygen became available^{38,56}. To avoid the overly precise and potentially misleading age estimates, we calibrated the upper limit of the Cyanobacteria root using the ages when the planet Earth formed and became habitable (C15-C38 in Table S1; see Section 2.3 in Supplemental Methods for details). Using this strategy, we show that the age of *Prochlorococcus* major genome reduction remains stable when non-oxygenic Cyanobacteria outgroups were included (Fig. S2B). Since including the non-oxygenic Cyanobacteria have consistently reduced the precision of posterior age estimates, manifested as the higher slopes of the regression line between HPD width and the posterior age estimates compared to those without including these lineages (C15-C38 versus C1-C14 in Fig. S3; also see Section 2.6 in Supplemental Methods for extended discussion), we focus on the crown oxygenic Cyanobacteria group dating (C7-C14) in the following discussions.

By comparing the width of the 95% highest posterior density (HPD) derived from each molecular clock analysis (Fig. S3), we inferred the most precise timeline of *Prochlorococcus* evolution (corresponding to calibration set C14 in Table S1; see Section 2.3 in Supplemental Methods). Our time estimates revealed that the LCA of *Prochlorococcus* HL, LLI and LLII/III clades diversified at 682 Mya [95% HPD

(=highest posterior density) 632-732 Mya], precisely dating the genome reduction event to this time. A 682 Mya date for the emergence of the LCA of *Prochlorococcus* HL, LLI and LLII/III clades places the large genome reduction that took place in this lineage firmly within the Cryogenian Period (~720 to 635 Mya; Fig. 1A) and implicates the Snowball Earth icehouse climate conditions eponymous with the Period in the corresponding *Prochlorococcus* population bottleneck. We therefore refer to this ancestor as SBE-LCA (see Fig. 1A), short for “Snowball Earth” LCA. The Neoproterozoic climate catastrophe culminated in the Sturtian (~717 to 659 Mya) and Marinoan (~645 to 635 Mya) glaciations (Fig. 1A), which stretched from the poles to sea level near the equators, possibly wrapping the entire Earth under a frozen skin¹⁹. This “Snowball Earth” persisted with the freezing temperature of seawater below the ice sheet lowered to -3.5 °C²⁰. Since all assayed *Prochlorococcus* strains, including those affiliated with the basal LL ecotypes, reach maximum growth rates at approximately 25 °C and rarely survive when temperature drops to ~10 °C (Fig. 1C)²¹, we propose that extreme climate cooling during the Neoproterozoic Snowball Earth events was likely the major driver of severe bottlenecks in early *Prochlorococcus* populations.

Survival of *Prochlorococcus* populations through the Cryogenian would have required refugia, the nature of which would have shaped continued *Prochlorococcus* evolution. A variety of biotic refugia have been identified during Snowball Earth intervals, including the sea-ice brine channels within ice grounding-line crack systems²² and cryoconite holes/ponds on the surface of the sublimation zone, which may have represented ~12% of the global sea glacier surface²³ (Fig. 1D). Despite providing the essential space for *Prochlorococcus* survival, these refugia would have presented a

number of environmental stresses to *Prochlorococcus* populations including: low temperature, low light, high and variable osmotic pressure, and limited nutrients²²⁻²⁴. *Prochlorococcus* thus evolved a number of adaptive mechanisms to cope with these stresses via gene gains and losses, which we assessed by reconstructing the evolutionary paths of imprints that the Snowball Earth climate left in extant *Prochlorococcus* genomes.

Among these stresses, the most prominent was likely lethally low temperature. Maintaining membrane fluidity is of paramount importance under low temperature conditions, which is largely achieved by the activities of fatty acid desaturase encoded by *desA* and *desC*. As a result, we inferred that these genes were retained in SBE-LCA (Fig. 1A). Lipopolysaccharide (LPS) in the outer membrane is known to provide the first line of defense against harsh environments²⁵, which contains the O-specific polysaccharide, the glycolipid anchor lipid A, and the polysaccharide core region. Based on our analyses, genes encoding the polysaccharide core region (*kdsABCD* for 3-deoxy-d-manno-octulosonate biosynthesis; Fig. 1A) were likely lost at SBE-LCA, while those encoding the other components were retained (*lpxABCD* and *rfbABC* for Lipid A precursor and O-specific LPS precursor biosynthesis; Fig. 1A). This inference is consistent with a previous conclusion that the loss of the LPS core region would increase the hydrophobicity and permeability of the cell envelope²⁶ to protect against the cold conditions²⁷. The amino sugar N-acetylglucosamine (GlcNAc) is used by bacteria such as *Corynebacterium glutamicum* as a carbon, energy and nitrogen source²⁸. GlcNAc enters bacteria in the form of GlcNAc-6-phosphate (GlcNAc-6-P). However, instead of being metabolized, the loss of *nagB* for GlcNAc-6-P deamination at SBE-LCA suggests that

185 GlcN6P is more likely to be involved in peptidoglycan (PG) recycling through the
186 cascade catalysis by GlmM and GlmU (Fig. 1A) to generate UDP-GlcNAc, which is an
187 essential precursor of cell wall PG and LPS²⁹. During cell turnovers, PG is continuously
188 broken down and reused through the PG recycling pathway to produce new PG, and in
189 some bacteria, PG recycling is critical for their long-term survival when growth is stalled
190 under nutrient limitation³⁰. Thus, such a recycling mechanism seems to be key for
191 maintenance of cell integrity in SBE-LCA under oligotrophic and lethally freezing
192 conditions²³. Another metabolic modification in SBE-LCA was related to heat shock
193 proteins (HSPs), which play crucial roles in tolerating environmental stresses including
194 thermal shocks. Typically, HSPs are tightly regulated, as they respond quickly to stress
195 and turn off rapidly once the stress disappears³¹. However, the HSP repressor protein
196 encoded by *hrcA* was inferred to be lost at SBE-LCA, which thus likely allowed the
197 organism to continuously express HSPs to cope with prolonged lethally low temperature.
198 In fact, constitutive expression of HSPs occurs in polar organisms such as the Antarctic
199 ciliate *Euplotes focardii*³² and the polar insect *Belgica antarctica* larvae³³. Extremely low
200 temperature also made substrate acquisition difficult due to increased lipid stiffness and
201 decreased efficiency and affinity of membrane transporters³⁴. Under such conditions,
202 bacteria may increasingly rely on substrates whose uptake shows lower dependence on
203 temperature. In sea-ice brines where less CO₂ is dissolved³⁵, elevated pH promotes the
204 conversion of ammonium to ammonia, which diffuses directly into cells without the aid
205 of transporters in the membrane. Accordingly, species of bacteria and microalgae show a
206 greater dependence on ammonium and ammonia at low temperatures and high pH than
207 nitrate^{36,37}, thereby reducing reliance on membrane transporters. In SBE-LCA, the

potentially efficient utilization of ammonia made other N acquisition genes dispensable, leading to the neutral loss of nitrite transporter (*nitM*), whereas glutamine synthetase (*glnA*) and glutamate synthase (*gltS*) responsible for the utilization of ammonia after its assimilation were conserved (Fig. 1A).

An additional stress to *Prochlorococcus* during Snowball Earth would have been variable osmotic pressure. Although multiple refugia might have supported bacterial survival during Snowball Earth, the schizohaline nature of these refugia must have imposed strong osmotic pressure on ice-trapped bacteria²³. As temperature dropped, salts would have become increasingly concentrated in brine channels²², whereas cryoconite ponds would have remained hyposaline, similar to modern Arctic and Antarctic cryoconite ecosystems³⁸. Glycine betaine (GB) is among the most important organic osmolytes in halophilic cyanobacteria, and is used as the major osmolyte in *Synechococcus* sp. WH8102³⁹. However, genes involved in glycine betaine biosynthesis and transport were lost at SBE-LCA, including *bsmB* for dimethylglycine N-methyltransferase, *gsmt* for glycine/sarcosine N-methyltransferase, and *proVWXP* for glycine betaine/proline transport system (Fig. 1A). Instead, several other organic osmolytes might have been used during the Snowball Earth, as their biosynthetic genes were retained at SBE-LCA. The first examples are the *ggpS* gene encoding glucosylglycerol phosphate synthase for glucosylglycerol (GG) synthesis and the *gpgS* encoding glucosyl-phosphoglycerate synthase for glucosylglycerate (GGA) synthesis (Fig. 1A). GG and GGA may be sufficient to provide osmotic tolerance under moderately saline conditions⁴⁰. Interestingly, the biosynthesis of GG and GGA requires less N compared to that of GB^{40,41}, and thus the potential use of GG/GGA instead of GB

appeared favorable to SBE-LCA, which had a reduced efficiency of membrane transporters and a low affinity for external nutrients at exceedingly low temperature.

Low light intensity during Snowball Earth was another formidable challenge to phototrophs including *Prochlorococcus*. In the Neoproterozoic, the Sun was still at least 6% dimmer than that at present⁴². Moreover, sea ice, especially when covered with snow, is an effective barrier to light transmission⁴³. This is in analogy to the deeper layers of today's polar snow and glacier ice where irradiation is reduced and photosynthetic organisms and activities are scarcely detectable⁴⁴. Consequently, photosynthetic organisms trapped in the brine channels or inhabiting waters below ice need to be physiologically geared to cope with low light. It was proposed that modification of the photosystem structure enables adaptation to the low light condition⁴⁵. We inferred a few changes in photosystem I and II (PSI/PSII) that occurred in SBE-LCA, including the gain of RC1 subunit PsaM, RC2 subunit PsbY, and an extra copy of the RC2 subunit PsbF, the loss of RC2 protein PsbU, and the replacement of RC2 subunit PsbX (Fig. 1A), but the molecular mechanism of these changes underlying low light adaptation is poorly understood. We also inferred an expansion of the *Prochlorococcus* antenna Pcb from two to six copies during the Snowball Earth (Fig. 1A), which may boost the light-harvesting capacity under low-light conditions⁴⁶.

Many cyanobacteria have a sophisticated circadian clock, which is essential in controlling global diel transcriptional activities of the cells. This circadian oscillator system requires only three components: KaiA, KaiB and KaiC⁴⁷. While all marine *Synechococcus* possess the three *kai* genes, most *Prochlorococcus* lack *kaiA* and, as a consequence, their circadian clocks rather behave like an “hourglass” which is reset every

morning⁴⁸⁻⁵⁰. Our analysis indicated that *kaiA* was lost at SBE-LCA (Fig. 1A). This is likely due to the prolonged darkness or low light conditions during the Snowball Earth, rendering the sophisticated circadian clock dispensable.

We argue that the genome reduction and metabolic adaptation events discussed above not only enabled *Prochlorococcus* to survive the Snowball Earth climate catastrophe, but also shaped the physiological characters and the biogeographic distribution of their descendants in the modern ocean. For example, the genome reduction that occurred in the early evolution of *Prochlorococcus* likely resulted in the reduced cell size and increased surface-to-volume ratio in their descendants, which may have enhanced their efficiency in nutrient acquisition⁵¹ and eventually led them to dominate the photosynthetic communities in the most oligotrophic regions of today's oceans². Likewise, new metabolic strategies that *Prochlorococcus* evolved to overcome the nutrient stresses during Snowball Earth, such as the recycling of cell wall components and the use of GG and GGA instead of nitrogen-rich GB as the organic osmolytes, decreased the nutrient requirements of the descendants' cells and thus contributed to their success in the modern oligotrophic nitrogen-limited oceans. On the other hand, modifications of some important metabolic pathways may also have imposed deleterious effects on *Prochlorococcus* descendants. For example, whereas the replacement of circadian clock with an hourglass-like mechanism might have facilitated the ancestral lineage to adapt to the prolonged dim light condition during the Snowball Earth catastrophe, it likely prevents the dispersal of *Prochlorococcus* to high latitude regions in the modern ocean, where the day length varies substantially across seasons. Normally, organisms with circadian rhythms deal with these changes by anticipating the changes of

light intensity and promptly regulating cellular processes such as DNA transcription and recombination via chromosome compaction, a known mechanism to protect DNA from UV radiation^{52,53}. In the absence of the circadian clock, however, species such as *Prochlorococcus* cannot synchronize the endogenous oscillation with the environmental cycles and thus are under high risks of cell damages⁵⁴.

Concluding Remarks

The Neoproterozoic Snowball Earth hypothesis was proposed decades ago, which claimed the entire extinction of the photosynthetic organisms¹⁹. In contrast to the original “hard” version of the hypothesis, a modified “soft” version of the Snowball Earth hypothesis was later proposed to include the likely persistence of refugia across the Cryogenian Period, which allowed for the survival of bacterial and simple eukaryotic lineages^{55,56}. Survivors of the Snowball Earth included photosynthetic microorganisms^{57,58}, which enabled continuous primary production across the interval^{55,59}. Like other autotrophic organisms at the base of a food web, the survival of *Prochlorococcus* was likely important in sustaining primary production, heterotrophy and carbon cycling, as well as broader ecosystem functioning during the Snowball Earth glaciations⁵⁵.

On the other extreme, a few studies have proposed that microbial communities might have been only mildly affected by the Snowball Earth climate catastrophe^{58,59}. These inferences were based on the microfossil and biomarker records, which, due to the lack of lineage-specificity, did not capture the nuances required to reconstruct effects on many ecologically important lineages like the *Prochlorococcus* studied here. Instead, we find that substantial disruptions to the Earth system, like the Neoproterozoic Snowball Earth,

leave indelible signatures in microbial genomes, such that these heritable changes allow us to reconstruct interactions between environmental change and biological evolution deep in Earth's history. By employing the accelerated genome-wide accumulation of the deleterious type mutations as a proxy for a rapid decrease in the population size of ancient lineages, we uncovered severe bottlenecks that shaped the early evolution of *Prochlorococcus* lineages. The precise molecular clock analyses as well as the ancestral genome reconstruction, furthermore, enabled us to link dynamics in ancestral population sizes to changes in metabolic potential and adaptation to icehouse climates through natural selection. Collectively, our findings demonstrate how paleomicrontological approaches can be used to connect large-scale dynamics in the Earth System to the genomic imprints left on extant microorganisms, which shape their ecological role and biogeographic distribution in the world today. They also illustrate how *Prochlorococci* acted as important “guardians of metabolism”¹², safeguarding photosynthetic metabolic potential across the Snowball Earth climate catastrophe.

Materials and Methods

Genomic sequences of Cyanobacteria were downloaded from public databases and manually annotated (Table S2; see Section 1 in Supplemental Methods). Divergence time of *Prochlorococcus* was estimated with MCMCTree v4.9e⁶⁰ on top of 27 genes (Table S3) previously proposed to be valuable to date bacterial divergence⁶¹ and *Cyanobacteria* phylogenomic trees. In previous studies, the LPP (*Leptolyngbya*, *Plectonema* and *Phormidium*) group of *Cyanobacteria* located either at the basal of the Microcyanobacteria group^{14,62} or at the basal of the Macrocyanobacteria group^{63,64}. Our

analysis showed that this controversy is likely caused by the inclusion of composition-heterogeneous proteins, and that using composition-homogeneous proteins led to consistent support for the former hypothesis (Fig. S4; Table S4; see Section 2.2 in Supplemental Methods). Since molecular dating analysis is known to be intrinsically associated with calibration points⁶⁵, we summarized the calibrations of Cyanobacteria used in previous studies, and modified them for our own analyses with caution. Moreover, we proposed a new strategy to use calibrations when non-oxygenic Cyanobacteria were used as outgroups (Table S1; see Section 2.3 in Supplemental Methods for justification). We further assessed the fitness of different molecular clock models implemented in MCMCTree by using the package “mcmc3r” v0.3.2, based on which we decided to use the independent rates model for further molecular clock analyses. For each molecular clock analysis, the software ran twice with a burn-in of 50,000 and a total of 500,000 generations. The convergence was assessed based on the correlations of posterior mean time of all ancestral nodes between independent runs (Fig. S5). By implementing statistical tests based on the “infinite-site” theory (Fig. S3; see Section 2.6 in Supplemental Methods) we were able to select the most precise estimates of *Prochlorococcus* evolutionary timeline for illustration (Fig. S6) and further discussion.

Evolution of genome content via gene gains and losses was inferred using two independent methods, AnGST⁶⁶ and BadiRate v1.35⁶⁷. The former assumes that the statistically supported topological differences between a gene tree and the species tree result from evolutionary events (gene loss, gene duplication, HGT, gene birth and speciation), and infers these evolutionary events by reconciling the topological incongruences under a generalized parsimony framework by achieving a minimum

number of the evolutionary events along the species tree, with penalties of an evolutionary event determined by the genome flux analysis (Fig. S7)⁶⁶. The latter does not rely on the tree topological incongruence information, but instead uses a full maximum-likelihood approach to determine the gene family turnover rates that maximizes the probability of observing the gene count patterns provided by the family size table. The BadiRate analyses were run using nine strategies each with a distinct turnover rate model and a distinct branch model. The likelihoods of different runs were compared (Fig. S8), and three strategies with highest likelihood values were used (Fig. S9). Further, results derived from AnGST (Fig. S10) and BadiRate were compared and summarized to determine the common patterns shared by the two software (Fig. S11), and important functional genes discussed were consistently inferred by these two methods. As the two methods inferred the qualitatively same pattern of genome size reduction on the branches leading to SBE-LCA, the number of gene gains and losses derived from the AnGST analysis was presented.

The inference of a potential change of selection efficiency on a given branch was performed by comparing the genome-wide d_R/d_C value across single-copy orthologous genes of the branch to that of the closest sister branch. The d_R/d_C value was calculated using RCCalculator (<http://www.geneorder.org/RCCalculator/>; see Section 4 in Supplemental Methods) based on two independent amino acid classification schemes (Table S5).

References

- 1 Biller, S. J., Berube, P. M., Lindell, D. & Chisholm, S. W. Prochlorococcus: the structure and function of collective diversity. *Nat Rev Microbiol* **13**, 13 (2015).
- 2 Johnson, Z. I. *et al.* Niche partitioning among Prochlorococcus ecotypes along ocean-scale environmental gradients. *Science* (80-) **311**, 1737-1740 (2006).
- 3 West, N. J. & Scanlan, D. J. Niche-partitioning of Prochlorococcus populations in a stratified water column in the eastern North Atlantic Ocean. *Appl Environ Microbiol* **65**, 2585-2591 (1999).
- 4 Hess, W. R. *et al.* The photosynthetic apparatus of Prochlorococcus: insights through comparative genomics. *Photosynth Res* **70**, 53-71 (2001).
- 5 Moore, L. R., Rocap, G. & Chisholm, S. W. Physiology and molecular phylogeny of coexisting Prochlorococcus ecotypes. *Nature* **393**, 464 (1998).
- 6 Batut, B., Knibbe, C., Marais, G. & Daubin, V. Reductive genome evolution at both ends of the bacterial population size spectrum. *Nat Rev Microbiol* **12**, 841 (2014).
- 7 Luo, H., Friedman, R., Tang, J. & Hughes, A. L. Genome reduction by deletion of paralogs in the marine cyanobacterium Prochlorococcus. *Mol Biol Evol* **28**, 2751-2760 (2011).
- 8 Hu, J. & Blanchard, J. L. Environmental sequence data from the Sargasso Sea reveal that the characteristics of genome reduction in Prochlorococcus are not a harbinger for an escalation in genetic drift. *Mol Biol Evol* **26**, 5-13 (2008).
- 9 Luo, H., Huang, Y., Stepanauskas, R. & Tang, J. Excess of non-conservative amino acid changes in marine bacterioplankton lineages with reduced genomes. *Nat Microbiol* **2**, 17091 (2017).
- 10 Zuckerkandl, E. P., Linus. in *Evolving genes and proteins* 97-166 (Elsevier, 1965).
- 11 Dayhoff, M. O. A model of evolutionary change in proteins. *Atlas of protein sequence and structure* **5**, 89-99 (1972).
- 12 Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive Earth's biogeochemical cycles. *Science* (80-) **320**, 1034-1039 (2008).
- 13 Kettler, G. C. *et al.* Patterns and implications of gene gain and loss in the evolution of Prochlorococcus. *PLoS Genet* **3**, e231 (2007).
- 14 Sánchez-Baracaldo, P. Origin of marine planktonic cyanobacteria. *Sci Rep* **5**, 17418 (2015).
- 15 Sánchez-Baracaldo, P., Ridgwell, A. & Raven, J. A. A neoproterozoic transition in the marine nitrogen cycle. *Current Biology* **24**, 652-657 (2014).
- 16 Marshall, C. R. Using the Fossil Record to Evaluate Timetree Timescales. *Front Genet* **10**, 1049 (2019).
- 17 Di Rienzi, S. C. *et al.* The human gut and groundwater harbor non-photosynthetic bacteria belonging to a new candidate phylum sibling to Cyanobacteria. *Elife* **2**, e01102 (2013).
- 18 Soo, R. M., Hemp, J., Parks, D. H., Fischer, W. W. & Hugenholtz, P. On the origins of oxygenic photosynthesis and aerobic respiration in Cyanobacteria. *Science* (80-) **355**, 1436-1440 (2017).
- 19 Hoffman, P. F., Kaufman, A. J., Halverson, G. P. & Schrag, D. P. A Neoproterozoic snowball earth. *Science* (80-) **281**, 1342-1346 (1998).
- 20 Ashkenazy, Y. *et al.* Dynamics of a Snowball Earth ocean. *Nature* **495**, 90 (2013).
- 21 Zinser, E. R. *et al.* Influence of light and temperature on Prochlorococcus ecotype distributions in the Atlantic Ocean. *Limnol Oceanogr* **52**, 2205-2220 (2007).
- 22 Thomas, D. & Dieckmann, G. Antarctic sea ice--a habitat for extremophiles. *Science* (80-) **295**, 641-644 (2002).
- 23 Hoffman, P. F. *et al.* Snowball Earth climate dynamics and Cryogenian geology-geobiology. *Sci Adv* **3**, e1600983 (2017).

414 24 Takeuchi, N. Optical characteristics of cryoconite (surface dust) on glaciers: the
415 relationship between light absorbency and the property of organic matter contained in the
416 cryoconite. *Annals of Glaciology* **34**, 409-414 (2002).

417 25 Benforte, F. C. *et al.* Novel role of the LPS core glycosyltransferase WapH for cold
418 adaptation in the Antarctic bacterium *Pseudomonas extremaustralis*. *PLoS One* **13**,
419 e0192559 (2018).

420 26 Wang, Z., Wang, J., Ren, G., Li, Y. & Wang, X. Influence of core oligosaccharide of
421 lipopolysaccharide to outer membrane behavior of *Escherichia coli*. *Mar Drugs* **13**, 3325-
422 3339 (2015).

423 27 Feller, G. & Gerday, C. Psychrophilic enzymes: hot topics in cold adaptation. *Nat Rev*
424 *Microbiol* **1**, 200 (2003).

425 28 Uhde, A. *et al.* Glucosamine as carbon source for amino acid-producing *Corynebacterium*
426 *glutamicum*. *Appl Microbiol Biotechnol* **97**, 1679-1687 (2013).

427 29 Park, J. T. & Uehara, T. How Bacteria Consume Their Own Exoskeletons (Turnover and
428 Recycling of Cell Wall Peptidoglycan). *Microbiology and Molecular Biology Reviews* **72**,
429 211-227 (2008).

430 30 Borisova, M. *et al.* Peptidoglycan Recycling in Gram-Positive Bacteria Is Crucial for
431 Survival in Stationary Phase. *MBio* **7** (2016).

432 31 Schumann, W. Regulation of bacterial heat shock stimulons. *Cell Stress and Chaperones*
433 **21**, 959-968 (2016).

434 32 La Terza, A., Papa, G., Miceli, C. & Luporini, P. Divergence between two Antarctic species
435 of the ciliate *Euplotes*, *E. focardii* and *E. nobilii*, in the expression of heat-shock protein 70
436 genes. *Mol Ecol* **10**, 1061-1067 (2001).

437 33 Rinehart, J. P. *et al.* Continuous up-regulation of heat shock proteins in larvae, but not
438 adults, of a polar insect. *Proc Natl Acad Sci U S A* **103**, 14223-14227 (2006).

439 34 Lawrence, R. P. & William, J. W. Temperature and substrates as interactive limiting factors
440 for marine heterotrophic bacteria. *Aquatic Microbial Ecology* **23**, 187-204 (2001).

441 35 Gleitz, M., v.d. Loeff, M. R., Thomas, D. N., Dieckmann, G. S. & Millero, F. J. Comparison
442 of summer and winter inorganic carbon, oxygen and nutrient concentrations in Antarctic
443 sea ice brine. *Mar Chem* **51**, 81-91 (1995).

444 36 Reay, D. S., Nedwell, D. B., Priddle, J. & Ellis-Evans, J. C. Temperature dependence of
445 inorganic nitrogen uptake: reduced affinity for nitrate at suboptimal temperatures in both
446 algae and bacteria. *Appl. Environ. Microbiol.* **65**, 2577-2584 (1999).

447 37 Raven, J. A., Wollenweber, B. & Handley, L. L. A comparison of ammonium and nitrate as
448 nitrogen sources for photolithotrophs. *New Phytologist* **121**, 19-32 (1992).

449 38 Webster-Brown, J. G., Hawes, I., Jungblut, A. D., Wood, S. A. & Christenson, H. K. The
450 effects of entombment on water chemistry and bacterial assemblages in closed cryoconite
451 holes on Antarctic glaciers. *FEMS Microbiol Ecol* **91** (2015).

452 39 Mao, X. *et al.* Computational prediction of the osmoregulation network in *Synechococcus*
453 *sp.* WH8102. *BMC Genomics* **11**, 291 (2010).

454 40 Scanlan, D. J. *et al.* Ecological genomics of marine picocyanobacteria. *Microbiol. Mol.*
455 *Biol. Rev.* **73**, 249-299 (2009).

456 41 Empadinhas, N. & da Costa, M. S. To be or not to be a compatible solute: bioversatility of
457 mannosylglycerate and glucosylglycerate. *Syst Appl Microbiol* **31**, 159-168 (2008).

458 42 Carver, J. H. & Vardavas, I. M. Precambrian glaciations and the evolution of the
459 atmosphere. *Ann Geophys* **12**, 674-682 (1994).

460 43 Raven, J. A., Kübler, J. & Beardall, J. Put out the light, and then put out the light. *Journal*
461 *of the Marine Biological Association of the United Kingdom* **80**, 1-25 (2000).

462 44 Simon, C., Wiezer, A., Strittmatter, A. W. & Daniel, R. Phylogenetic diversity and

463 metabolic potential revealed in a glacier ice metagenome. *Appl Environ Microbiol* **75**,
464 7519-7526 (2009).

465 45 Kouril, R., Wientjes, E., Bultema, J. B., Croce, R. & Boekema, E. J. High-light vs. low-
466 light: effect of light acclimation on photosystem II composition and organization in
467 *Arabidopsis thaliana*. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1827**, 411-419
468 (2013).

469 46 Bibby, T., Mary, I., Nield, J., Partensky, F. & Barber, J. Low-light-adapted *Prochlorococcus*
470 species possess specific antennae for each photosystem. *Nature* **424**, 1051 (2003).

471 47 Dong, G. & Golden, S. S. How a cyanobacterium tells time. *Curr Opin Microbiol* **11**, 541-
472 546 (2008).

473 48 Holtzendorff, J. *et al.* Genome streamlining results in loss of robustness of the circadian
474 clock in the marine cyanobacterium *Prochlorococcus marinus* PCC 9511. *J Biol Rhythms*
475 **23**, 187-199 (2008).

476 49 Axmann, I. M. *et al.* Biochemical evidence for a timing mechanism in *prochlorococcus*. *J*
477 *Bacteriol* **191**, 5342-5347 (2009).

478 50 Mella-Flores, D. *et al.* *Prochlorococcus* and *Synechococcus* have Evolved Different
479 Adaptive Mechanisms to Cope with Light and UV Stress. *Front Microbiol* **3**, 285 (2012).

480 51 Giovannoni, S. J., Thrash, J. C. & Temperton, B. Implications of streamlining theory for
481 microbial ecology. *ISME J* **8**, 1553 (2014).

482 52 Warters, R. L. & Lyons, B. W. Variation in radiation-induced formation of DNA double-
483 strand breaks as a function of chromatin structure. *Radiat Res* **130**, 309-318 (1992).

484 53 Simons, M. J. The evolution of the cyanobacterial posttranslational clock from a primitive
485 “phoscillator”. *J Biol Rhythms* **24**, 175-182 (2009).

486 54 Mullineaux, C. W. & Stanewsky, R. The rolex and the hourglass: a simplified circadian
487 clock in *Prochlorococcus*? *J Bacteriol* **191**, 5333-5335 (2009).

488 55 Moczyłowska, M. The Ediacaran microbiota and the survival of Snowball Earth
489 conditions. *Precambrian Res* **167**, 1-15 (2008).

490 56 Hoffman, P. F. & Schrag, D. P. The snowball Earth hypothesis: testing the limits of global
491 change. *Terra nova* **14**, 129-155 (2002).

492 57 Allison, C. W. & Awramik, S. M. Organic-walled microfossils from earliest Cambrian or
493 latest Proterozoic Tindir Group rocks, northwest Canada. *Precambrian Res* **43**, 253-294
494 (1989).

495 58 Corsetti, F. A., Awramik, S. M. & Pierce, D. A complex microbiota from snowball Earth
496 times: microfossils from the Neoproterozoic Kingston Peak Formation, Death Valley, USA.
497 *Proc Natl Acad Sci USA* **100**, 4399-4404 (2003).

498 59 Olcott, A. N., Sessions, A. L., Corsetti, F. A., Kaufman, A. J. & De Oliveira, T. F. Biomarker
499 evidence for photosynthesis during Neoproterozoic glaciation. *Science (80-)* **310**, 471-474
500 (2005).

501 60 Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-
502 1591 (2007).

503 61 Battistuzzi, F. U. & Hedges, S. B. A major clade of prokaryotes with ancient adaptations to
504 life on land. *Mol Biol Evol* **26**, 335-343 (2008).

505 62 Shih, P. M. *et al.* Improving the coverage of the cyanobacterial phylum using diversity-
506 driven genome sequencing. *Proc Natl Acad Sci USA* **110**, 1053-1058 (2013).

507 63 Blank, C. & Sanchez-Baracaldo, P. Timing of morphological and ecological innovations in
508 the cyanobacteria—a key to understanding the rise in atmospheric oxygen. *Geobiology* **8**, 1-
509 23 (2010).

510 64 Uyeda, J. C., Harmon, L. J. & Blank, C. E. A comprehensive study of cyanobacterial
511 morphological and ecological evolutionary dynamics through deep geologic time. *PLoS*

512 *One* **11**, e0162539 (2016).
513 65 Schirrmeister, B. E., Sanchez-Baracaldo, P. & Wacey, D. Cyanobacterial evolution during
514 the Precambrian. *Int J Astrobiol* **15**, 187-204 (2016).
515 66 David, L. A. & Alm, E. J. Rapid evolutionary innovation during an Archaean genetic
516 expansion. *Nature* **469**, 93 (2011).
517 67 Librado, P., Vieira, F. & Rozas, J. BadiRate: estimating family turnover rates by likelihood-
518 based methods. *Bioinformatics* **28**, 279-281 (2011).
519

Acknowledgments

We thank Allison Coe, Erik Zinser, and Zackary Johnson for providing the data of *Prochlorococcus* growth rates, Sishuo Wang, Tianhua Liao and Xiaoyuan Feng for their suggestions on molecular dating analyses. This work is supported by the National Natural Science Foundation of China (92051113), the Hong Kong Research Grants Council General Research Fund (14110820), the Hong Kong Research Grants Council Area of Excellence Scheme (AoE/M-403/16), HKU FoS funds to SAC, and the Direct Grant of CUHK (4053257 and 3132809).

Author Contributions

H.L. conceived and directed the study. H.Z. and Y.S. performed the bioinformatics. All authors contribute to the interpretation of the results. H.Z., Y.S., S.A.C and H.L. wrote the paper.

Conflict of interests

The author declares no competing interests.

Fig. 1

Fig. 1 Evolution of *Prochlorococcus* during the Neoproterozoic Snowball Earth events.

(A) (Left) Chronogram of the evolutionary history of *Prochlorococcus* estimated by MCMCTree. The evolutionary tree shown here is part of the species tree constructed with MrBayes based on 90 compositionally homogenous gene families shared by 159 cyanobacterial genomes (Fig. S4 D). Divergence time is estimated based on 27 gene sequences under calibration set C9 (Table S1). The vertical bars represent the estimated time of the Neoproterozoic glaciation events. The flanking horizontal blue bars on ancestral nodes represent the posterior 95% highest probability density (HPD) interval of the estimated divergence time. The pie chart on the ancestral branches leading to the node SBE-LCA provides the proportion of reconstructed genomic events including gene gain, gene loss, gene replacement, gene duplication and gene vertical inheritance. (Right) Phyletic pattern of key gene families that potentially enabled *Prochlorococcus* to survive harsh conditions during the Neoproterozoic Snowball Earth (at the ancestral node ‘SBE-LCA’). Solid square, solid circle and open circle next to each extant taxon represent multi-copy gene family, single-copy gene family, and absence of the gene family, respectively, in the genome. (B) (Left) The diagram helps understand how the d_R/d_C was calculated. In this context, the ‘Target’ group includes all genomes of all HL clades, LLI and LLII/III, the ‘Control’ group includes all genomes of LLIV, and the ‘reference’ group includes all genomes of Syne 5.1. The d_R/d_C for the ‘Target’ group (shown in Middle & Right) is calculated by comparing a genome from the ‘Target’ group to a genome from the ‘reference’ group (marked with red), followed by averaging the value across all possible genome pairs. Likewise, the d_R/d_C for the ‘Control’ group (shown in Middle & Right) is calculated by comparing a genome from the ‘Control’ group to a

559 genome from the ‘reference’ group (marked with green) and then by averaging the value
560 across all possible genome pairs. (Middle & Right) The genome-wide means of d_R/d_C
561 values at the ancestral branch leading to SBE-LCA and that at its sister lineage. They
562 were classified based on the physicochemical classification of the amino acids by charge
563 or by volume and polarity, and were either GC-corrected by codon frequency (blue), GC-
564 corrected by amino acid (AA) frequency (red) or uncorrected (gray). Error bars of d_R/d_C
565 values represent the standard error of the mean. (C) Diagram of putative bacterial refugia
566 including cryoconite holes and sea ice brine channels in Neoproterozoic Snowball Earth,
567 which were featured with a number of stresses such as low temperature, low light, and
568 variable osmotic pressure. (D) The average growth rate of *Prochlorococcus* ecotypes at
569 different temperatures. Replicate cell cultures were grown in a 14:10 light: dark cycle at
570 $66 \pm 1 \mu\text{mol m}^{-2}\text{s}^{-1}$. The growth data used for plotting are collected from Johnson et al.
571 2006 and Zinser et al. 2007.