Cooperative herbivory between two important pests of rice

Qingsong Liu^{1,2†}, Xiaoyun Hu^{1†}, Shuangli Su^{1†}, Yufa Peng¹, Gongyin Ye³, Yonggen Lou³,
Ted C. J. Turlings⁴, and Yunhe Li^{1*}

¹State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China.

²College of Life Sciences, Xinyang Normal University, Xinyang 464000, China.

³Institute of Insect Sciences, Zhejiang University, Hangzhou 310058, China.

⁴Laboratory of Fundamental and Applied Research in Chemical Ecology, University of Neuchâtel, Neuchâtel 2000, Switzerland.

[†]These authors contributed equally to this work.

***To whom correspondence should be addressed.**

Yunhe Li

No. 2 West Yuanmingyuan Road, Beijing, 100193, China

E-mail: liyunhe@caas.cn

21 **Abstract**

22 Normally, when different species of herbivorous arthropods feed on the same
 23 plant this leads to fitness reducing competition. We found this to be uniquely
 24 different for two of Asia's most destructive rice pests, the brown planthopper
 25 and the rice striped stem borer. Both insects directly and indirectly benefit from
 26 jointly attacking the same host plant. Double infestation improved plant quality,
 27 particularly for the stemborer because the planthopper fully suppresses
 28 caterpillar-induced production of proteinase inhibitors. It also drastically
 29 reduced the risk of egg parasitism, due to diminished parasitoid attraction.
 30 Females of both pests have adapted their oviposition behaviour accordingly.
 31 Their strong preference for plants infested by the other species even overrides
 32 their avoidance of plants already attacked by conspecifics. This uncovered
 33 cooperation between herbivores is telling of the exceptional adaptations
 34 resulting from the evolution of plant-insect interactions, and points out
 35 mechanistic vulnerabilities that can be targeted to control two major pests.

36

37 **Keywords:** *Nilaparvata lugens*, *Chilo suppressalis*, rice defense suppression,
 38 conventional interspecific competition theory, collaborative interaction

39

40

Introduction

Species that feed on the same resource are commonly regarded to be competitors¹⁻³. In the great majority of cases, species that share a food resource negatively affect each other's performance, and the conventional interspecific competition theory is widely recognized for a diverse range of taxonomic groups including plants, birds, reptiles, marine invertebrates, insects and microbes^{1,4,5}. Yet, there are instances of mutually beneficial interactions between species that feed on the same food source. This is mostly known for microbes that can assist other organisms in various ways to help reach, convert and digest food⁶⁻⁸. In very rare cases, higher organisms have also evolved cooperative interactions to be better exploit and share resources, as, for instance, has been shown for predators with complementary hunting tactics⁹⁻¹¹.

Amongst arthropods, mutually beneficial interactions have been reported mainly for social insects with food providing partners¹². The classic example is the symbiotic relationship between ants and aphids, in which aphids produce sugar-rich honeydew that is collected by the ants, and in exchange ants care for and protect the aphids from predators and parasitoids^{13,14}. In none of these associations the arthropods share the same food sources, and we are not aware of any example of mutually beneficial exploitation of the same resources by arthropods.

The most commonly food sources shared among arthropods are plants, with virtually all plants being attacked by a number of different species. This has so far assumed to always lead to competition^{1,3}. The extent of this competition can vary and the consequences can be highly asymmetrical, but in all reported examples the effects of feeding on the same plant are negative for at least one of the herbivores, independently of whether they are phylogenetically close, have the same mode of feeding, or feed on the same tissues^{1,15-18}. In certain cases, one herbivore species can benefit from the presence of another herbivore species; for instance, by causing physiological changes in the plants

that make these plants less toxic and/or more nutritious^{16,17,19-22} or by masking the (volatile) foraging cues used by natural enemies²³⁻²⁷. To our knowledge, however, there are no known examples of two species of herbivores both consistently benefitting from presence of each other on the same host plants.

It has been postulated that mutually beneficial interactions among species of insect herbivores must exist, but no specific examples have been uncovered yet^{20,28}. It is one thing to demonstrate that two herbivores benefit from jointly feeding on the same plant, it is another to conclude that they actively pursue the plant mediated-benefits. Here we propose such a coevolved partnership between the brown planthopper (BPH), *Nilaparvata lugens* and the rice striped stem borer (SSB), *Chilo suppressalis*, two of the most devastating pests of rice²⁹. Our hypothesis that both insects benefit from attacking the same plant was prompted by our recent finding that BPH can escape parasitism of its eggs by preferentially ovipositing on rice plants that are already infested by SSB²⁵. The apparent reason for this oviposition strategy is that, *Anagrus nilaparvatae*, the most common egg parasitoid of BPH, uses volatiles emitted by BPH-infested plants to locate plants with eggs. Plants that are co-infested by SSB release a different blend of volatiles that is not attractive to the parasitoid²⁵. Previous work also indicates that SSB larvae perform poorly on rice plants that are already infested by conspecifics due to induced plant resistance, and that females show a strong oviposition preference for uninfested rice plants relative to SSB-infested rice plants, in accordance with the ‘mother knows best’ principle³⁰. BPH has been shown to suppress certain defenses in rice^{31,32}. This raises the question whether sharing host plants with BPH can help SSB to counter the direct defenses of rice plants and possibly mitigate the defense responses to SSB infestation, and, if so, whether it also has adapted its oviposition behavior accordingly.

To answer these questions, we tested the performance of SSB larvae on uninfested rice plants, and plants infested either by BPH only, by SSB only, or by both species. We further tested if the oviposition preferences of SSB moths

101 matched the measured performances on the differently pre-infested plants.
 102 The results prompted us to investigate the potential molecular and biochemical
 103 mechanisms that may be involved, with a focus on protease inhibitors. To also
 104 address how BPH may affect the indirect defenses of rice plants against SSB,
 105 we further tested the odor preferences of *Trichogramma japonicum*, an
 106 important egg parasitoid of SSB, for differently treated plants. With these
 107 olfactometer assays and additional cage experiments we tested whether the
 108 presence of BPH can also reduce the risk of SSB moth eggs to be parasitized.
 109 The combined results support our hypothesis of a co-evolve cooperation
 110 between two intimately interacting herbivorous species that feed on the same
 111 plants, which exceptionally goes against conventional competition theory of
 112 interspecific interactions among phytophagous insects¹. The elucidation of the
 113 underlying plant-mediated mechanisms that facilitate this cooperation can be
 114 the basis for plant breeding strategies to control the two exceedingly important
 115 rice pests.

116

117 **Results**

118 **Performance of SSB caterpillars on herbivore-infested rice plants**

119 When *C. suppressalis* larvae were allowed to feed for 7 days on rice plants
 120 that were either uninfested (control), infested by SSB larvae alone (SSB), BPH
 121 nymphs alone (BPH), or both SSB larvae and BPH nymphs (SSB/BPH), the
 122 body weight of SSB caterpillars was significantly different among the
 123 treatments ($F_{3,165} = 8.462$, $P < 0.001$) (Fig. 1a). The body weight of SSB larvae
 124 was significantly lower when fed on plants that had been infested by SSB
 125 larvae than on all other plant treatments (all $P < 0.01$). Importantly, there was
 126 no difference in larval weight among the other three treatments (uninfested
 127 plants, feeding on BPH-infested plants or SSB/BPH-infested plants; $P > 0.05$).
 128 These results imply that additional infestation by BPH fully eliminated the
 129 negative effects of SSB-infestation on successively feeding conspecifics. And
 130 this effect seems to be independent of infestation sequence, as weight gain

131 was only marginally different between SSB larval that were placed on dual
132 infested plants, infested, with SSB treatment occurring either before or after
133 BPH infestation ($P = 0.051$) (Fig. 1b).

134

135

136 **Figure 1. Weight of SSB larvae after seven days of feeding on**
137 **differentially infested rice plants.**

138 **a** Neonates of SSB were individually placed on rice plants that were either
139 uninfested (Control), infested by SSB larvae only (SSB), BPH only (BPH), or
140 both SSB and BPH (SSB/BPH, two species were simultaneously introduced to
141 the plants); **b** Neonates of SSB were individually placed on rice plants that
142 were either uninfested (Control), infested by SSB only, or both SSB and BPH
143 in sequencing order (SSB→BPH, plants were infested with SSB larvae for 24 h
144 then BPH were added for another 24 h; BPH→SSB, plants were infested with
145 BPH for 24 h then SSB larvae were added for another 24 h). Bars indicate
146 mean $\pm$ SE. Data was analyzed using Two-way analysis of variance followed
147 by LSD post hoc test. Small letters indicate significant differences between
148 treatments ($P < 0.05$) ($N = 30-46$).

149

150

Oviposition preferences of SSB females

Oviposition preference in greenhouse

When given a choice between uninfested and SSB-infested plants, SSB females laid significantly fewer eggs on SSB-infested plants than on uninfested plants (RT-test applied to a GLMM, Poisson distribution error; $P < 0.001$) (Fig. 2a). However, compared to uninfested plants, the females strongly preferred to lay eggs on BPH-infested plants (RT-test applied to a GLMM, Poisson distribution error; $P = 0.007$) (Fig. 2b) or on SSB/BPH-infested plants (RT-test applied to a GLMM, Poisson distribution error; $P = 0.03$) (Fig. 2c). As expected, SSB females also laid significantly more eggs on BPH-infested or SSB/BPH-infested plants (RT-test applied to a GLMM, Poisson distribution error; both $P < 0.001$) relative to SSB-infested plants (Fig. 2e, f), while they laid similar numbers of eggs on BPH-infested and SSB/BPH-infested plants (Fig. 2g).

When SSB females were offered the 4 types of rice plants simultaneously, SSB females laid significantly different numbers of eggs among treatments (Fig. 2h; RT-test applied to a GLMM, Poisson distribution error; $P < 0.001$), with the most eggs on plants infested by both SSB and BPH, and slightly less on BPH-infested plants and uninfested plants. Importantly, SSB females laid only very few eggs on SSB-infested plants, significantly less than on the other three treatment (all $P < 0.001$) (Fig. 2h).

Figure 2. Oviposition preference of SSB female moths in greenhouse experiments.

Number of eggs laid by female *C. suppressalis* on rice plants that were either uninfested (Control), infested by SSB larvae only (SSB), BPH only (BPH), or by both SSB and BPH (SSB/BPH). **a** Schematic drawing of the oviposition experiments. **b** Control versus SSB larva-infested plants. **c** Control versus BPH-infested plants. **d** Control versus dually infested plants. **e** SSB larva-infested plants versus BPH-infested plants. **f** SSB larva-infested plants versus dually infested plants. **g** BPH-infested plants versus dually infested plants. **h** SSB female moths were exposed to 4 types of rice plants together. LR tests applied to a GLMM were conducted for the number of eggs (Poisson distribution error). Each bar represents the mean $\pm$ SE. Data columns with asterisks (***) $P < 0.001$, ** $P < 0.01$, * $P < 0.05$, or with small letters ($P < 0.05$) indicate significant differences between treatments; n.s. indicates a non-significant difference ($P > 0.05$) (N = 9–11).

Oviposition preferences under field conditions

Consistent with the results from the greenhouse experiments, far more eggs were laid on uninfested plants than on SSB-infested plants (RT-test applied to a GLMM, Poisson distribution error; $P < 0.001$) (Fig. 3a). Compared to uninfested rice plants, the females preferred to lay eggs on BPH-infested plants ($P < 0.001$) or SSB/BPH-infested plants ($P = 0.03$) (Fig. 3b, c). When given a choice between SSB-infested plants and SSB/BPH-infested plants, the females preferred to lay eggs on the co-infested plants, as expected (RT-test applied to a GLMM, Poisson distribution error; $P < 0.001$) (Fig. 3d).

Figure 3. Oviposition preference of SSB female moths in field cage experiments.

Number of eggs laid by female *C. suppressalis* on rice plants that were either uninfested (Control), infested by SSB larvae only (SSB), BPH only (BPH), or both SSB and BPH (SSB/BPH). **a** Control versus SSB larva-infested plants. **b** Control versus BPH-infested plants. **c** Control versus dually infested plants. **d** SSB larva-infested plants versus dually infested plants. LR tests applied to a GLMM were conducted for the number of eggs (Poisson distribution error). Each bar represents the mean $\pm$ SE. Data columns with asterisks (***) $P < 0.001$, * $P < 0.05$) indicate significant differences between treatments (N = 3–4).

Rice plant defense responses to herbivore infestation

Gene expression changes

RNA-seq analysis was carried out to assess gene expression changes in response to infestation by SSB, BPH or both species. A partial least squares discriminant analysis on all 12 transcriptomic datasets provided a global view of the total gene expression across the four treatments. As shown in Fig. 4a, the first two principal components (PCs) explained 39.5% (PC1), and 13.3% (PC2) of the total variance, respectively. PC1 revealed a clear separation of samples with SSB infestation (SSB and SSB/BPH) from others (BPH and control). And PC2 separated samples with BPH infestation (BPH and SSB/BPH) from others (SSB and control). Taken together, these results suggest that each infestation regime had distinctly different effects on the gene expression of rice plants.

233 The gene expression analyses showed that feeding by SSB resulted in the
234 differential expression of 12512 genes (absolute $\log_2(\text{fold change}) > 0$ and $P <$
235 0.05), of which 6533 genes were up-regulated and 5979 genes
236 down-regulated. Infestation by BPH alone induced a relative weak response of
237 rice plant, that is, 2523 differentially expressed genes including 1292
238 upregulated genes and 1231 downregulated genes. Co-infestation by the two
239 species induced the upregulation of 3640 genes and the downregulation of
240 4082 genes (Fig. 4b). Interestingly, compared to SSB-infested plants, 992
241 genes were downregulated when plants were co-infested with BPH and SSB,
242 and the gene ontology (GO) enrichment analysis showed that these repressed
243 genes were largely associated with defense responses, such as JA-related
244 process, and enzyme inhibitor activity (Supplementary Data 1).

245

246 ***JA and SA associated gene expression and their accumulation***

247 A total of 10392 *Arabidopsis* orthologs of rice genes were included in
248 phytohormone-related gene expression signatures (Supplementary Data 2).
249 The results showed that, when compared to uninfested plants, genes
250 associated with JA and SA pathways were generally upregulated in plants
251 infested by SSB larvae, BPH nymphs, or both species (Fig. 4c). However, dual
252 infestation by SSB and BPH resulted in an apparent downregulation of both JA
253 and SA pathways in rice, as when compared to SSB-infested plants.
254 Specifically, four genes involved in JA biosynthesis, *OsLOX9*, *OsJAR1;2*,
255 *OsDAD1;3*, and *OsAOC*, were significantly induced upon SSB infestation but
256 were not induced when plants were co-infested by both BPH and SSB
257 (Supplementary Data 3). More surprisingly, two genes involved in JA pathway,
258 *OsLOXL-2* and *OsAOS3*, and nine SA-responsive genes (*OsPR2*, *OsPR4*,
259 *OsPR4B*, *OsPR4C*, *OsPR4D*, *OsPR6*, *OsPR10*, *OsPR10A*, and *OsPR10B*)
260 were activated by SSB infestation, but were suppressed by dual infestation
261 (Supplementary Data 4).

262 Consistent with the phytohormone-related gene expression results, SSB

infestation induced a significant increase in the levels of both JA (12 hr vs. 0 hr, $P < 0.001$; 24 hr vs 0 hr, $P < 0.001$) and SA (12 hr vs. 0 hr, $P < 0.001$; 24 hr vs 0 hr, $P = 0.004$; Fig. 4d, e). However, when BPH nymphs were added after 12 hr infestation by SSB alone, the levels of JA and SA in rice plants significantly decreased (JA: $P = 0.02$; SA: $P < 0.001$; Fig. 4d, e).

Protease inhibitor-associated gene expression and their accumulation

When determining GO terms that were significantly enriched ($P_{adj} < 0.05$) in the set of 992 downregulated DEGs comparing dual infestation samples and SSB infestation samples, we identified several molecular function terms that were associated with protease inhibitor activity including serine-type endopeptidase inhibitor activity (GO:0004867), enzyme inhibitor activity (GO:0004857), endopeptidase inhibitor activity (GO:0004866), peptidase inhibitor activity (GO:0030414) (Supplementary Data 1). After screening for genes involved in these categories, we found 11 genes related to protease inhibitors that were highly induced by SSB infestation, but were significantly attenuated by the additional infestation of BPH nymphs (Fig. 4f). The expression of nine genes was validated by qRT-PCR, showing similar expression patterns among the four treatments as obtained with RNA-seq (Supplementary Fig. 1), confirming the reliability of the RNA-seq data.

Prompted by the observed changes in protease inhibitor-associated gene expression, we further measured the contents of TPIs in rice plants responding to different herbivore infestations. The results showed a significant increase in TPIs content after 72 hr SSB infestation compared to uninfested plants ($P = 0.03$). This TPIs content increase was not observed for BPH-infested plants or SSB/BPH-infested plants (Fig. 4g).

Figure 4. Rice plant responses to herbivore infestation.

a Partial least squares discriminant analysis (PLS-DA) of all detected genes in rice plants that were either untreated (control), infested by SSB, BPH, or both herbivores (SSB+BPH) for 48 h (three biological replicates per treatment). The percentage of variation of the data explained by principal component 1 (PC1) and PC2 is in parentheses (39.5 and 13.3%, respectively). The score plot displays the grouping pattern according to the first two PCs and the ellipse defines the Hotelling's T² confidence interval (95%) for the observations. **b** Differentially expressed genes among differently treated rice plants. **c** Hormonometer analyses for JA and SA signatures based on transcriptomic responses of rice to herbivory. The colors indicate similarity between herbivore infestation and a particular hormone response (blue and red for negative and positive correlations, respectively, see bottom). MJ, methyl jasmonate; SA: salicylic acid. The contents of endogenous JA (**d**) and SA (**e**) in rice plants subjected to SSB infestation or SSB plus BPH in sequence. Letters indicate significant differences among SSB treatment with different time points; asterisks indicate significant differences between SSB and SSB→BPH treatments ($P < 0.01$; $N = 3$). Bars indicate mean $\pm$ SE. **f** Heatmap of the expression of 10 enzyme inhibitors genes. Log₂-transformed RPKM values are plotted. BBI 3-1, Bowman-Birk inhibitor 3-1; BBI 3-2, Bowman-Birk inhibitor 3-2;

314 BBI3-3, Bowman-Birk inhibitor 3-3; OCPI1, *Oryza sativa* chymotrypsin
 315 inhibitor-like 1; OCPI2, *Oryza sativa* chymotrypsin inhibitor-like 2; PME1 15,
 316 pectin methylesterase inhibitors 15; PME1 16, pectin methylesterase inhibitors
 317 16; PR6, pathogenesis-related proteins 6; RASI, rice alpha-amylase/subtilisin
 318 inhibitor; SCI2A, subtilisin-chymotrypsin inhibitor-2A; WIP1, wound-induced
 319 protease inhibitor. FPKM, fragments per kilobase of transcript per million
 320 fragments mapped. **g** Time course of the contents of trypsin in rice plants that
 321 were either uninfested (Control), infested by SSB, BPH, or both (N=3). Bars
 322 indicate mean $\pm$ SE. Data was analyzed using one-way ANOVA followed by
 323 LSD post hoc test. Letters indicate significant differences between treatments
 324 ($P < 0.05$). n.s., not significant ($P > 0.05$).

325

326 **Volatile profiles of rice plants and their effects on the oviposition** 327 **behavior of SSB females**

328 A total of 61 compounds were detected in the headspace of the four plant
 329 treatments (Supplementary Data 5). Plants infested by both SSB and BPH
 330 emitted the highest amounts of volatiles, followed by plants infested by SSB,
 331 plants infested by BPH, whereas control plants emitted the lowest amounts of
 332 volatiles. A projection to partial least squares-discriminant analysis (PLS-DA)
 333 using the contents of all detected volatiles showed a clear separation among
 334 the four treatments (Fig. 5a). The first two significant principle components
 335 (PCs) explained 33.8% and 7.44% of the total variance, respectively.
 336 Consistent with gene expression data, the first PC showed a clear separation
 337 between plant infested with SSB (SSB and SSB/BPH) from others (BPH and
 338 control), and the second PC separated BPH-infested samples and
 339 SSB-infested samples (BPH and SSB) from others (SSB/BPH and control).

340 Using volatiles that had been collected from plants subjected to the four
 341 types of treatments as odor sources (applied on filter paper strips), we tested
 342 the oviposition preference of SSB females. They differently distributed their
 343 eggs among the four treatments (Fig. 5b, RT-test applied to a GLMM, Poisson
 344 distribution error; $P < 0.001$), with highest number of eggs observed on or near
 345 filter paper that had been treated with volatiles collected from
 346 SSB/BPH-infested plants, which was statistically similar to the numbers of

eggs laid on BPH-infested plants and uninfested plants. However, SSB females laid significantly lower numbers of eggs on filter paper treated with volatiles collected from SSB-infested plants compared to any of the other treatment (all $P < 0.001$) (Fig. 5b).

Figure 5. Volatiles released by rice plants and effect on oviposition behavior of *C. suppressalis* female.

a Projection to latent structures-discriminant analysis (PLS-DA) of volatile emissions produced by rice plants that were either untreated (Control), infested by SSB, BPH, or both herbivores (SSB/BPH) for 48 h. The score plot display the grouping pattern according to the first two components and the ellipse defines the Hotelling's T2 confidence interval (95%) for the observations. **b** Number of eggs laid by female *C. suppressalis* on filter paper that had been treated with volatiles collected from 4 types of differently treated rice plants. LR tests applied to a GLMM were conducted for the number of eggs (Poisson distribution error). Each bar represents the mean $\pm$ SE. Letters above bars indicate significant differences between treatments (RT-test applied to a GLM, Poisson distribution error; $P < 0.05$; $N = 8$).

367

368 **Responses of *T. japonicum* wasps to herbivore-infested rice plants**

369 In a dual-choice assay, the *T. japonicum* wasps showed a strong preference for
 370 plants infested by SSB eggs ($P = 0.004$) over uninfested intact plants (Fig. 6a).
 371 When offered plants carrying SSB eggs in both sides of the olfactometer, the
 372 wasps exhibited a significant preference for plants that were additionally
 373 infested by SSB larvae relative to plants infested by SSB eggs alone ($P =$
 374 0.002). In contrast, the wasps were significantly more attracted to plants with
 375 eggs alone than plants that were additional infested by BPH nymphs ($P <$
 376 0.001) or both SSB larvae and BPH nymphs ($P = 0.001$). We further tested the
 377 preferences of wasps that were offered plants with combinations of herbivore
 378 and egg infestation. The results (Fig. 6a) showed that *T. japonicum* females
 379 significantly preferred to rice plants with SSB and eggs over plants coinfecting
 380 by BPH and eggs ($P < 0.001$) or plants with SSB, BPH, and eggs ($P = 0.01$).
 381 Finally, as expected, the wasps showed a significant preference for the odor of
 382 plants with the combination of SSB, BPH, and eggs over the odor of plants
 383 with BPH and eggs ($P = 0.009$). These results imply that additional infestation
 384 by BPH resulted in the repellence of *T. japonicum* wasps.

385

386 **Parasitism rates of SSB eggs by *T. japonicum* wasps**

387 In the greenhouse cages with the four plant treatments, the rate of parasitism
 388 of SSB eggs by *T. japonicum* wasps were highest on plants infested with SSB
 389 larvae only (Fig. 6b), which was significantly higher than on plants infested with
 390 BPH only ($P < 0.001$) or plants infested with both herbivore species ($P = 0.016$).
 391 The lowest parasitism rate of SSB eggs was observed on plants infested by
 392 BPH only. Although parasitism was lower on plants infested by both species,
 393 this was not significant different from parasitism on control plants ($P = 0.28$).
 394 The parasitism rates of SSB eggs in the cages nicely reflected the trends of
 395 responses of the parasitoids in the olfactometer (Fig. 6b).

396

benefits for SSB when infesting plants that are already under attack by BPH. The feeding assays show that SSB infestation induces rice defense responses that cause significantly reduced fitness in SSB larvae that subsequently feed on the same plants (Fig. 1). Remarkably, additional infestation by BPH completely neutralized the negative effects of SSB defense induction; SSB caterpillars placed on plants already infested by conspecifics grew considerably better on plants that were also attacked by BPH, independent of order of infestation. On BPH-infested plants the caterpillars performed just as well as on previously uninfested control plant (Fig. 1)

RNA-seq and biochemical analyses showed that infestation by BPH suppresses a broad spectrum of defense related genes (Fig. 4f, Supplementary Data 1), with one of the main consequences being a significant reduction in SSB-induced production of proteinase inhibitors (Fig. 4g). These are key defensive compounds of rice plants that are particularly effective against chewing herbivores, including SSB^{35,36}. We also found that co-infestation with BPH suppresses the expression of SA- and JA-associated genes that are normally upregulated by SSB infestation (Fig. 4c, Supplementary Data 4). This suppression led to reduced levels of JA and SA in the plants (Fig. 4d, e). The JA signal transduction pathway is responsible for the production of TPIs in rice plant, and it is known that SSB performs better on JA-deficient mutant rice lines mainly due to reduced TPIs levels^{36,37}. Collectively, this insight explains why BPH feeding neutralizes the negative effects of SSB defense induction.

Insect attack typically induces defense responses in plants³⁸⁻⁴¹, but it is also increasingly evident that various insect herbivores have the ability to, at least partially, suppress plant defenses by interfering with JA or SA biosynthesis and that this can enhance their own performance and that of conspecifics^{28,42-45}. This can also benefit certain other species that feed on the same plants. For example, the silverleaf whitefly (*Bemisia tabaci*) activates SA signaling and represses JA-regulated defenses, leading to enhanced nymphal

development of this insect²⁸, which also benefits spider mites²⁶. None of these studies appear to show reciprocal benefits for species feeding on the same plant.

Consistent with ‘mother knows best’ hypothesis⁴⁶, SSB females avoid to lay eggs on rice plants that are already infested by conspecifics, thus ensuring that their offspring evade the negative effects that SSB-induced defenses³⁰. Here we show that SSB females have adapted their oviposition behavior to preferentially oviposit on BPH-infested rice plants, independent of whether SSB larvae are already present or not, as compared to healthy plants (Figs. 2 and 3). By doing so, they benefit from the BPH-mediated suppression of rice defense responses (Fig. 1a). However, the performance of SSB larvae was not any better on rice plants that were already co-infested by SSB plus BPH compared to their performance on healthy plants. So why did SSB females prefer to lay eggs on dual-infested plants rather than on healthy plants (Fig. 2d)? The experiments with the egg parasitoid *T. japonicun* provide a plausible explanation, as they showed that the presence of BPH significantly reduced the attractiveness of rice plants to the wasp (Fig. 6a). The cage experiment confirmed that the presence of BPH decreased the risk for SSB eggs to be parasitized, implying that the oviposition strategy of SSB females is highly adaptive (Fig. 6b).

It appears that certain well-adapted herbivores can also manipulate the emission of volatiles of their host plants⁴⁷. In most such cases, herbivore infestation suppresses certain key volatile compounds and thereby possibly reduce the attractiveness of plants to certain natural enemies^{26,27,42,48,49}. Other herbivores can benefit from this as well. Simultaneous feeding by slugs, for instance, suppress caterpillar-induced volatiles in cabbage plants, thereby reducing the attractiveness of the plants to parasitoids²⁷. Similarly, when spider mite-infested Lima bean plants are also infested by whiteflies this leads to a reduced emission of the volatile (E)- β -ocimene compared to plants infested by spider mites only, resulting in a reduced attraction of predatory

484 mites²⁶. In other cases, double infestation may actually lead to higher
 485 quantities of volatiles being emitted, but the blend is altered in a way that it is
 486 no longer attractive to parasitoids²³. This is also the case for our study system.
 487 We previously showed that BPH preferentially oviposits on SSB-infested rice
 488 plants, thereby avoiding the attraction of the egg parasitoid *Anagrus*
 489 *nilaparvatae*²⁵. These various examples confirm that insect herbivores not
 490 only can evolve the ability to use volatiles to identify host plants of better
 491 nutritional quality, but also plants where their offspring can escape natural
 492 enemies⁵⁰.

493 Our combined results, including the additional tests showing that SSB
 494 infestation of rice plants significantly increases the performance of BPH
 495 nymphs (Supplementary Fig.2) strongly supported the conclusion of mutually
 496 beneficial interaction between SSB and BPH. This goes against the ingrained
 497 notion of competition between phytophagous insects that share a common
 498 host plant, and how this competition shapes insect assemblages^{1,15,17,51-53}.
 499 The resource-based competition theory has been challenged before, but the
 500 examples involve specific asymmetric beneficial plant-mediated interactions,
 501 meaning that only one of the herbivore species benefits from the presence of
 502 another^{20,21,54-58}. In these cases, the benefit is never reciprocal, nor do they
 503 seem to represent tightly coevolved interactions with specific behavioral
 504 adaptations as found for BPH and SSB in the current study and certain
 505 vertebrates⁹⁻¹¹.

506 The interaction between SSB and BPH reported here seems to represent a
 507 highly evolved collaboration to cope with and exploit the direct and indirect
 508 defense responses of rice plants. The two species share the same host plants
 509 and have a similar spatial and temporal distribution throughout Asia's rice
 510 paddy area (²⁵; <https://www.cabi.org/isc/>). Why has their interaction evolved
 511 into collaboration rather than competition? It is likely because of their
 512 differential feeding strategies. Although the two insects occur side-by-side on
 513 rice plants, SSB is a stemborer insect that feeds inside the rice plants, while

514 BPH is a phloem-sucking insect that feeds at the surface on leaf blades and
 515 leaf sheaths ^{25,59}. Hence, there is usually no direct physical interaction
 516 between the two species. During the coevolutionary arms race between
 517 herbivores and host plants, both sides may evolve multiple defense
 518 mechanisms against the other. We speculate that the cooperative relationship
 519 between SSB and BPH may be the result of two opposing coevolutionary arms
 520 races that in combination benefit both herbivores.

521 Although the mutually beneficial interaction between the stemborer and
 522 planthopper bears no resemblance to any of the known interactions between
 523 other herbivore species attacking a same plant ^{1,15}, the reported type of
 524 cooperation is unlikely to be unique. We postulate that agricultural pests are
 525 especially prone to rapid evolutionary changes that allow such cooperation to
 526 emerge. As pest populations build up in vast monocultures their main
 527 challenges revolve around coping with plant defenses and avoiding and
 528 resisting their specific natural enemies, whereas finding host plants is no
 529 longer a challenge. In such scenarios, different insect species will encounter
 530 each other frequently. Unlike the cultivated plants, the insects are subject to
 531 natural selection and can evolve traits to jointly overcome plant defenses,
 532 without the cultivated plants being able to coevolve to resist these traits. The
 533 plants are at the mercy of human selection, which is focused at traits that favor
 534 yield and nutritional value, often at the cost of reduced resistance against
 535 pests and diseases ^{60,61}. Yet, as we discover and unravel the intricate
 536 adaptations in the insects we can start steering this human selection in favor of
 537 potent pest resistance traits. For the specific example uncovered in our study,
 538 interfering with the ability of BPH to suppress the biosynthesis of proteinase
 539 inhibitors could be a highly effective and sustainable strategy to control two of
 540 rice's most common and most harmful pests.

541 In summary, the current study reveals a highly adaptive, mutually beneficial
 542 relationship between rice planthoppers and stem borers that is mediated by
 543 opposing rice plant defense responses. The findings represent a unique

example of a cooperative interaction that challenges traditional interspecific competition theory. The two insect species take advantage of the rice defense responses induced by each other in a manner that suggests that together they are the tentative winners in the arms race with rice plants. The results are also illustrative of the complexity and intricate dynamics of the interaction between plants and insects, and challenge the conventional paradigms of interspecific competition. Future work should further unravel more details about the molecular mechanisms underlying the insect-controlled interactions, which might lead the development of rice varieties that disrupt the cooperative interaction as potential strategy to control the two pests.

Methods

Plants and insects

Rice plants (*Oryza sativa*, cultivar Minghui63) were grown in a greenhouse at 27 ± 3 °C with $75 \pm 10\%$ RH (relative humidity) and a photoperiod of 16:8 hr L:D (light:dark). The cultivation of rice plants followed the same procedure as described previously²⁵. Plants were used for experiments when they were at the tillering stage, which occurred about 44–49 days after sowing.

C. suppressalis larvae were reared on an artificial diet as described⁶². Ten percent honey water solution was provided to supply nutrition for the adults. *N. lugens* were maintained on a BPH-susceptible rice variety Taichung Native 1 (TN1)³⁴. *T. japonicun* were obtained from Keyun Industry Co., Ltd (Jiyuan, China). Newly emerged adult wasps were maintained in glass tubes (3.5 cm diameter, 20 cm height) and supplied with 10% honey water solution as a food source and were maintained for at least 6 hr to ensure freely mating, before females were used for the following experiments. All three species were maintained in climatic chambers at 27 ± 1 °C, $75 \pm 5\%$ RH, and a photoperiod of 16:8 hr L:D.

Performance of caterpillars on insect-infested rice plants

Multiple types of rice plants were prepared: i) uninfested plants, meaning that potted rice plants remained intact without insect infestation; (ii) SSB-infested plants, each potted rice plant was artificially infested with one 3rd instar SSB larva that had been starved for > 3 hr for 48 hr; iii) BPH-infested plants, each potted rice plant was artificially infested with a mix of fifteen 3rd and 4th instars BPH nymphs for 48 hr; iv) SSB/BPH-infested plants, each potted rice plant was simultaneously infested with one SSB larva and 15 BPH nymphs for 48 hr; (v) SSB→BPH-infested plants, each potted rice plant was artificially infested with one SSB larvae alone for the first 24 hr, then 15 BPH nymphs were additionally introduced for another 24 hr; vi) BPH→SSB-infested plants, namely each potted rice plant was artificially infested with 15 BPH nymphs for the first 24 hr, then one SSB larvae were additionally introduced for another 24 hr. Plant treatments were conducted as described in detail in our previous study²⁵. During herbivory treatment, the uninfested plants were placed in a separate room to avoid possible volatile-mediated interference. During the subsequent bioassays, both SSB caterpillar and BPH nymphs remained in or on the rice plants.

Two bioassays were conducted to test the performance of *C. suppressalis* larvae feeding on differently treated rice plants. The first bioassay included the plants treatments i, ii, iii, and vi, and the second bioassay included the plants treatments i, ii, v and vi. Three 2-day-old larvae of *C. suppressalis* were gently introduced onto the middle stem of each rice plant using a soft brush. The infested rice plants were then placed in climatic chambers at $27 \pm 1^\circ\text{C}$, $75 \pm 5\%$ relative humidity, and a photoperiod of 16:8 hr L:D. The *C. suppressalis* larvae were retrieved from the rice plants after 7 days feeding, and they were weighed on a precision balance (CPA2250, Sartorius AG, Germany; readability = 0.01 mg). The mean weight of the three caterpillars on each plant was considered as one biological replicate. The experiment was repeated four times using different batches of plants and herbivores, resulting in a total of 30–46 biological replicates for each treatment.

604

605 **Oviposition-preferences of *C. suppressalis* females choosing among** 606 **differently infested rice plants**

607 ***Greenhouse experiment***

608 In the greenhouse, seven choice tests were conducted with *C. suppressalis*
609 females including i) SSB-infested plants versus uninfested plants; ii)
610 BPH-infested plants versus uninfested plants; iii) SSB/BPH-infested plants
611 versus uninfested plants; iv) SSB-infested plants versus BPH-infested plants; v)
612 SSB-infested plants versus SSB/BPH-infested plants; vi) BPH-infested plants
613 versus SSB/BPH-infested plants; and vii) the test in which *C. suppressalis*
614 females were exposed to all four types of rice plants. The experiments were
615 performed as described in detail by ³⁰. In brief, four potted plants were
616 positioned in the 4 corners of a cage (80 cm x 80 cm x 100 cm) made of
617 80-mesh nylon nets for each test. For paired comparisons, two potted plants
618 belonging to the same treatment were placed in opposite corners of each cage,
619 and in the test with 4 types of rice plants, each type of plant was positioned in
620 one of the four corners of each cage. Five pairs of freshly emerged moths (less
621 than 1 day) were released in each cage, and a clean Petri dish (9 cm diameter)
622 containing a cotton ball soaked with a 10% honey solution was placed in the
623 center of the cage as food source. After 72 h, the number of individual eggs on
624 each plant were determined. The experiment was conducted in a greenhouse
625 at 27 ± 3°C, 65 ± 10% RH, and a photoperiod of 16:8 hr L:D. Each choice test
626 was repeated with 9–11 times (replicates).

627 ***Field cage experiment***

628 The oviposition preference of SSB females was further assessed in a field
629 near Langfang City (39.58° N, 116.48° E), China. Four choice tests were
630 conducted: i) SSB-infested plants versus uninfested plants; ii) BPH-infested
631 plants versus uninfested plants; iii) SSB/BPH-infested plants versus uninfested
632 plants; and iv) SSB/BPH-infested plants versus SSB-infested plants. The
633 treated rice plants were prepared as described above and were transplanted

634 into experimental plots (1.5 m × 1.5 m). For each pairwise comparison, six
635 plots of rice plants were covered with a screened cage (8 m × 5 m × 2.5 m)
636 made of 80-mesh nylon net to prevent moths from entering or escaping. Each
637 of the six plots contained 9 rice plants of a particular treatment, with 3 plots per
638 cage representing the same treatment. Plots were separated by a 1-m buffer
639 and they were alternately distributed in a 3 × 2 grid arrangement in each cage
640 (Supplementary Fig. 1). Approximately 50 mating pairs of newly emerged *C.*
641 *suppressalis* adults (< 24 hr) were released into each cage. After 72 hr, the
642 number of individual eggs on each plant were determined. The total number of
643 eggs of three plots in each cage was regarded as one replicate, 3–4 replicates
644 were conducted for each pairwise comparisons.

645

646 **Rice plant response to herbivore infestation**

647 ***RNA-seq and data analysis***

648 To explore the molecular mechanisms underlying the rice plant-mediated
649 interaction between BPH and SSB, gene expression changes in rice response
650 to infestation by SSB, BPH or both were analyzed by RNA-seq. The rice plants,
651 uninfested (control) or infested, were prepared as described above. After 48h,
652 the stems of the plants were harvested and frozen in liquid nitrogen. Samples
653 from 5 individual plants of the same treatment were pooled together as one
654 biological replicate, and three replicates were collected for each treatment.

655 RNA-seq analyses were performed as described previously⁶³. In brief, total
656 RNA was extracted using the TRIzol reagent (Invitrogen, Carlsbad, CA, USA)
657 and treated with RNase-free DNase I (NEB, Ipswich, MA, USA) to remove any
658 genomic DNA. Library preparation and RNA-seq were performed by Novogene
659 (Beijing, China) using an Illumina Hiseq 4000 system, resulting in ~45–55
660 million raw reads per sample. Raw reads were subjected to quality checking
661 and trimming to remove adapters, poly-N sequences, and low-quality bases
662 (Phred quality score Q < 20). The yield clean data of each sample were
663 aligned to the rice reference genome IRGSP-1.0 (<https://rapdb.dna.affrc.go.jp>)

664 using HISAT2 (v2.09) ⁶⁴, and the number of reads mapped to each gene was
665 counted with featureCounts (v1.5.0-p3) ⁶⁵. The expression level of each gene
666 was calculated as FPKM (fragments per kilobase of transcript per million per
667 million fragments mapped) according to an established protocol ⁶⁶. Expression
668 differentiation analyses were conducted DESeq2 R package (v. 1.18.0) ⁶⁷.
669 Genes with absolute value of $\log_2(\text{fold change}) > 0$ and $P\text{-value} < 0.05$ were
670 defined as differentially expressed genes (DEGs). The enriched functions of
671 DEGs in RNA-seq data sets were annotated with the Gene Ontology (GO)
672 function using the clusterProfiler R package ⁶⁸, and GO terms with Benjamini–
673 Hochberg false discovery rate (FDR) adjusted $P\text{-value} (Padj) < 0.05$ were
674 considered significantly enriched. The transcriptional signatures of hormonal
675 responses of rice plant to herbivory relative to gene expression in *Arabidopsis*
676 induced by diverse phytohormones was analyzed using Hormonometer
677 program ⁶⁹. Since trypsin protease inhibitors (TPIs) serve as indicators of
678 induced resistance in rice plants, especially against chewing herbivores such
679 as SSB ^{35,36}, the analyses focused on expression profiles of TPIs-related
680 genes among the four plant treatments. The expression of nine selected TPIs
681 genes were validated by quantitative real-time PCR (qRT-PCR) analyses as
682 previously described ⁷⁰. qRT-PCR was conducted on a Bio-RadCFX96 Touch
683 Real-time PCR Detection System instrument (Bio-Rad, Hercules, CA, USA)
684 using TransStart[®] Top Green qPCR SuperMix (TransGen Biotech, Beijing,
685 China). The rice *ubiquitin 5* gene was used as the internal standard to
686 normalize the variations in gene expression. The primers used are listed in
687 Supplementary Table 1.

688

689 ***Quantification of endogenous jasmonic and salicylic acid***

690 Our RNA-seq results suggested that additional infestation by BPH significantly
691 suppressed the expressions of genes related to the defense hormones
692 jasmonic acid (JA) and salicylic acid (SA). Both types of genes are highly
693 upregulated in response to SSB infestation ⁷¹. To confirm this, we quantified

the JA and SA levels in rice plants with two treatments: i) rice plants that were infested with one third-instar SSB larva alone for 24 hr; ii) rice plants that were first infested with one third-instar SSB larva for 12 hr and then also with 15 BPH female adults for another 12 hr. Rice stems were harvested at three time points: 0 hr (uninfested control plants), as well as 12 hr and 24 hr after infestation. For each treatment, stems from 5 individual plants were harvested and pooled together as one biological replicate, and three replicates were collected for each time point.

Endogenous measurements of JA and SA were performed by the plant hormone platform at the Institute of Genetics and Developmental Biology, Chinese Academy of Sciences as previously described⁷².

Quantification of TPIs

We further measured the accumulation of TPIs in rice plants subjected to insect infestation. These experiments were prompted by the RNA-seq results indicating that the upregulation of TPIs-related genes in response to SSB infestation is significantly suppressed after co-infestation with BPH. The same plant treatments were included as used for RNA-seq but with new batches of plants. Samples were collected at 48 hr and 72 hr of insect infestation. Intact rice plants that served as controls were also sampled at the same time points. Samples from five individual plants were pooled together as one biological replicate, and three replicates were collected for each treatment. All samples were immediately frozen in liquid nitrogen and stored at -80°C until further analyses.

TPIs contents was determined using enzyme linked immunosorbent assay (ELISA) kits (J&L Biological, Shanghai, China). The stem samples were ground into a fine power in liquid nitrogen using a mortar and pestle, and each sample (0.1 g) was homogenized in 0.01 M PBS (Phosphate Buffered Saline) buffer (pH = 7.4) (Sigma-Aldrich, St. Louis, MO, USA) with a sample–PBS proportion of 1:9 (1 g plant sample/9 ml of PBS). Samples were centrifuged at 4000 *g* for 15 min at 4°C and the supernatant was collected. The ELISA

experiments were performed following the protocols provided with the kits. The optical density values were recorded at 450 nm using a microplate spectrophotometer (PowerWave XS2, BioTek, Winooski, VT, USA). The protein concentrations in plant samples were measured using a bicinchoninic acid (BCA) protein assay kit (Aidlab Biotechnologies Co., Ltd., Beijing, China) according to the manufacturer's instructions. The amount of protease inhibitor was calculated based on a standard curve, and results were expressed as μg protease inhibitor per mg protein.

Effect of insect-induced volatiles on the oviposition behavior of SSB moths

Collection and analysis of rice plant volatiles

Individual rice plants were either uninfested or infested with SSB larvae alone, BPH nymphs alone, or both species simultaneously for 48 hr using the method described above. The emitted volatiles were trapped for 3 hr (21:00–24:00, the time period that SSB lay their eggs), and then analyzed and identified as described²⁵. The compounds were quantified as a percentage of peak areas relative to the internal standard (nonyl acetate) per 3 hr of trapping for one plant. For each treatment, collections and analyses were repeated 7–9 times.

Odor preferences of SSB females

The response of SSB females to volatiles released from differently treated rice plants were investigated to better understand the mechanism underlying the moth's oviposition preferences. The total volatiles emitted from uninfested plants, SSB-infested plants, BPH-infested plants and SSB/BPH-infested plants were collected for this experiment. Plant treatments and volatiles collections were the same as described above but without the addition of the internal standard. The collected volatiles were diluted in paraffin oil (purity 99%; Sigma-Aldrich, St. Louis, MO, USA) at 1:4 (v/v) and were stored at $-80\text{ }^{\circ}\text{C}$ before use.

One ml of each of the four types of volatile solutions were separately

754 pipetted on the center of a filter paper strip (4 cm × 21 cm), which were then
755 hung from the four corners of a cage (45 cm × 45 cm × 45 cm) made of
756 80-mesh nylon net. Five pairs of freshly emerged SSB moths (< 24 hr) were
757 released in each cage. After 72 hr, the number of eggs deposited on the filter
758 paper strips and the surface of the nylon nets near each paper strip were
759 determined. This oviposition choice test was repeated 8 times.

760

761 **Response of the egg parasitoid *T. japonicum* to herbivore-infested rice** 762 **plants**

763 Multiple types of herbivore-infested rice plants were prepared: i) uninfested
764 plants (control); ii) SSB-infested plants; iii) BPH-infested plants; iv)
765 SSB/BPH-infested plants; v) plants infested with SSB eggs (referred to as
766 egg-infested plants); vi) plant infested with SSB larvae and their eggs (referred
767 to as SSB/egg-infested plants); vii) plants infested with BPH nymphs and SSB
768 eggs (BPH/egg-infested plants); and viii) plants infested with both SSB larvae,
769 BPH nymphs and SSB eggs (referred to as SSB/BPH/egg-infested plants). To
770 prepare these treatments, plants were first artificially infested with herbivores
771 for 48 hr as described above, then some of them were subjected to SSB eggs
772 deposition. For that, two potted rice plants of the same type were placed in a
773 cage (45 cm × 45 cm × 45 cm) made of 80-mesh nylon nets, then 30 pairs of
774 freshly emerged moths (< 24 hr) were released in each cage to mate and lay
775 eggs. After 24 hr, the plants were removed from the cage and those that
776 carried 200-250 eggs were used as odor sources. During the period of egg
777 deposition and the subsequent olfactometer experiments with the parasitoid,
778 all insects remained in or on the rice plants.

779 To test the behavioral responses of *T. japonicum* to differently treated rice
780 plants, they were offered the following pairs of odor sources: i) uninfested
781 plants versus egg-infested plants; ii) uninfested plants versus SSB-infested
782 plants; iii) uninfested plants versus BPH-infested plants; iv) egg-infested plants
783 versus SSB/egg-infested plants; v) egg-infested plants versus BPH/eggs

infested plants; vi) SSB/egg-infested plants versus BPH/egg-infested plants;
vii) egg-infested plants versus SSB/BPH/egg-infested plants; viii)
SSB/egg-infested plants versus SSB/BPH/egg-infested plants; and ix)
BPH/egg-infested plants versus SSB/BPH/egg-infested plants.

Responses of *T. japonicun* females to these odor sources were investigate
in a Y-tube olfactometer as described ²⁵. Newly emerged adult wasps were
maintained in glass tubes (3.5 cm diameter, 20 cm height) for at least 6 hr to
ensure that they would mate, before females were used for the experiments.
Two rice plants of the same treatment were enclosed in a glass bottle and
used as one odor source, and each pair of odor sources was replaced after ten
parasitic wasps were tested. For each treatment, a total of 64–88 female
wasps were tested. The experiments were conducted between 10:00 and
16:00 on several consecutive days.

Parasitism rates of *C. suppressalis* eggs by *T. japonicun* wasps

In a cage experiment, we further tested if the differences in parasitoid
attraction observed in the olfactometer for the differentially infested plants can
result in differences in parasitism rates of SSB eggs under realistic conditions.
The following herbivore-treated plants were prepared as described above:
SSB eggs on uninfested, SSB-infested, BPH-infested and SSB/BPH-infested
plants. The four types of plants were placed in the four corners of a cage
(60cm × 60 cm × 60 cm) made of 80-mesh nylon nets, respectively.
Subsequently, 40 pairs of newly emerged wasps (<1 day old) were released
into the cage. After 48 hr, the rice leaves with SSB eggs were collected, and
the total number of SSB eggs on each plant was counted and their
parasitization status was determined under a microscope two days later; the
eggs turned black after being parasitized for 3 days. The experiment was
replicated 12 times. The experiment was performed in a greenhouse at 27 ±
3°C and with 75 ± 10% RH and a photoperiod of 16:8 hr L:D.

814 **Statistical analyses**

815 All data were checked for normality and equality of variances prior to statistical
816 analysis. Likelihood ratio test (LR test) applied to a generalized lineal mixed
817 model (GLMM) for overdispersion and grouped design were conducted to
818 compare the number of eggs laid by SSB females on rice plants (Poisson error
819 structure with log link function). Likelihood ratio test (LR test) applied to a
820 generalized lineal model (GLM) were conducted to compare the parasitism
821 rates of SSB eggs by *T. japonicun* (normal distribution error) with the cage
822 treated as a random factor; the percentage data of parasitism rates were
823 arcsin square-root transformed prior to analyses. Two-way analysis of variance
824 (ANOVA) followed by least significant difference (LSD) test was used to
825 compare the body weight increases of the SSB larvae on different plant
826 treatments. The contents of JA and SA in different samples was analyzed
827 using one-way ANOVA followed by Tukey honest significant difference (HSD)
828 test or two-sided Student's *t*-test. Behavioral responses of *T. japonicun* in
829 Y-tube assays were analyzed using binomial test with an expected response of
830 50% for either olfactometer arm; parasitoids that did not make a choice were
831 excluded from the analysis. Differences in volatile emission and in gene
832 expression were analyzed by partial least squares-discriminant analysis
833 (PLS-DA)^{25,55} using SIMCA 14.1 software (Umetrics, Umeå, Sweden). The
834 omics data were normalized by medians, log-transformed, and then
835 auto-scaled (mean centered and divided by the standard deviation of each
836 variable) using Metaboanalyst 4.0 software⁷³ before they were subjected to
837 PLS-DA. All statistical analyses were conducted with SPSS 22.0 (IBM SPSS,
838 Somers, NY, USA), except for the PLS-DA were performed using SIMCA 14.1
839 software.

840

841 **Data availability**

842 RNA-seq raw data have been deposited to the Gene Expression Omnibus
843 (GEO) database in the National Center for Biotechnology Information (NCBI)

(note: accession number is not open yet). All other relevant data are available from the corresponding author upon reasonable request.

846

847 **Acknowledgements**

848 The study was supported by the National Natural Science Foundation of China
849 (31972984 and 31901896). The contribution by T.C.J.T. was supported by
850 European Research Council Advanced Grant 788949.

851

852 **Author contributions**

853 Y.Li conceived and directed the project. Y.Li, Q.L., X.H., and T.C.J.T. designed
854 the study. X.H. and S.S. performed the experiments. Q.L., X.H., S.S., Y.Li, and
855 T.C.J.T analyzed the data. Q.L., X.H., S.S., Y.P., G.Y., Y.Lou, T.C.J.T., and Y.Li
856 wrote the manuscript. All authors have read and approved the manuscript for
857 publication.

858

859 **Competing interests**

860 The authors declare no competing financial interests.

861

862 **Additional information**

863 **Supplementary information** The online version contains supplementary
864 material available.

865 **Correspondence** and requests for materials should be addressed to Y.Li.

866

867

868 **References**

- 869 1 Kaplan, I. & Denno, R. F. Interspecific interactions in phytophagous
870 insects revisited: a quantitative assessment of competition theory. *Ecol.*
871 *Lett.* **10**, 977-994 (2007).
- 872 2 Ohgushi, T. in *Trait-mediated indirect interactions: ecological and*
873 *evolutionary perspective* (eds T. Ohgushi, O. Schmitz, & R.D. Holt)
874 161-185 (Cambridge University Press, 2012).
- 875 3 Yamauchi, A., Ikegawa, Y., Ohgushi, T. & Namba, T. Density regulation

- 876 of co-occurring herbivores via two indirect effects mediated by biomass
877 and non-specific induced plant defenses. *Theor. Ecol.* Doi:
878 10.1007/s12080-020-00479-2 (2020).
- 879 4 Tilman, D. *Resource competition and community structure*. (Princeton
880 university press, 1982).
- 881 5 Foster, K. R. & Bell, T. Competition, not cooperation, dominates
882 interactions among culturable microbial species. *Curr. Biol.* **22**,
883 1845-1850 (2012).
- 884 6 Elias, S. & Banin, E. Multi-species biofilms: living with friendly neighbors.
885 *FEMS Microbiol. Rev.* **36**, 990-1004 (2012).
- 886 7 Little, A. E., Robinson, C. J., Peterson, S. B., Raffa, K. F. & Handelsman,
887 J. Rules of engagement: interspecies interactions that regulate
888 microbial communities. *Annu. Rev. Microbiol.* **62**, 375-401 (2008).
- 889 8 Lohberger, A. et al. Effect of organic carbon and nitrogen on the
890 interactions of *Morchella* spp. and bacteria dispersing on their mycelium.
891 *Front. Microbiol.* **10**, 124 (2019).
- 892 9 Bshary, R., Hohner, A., Ait-el-Djoudi, K. & Fricke, H. Interspecific
893 communicative and coordinated hunting between groupers and giant
894 moray eels in the Red Sea. *PLoS Biol.* **4**, e431 (2006).
- 895 10 Vail, A. L., Manica, A. & Bshary, R. Referential gestures in fish
896 collaborative hunting. *Nat. Commun.* **4**, 1765 (2013).
- 897 11 Sfarra, A. *10 amazing acts of cooperation between different animal*
898 *species*,
899 <[https://listverse.com/2015/02/23/10-amazing-cooperations-between-di](https://listverse.com/2015/02/23/10-amazing-cooperations-between-different-animal-species/)
900 [fferent-animal-species/](https://listverse.com/2015/02/23/10-amazing-cooperations-between-different-animal-species/)> (2015).
- 901 12 Parmentier, T., Yéo, K., Dekoninck, W. & Wenseleers, T. An apparent
902 mutualism between Afrotropical ant species sharing the same nest.
903 *Behav. Ecol. Sociobiol.* **71** (2017).
- 904 13 Yao, I. Costs and constraints in aphid-ant mutualism. *Ecol. Res.* **29**,
905 383-391 (2014).
- 906 14 Debbie, H. *How ants and aphids help each other*, <
907 <https://www.thoughtco.com/aphid-herding-ants-1968237>>(2019).
- 908 15 Denno, R. F., McClure, M. S. & Ott, J. R. Interspecific Interactions in
909 Phytophagous Insects: Competition Reexamined and Resurrected.
910 *Annu. Rev. Entomol.* **40**, 297-331 (1995).
- 911 16 Poelman, E. H. & Dicke, M. in *Annual Plant Reviews: Insect - Plant*
912 *Interactions* Vol. 47 (eds C. Voelckel & G. Jander) 309-338
913 (Wiley-Blackwell, 2014).
- 914 17 Stam, J. M. et al. Plant interactions with multiple insect herbivores: from
915 community to genes. *Annu. Rev. Plant Biol.* **65**, 689-713 (2014).
- 916 18 Li, X. et al. Reduced *Mythimna separata* infestation on *Bt* corn could
917 benefit aphids. *Insect Sci.* doi: 10.1111/1744-7917.12833 (2020).
- 918 19 Nakamura, M., Miyamoto, Y. & Ohgushi, T. Gall initiation enhances the
919 availability of food resources for herbivorous insects. *Funct. Ecol.* **17**,

- 851-857 (2003).
- 20 Soler, R. et al. Plant-mediated facilitation between a leaf-feeding and a phloem-feeding insect in a brassicaceous plant: from insect performance to gene transcription. *Funct. Ecol.* **26**, 156-166 (2012).
- 21 Su, Q. et al. A non-vector herbivore indirectly increases the transmission of a vector-borne virus by reducing plant chemical defences. *Funct. Ecol.* **34**, 1091-1101 (2020).
- 22 Utsumi, S. & Ohgushi, T. Host plant variation in plant-mediated indirect effects: moth boring - induced susceptibility of willows to a specialist leaf beetle. *Ecol. Entomol.* **33**, 250-260 (2008).
- 23 Shiojiri, K., Takabayashi, J., Yano, S. & Takafuji, A. Oviposition preferences of herbivores are affected by tritrophic interaction webs. *Ecol. Lett.* **5**, 186-192 (2002).
- 24 Desurmont, G. A., Guiguet, A. & Turlings, T. C. J. Invasive insect herbivores as disrupters of chemically-mediated tritrophic interactions: effects of herbivore density and parasitoid learning. *Biol. Invasions* **20**, 195-206 (2017).
- 25 Hu, X. et al. Caterpillar-induced rice volatiles provide enemy-free space for the offspring of the brown planthopper. *eLife* **9**, e55421 (2020).
- 26 Zhang, P. J. et al. Whiteflies interfere with indirect plant defense against spider mites in Lima bean. *Proc Natl Acad Sci USA* **106**, 21202-21207 (2009).
- 27 Desurmont, G. A., Zemanova, M. A. & Turlings, T. C. The gastropod menace: slugs on Brassica plants affect caterpillar survival through consumption and interference with parasitoid attraction. *J. Chem. Ecol.* **42**, 183-192 (2016).
- 28 Zarate, S. I., Kempema, L. A. & Walling, L. L. Silverleaf whitefly induces salicylic acid defenses and suppresses effectual jasmonic acid defenses. *Plant Physiol.* **143**, 866-875 (2007).
- 29 Chen, M., Shelton, A. & Ye, G. Y. Insect-resistant genetically modified rice in China: from research to commercialization. *Annu. Rev. Entomol.* **56**, 81-101 (2011).
- 30 Jiao, Y. et al. *Bt* rice plants may protect neighbouring non-*Bt* rice plants against the striped stem borer, *Chilo suppressalis*. *Proc. R. Soc. B* **285**, 20181283 (2018).
- 31 Ye, M. et al. Suppression of a leucine-rich repeat receptor-like kinase enhances host plant resistance to a specialist herbivore. *Plant Cell Environ.* **43**, 2571-2585 (2020).
- 32 Lu, H. P. et al. Resistance of rice to insect pests mediated by suppression of serotonin biosynthesis. *Nat. Plants* **4**, 338-344 (2018).
- 33 Barker, J. L. et al. Synthesizing perspectives on the evolution of cooperation within and between species. *Evolution* **71**, 814-825 (2017).
- 34 Wang, X. et al. *Bt* rice could provide ecological resistance against nontarget planthoppers. *Plant Biotechnol. J.* **16**, 1748-1755 (2018).

964 35 Lu, J. et al. Contrasting effects of ethylene biosynthesis on induced
965 plant resistance against a chewing and a piercing-sucking herbivore in
966 rice. *Mol. Plant* **7**, 1670-1682 (2014).

967 36 Zhou, G. et al. Silencing *OsHI-LOX* makes rice more susceptible to
968 chewing herbivores, but enhances resistance to a phloem feeder. *Plant*
969 *J.* **60**, 638-648 (2009).

970 37 Ye, M. et al. silencing *COI1* in rice increases susceptibility to chewing
971 insects and impairs inducible defense. *PloS one* **7**, e36214 (2012).

972 38 Wu, J. & Baldwin, I. T. New insights into plant responses to the attack
973 from insect herbivores. *Annu. Rev. Genet.* **44**, 1-24 (2010).

974 39 Erb, M. & Reymond, P. Molecular interactions between plants and
975 insect herbivores. *Annu. Rev. Plant Biol.* **70**, 527-557 (2019).

976 40 Farmer, E. E. *Leaf defence*. (Oxford University Press, 2014).

977 41 Mao, Y. B. et al. Jasmonate response decay and defense metabolite
978 accumulation contributes to age-regulated dynamics of plant insect
979 resistance. *Nat. Commun.* **8**, 13925 (2017).

980 42 Sarmento, R. A. et al. A herbivore that manipulates plant defence. *Ecol.*
981 *Lett.* **14**, 229-236 (2011).

982 43 De Lange, E. S. et al. *Spodoptera frugiperda* caterpillars suppress
983 herbivore-induced volatile emissions in maize. *J. Chem. Ecol.* **46**,
984 344-360 (2020).

985 44 Walling, L. L. Avoiding effective defenses: strategies employed by
986 phloem-feeding insects. *Plant Physiol.* **146**, 859-866 (2008).

987 45 Will, T., Tjallingii, W. F., Thonnessen, A. & van Bel, A. J. Molecular
988 sabotage of plant defense by aphid saliva. *Proc Natl Acad Sci USA* **104**,
989 10536-10541 (2007).

990 46 Gripenberg, S., Mayhew, P. J., Parnell, M. & Roslin, T. A meta-analysis
991 of preference-performance relationships in phytophagous insects. *Ecol.*
992 *Lett.* **13**, 383-393 (2010).

993 47 Turlings, T. C. J. & Erb, M. Tritrophic interactions mediated by
994 herbivore-induced plant volatiles: mechanisms, ecological relevance,
995 and application potential. *Annu. Rev. Entomol.* **63**, 433-452 (2018).

996 48 Tooker, J. F., Rohr, J. R., Abrahamson, W. G. & De Moraes, C. M. Gall
997 insects can avoid and alter indirect plant defenses. *New Phytol.* **178**,
998 657-671 (2008).

999 49 Hogenhout, S. A. & Bos, J. I. Effector proteins that modulate
1000 plant-insect interactions. *Curr. Opin. Plant Biol.* **14**, 422-428 (2011).

1001 50 Zu, P. et al. Information arms race explains plant-herbivore chemical
1002 communication in ecological communities. *Science* **368**, 1377-1381
1003 (2020).

1004 51 Gause, G. F. *The struggle for existence*. (Williams & Wilkins, 1934).

1005 52 Park, T. Experimental studies of interspecies competition. 1.
1006 Competition between populations of the flour beetles, *Tribolium*
1007 *confusum* Duval and *Tribolium castaneum* Herbst. *Ecol. Monogr.* **18**,

1008 265-307 (1948).
1009 53 Janzen, D. H. Host plants as islands. II. Competition in evolutionary and
1010 contemporary time. *Am. Nat.* **107**, 786-790 (1973).
1011 54 Kant, M. R. et al. Mechanisms and ecological consequences of plant
1012 defence induction and suppression in herbivore communities. *Ann. Bot.*
1013 **115**, 1015-1051 (2015).
1014 55 Zhang, P. J. et al. Jasmonate and ethylene signaling mediate
1015 whitefly-induced interference with indirect plant defense in *Arabidopsis*
1016 *thaliana*. *New Phytol.* **197**, 1291-1299 (2013).
1017 56 Ali, J. G., Agrawal, A. A. & Fox, C. Asymmetry of plant-mediated
1018 interactions between specialist aphids and caterpillars on two
1019 milkweeds. *Funct. Ecol.* **28**, 1404-1412 (2014).
1020 57 Ray, S. et al. Asymmetry in herbivore effector responses: caterpillar
1021 frass effectors reduce performance of a subsequent herbivore. *J. Chem.*
1022 *Ecol.* **46**, 76-83 (2020).
1023 58 Chisholm, P. J., Eigenbrode, S. D., Clark, R. E., Basu, S. & Crowder, D.
1024 W. Plant-mediated interactions between a vector and a non-vector
1025 herbivore promote the spread of a plant virus. *Proc. Biol. Sci.* **286**,
1026 20191383 (2019).
1027 59 Jiang, Y., Zhang, C. X., Chen, R. & He, S. Y. Challenging battles of
1028 plants with phloem-feeding insects and prokaryotic pathogens. *Proc*
1029 *Natl Acad Sci USA* **116**, 23390-23397 (2019).
1030 60 Johnson, S. N., Karley, A. J., Gregory, P. J. & Brennan, R. M. Editorial:
1031 Crop traits for defense against pests and disease: durability, breakdown
1032 and future prospects. *Front. Plant Sci.* **8**, 209 (2017).
1033 61 Chen, Y. H., Gols, R. & Benrey, B. Crop domestication and its impact on
1034 naturally selected trophic interactions. *Annu. Rev. Entomol.* **60**, 35-58
1035 (2015).
1036 62 Han, L., Li, S., Liu, P., Peng, Y. & Hou, M. New artificial diet for
1037 continuous rearing of *Chilo suppressalis* (Lepidoptera: Crambidae). *Ann.*
1038 *Entomol. Soc. Am.* **105**, 253-258 (2012).
1039 63 Liu, Q. et al. Plant breeding involving genetic engineering does not
1040 result in unacceptable unintended effects in rice relative to conventional
1041 cross-breeding. *Plant J.* **103**, 2236-2249 (2020).
1042 64 Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner
1043 with low memory requirements. *Nat. Methods* **12**, 357-360 (2015).
1044 65 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general
1045 purpose program for assigning sequence reads to genomic features.
1046 *Bioinformatics* **30**, 923-930 (2014).
1047 66 Garber, M., Grabherr, M. G., Guttman, M. & Trapnell, C. Computational
1048 methods for transcriptome annotation and quantification using RNA-seq.
1049 *Nat. Methods* **8**, 469-477 (2011).
1050 67 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change
1051 and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550

1052 (2014).
1053 68 Yu, G., Wang, L., Han, Y. & He, Q. clusterProfiler: an R package for
1054 comparing biological themes among gene clusters. *OMICS: J.*
1055 *Integrative Biol.* **16**, 284-287 (2012).
1056 69 Volodarsky, D., Leviatan, N., Otcheretianski, A. & Fluhr, R.
1057 HORMONOMETER: a tool for discerning transcript signatures of
1058 hormone action in the *Arabidopsis* transcriptome. *Plant Physiol.* **150**,
1059 1796-1805 (2009).
1060 70 Wang, Y., Liu, Q., Du, L., Hallerman, E. M. & Li, Y. Transcriptomic and
1061 metabolomic responses of rice plants to *Cnaphalocrocis medinalis*
1062 caterpillar infestation. *Insects* **11**, 705 (2020).
1063 71 Liu, Q. et al. Combined transcriptome and metabolome analyses to
1064 understand the dynamic responses of rice plants to attack by the rice
1065 stem borer *Chilo suppressalis* (Lepidoptera: Crambidae). *BMC Plant*
1066 *Biol.* **16**, 259 (2016).
1067 72 Fu, J., Chu, J., Sun, X., Wang, J. & Yan, C. Simple, rapid, and
1068 simultaneous assay of multiple carboxyl containing phytohormones in
1069 wounded tomatoes by UPLC-MS/MS using single SPE purification and
1070 isotope dilution. *Anal. Sci.* **28**, 1081-1087 (2012).
1071 73 Chong, J. et al. MetaboAnalyst 4.0: towards more transparent and
1072 integrative metabolomics analysis. *Nucleic Acids Res.* **46**, W486-W494
1073 (2018).
1074