1 In-land long-term culture of colonial ascidians

2 Wawrzyniak M.¹, Matas Serrato L.A.¹, Blanchoud S.^{1*}

3

4 1. Department of Biology, University of Fribourg, Chemin du Musée 10, 1700 Fribourg,
5 Switzerland

6 * corresponding author: simon.blanchoud@unifr.ch

7

8 Highlights:

- 9 • First in-land recirculating aquaculture for colonial ascidians
- 10 • Over 3 years of continuous breeding
- 11 • Semi-automated setup with minimized maintenance
- 12 • Good biomass production for strain propagation
- 13 • 4 different culture conditions for optimized breeding of species of interest

14

15 Abstract

16 Tunicates are highly diverse marine invertebrate filter-feeders that are vertebrates' closest
17 relatives. These organisms, despite a drastically different body plan during their adulthood,
18 have a tissue complexity related to that of vertebrates. Ascidians, which compose most of the
19 Tunicata, are benthic sessile hermaphrodites that reproduce sexually through a motile tadpole
20 larval stage. Over half of the known ascidians species are able to reproduce asexually by
21 budding, typically leading to the formation of colonies where animals, called zooids, are
22 interconnected through an external vascular system. In addition, colonial ascidians are
23 established models for important biological processes including allrecognition,

immunobiology, aging, angiogenesis and whole-body regeneration. However, the current paucity in breeding infrastructures limits the study of these animals to coastal regions.

To promote a wider scientific spreading and popularity of colonial ascidians, we have developed a flexible recirculating husbandry setup for their long-term in-lab culture. Our system is inspired both by the flow-through aquariums used by coastal ascidian labs, as well as by the recirculating in-lab systems used for zebrafish research. Our hybrid system thus combines colony breeding, water filtering and food culturing in a semi-automated system where specimens develop on hanging microscopy glass slides. Temperature, light/dark cycles, flow speed and feeding rates can be controlled independently in four different breeding environments to provide room for species-specific optimization as well as for running experiments. This setup is complemented with a quarantine for the acclimatization of wild isolates.

Herein we present our success in breeding *Botrylloides diegensis*, a species of colonial ascidians, for more than 3 years in recirculating artificial seawater over 600 km away from their natural habitat. We show that colonies adapt well to in-lab culturing and that a specific strain can be isolated, propagated and efficiently used for research over prolonged periods of time. The flexible and modular structure of our system can be scaled and adapted to the needs of specific species as well as of particular laboratory spaces. Overall, we show that *Botrylloides diegensis* can be proficiently bred in-land and suggest that our results can be extended to other species of colonial ascidians to promote research on these fascinating animals.

Keywords: Aquaculture, colonial tunicates, recirculating system, *Botrylloides*, *Botryllus*, breeding

46 1. Introduction

47 Tunicates are a group of over 3000 known extent highly diverse worldwide marine
 48 invertebrates, the majority of which are sub-tidal suspension-feeding hermaphrodites (Holland,
 49 2016). Tunicata is separated into three classes (WoRMS Editorial Board, 2017): Appendicularia,
 50 Thaliacea and Ascidiacea. Appendicularia has 68 identified species of planktonic free-
 51 swimming organisms that possess chordate traits common to most tunicate larvae including a
 52 notochord, neural tube and pharyngeal slits (Holland, 2016). These social animals form
 53 communities where each individual is enclosed inside a special external mucous structure,
 54 termed house, which concentrates and funnels their food (Lambert, 2005). Thaliacea is
 55 composed of 77 species of planktonic pelagic animals forming free-floating compound
 56 colonies (Lambert, 2005). These organisms alternate between sexual reproduction to initiate
 57 new colonies, and asexual reproduction by budding to increase the size of the colony. With
 58 2935 currently identified species, of which 1730 are colonial species (Shenkar and Swalla,
 59 2011), Ascidiacea is by far the most diverse class of tunicates (WoRMS Editorial Board, 2017).
 60 Ascidians are sessile benthic organisms that are present in all marine habitats, from sandy
 61 beds to rocky shores and from coral reefs to kelp forests (Shenkar and Swalla, 2011). In
 62 addition, ascidians very efficiently colonize artificial substrata and have thus become a major
 63 source of biofouling, with a negative impact on economical sectors such as shellfish farming
 64 and marine transport (Aldred and Clare, 2014; Comeau et al., 2015).

65 However, ascidians also represent great opportunities for research as well as for the
 66 industry. First, Tunicata's phylogenetic position as vertebrates' closest relatives (Delsuc et al.,
 67 2006) renders them key models for evolutionary research, and to study the emergence of
 68 chordate-specific processes in particular. Second, colonial ascidians (**Fig. 1A**) are the only

known chordates capable of undergoing whole-body regeneration whereby a fully functional adult is restored from a minute fragment of vasculature in as little as 10 days (Blanchoud et al., 2018a). Third, their recently established application as bioindicators of micro-pollution (Navon et al., 2020; Tzafriri-Milo et al., 2019; Vered et al., 2019) combined with their ubiquitous localization makes these animals a perfect tool for environmental bio-surveillance. Fourth, some colonial ascidians have emerged as model organisms for an increasing number of research fields, including immunobiology, allorecognition, angiogenesis and aging (Blanchoud et al., 2018a). In addition, ascidians are a particularly promising source of marine natural products (MNP) with already over 1000 isolated molecules (Chen et al., 2019) and numerous reported applications including antibacterial, anti-fungal, anti-viral, anti-inflammatory and antitumoral (Palanisamy et al., 2017). There is thus a tremendous scientific interest in the study of ascidians. In our lab, we are particularly interested in *Botrylloides diegensis* because it is the only known colonial tunicate that combines a high quality genome assembly (Blanchoud et al., 2018b), a constant WBR ability throughout its developmental cycle (Blanchoud et al., 2018a) and an identified population of stem-cells responsible for its capacity to undergo WBR (Kassmer et al., 2020). Consequently, we set out to develop a suitable husbandry setup in our research institute, located over 5 hours of travel from the closest sea.

Ascidian research is a particularly active community with numerous groups working worldwide using three main culturing methods. One most straightforward technique is onsite processing (Roberts et al., 2007; Sheets et al., 2016; Zondag et al., 2016; Ueki et al., 2018; Hiebert et al., 2019) (**Fig. 1B**). No long-term culture is involved, animals are sampled in their wild natural environment and directly processed for downstream analysis. This approach is thus well suited for studying *in situ* physiological aspects of Tunicates but it requires a reasonably

close-by sampling site and can be influenced by the potential seasonal availability of some species in some regions of the globe (Coma et al., 2000). A second prominent approach is the culture in a flow-through system (Hirose et al., 1995; Taketa et al., 2015; Blanchoud et al., 2017; Scelzo et al., 2019) (**Fig. 1C**), typically with filtered natural seawater from the site of sampling, which provides close to natural environmental conditions including in terms of feeding diet. This method allows to isolate and maintain strains on the long-term in confined and hazards-free conditions. It does require either a close proximity to a source of seawater or a consequent organization to provide for the necessary amounts of seawater. A third common approach is the culture in a closed system with periodic water changes (Brown et al., 2009; Epelbaum, 2009; Rinkevich et al., 2010) (**Fig. 1D**). This setup is most typically used with water from the sampling site for short-term culture of animals, although it has been used for extensively long studies too (Rinkevich and Shapira, 1998), as well as with artificial sea water (Nagar and Shenkar, 2016). The main limitations of this approach is the maintenance it requires on the long term, and the amount of water used when culturing a large number of colonies simultaneously. Consequently, all the labs that have the opportunity to do so combine these three approaches to benefit from their specific advantages. Yet, it consistently requires a routine access to either wild specimens or natural seawater and thus restricts their application to coastal locations. The only exception to this spatial limitation that we are aware of is that of Joly et al. who developed a recirculating system for breeding transgenic lines of the solitary ascidian *Ciona intestinalis* (Joly et al., 2007). Consequently, there is currently, to our knowledge, no recirculating husbandry system for the culture of colonial ascidians, which spatially limits their study and thus hinders their popularity as model organisms.

Here, we present a husbandry system designed for the in-land breeding of sessile colonial tunicates (**Fig. 2**). We have been successfully using this setup for over 3 years with the continuous presence of some of our strains, combined with a periodic addition of new wild isolates. Our recirculating system runs on artificial seawater and uses commercial algae pastes as feeding diets. Our aquarium is designed to permit up to 4 different culture conditions in parallel to optimize the breeding of the species of interest. In this study, we have been growing both *Botrylloides diegensis* as well as *Botryllus schlosseri* on microscopy glass slides. We present our developed aquaculture system, the maintenance setup used for the colonies, our long-term results in the clonal expansion of colonial ascidians and an example of a comparative breeding experiment. Our data shows that we are able on the long-term to abolish any spatial constraints with respect to the natural environment of colonial ascidians. We believe that this setup will enable research on colonial tunicates to be undertaken in novel places and will thus promote as well as popularize these fascinating animals as model organisms.

127

128 2. Materials and Methods

129 2.1 Aquarium structure

The complete list of material and equipment used to assemble our system is provided in **Table 1 (Supplementary Figures S1-S4)**. Briefly, the structure of the system is a SS 316 40x40 and 30x30 mm powder coated stainless steel stand (LxWxH 2.0 x 0.9 x 2.2 m), custom manufactured PE/PP 40 L specimen tanks, a custom manufactured PE/PP 200 L filtration sump as well as four custom manufactured PE/PP 10 L food culturing vessels. The system is equipped with the following materials: fountain pump (universal 300, Eheim, DE), air blower (Medo LA-60B, Nitto Kohki, DE), cooling unit (Titan 4000, Aqua Medic, DE), heaters (titanium D-

137 D 600W, Schego, DE), motor driven protein skimmer (DC runner 1.1, AquaMedic GmbH, DE),
 138 circulation pump (Titan-Tec 10000, Messner GmbH, DE), UV steriliser (Vecton2, TMC Aquarium,
 139 US), lights (AquaBeam 1500 Ocean blues, TMC Aquarium, US). The construction of our
 140 prototype was outsourced to a company specialized in aquatic research equipment (Zebcare,
 141 Fleuren & Nooijen BV, NL).

142 A quarantine area composed of two flow-through 30 L tanks (Gripline, Semadeni AG, CH),
 143 aerated by one pump each (Wavemaker Jecod SW-2, Jebao, CN) and illuminated by a led rack
 144 each (Marine hybrid powerLED+, Eheim, DE) was added to the main system by our
 145 departmental workshop. Our quarantine tanks are placed on a heavy-duty plastic stand
 146 (Camshelving Premium, Cambro, US). Staining plastic racks (1233, Semadeni AG, CH) were used
 147 for hosting the animals on their glass slides.

148 The entire setup was equipped with a range of sensors to obtain a fully automatized
 149 control on the environment of the aquarium using the Apex (Apex 2016, Neptune Systems, US)
 150 controller, a custom micro-controller (Pico, Raspberry Pi, UK) as well as light controllers
 151 (BioLumen, TMC Aquarium, US). In particular, salinity, pH, day/night cycles, temperature and
 152 changes of water are performed automatically and reproducibly.

153

154 2.2 Artificial sea water production and filtration

155 All kits for water quality as well as syn-biotic sea salt were purchased from Tropic Marin
 156 (DE). Seawater refractometer was purchased from Red Sea (US), Optic USB base station and
 157 Hobo Pendant temperature/light data logger were purchased from Onset Computer
 158 Corporation (US). Filtration is performed by a nitrate filter (Sulphur BioDenitrator, Korallin, DE),
 159 ceramic beads (Tetra, DE), nitrifying bacteria (Bio reefclear, Sera, DE), stirred plastic moving bed

160 (K+Media, Evolution Aqua, UK), an algae scrubber (Medium Turf Scrubber, IceCap, US) and
 161 phosphate absorption resin (Aqua Crown Phosphatadsorber, Deltec, DE). Live stones cured
 162 with Coralline algae as well as macroalgae were bought from Aqua-Home (CH). Concentrated
 163 marine biota was made by sterile filtering 10 L of gravity filtered natural seawater using 0.22
 164 µm Nalgene filters and resuspending the content of the filters in 50 mL of gravity filtered natural
 165 seawater. Other chemicals and small laboratory equipment were purchased from Thermo
 166 Fisher Scientific (US).

167

168 2.3 Animal breeding and handling

169 Numerous colonial ascidians (*Botrylloides diegensis* in particular) were collected in the
 170 coastal town of Chioggia (IT) and successfully brought back to the lab with their original
 171 substrate in closed boxes filled with sea water. Half a dozen mussels were also collected to
 172 inoculate the husbandry system. Animals were then acclimatized in our quarantines and
 173 transferred to 76x26x1 mm microscopic slides (Marienfeld, DE). After the colonies had properly
 174 settled on the slide, they were transferred to the recirculating system where they developed
 175 vertically inside the specimen tanks using a custom attachment setup using cable ties
 176 (Netztech Handels AG, CH) and a silicone badge gripper (Specialist ID, US). Colonies were fed
 177 using a blend of commercial algae concentrates from Instant algae (Shellfish Diet 1800,
 178 RotiGrow Plus, US), complemented with a planktonic substitute (Zootonic, Tropic Marin, DE)
 179 and commercial bakery yeast (Rinkevich and Shapira, 1998). Feeding was controlled by 12V
 180 normal closed solenoid valves (CJV23, Conjoin, CN) controlled by a customized microcontroller
 181 (Pico, Raspberry Pi, UK, **Supplementary Material 1**). Colonies were monitored weekly by
 182 macrophotography using a Canon 5D Mark II with objective Sigma 50 mm f/2.8 EX DG macro

183 and a flash Canon Speedlite 430EX II. Statistical significance between growth rates was
184 measured using two-tailed Student's *t*-tests. Salinity was kept constant at 35 g/L of seasalt,
185 temperature was set at 21 °C all year round.

186 Live L-type rotifers (Planktovie, FR) were purchased in brackish 15 g/L seawater. We thus
187 acclimatized them to our system's 35 g/L by slowly increasing the salinity of their medium by 5
188 g/L every week. Rotifers were cultured in 1 L Erlenmeyer flasks without agitation, fed with RG
189 Complete every day and their water was changed every 2 days. Once the population became
190 sufficiently dense, typically every other day, they were splitted in half. One half serving for
191 continuing their expansion, the other half being fed to the specimen tanks.

192 3. Results

193 3.1. Description of the system

194 We designed a custom recirculating system for the in-land culture of colonial tunicates
 195 (**Fig. 2A**), which allows to reduce periodic water changes to minimal amounts. In such closed
 196 systems, the wastes produced by the animals (e.g. NH_4^+ , NO_2^- , NO_3^-) accumulate over time and
 197 need to be continuously purified. Our modular aquaculture system is structured in three levels
 198 around a stainless stand to maximize the use of gravity for the circulation of the artificial sea
 199 water (ASW) in the aquarium. On the top, food culturing tanks distribute their content into the
 200 specimen tanks located in the middle, which overflow into the water filtration units (sump)
 201 placed at the bottom of the apparatus (**Fig. 2B**). The filtration unit combines mechanical and
 202 biological filters to reach a minimal weekly water change of 5 % of the total 450 L of ASW
 203 present in our system. In particular, filtration includes a particle filter, a so-called moving bed
 204 filter (or upflow filter), a nitrate filter, an algae scrubber, a protein skimmer, live cured stones and
 205 a UV-C unit (**Fig. 2C**).

206 To breed our animals in optimal conditions, we specifically aimed at using a maximum of
 207 food-grade plastic material, to avoid rust, wear and chemical contamination as much as
 208 possible. Four 40 L specimen tanks were custom made from 15/20mm PE/PP material. Tanks
 209 have one compartment and each has a heater which is placed at the bottom of the tank and is
 210 controlled by a temperature sensor placed on the side of the tank. Aeration and mixing of the
 211 water are ensured by a circulation pump placed on the back side of each tank. Tanks are
 212 completely closed to make a light-isolated environment, with a convenient front opening (**Fig.**
 213 **2D**) for easy access and cleaning. Water falls into the tank through a silicone tube, which flow
 214 rate can be regulated to modulate the amount of flow-through, and leaves the tank from an

215 outlet located in the back of the tank. A fleece is placed at the bottom of the tanks to collect
 216 solid wastes. Each tank is equipped with two lights sunken in the lid of the tank in such a way
 217 that the heat sink at the back of the LED tile is outside the tank for optimal heat transfer (**Fig.**
 218 **2D**). The intensity and photoperiod of each LED unit can be controlled independently through a
 219 central control unit. Animals are placed in the tanks on a custom system of hangers made
 220 using plastic tubes, plastic hooks and badge grippers (**Fig. 2E**).

221 The central filtration unit (sump) has been placed on the ground (**Supplementary Figure**
 222 **S1**). Water from the tanks is drained through a fleece filter into an aerated moving-bed biofilter,
 223 through a ceramic-beads biofilter and into the main sedimentation tank (**Fig. 2C**). The
 224 sedimentation tank is connected to an external protein skimmer with its submerged pump, a
 225 nitrate filter, an algae scrubber and a refugium populated with live cured stones and macro-
 226 algae. The refugium and the algae scrubber are constantly illuminated by two LED units. The
 227 sedimentation tank is connected to the main circulation pump, which provides system water to
 228 the cooling unit, the algae scrubber and the refugium in a closed loop, as well as to the
 229 specimen tanks through a UV sterilizer. In addition, periodic water changes are controlled by a
 230 timer-activated submerged fountain pump. This pump discards system water into the sewage
 231 through the sump's overflow drain to induce its replacement with fresh seawater. This pump
 232 can also be used for complete drainage of the system. In order to reduce evaporation, the
 233 whole filtration tank is fitted with a clear plastic lid.

234 Above the specimen tanks, we have installed four 10 L cylindrical feeding tanks
 235 (**Supplementary Figure S2**). Each tank is made of non-transparent PVC pipes 200 mm in
 236 diameter and 300 mm long. An external water level made of clear PVC hose and a visit window
 237 allow to monitor the content of each tank. Content of the tanks is mixed by injecting air through

238 a microporous tube placed at the bottom of the tank, and lighting is done by two LED rails
239 placed inside a clear pipe through the center of the non-air tight lid of the tank. Food is drained
240 outside of each tank by gravity through an electronically controlled micro-valve connected to
241 the outlet of the tank (**Supplementary Figure S3**). Air is supplied via 4 mm silicone hoses
242 directly from the main air pump through a custom-built manifold (**Supplementary Figure S4**).

243 Besides the main system, we also maintain two quarantine containers for new comers
244 and new subclones (**Fig. 2F**). This part has been designed and built up on site and is on
245 purpose not recirculating into the main system. However, both tanks can be filled with system
246 water, and system water can be collected there for experiments using handy garden hose
247 nozzles. Each container can be filled with up to 30 L of saltwater, aerated and mixed by a pump
248 placed in the back of the tank. Each tank is illuminated with photoperiod-controlled lights.
249 Animals are placed in standing plastic staining racks (**Fig. 2G**) inside the quarantine tanks.

250 The conditions in the system are monitored and logged automatically (**Fig. 2H**) using an
251 APEX system from Neptune Systems (**Supplementary Figure S3**), connected to a number of
252 sensors and actuators. Temperature, salinity, pH and ORP (oxidation reduction potential) are
253 measured every 200 s in the main sedimentation tank. Temperature is controlled by the cooling
254 unit and a heater, salinity can be increased by pumping 270 g/L brine water located in a 75 L
255 tank placed behind the feeding tanks using two peristaltic pumps, also controlled by Apex.
256 Brine tank is mixed four times per day using a heavy duty concrete mixer controlled by a timer
257 (**Supplementary Figure S4**). An alert email is sent if some values deviate from their defined ideal
258 range (7.9-8.5 pH, 27-35 salt, 14-24 °C, 300-400 ORP). Water level is monitored using two
259 optical sensors driving an automatic top-off kit that delivers deionized water directly in the main
260 sedimentation tank by opening a solenoid valve connected to the University's water system

(Supplementary Figure S1). Temperature is also measured individually in each specimen tank, and salinity is measured in one of the tanks for redundancy. Leaks and water loss are monitored using a floor sensor and an optical sensor placed at the bottom of the sedimentation tank, both leading to the shutdown of the whole system in case of emergency. Additional water quality parameters (pH, NH_4^+ , NO_2^- , NO_3^- , PO_4^{3-}) and the conditions in the quarantine are monitored manually.

267

3.2. Initialization and maintenance of the system

Before its first usage, the aquaculture system was cured over two weeks. First, the system was filled with tap water, run for 48 h and completely emptied. This wash was repeated two more times to remove all traces of chemicals from the system. During this time, the wild mussels were cured and cleaned in our quarantine using ASW changed once per week and fed using 1 mL/day of Shellfish Diet 1800. The cleaned aquarium was then filled up with ASW and curing was started. Ammonium chloride was added (3 mg/L) together with the nitrifying bacteria, the concentrated marine biota, the mussels and the live rocks, and the system was left to run for 24 hours. 5 grams of commercial fish food mix was added in the sedimentation tank followed by 7 g every week. Water quality (pH, NH_4^+ , NO_2^- , NO_3^-) was measured every other day until ammonium concentration started to decrease (1.5 weeks post inoculation). 1.5 mg/L of ammonium was then added and the monitoring continued. When the nitrite concentration started to decrease (4 weeks post inoculation), the nitrate filter was started following the manufacturer's instructions. 0.5 mg/L of ammonium was added and the system monitored until the water quality was within the range defined by commercial kits (6 weeks post

inoculation in total). System was then used following our routine maintenance as described below.

Maintenance of our aquaculture system was reduced to a minimum by automatizing as many steps as possible, the bulk of the remaining manual labor being dedicated to tending to the various needs of the animals. Weekly duties include only the cleaning of the protein skimmer and every other week the replacement of the fleece placed at the bottom of the specimen tanks for trapping solid waste as well as the cleaning of the algae scrubber. Once a month, the excess of algae growing in the refugium is removed, the live stones scrubbed clean and the Apex system checked. Every 2-3 months depending on our use of system water, the brine tank needs refilling, and once a year a general maintenance is done where all the tubing pumping brine water are thoroughly cleaned.

In our hands, the most important factors for a reliable performance of our recirculation system is the quality of the water; not inasmuch as having perfectly low concentrations for all the noxious compounds in the system water, but in preventing these compounds from accumulating to dangerously high concentrations. Consequently, we manually measured on a weekly basis the four most important ions (NH_4^+ , NO_2^- , NO_3^- , PO_4^{3-} ; **Fig. 2H**) as well as pH and salinity to confirm the readings from our controller. In addition, we tested monthly the following other parameters: KH, O_2 , I, Ca_2^+ , Mg_2^+ , Si and K^+ .

301

3.3. Breeding colonial tunicates in-land

Wild isolates were collected twice a year (Spring/Autumn) in the Venetian lagoon (**Fig. 1B**). Animals were transferred on their original substrate into 1 L water-tight wide-mouth cylindrical containers. Animals were then transported to the laboratory, which typically took

306 between 8 h and 10 h. Important parameters for successful transportation of the collected
 307 samples are first a large animal to volume ratio ($> 20X$), regular water changes during
 308 transportation (every 2-3 h), as little air in the container as possible and minimizing the
 309 turbulences during transport. However, the best predictor of a colony that will well tolerate
 310 being transferred was its size, namely medium-sized colonies composed of 50-100 zooids
 311 were typically fitter once in our laboratory. Another important aspect to consider when
 312 collecting is the type of substrate the animal is growing on, and its shape. Colonies growing flat
 313 on shellfish were the most appropriate for later transfer to our glass slides, while those
 314 intermingled with brittle algae were very difficult to isolate once in the lab.

315 Upon arrival, wild specimens were transferred to 2 L tanks filled with ASW adapted to the
 316 salinity and pH of the natural sea water they were transported in. Animals were cleaned, their
 317 substrate reduced to a minimum and the ones without any sign of filter-feeding activity
 318 discarded. Animals were placed in our quarantine tanks, fed twice daily with 1 mL of feeding
 319 mix and left 24 h to adapt. On the next day, colonies were subcloned to numbered microscopy
 320 glass slides using a single-edge razor blade following a customized protocol (Bugada and
 321 Blanchoud, 2021). Slides were placed in plastic staining racks at a density of 7 slides per rack
 322 standing at the bottom of the quarantine tank. Colonies were left for 48 h to attach to the slides,
 323 carefully cleaned (Blanchoud and Wawrzyniak, 2021), the decaying pieces of tissue discarded
 324 and the remaining attached colonies placed in the second quarantine tank. Water in the
 325 quarantines was changed when the NH_4^+ levels rose above 1.5 mg/L, or at least once a week.
 326 We have observed that in our 30 L setup, with up to two dozen colonies, this will take around 7
 327 days. The colonies that detached were subcloned again into the first quarantine and the
 328 process repeated until all colonies either attached or decayed. Colonies were then acclimatized

329 to the salinity of our recirculating system by lowering the salinity of our quarantine by 1 g/L per
 330 week. Once the quarantines' salinity was identical to that of our system, and the colonies had
 331 spent at least 2 weeks in quarantine under weekly monitoring against diseases and parasites,
 332 they were transferred to the main system where they proliferated actively (**Fig. 3A, Movie 1**).
 333 The lineage of all strains, their clonal offspring, lifespan, position in the system and treatments
 334 were recorded in an online database. Approximately 50 % of the collected specimens could be
 335 transferred successfully to our recirculating system.

336 Over the 3 years of this experiment, we transferred around 100 strains of colonial
 337 ascidians, and of *Botrylloides diegensis* in particular, from their natural environment to our
 338 recirculating system. We repeatedly observed that some strains declined in the first few
 339 months after their acclimatization and ultimately went extinguished without any apparent
 340 reason, while neighboring slides with different animals propagated totally fine. We could not
 341 find any pattern in the premature disappearance of these isolates. However, some strains have
 342 been much more productive than others (**Fig. 3B**) and vastly compensated these losses.
 343 Currently, 6 strains have ended up constituting over 95 % of all the colonies present in our
 344 aquaculture. Although we could have promoted a higher genetic diversity in our lab, this was
 345 not an objective in this study. We rather favored fast propagating animals to identify candidate
 346 reference strains for in-lab studies where biological replicates and large sample numbers are
 347 important.

348 In our system, colonies are cultured on numbered microscopy glass slides hanging
 349 vertically in the water, which reproduces somewhat their natural environment where we mainly
 350 collected them from the lateral sides of floating pontoons (**Fig. 1B**). This breeding method has
 351 multiple advantages, first for microscopy and general handling of the animals in the lab, second

for the ease of use of the badge gripper and the total absence of metal parts (**Fig. 2E**), and finally for the identification and localization of specific colonies within the system. Each slide hangs onto a hook placed onto a PVC tube, itself fixed in the top part of the specimen tanks. Given the space constrains of our system to allow manipulation of each animal individually, each specimen tank can contain up to 32 slides (**Fig. 2D**).

Colonies are checked upon at least once every two weeks, and often daily for slides participating in specific studies. In this occasion, colonies are taken out of the system, cleaned and subcloned if necessary. Subcloning was undertaken for three main reasons. First, for maintenance to compensate the propagation of the animals away from the center of the slide in particular, because this impedes its ease of manipulation and observation. Second to produce clonal biological replicates for experiments. Third, to fuse smaller clones from the same strain to obtain bigger samples for studies where large amounts of tissue were required.

Excitingly, several *B. diegensis* strains have been observed to become gravid (**Fig. 3C-D**). Some of these oocytes even got fertilized as we could identify oozoids developing later on in our system (**Fig. 3E**). Although this result was neither expected nor aimed for, it confirms that our colonies adapted well to our recirculating system and it shows that induction of sexual reproduction is possible under controlled artificial conditions.

3.4. Differential breeding of colonial tunicates

Colonies in our recirculating system were bred under continuous feeding following preliminary indications (Blanchoud, unpublished) that such delivery was promoting growth. In addition, this type of feeding has two advantages, first a reduced contamination of the colonies and their slides because of the overall lower concentration of food present in the water, and

375 second a lower daily maintenance as food needs to be prepared only once for multiple days.
 376 Continuous feeding was implemented using electronically controlled valves (**Supplementary**
 377 **Material 1**). We optimized the frequency and duration of the opening of each valve to reach
 378 uniform amounts between all four specimen tanks. On average, valves opened 1.5 s every 3
 379 min, thus delivering around 2 L of diluted food per day. Food was prepared every 4 days to
 380 guarantee food quality after culture in our feeding tanks and concentration was adapted to
 381 minimize clotting.

382 Animals were fed with a custom food mix composed of equal parts of two commercial
 383 blends of concentrated micro-algae (Shellfish Diet 1800, RotiGrow Plus), supplemented with
 384 live rotifers (cultured in-lab), enriched with 1 % (v/v) bakery yeast and 2 % (v/v) of planktonic
 385 supplement. To monitor the well-being of our animals, 4 slides representative of the same 4
 386 strains were imaged per tank once a week. We manually counted the number of zooids in each
 387 picture to estimate the growth rate of our animals and the impact of the different breeding
 388 conditions (**Fig. 3F**).

389 The favorable environmental culturing conditions presented here were identified thanks to
 390 multiple differential breeding experiments. Such approach could be used to test the impact of
 391 environmental conditions on specific aspects of the specimens' development, and to adapt our
 392 culturing conditions to different species for their breeding. Herein we show one example of
 393 such experiments. Animals were fed with four different food mixes: A) 1.6 mL/day ShellFish
 394 Diet 1800, 1.6 mL/day RotiGrow Plus, B) 1.1 mL/day ShellFish Diet 1800, 1.1 mL/day RotiGrow
 395 Plus, C) 3.2 mL/day RotiGrow Plus, D) 3.2 mL/day Shellfish Diet 1800. All food mixes were
 396 complemented with bakery yeast and planktonic substitute at the same proportion. Animals
 397 were bred under these conditions for four consecutive weeks while being monitored weekly. To

398 compensate for any possible bias in specific specimen tanks, we swapped the feeding diets
399 after 4 weeks, and monitored the same colonies for another 4 weeks.

400 Animals fed with food mix A and D outperformed those fed with B and C (Fig. 3G).
401 Moreover, animals fed with food mix A and D had positive growth rates (> 1), meaning that each
402 generation produces more animals than the previous one thus leading to exponential growth.
403 Although the differences were not significant in the first set of experiments, we measured the
404 same trends in both sets of feeding experiments. This confirms that the impact of tank specific
405 biases, if any, did not significantly influence our measurements. The low breeding measured
406 with mix B indicates less food than our baseline was not sufficient to sustain a positive growth
407 rate. This confirms that we are not overfeeding our animals. The low breeding measured with
408 mix C indicated that feeding on a blend of two type of algae mix was more beneficial as
409 previously published (Rinkevich and Shapira, 1998). However, the good performance of feeding
410 with D, also a single algae mix, suggests that this food source (Shellfish Diet 1800) is the
411 source of most growth in our species. Overall, this experiment exemplifies how differential
412 breeding can be used in our in-land aquaculture system to study how feeding influences the
413 growth of our colonial ascidians.

414 4. Discussion & Conclusions

415 We here present a hybrid flexible recirculating husbandry setup for the long-term in-lab
 416 culture of colonial ascidians. We developed this prototype to sustain our research on the
 417 exceptional capacity of *Botrylloides* to undergo whole-body regeneration (Blanchoud et al.,
 418 2018a). Yet, in this endeavor we also set out to establish a system that would be suitable for
 419 promoting as well as popularizing these colonial ascidians as model organisms more globally.
 420 Our achievements in constantly breeding *Botrylloides* over more than 3 years shows that this is
 421 indeed possible, and we do hope that it will initiate a renewed interest in studying these
 422 fascinating animals in novel places, including in non-coastal ones.

423 As pioneers in the in-land breeding of ascidians with limited experience in saltwater
 424 aquaculture, we came across some difficulties with the maintenance of the quality of our
 425 recirculating water. Accumulation of noxious ions is the main challenge for such a system.
 426 Nitrates in particular tend to accumulate quickly and on the long term because the anaerobic
 427 bacteria that can process this ion develop much more slowly than the others involved in the
 428 rest of the nitrogen cycle (Arrigo, 2005). Although our nitrate filter solved this particular issue
 429 relatively well, the most common solution to this accumulation remains to increase water
 430 changes (Zebcare, personal communications). Consequently, the main strategies available to
 431 maintain the quality of our ASW were to slowly vary the biological load of our system (i.e.
 432 animals and feeding), to use periodic treatments with removable media (e.g. phosphate binding
 433 resin), and to perform large water changes (50 % of the total water amount) when needed. To
 434 avoid sudden changes in water quality, we thus focused on reducing the variations in biological
 435 load on our system. Indeed, the three major regression events we experienced (**Fig. 3A-B**
 436 **shaded areas**) followed a sudden increase in the number of colonies that we naively combined

437 with a corresponding increase in feeding to try supporting these additional animals. We
 438 experienced a similar issue three times consecutively because we were initially not monitoring
 439 all the noxious ions in the system's water, and thus wrongly believed that we had solved the
 440 issue at the root of the decline. Instead, the accumulation was transferred to different noxious
 441 ions. The third and more drastic regression (**Fig. 3A-B blue area**) was unfortunately concurrent
 442 with a sanitary confinement that prevented us from tending properly to our aquaculture.
 443 Consequently, the latent issue lasted longer than the two previous ones. Nevertheless, colonies
 444 recovered particularly quickly and repopulated our culture setup very efficiently. Moreover, we
 445 resorted only twice over the 3 years to large water changes, confirming that the quality of our
 446 water is stable.

447 An important point in controlling the quality of our ASW was to understand that although
 448 a particular food source might be beneficial to the growth of our species, it might be
 449 imbalanced with respect to our target water parameter values. For instance, we observed that
 450 our food mix has a higher pH than our ideal pH 8.2. Although the important dilution that takes
 451 place during feeding renders this increase quite minimal, it similarly accumulated over time to
 452 yield to a noticeable increase in the whole system (**Fig. 2H**). Properly balancing the pH of the
 453 food mix before feeding it to the specimen tanks solved the problem at its root. Consequently,
 454 we advise to measure all the parameters of the food mix being delivered to the animals before
 455 starting a new feeding experiment. Feeding using live algae (Joly et al., 2007; Brown et al., 2009;
 456 Blanchoud et al., 2017) or using home-made blends of vegetable pastes (Kowarsky et al., 2021)
 457 might be more appropriate in the context of an in-lab monoculture as presented here and will
 458 be experimented in the future.

459 Colonies of *Botrylloides diegensis* have been cultured in our laboratory for more than 3
 460 years. We have established an in-lab all-year-round stable and fully monitored system which
 461 can be adapted to the needs of other tunicates to further popularize this subphylum. In parallel,
 462 we have introduced over the last year another colonial ascidian in our system, *Botryllus*
 463 *schlosseri*. While this species has so far well adapted to our recirculating ASW aquaculture, it
 464 grows much slower than *B. diegensis*. Yet, colonies look healthy and they do propagate. One
 465 potential explanation for this behavior is that *Botryllus schlosseri* colonies thrive at
 466 temperatures lower than 20 °C (Boyd et al., 1986), above which they tend to become gravid and
 467 slow down their asexual reproduction. Our system allows to set different water temperatures in
 468 different specimen tanks and optimizing *Botryllus* culture will be attempted in this direction.

469 While gravid colonies confirm the good status of our colonies, the observed sexual
 470 reproduction occurred without apparent relation to any specific environmental condition. In
 471 addition, the open configuration of our specimen tanks together with our important flow-
 472 through is not adapted to handling spawning of colonial tunicates. Consequently, we will
 473 explore adapting our in-land aquaculture with published methods used in other species of
 474 colonial ascidians (Fletcher and Forrest, 2011; Joly et al., 2007), both for the induction of
 475 spawning as well as for the retrieval and settling of the motile larvae.

476 A surprising result from our differential breeding experiment is that only one set of data
 477 shows statistically significant effects, while the same trends can be observed in the other one.
 478 One potential explanation is that some remnant effect of the previous set of feeding
 479 experiment was influencing the monitored colonies. Indeed, because *Botrylloides* have three
 480 generations co-existing constantly within the colony (Berrill, 1947), the number of adults
 481 observed in the following generation was determined based on the breeding conditions that

were in place when this new generation budded, which means around two weeks in our 21 °C system. Consequently, differential breeding experiments might need to be monitored over periods that include 3 additional weeks for buffering between experiments. Alternatively, incorporating only the data from the last week of the experiment could remove this issue, but more colonies would need to be monitored to yield statistically significant results.

Overall, we show that *Botrylloides diegensis* can be proficiently bred in-land and our preliminary results with *Botryllus schlosseri* suggest that our system for controlled breeding can provide the necessary platform for the worldwide spreading of colonial ascidians.

490

491 5. Acknowledgments

We would like to thank Dr. Megan Wilson for sharing her experience with breeding *Botrylloides*, Pr. Miles Lamar for providing space for our trials when setting up a flow-through system at the University of Otago, Pr. Lorian Ballarin for his extensive experience on breeding *Botryllus* in closed systems, Dr. Fabio Gasparini for helping us with the collection of our wild isolates, all the members of these laboratories for useful discussions on the breeding of colonial ascidians as well as Laura Bugada, Mohamed Abshir Mohamed, Margaux de Raemy, Lionel Page, Jeanne Bruehlhart and Silvia Moreno Forero for their help in taking care of the husbandry setup.

We thank the people from our institute's workshops for helping us, teaching us and showing us how to modify, improve and develop an aquaculture system. In particular, Ruben Pooley, Alain Werro, Jean-Daniel Niederhauser, Bertrand Dubois and François Zosso.

503 We also thank Dr. Megan Wilson and Dr. Arzu Karahan for kindly providing the pictures of
504 their respective husbandry system displayed in Fig. 1, and Aude Blanchoud for comments on
505 the manuscript.

506

507 6. Author contributions

508 Simon Blanchoud: Conceptualization, Funding acquisition, Project administration
509 Visualizaton. Marta Wawrzyniak: Writing-original draft. Marta Wawrzyniak, Lluís Matas Serrato,
510 Simon Blanchoud: Investigation, Methodology, Validation, Writing-review & editing.

511

512 7. Funding

513 This work was funded by the Swiss National Science Foundation (SNF) [grant number
514 PZ00P3_173981]. The SNF had no involvement in the conduct and preparation of this work.

515

516 8. Declaration of Interests

517 The authors have declared no conflict of interest.

518 9. References

- 519 Aldred, N., Clare, A.S., 2014. Mini-review: Impact and dynamics of surface fouling by solitary and
520 compound ascidians. *Biofouling* 30, 259–270.
521 <https://doi.org/10.1080/08927014.2013.866653>
- 522 Arrigo, K.R., 2005. Marine microorganisms and global nutrient cycles. *Nature* 437, 349–355.
523 <https://doi.org/10.1038/nature04159>
- 524 Berrill, N.J., 1947. The developmental cycle of *Botrylloides*. *Q J Microsc Sci* 88, 393–407.
- 525 Blanchoud, S., Rinkevich, B., Wilson, M.J., 2018a. Whole-Body Regeneration in the Colonial
526 Tunicate *Botrylloides leachii*. *Results Probl Cell Differ* 65, 337–355.
527 https://doi.org/10.1007/978-3-319-92486-1_16
- 528 Blanchoud, S., Rutherford, K., Zondag, L., Gemmell, N.J., Wilson, M.J., 2018b. *De novo* draft
529 assembly of the *Botrylloides leachii* genome provides further insight into tunicate
530 evolution. *Sci Rep* 8, 5518. <https://doi.org/10.1038/s41598-018-23749-w>
- 531 Blanchoud, S., Wawrzyniak, M., 2021. Cleaning colonial ascidians. *Protocols.io*.
532 <https://doi.org/10.17504/protocols.io.brxiem7je>
- 533 Blanchoud, S., Zondag, L., Lamare, M.D., Wilson, M.J., 2017. Hematological Analysis of the
534 Ascidian *Botrylloides leachii* (Savigny, 1816) During Whole-Body Regeneration. *The*
535 *Biological Bulletin* 232, 143–157. <https://doi.org/10.1086/692841>
- 536 Boyd, H.C., Brown, S.K., Harp, J.A., Weissman, I.L., 1986. Growth and sexual maturation of
537 laboratory-cultured monterey *Botryllus schlosseri*. *The Biological Bulletin* 170, 91–109.
538 <https://doi.org/10.2307/1541383>
- 539 Brown, F.D., Keeling, E.L., Le, A.D., Swalla, B.J., 2009. Whole body regeneration in a colonial
540 ascidian, *Botrylloides violaceus*. *J. Exp. Zool.* 312B, 885–900.
541 <https://doi.org/10.1002/jez.b.21303>
- 542 Bugada, L., Blanchoud, S., 2021. Subcloning colonial ascidians. *Protocols.io*.
543 <https://doi.org/10.17504/protocols.io.brw9m7h6>
- 544 Chen, L., Wang, X.-N., Fu, C.-M., Wang, G.-Y., 2019. Phylogenetic Analysis and Screening of
545 Antimicrobial and Antiproliferative Activities of Culturable Bacteria Associated with the
546 Ascidian *Styela clava* from the Yellow Sea, China. *BioMed research international* 2019,
547 7851251. <https://doi.org/10.1155/2019/7851251>
- 548 Coma, R., Ribes, M., Gili, J.-M., Zabala, M., 2000. Seasonality in coastal benthic ecosystems.
549 *Trends in Ecology & Evolution* 15, 448–453. [https://doi.org/10.1016/S0169-5347\(00\)01970-4](https://doi.org/10.1016/S0169-5347(00)01970-4)
- 551 Comeau, L.A., Filgueira, R., Guyondet, T., Sonier, R., 2015. The impact of invasive tunicates on
552 the demand for phytoplankton in longline mussel farms. *Aquaculture* 441, 95–105.
553 <https://doi.org/10.1016/j.aquaculture.2015.02.018>

554 Delsuc, F., Brinkmann, H., Chourrout, D., Philippe, H., 2006. Tunicates and not cephalochordates
555 are the closest living relatives of vertebrates. *Nature* 439, 965–8.
556 <https://doi.org/10.1038/nature04336>

557 Epelbaum, A., 2009. Botryllid tunicates: Culture techniques and experimental procedures. *AI* 4,
558 111–120. <https://doi.org/10.3391/ai.2009.4.1.12>

559 Fletcher, L., Forrest, B., 2011. Induced spawning and culture techniques for the invasive
560 ascidian *Didemnum vexillum* (Kott, 2002). *AI* 6, 457–464.
561 <https://doi.org/10.3391/ai.2011.6.4.11>

562 Hiebert, L.S., Vieira, E.A., Dias, G.M., Tiozzo, S., Brown, F.D., 2019. Colonial ascidians strongly
563 preyed upon, yet dominate the substrate in a subtropical fouling community. *Proc. R.*
564 *Soc. B.* 286, 20190396. <https://doi.org/10.1098/rspb.2019.0396>

565 Hirose, E., Saito, Y., Watanabe, H., 1995. Regeneration of the tunic cuticle in the compound
566 ascidian, *Botrylloides simodensis*. *Developmental & Comparative Immunology* 19, 143–
567 151. [https://doi.org/10.1016/0145-305X\(94\)00062-K](https://doi.org/10.1016/0145-305X(94)00062-K)

568 Holland, L.Z., 2016. Tunicates. *Current Biology* 26, R146–R152.
569 <https://doi.org/10.1016/j.cub.2015.12.024>

570 Joly, J.-S., Kano, S., Matsuoka, T., Auger, H., Hirayama, K., Satoh, N., Awazu, S., Legendre, L.,
571 Sasakura, Y., 2007. Culture of *Ciona intestinalis* in closed systems. *Dev Dyn* 236, 1832–
572 1840. <https://doi.org/10.1002/dvdy.21124>

573 Kassmer, S.H., Langenbacher, A.D., De Tomaso, A.W., 2020. Integrin- α -6+ Candidate stem
574 cells are responsible for whole body regeneration in the invertebrate chordate
575 *Botrylloides diegensis*. *Nat Commun* 11, 4435. <https://doi.org/10.1038/s41467-020-18288-w>

577 Kowarsky, M., Anselmi, C., Hotta, K., Burighel, P., Zaniolo, G., Caicci, F., Rosental, B., Neff, N.F.,
578 Ishizuka, K.J., Palmeri, K.J., Okamoto, J., Gordon, T., Weissman, I.L., Quake, S.R., Manni,
579 L., Voskoboynik, A., 2021. Sexual and asexual development: two distinct programs
580 producing the same tunicate. *Cell Reports* 34, 108681.
581 <https://doi.org/10.1016/j.celrep.2020.108681>

582 Lambert, G., 2005. Ecology and natural history of the protochordates. *Canadian Journal of*
583 *Zoology* 83, 34–50. <https://doi.org/10.1139/Z04-156>

584 Nagar, L., Shenkar, N., 2016. Temperature and salinity sensitivity of the invasive ascidian
585 *Microcosmus exasperatus* Heller, 1878. *AI* 11, 33–43.
586 <https://doi.org/10.3391/ai.2016.11.1.04>

587 Navon, G., Kaplan, A., Avisar, D., Shenkar, N., 2020. Assessing pharmaceutical contamination
588 along the Mediterranean and Red Sea coasts of Israel: Ascidians (Chordata, Ascidiacea)
589 as bioindicators. *Marine pollution bulletin* 160, 111510.
590 <https://doi.org/10.1016/j.marpolbul.2020.111510>

591 Palanisamy, S.K., Rajendran, N.M., Marino, A., 2017. Natural Products Diversity of Marine
592 Ascidiaceae (Tunicates; Ascidiaceae) and Successful Drugs in Clinical Development. Nat.
593 Prod. Bioprospect. 7, 1–111. <https://doi.org/10.1007/s13659-016-0115-5>

594 Rinkevich, B., Shapira, M., 1998. An improved diet for inland broodstock and the establishment
595 of an inbred line from *Botryllus schlosseri*, a colonial sea squirt (Ascidiaceae). Aquatic
596 Living Resources 11, 163–171. [https://doi.org/10.1016/S0990-7440\(98\)80113-7](https://doi.org/10.1016/S0990-7440(98)80113-7)

597 Rinkevich, Y., Rosner, A., Rabinowitz, C., Lapidot, Z., Moiseeva, E., Rinkevich, B., 2010. *Piwi*
598 positive cells that line the vasculature epithelium, underlie whole body regeneration in a
599 basal chordate. Developmental Biology 345, 94–104.
600 <https://doi.org/10.1016/j.ydbio.2010.05.500>

601 Roberts, B., Davidson, B., MacMaster, G., Lockhart, V., Ma, E., Wallace, S.S., Swalla, B.J., 2007. A
602 complement response may activate metamorphosis in the ascidian *Boltenia villosa*. Dev
603 Genes Evol 217, 449–458. <https://doi.org/10.1007/s00427-007-0157-0>

604 Scelzo, M., Alié, A., Pagnotta, S., Lejeune, C., Henry, P., Gilletta, L., Hiebert, L.S., Mastrototaro, F.,
605 Tiozzo, S., 2019. Novel budding mode in *Polyandrocarpa zorritensis*: a model for
606 comparative studies on asexual development and whole body regeneration. EvoDevo 10,
607 7. <https://doi.org/10.1186/s13227-019-0121-x>

608 Sheets, E.A., Cohen, C.S., Ruiz, G.M., da Rocha, R.M., 2016. Investigating the widespread
609 introduction of a tropical marine fouling species. Ecol Evol 6, 2453–2471.
610 <https://doi.org/10.1002/ece3.2065>

611 Shenkar, N., Swalla, B.J., 2011. Global Diversity of Ascidiaceae. PLoS ONE 6, e20657.
612 <https://doi.org/10.1371/journal.pone.0020657>

613 Taketa, D.A., Nydam, M.L., Langenbacher, A.D., Rodriguez, D., Sanders, E., De Tomaso, A.W.,
614 2015. Molecular evolution and in vitro characterization of *Botryllus* histocompatibility
615 factor. Immunogenetics 67, 605–623. <https://doi.org/10.1007/s00251-015-0870-1>

616 Tzafriri-Milo, R., Benaltabet, T., Torfstein, A., Shenkar, N., 2019. The Potential Use of Invasive
617 Ascidiaceae for Biomonitoring Heavy Metal Pollution. Front. Mar. Sci. 6, 611.
618 <https://doi.org/10.3389/fmars.2019.00611>

619 Ueki, T., Koike, K., Fukuba, I., Yamaguchi, N., 2018. Structural and Mass Spectrometric Imaging
620 Analyses of Adhered Tunic and Adhesive Projections of Solitary Ascidiaceae. Zoological
621 Science 35, 535–547. <https://doi.org/10.2108/zs180051>

622 Vered, G., Kaplan, A., Avisar, D., Shenkar, N., 2019. Using solitary ascidiaceae to assess
623 microplastic and phthalate plasticizers pollution among marine biota: A case study of
624 the Eastern Mediterranean and Red Sea. Marine pollution bulletin 138, 618–625. <https://doi.org/10.1016/j.marpolbul.2018.12.013>

626 WoRMS Editorial Board, 2017. World Register of Marine Species. Available from
627 <http://www.marinespecies.org> at VLIZ. Accessed 2020-09-29.
628 <https://doi.org/10.14284/170>

629 Zondag, L.E., Rutherford, K., Gemmell, N.J., Wilson, M.J., 2016. Uncovering the pathways
630 underlying whole body regeneration in a chordate model, *Botrylloides leachi* using *de*
631 *novo* transcriptome analysis. BMC Genomics 17, 114. [https://doi.org/10.1186/s12864-](https://doi.org/10.1186/s12864-016-2435-6)
632 016-2435-6
633

634 **10. Figure legends (all figures should be printed in color)**

635

636 **Figure 1. Current status of colonial ascidian husbandry.** (A) A typical colony of *Botrylloides*
637 *diegensis* settled on a glass slide. (B) The sampling site where colonies were collected
638 (Chioggia, IT). (C) Example of a flow-through system (Wilson lab, University of Otago, NZ). (D)
639 Example of a closed system (Karahan lab, METU, Turkey).

640

641 **Figure 2. The colonial ascidian husbandry setup.** (A) Schematic blueprint of our system. (B)
642 Picture of the developed prototype. The main system is composed of a shelf of tanks for food
643 flowing into the specimens' tanks recycling through the sump (C). View inside the sump where
644 water is filtered and rectified automatically. (D) Vertical culture of colonial ascidians – each
645 specimens' tank is equipped with racks of hooks. (E) Colony hanger used to hold the glass
646 slides vertically. (F) Two quarantine containers. (G) Staining racks are used in our quarantine
647 tanks. (H) Parameters of the water quality measured in our system between March 2018 and
648 March 2021. From left to right: ORP – oxidation reduction potential (mV); salinity (g/l); water
649 temperature in the sump and tanks 1 – 4 (°C); pH in the sump and measured with a kit; levels of
650 noxious substances: NH_4^+ , NO_2^- , NO_3^- and PO_4^{3-} (mg/l).

651

652 **Figure 3. Breeding of colonial ascidians.** (A) Lineage of a successful strain (S131) showing its
653 propagation by subcloning (from February 2019 till March 2021). Major recession events are
654 pointed at by shaded areas (grey and blue), the one concurrent with a sanitary confinement is
655 highlighted in blue. (B) Evolution of the number of clones bred in our system shown from March
656 2019 till February 2021, color-coded by strain ID. Major recession events are depicted as in A.

657 (C) Picture of a relatively large colony composed of over 300 zooids, which became gravid in
 658 2019. (D) Picture of a gravid bud with its oocyte visible (arrowhead). (E) Picture of a young
 659 oozoid recently metamorphosed. Visible organs including its oral siphon (s), ampullae (a) and
 660 bud (b). (F) Sample pictures taken on the specified days during our differential feeding
 661 experiment. Depicted pictures correspond to the same colony fed with the combination Mix A /
 662 Mix B. (G) Relative mean number of zooids counted per colony (n=3) averaged over the
 663 indicated number of weeks. $P^* < 0.05$; $** < 0.01$; $*** < 0.001$.

664

665 **Table 1. Components use in our husbandry system for colonial ascidians.** Most components
 666 are visible in **Supplementary Figures S1-S4**. Note that the given prices have been gathered from
 667 a variety of different providers and currencies and will likely vary. Transportation, taxes and
 668 assembly costs were not taken into account. a) these components were custom-build and their
 669 costs are thus estimates only. b) these components are consumables and the given price
 670 corresponds to what is required for assembling one complete setup.

671

672 **Movie 1. In-land breeding of *B. diegensis*.** Time-lapse imaging of a colony developing over a
 673 period of 25 days.

674

675 **Supplementary Figure S1. Detailed structure of the sump unit.** (A) Moving-bed filter, (B)
 676 refugium and (C) main sedimentation tank composing the sump of our husbandry system. (D)
 677 Fleece filter separated from its underlying moving-bed unit shown in A. (E) Components of the
 678 refugium filter and of the ceramic beads filtration unit visible in B. (F) Area between the sump
 679 and the chiller where the main circulation pump, the nitrate filter and the leak detection probe

are located. The main components visible in the panels are identified using their ID as listed in **Table 1**.

Supplementary Figure S2. Detailed structure of the feeding unit. (A) Feeding tank connected to the delivery valve for dispensing food in the specimen tank located underneath. Each feeding tank is composed of (B) a 10 L cylindrical container, (C) a micro-porous air diffuser and (D) a lid with a clear compartment for hosting the lighting bars. Components ID are specified in **Table 1**.

Supplementary Figure S3. Detailed structure of the controlling units. (A) Main control panel with the Apex controller and its various modules. (B) Behind the husbandry system where the solenoid valve controls the input of deionized water for topping-up the sump. (C) Components of the custom micro-controller unit for continuous feeding of our colonies. (D) Lights and feeding control units are installed behind the feeding tanks. Components ID are specified in **Table 1**.

Supplementary Figure S4. Detailed structure of the air and salts units. (A) The main air pump is isolated from the husbandry system and connects to a custom manifold. The pH unit is located on the left side of the system. (B) The manual control valve for the pH buffer is accessible from behind the main system. (C) The brine tank is located on the right side of the system and is pumped automatically by the two peristaltic pumps visible in A. Components ID are specified in **Table 1**.

702 **Supplementary File 1. Design and control of our custom micro-controller for continuous**
 703 **feeding.** This archive contains all the documents required to reproduce our micro-controller.
 704 The file used to design and produce the custom PCB board (controller_design.fzz) using
 705 Fritzing, a dedicated Open Source software. A schematic representation of that PCB board
 706 (controller_design_scheme.pdf) and two vectorized visualizations of the PCB connections, one
 707 for each side of the board (controller_design_pcb_top.pdf, controller_design_pcb_bottom.pdf). A
 708 list of the electronic components required to assemble the PCB board
 709 (controller_components.txt) and the MicroPython code used to control the valves
 710 (controller_code.py).

Figure 1

Figure 2

bioRxiv preprint doi: <https://doi.org/10.1101/2021.01.31.429038>; this version posted March 12, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC 4.0 International license.

Figure 3

bioRxiv preprint doi: <https://doi.org/10.1101/2021.01.31.429038>; this version posted March 12, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC 4.0 International license.

ID	Component	Manufacturer	Num.	Total (€)	
1 Specimen compartments					
1.01	Stainless frame	Fleuren&Nooijen	1	1000	a
1.02	Specimen tanks 40 L	Fleuren&Nooijen	4	1200	a
1.03	Flow regulation valve (VM DN 8)	FIP	4	112	
1.04	Heaters (titanium D-D)	Schego	4	196	
1.05	Wavemaker pump (Jecod SW-2)	Jebao	4	240	
1.06	Light AquaBeam 1500 Ocean blue	TMC Aquarium	8	800	
1.07	Light BioLumen Power Pod 2	TMC Aquarium	4	392	
1.08	Stainless support bar with stoppers	Fleuren&Nooijen	8	80	a
1.09	Plastic tubing for colony racks	Fleuren&Nooijen	16	8	a
1.10	Plastic J-hooks for hanging colonies	Sinopos	128	5	
1.11	Silicone badge gripper	Specialist ID	128	141	
1.12	Various cable ties	Netztech Handels	200	6	
1.13	Filter wool	Finest aquatics	2 m/w	6	b
2 Quarantine compartments					
2.01	Heavy-duty plastic stand	Cambro	1	700	a
2.02	Quarantine container 45 L (Gripline)	Semadeni AG	2	18	
2.03	Plastic staining rack (1233)	Semadeni AG	10	54	
2.04	Wavemaker pump (Jecod SW-2)	Jebao	2	120	
2.05	360 mm marine hybrid LED (powerLED+)	Eheim	2	218	
2.06	Wireless LED controller (control+)	Eheim	1	122	
2.07	LED power supply	Eheim	1	33	
3 Feeding compartments					
3.01	Feeding tanks 10 L	Fleuren&Nooijen	4	800	a
3.02	Light Aquabar T-series Colorplus 438mm	TMC Aquarium	8	624	
3.03	Aquabar 2 channel controller	TMC Aquarium	1	58	
3.04	4.5 mm barbed micro-valve (AC42155)	Antelco	4	4	
3.05	4 mm inner diameter silicone tubing	Cortailod	20	32	
3.06	Micro-porous tubing	Gardena	4	10	

bioRxiv preprint doi: <https://doi.org/10.1101/2021.01.31.429038>; this version posted March 12, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC 4.0 International license.

3.07	Polyamide cable gland	Wiska	8	6	
3.08	4.5 mm joiners barbed	Artec	10	2	
3.09	500 g waterproof weight	Tunturi	4	27	
4	Water compartment				
4.01	Sump tank 200 L	Fleuren&Nooijen	1	600	a
4.02	Fountain pump (universal 300)	Eheim	1	55	
4.03	Digital timer IP20	Steffen AG	1	17	
4.04	Circulation pump (Titan-Tec 10000)	Messner	1	700	
4.05	UV steriliser (Vecton2 600)	TMC Aquarium	1	78	
4.06	Bi-stop ball valve	Plasson	2	74	
4.07	Cooling unit Titan 4000	Aqua Medic	1	3400	
4.08	Heaters (titanium D-D)	Schego	1	49	
4.09	Protein skimmer (DC runner 1.1)	Aqua Medic	1	205	
4.10	Filter wool	Finest aquatics	0.2 m/w	1	b
4.11	Moving bed (K+media)	Evolution aqua	10 L	19	
4.12	Cermanic beads	TetraTec	3	34	
4.13	Bacteria (Bio reefclear)	Sera	100 mL	11	
4.14	Algae scrubber	IceCap	1	240	
4.15	Denitrifyer	Korallin	1	250	
4.16	25/30/40 mm PVC tubing	Van de Lande	20	80	
4.17	Various PVC connectors and tees	Van de Lande	25	75	
4.18	19 mm tubing	Angst + Pfister AG	2	16	
4.19	4-channels water distributor (8194-20)	Gardena	1	36	
4.20	Various hose connectors and fittings	Gardena	15	75	
4.21	Water trigger nozzle (18301-20)	Gardena	2	56	
5	Refugium compartment				
5.01	Light Aquabar T-series Colorplus 438mm	TMC Aquarium	2	156	
5.02	Aquabar Dimmer Switch	TMC Aquarium	1	24	
5.03	Meshed filter box	Semadeni	1	8	
5.04	Live rock with Coralline algae	Aqua Home	8 kg	182	
5.05	Macroalgae (Chaetomorpha sp.)	Aqua Home	1	9	

6 Air compartment

bioRxiv preprint doi: <https://doi.org/10.1101/2021.01.31.429038>; this version posted March 12, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC 4.0 International license.

6.01	Air blower (Medo LA-60B)	Angst + Pfister AG	3 m	24
6.02	18 mm tubing	Angst + Pfister AG	3 m	24
6.03	18 mm to 4 mm air manifold	Fleuren&Nooijen	1	100
6.04	4 mm inner diameter tubing	Cortailod	20 m	33
6.05	Air stones	Sera	3	3

7 Salt compartments

7.01	Mortar mixer	Einhell AG	1	80	
7.02	Digital timer IP20	Steffen AG	1	17	
7.03	Salt tank 75 L	Fleuren&Nooijen	1	180	
7.04	Seawater salt (SYN-BIOTIC)	Tropic Marin	260 g/L	57	b
7.05	pH tank 20 L	Fleuren&Nooijen	1	100	
7.06	Sodium bicarbonate	Fisher Chemicals	16 g/L	3	b
7.07	4.5 mm barbed micro-valve (AC42155)	Antelco	1	1	

8 Controller systems

8.01	Apex 2016 Lab Grade Kit	Neptune Systems	1	680
8.02	Magnetic holder	Neptune Systems	1	41
8.03	Automatic top-off kit	Neptune Systems	1	200
8.04	Solenoid valve	Neptune Systems	1	31
8.05	Optical sensor	Neptune Systems	1	36
8.06	Leak detection probe	Neptune Systems	1	30
8.07	Temperature probe	Neptune Systems	4	112
8.08	pH/ORP/Redox module	Neptune Systems	2	194
8.09	Salinity module	Neptune Systems	2	194
8.10	Dosing system	Neptune Systems	1	311
8.11	Conductivity probe	Neptune Systems	1	132
8.12	Microcontroller (Pico RP2040)	Raspberry Pi	1	4
8.13	Custom valve controller	Aisler	1	50
8.14	12V normal closed liquid valve (CJV23)	Xiamen Conjoin	4	8
8.15	Rocker switch (16Q7522)	Hornbach	1	3
8.16	Biolumen control unit	TMC Aquarium	1	413

Table 1. Components use in our husbandry system for colonial ascidians. Most components are visible in **Supplementary Figures S1-S4**. Note that the given prices have been gathered from a variety of different providers and currencies and will likely vary. Transportation, taxes and assembly costs were not taken into account. **a)** these components were custom-build and their costs are thus estimates only. **b)** these components are consumables and the given price corresponds to what is required for one complete setup.

95

55

5 mm

12:35 - 21/08/2020