Rediscovering the musician's brain: a systematic review and meta-analysis

Criscuolo, Antonio^{1,2+}, Pando-Naude, Victor^{2+*}, Bonetti, Leonardo², Vuust, Peter² & Brattico, Elvira²

¹Department of Neuropsychology & Psychopharmacology, Faculty of Psychology and Neuroscience, Maastricht University

²Center for Music in the Brain, Department of Clinical Medicine, Aarhus University & The Royal Academy of Music Aarhus/Aalborg, Denmark

+Shared first-authorship

***Corresponding author:**

Victor Pando-Naude, M.D., M.Sc., PhD Fellow

Center for Music in the Brain (MIB)

Department of Clinical Medicine, Aarhus University

Universitetsbyen, 3-0-17

8000 Aarhus C, Denmark

Email: pandonaude@clin.au.dk

Phone: +4552805306

Abstract.

Musical expertise is a model of neuroplasticity associated with pervasive, long-lasting training effects. Indeed, decades of cognitive neuroscience widely investigated brain functional and structural changes associated with musical training, providing a widespread and variegated set of findings. However, several controversial results emerged, leading the neuroscientific community to lack a well-defined neuro-functional-anatomy of musical expertise. Here, we performed a systematic review and meta-analysis of publications investigating brain functional and structural differences between musicians and non-musicians. Eighty-four publications were included in the qualitative synthesis. Coordinate-based meta-analyses were conducted using the anatomic/activation likelihood estimation (ALE) method implemented in GingerALE, with a total of 675 foci, 79 experiments and 2780 participants. Results showed a widespread and complex array of functional and structural changes in musicians' brains, revealing for the first time a comprehensive picture of the brain plasticity associated with musical training.

1. Introduction.

A key to survival is the ability to adapt to the environment as biologically reflected in the neuroplastic changes associated with experience¹. Thus, understanding the mechanisms behind neuroplasticity has always represented a primary and fascinating challenge in neuroscience². At a circuit level, studies employing animal models showed how small neural networks are shaped by experience and learning, highlighting the role of key brain mechanisms such as long-term potentiation (LTP) and long-term depression (LTD)³.

Yet, the whole-brain neuroplasticity associated with the complexity of human life has remained more elusive and difficult to investigate. Studying the musician's brain has been suggested as an ideal model for tackling this complex challenge⁴, allowing for both cross-sectional and longitudinal investigations^{5,6}.

Indeed, a long-lasting musical training promotes highly-specialized perceptual, motor, emotional and higher-order cognitive abilities⁷, as well as the capacity of integrating multimodal auditory, visual and motor information⁸. Musical training also requires the development of fine-grained skills in auditory perception, kinesthetic control, visual perception and pattern recognition⁹. Further, it stimulates mnemonic processes, fine-grained auditory perception (absolute pitch)¹⁰ and manual coordination. All these specialized abilities acquired through an extensive professional training have a neural counterpart, represented by a wide array of brain structural and functional neuroplastic changes associated with musical expertise.

In cognitive neuroscience, structural neuroplasticity is usually quantified in terms of decreased or increased volumes of gray and white matter (GM and WM, respectively) as measured by magnetic resonance imaging (MRI). GM volumes are typically derived from cortical thickness (CT) and voxel-based morphometry (VBM) analyses, and results are interpreted as differences in GM concentration¹¹. WM volumes can be measured with VBM analysis as well, although the study of fractional anisotropy (FA) values obtained by diffusion tensor imaging (DTI) represents a valid and more informative alternative for investigating WM¹². Differently, functional neuroplasticity can be examined in task-related studies by using metabolic measures of oxygen consumption as recorded by functional (f) MRI. Hence, the time-series of blood-oxygen-level-dependent (BOLD) signals are usually correlated to the tasks performed during image acquisition to infer region-specific or network-level activity patterns¹³.

Employing these techniques, decades of studies on musical expertise have provided a plethora of variegated findings which, while abundant, are scattered and often controversial. For instance, while some works reported an overall increase of GM volume when comparing musicians' to non-musicians' brains^{14,15}, other research mainly highlighted negative correlations between GM volumes and musical expertise¹⁶. Similarly, fMRI studies reported a large and sometimes non-consistent range of brain areas that were differently active during tasks when comparing musicians vs non-musicians¹⁷. Moreover, previous literature has mainly relied on cross-sectional comparisons rather than longitudinal designs, resulting in reduced consistency across studies and showing a lack of a well-defined neuro-functional-anatomy of musical expertise.

Strikingly, although the musician's brain has even been indicated as a "model of neuroplasticity"⁴, the neuroscientific community lacks meta-analytic efforts of summarizing the findings on brain differences between music experts and laypersons. Thus, for the first time, we conducted a coordinate-based meta-analysis of a large pool of MRI studies using state-of-the-art methods such as anatomic/activation likelihood estimation (ALE)¹⁸, to investigate the neuroanatomical and neurofunctional changes associated with musical training in healthy humans.

2. Results.

A total of 1169 records was identified through database searching, and after removing duplicates, 679 records were initially screened by title and abstract. A total of 145 articles was assessed for eligibility in the full-text screening stage. From these, 84 studies fulfilled criteria for eligibility and were thus included in the qualitative synthesis. Finally, 58 studies reported results in stereotactic coordinates (either Talairach or Montreal Neurological Institute (MNI) three-dimensional-coordinate system) and were therefore included in the quantitative synthesis (ALE meta-analyses) (**Supplementary Figure 1**).

2.1. Study characteristics.

The characteristics of all studies included in our work are shown in **Table 1**. Eighty-four studies met inclusion criteria and were included in the qualitative synthesis. The total number of participants was 3048, with 1581 musicians (M), and 1424 non-musicians (NM). Eighteen studies (21%) included amateur musicians, and only 7 studies (8.3%) reported absolute pitch possessors ($n = 97$). Musical instruments were reported in the majority of the studies (81%): piano or keyboard (62%), string instruments (41%), wind instruments (26%), percussion instruments (17%), voice (8%), and 19% studies failed to report musicians' instrument. Years of education was described only in 8% of the included studies. Years of musical training was reported in 63% of the studies, with a mean of 15.6 ± 5.9 years. The age of onset of musical training was reported in 49% of the studies, with a mean of 7.4 ± 2.3 years old. Weekly hours of training were reported in 32% of the studies, with a mean of 16.7 ± 8.9 hours per week.

Neuroimaging data was acquired in either 1.5 T (39%), or 3 T (56%) scanners, while 5% of studies did not report the magnetic field strength. MRI scanners included Siemens (40%), General Electric (25%), Philips (21%), Bruker (5%), while 7% did not report it. Analysis methods included fMRI (52%), VBM (29%), DTI (18%), and CT (10%). Finally, 84% of studies included GM analyses, while 23% included WM analyses (**Supplementary Tables 1 and 2**).

Insert Table 1.

2.2. MRI quality.

MRI quality of the included studies in the meta-analysis was assessed following a set of guidelines for the standardized reporting of MRI studies (**Supplementary Table 3**). All studies reported their MRI design, software package and image acquisition, processing and analyses. Overall, good MRI practices were performed in the included studies.

2.3. Primary outcome: ALE meta-analysis.

The first ALE meta-analysis included all foci extracted from the screened studies (both functional and anatomical; 675 foci, 79 experiments, and 2,780 participants). The analysis returned significant clusters peaking in the following areas (alphabetical order): anterior cingulate cortex, caudate, claustrum, inferior frontal gyrus, insula, inferior parietal lobule, middle frontal gyrus, middle temporal gyrus, parahippocampal gyrus, postcentral gyrus, precentral gyrus, putamen, supramarginal gyrus, superior parietal lobule and superior temporal gyrus. The comparison $M > NM$ (higher volume/activity in musicians) resulted in significant peak clusters of both left and right superior temporal gyri, extending to middle temporal gyri, middle and inferior frontal gyri, insula and pre-post central gyri. The comparison $NM > M$ (lower volume/activity in musicians) resulted in significant peak clusters located in the superior and inferior parietal lobules, pre-post central gyri, caudate, and cerebellum (**Figure 1, Table 2**).

The comparison M vs NM was further analyzed for structural and functional foci, independently. Structurally, the comparison M > NM in gray matter (GM) resulted in significant peak clusters of the superior temporal gyrus (BA22), including the Heschl's gyrus and planum temporale, postcentral gyrus, supramarginal gyrus, insula, hippocampus, thalamus, cerebellum and brainstem; and white matter (WM) tracts of the internal capsule bundle, bilaterally, corona radiata, corpus callosum, optic radiation and anterior thalamic radiations. Conversely, the comparison NM > M in GM resulted in significant peak clusters of the precentral gyrus, posterior cingulate cortex and supramarginal gyrus. No significant results were found in this comparison for WM. Functionally, M > NM resulted in significant peak clusters of the inferior frontal gyrus, superior temporal gyrus, and middle frontal gyrus. The comparison NM > N resulted in significant peak clusters of the inferior parietal lobule, precentral gyrus, superior frontal gyrus, superior parietal lobule and cerebellum (**Figure 1, Supplementary Table 4**).

Insert Figure 1.

Insert Table 2.

2.4. Contrast analysis.

A contrast ALE meta-analysis was conducted to compare foci from structural (e.g., VBM) and functional (e.g., fMRI) studies, by means of conjunction (similarity) and subtraction (difference). The purpose of this analysis was to identify common and distinct music-related brain areas between the neuroimaging modalities, thus, suggesting a more objective relationship between a function, and structural neuroadaptations produced by the repetition of that function. The superior temporal gyrus, precentral gyrus and insula are present in both modalities. The postcentral gyrus (BA3), supramarginal gyrus and transverse temporal gyrus are more present in structural imaging, while the inferior frontal gyrus, postcentral gyrus (BA4), middle frontal gyrus, middle temporal gyrus, and parietal lobules are more present in functional imaging (**Figure 2, Supplementary Table 5**).

Insert Figure 2.

2.5. Subgroup analysis.

A series of *post hoc* subgroup analyses were conducted to identify brain regions related to specific musical and cognitive features explored by the previous studies. Such features included improvisation, language, melody, memory, sensorimotor, pitch, rhythm and timbre (**Figure 3, Supplementary Table 6**).

The improvisation subgroup resulted in significant clusters of the right superior temporal gyrus extending to the insula, and precentral gyrus, bilaterally. Such results suggest the engagement of primary auditory and motor areas, presumably modulated by limbic processes. Motor-related foci converged in the precentral gyrus, inferior parietal lobule and medial frontal gyrus. Notably, motor, premotor and supplementary motor areas were expected. The language subgroup resulted in significant clusters in supramarginal gyrus, inferior parietal lobule, anterior cingulate cortex and middle frontal areas. Such areas suggest the activity of a network involved in language processing, and executive processing. Memory-related subgroup resulted in significant clusters in pre-postcentral gyri, middle, medial and superior frontal gyri. Interestingly, no areas from the parahippocampal gyrus were significant. The melody subgroup resulted in significant clusters of the cingulate gyrus, insula and superior temporal gyrus. Pitch subgroup resulted in convergence of foci in the insula extending to claustrum, inferior frontal gyrus and putamen. Rhythm-related foci converged in the supramarginal gyrus, angular gyrus, middle frontal gyrus, and somatosensory cortices. Timbre subgroup showed the inferior frontal gyrus, insula and inferior parietal lobule. Such physical auditory features in music seem

to rely on the function of primary cortical areas for audition, somatosensation, language and emotion perception.

Insert Figure 3.

2.6. Meta-analytic connectivity modelling (MACM) and functional characterization.

Meta-analytic connectivity modelling (MACM) was performed to analyze co-activation patterns of the regions-of-interest (ROI) resulting from the primary outcome, aiming to functionally segregate each region's contribution to behavioral domains and paradigm classes. A total of 30 music expertise-related ROIs that resulted from the primary outcome were included. The ROIs were input into the BrainMap database separately, to identify studies that reported activation within each ROI boundary. Then, whole-brain coordinates of activations were downloaded as new foci files, and ALE meta-analyses were performed on the foci to identify areas of convergence. Finally, functional characterization histograms were used to categorize each area's participation in behavioral domains and paradigm classes (**Supplementary Table 7** for MACM, and **Supplementary Table 8** for functional characterization).

3. Discussion.

Accumulating evidence has demonstrated that musical expertise is accompanied by several neuro-anatomical and -functional changes, electing the long-lasting training of professional musicians as a privileged tool to investigate whole-brain neuroplasticity in humans. However, discrepancies between findings challenged our understanding of such phenomenon. Thus, here we provide for the first time a coordinate-based anatomic/activation likelihood estimation (ALE) meta-analysis aiming to depict a unified picture of the neuro-functional-anatomy of musical expertise. Besides confirming that music experts possess highly developed auditory, sensorimotor, and subcortical regions, we highlight convergence towards bilateral and widely distributed networks. Such results challenge the simplistic and most commonly accepted view of left-lateralization for music-related functions and structures. Rather, this meta-analysis strengthens the view that musical training is a beneficial complex and multisensory experience which stimulates a wide variety of cognitive functions.

3.1. Characteristics of the included studies.

In this systematic review and meta-analysis, the included publications reported a clear research question, inclusion and exclusion criteria for participants, description of methods and explicit results. Most of the studies used state-of-the-art techniques and computational MRI-tools, important for the support of standardization and reproducibility of neuroimaging studies. However, we found that some of the studies lacked important demographic data such as the years of education, age of musical training onset, and current time of musical practice, which may have an effect on behavioral tasks and neuroimaging data.

3.2. Structural and functional neuroplasticity in musical expertise.

Our results highlight that expert musicians exhibit higher GM volume in bilateral superior temporal gyri, postcentral gyrus, medial frontal gyrus, supramarginal gyrus, hippocampus, insula, thalamus and left cerebellum; and greater WM volume in bilateral internal capsule bundle, anterior corpus callosum, longitudinal fasciculus and anterior thalamic radiations as compared to non-musicians. Additionally, musicians exhibit higher activity of the inferior frontal gyrus and superior temporal gyrus related to auditory tasks. On the other hand, musicians exhibit lower GM volume in areas of the sensorimotor cortex and cingulate cortex. No WM structure was found to have larger volume in non-musicians as compared to musicians. Finally, musicians exhibit lower activation of the parietal lobes and cerebellum.

The auditory system.

One of our main findings shows enlargement of GM volume in musicians located in medial and posterior superior temporal regions, with clusters extending into primary and secondary auditory cortices. These regions include neuronal assemblies dedicated to encoding of spectro-temporal features of sounds relevant to music¹⁹, such as the discrete pitches forming the Western chromatic scale and fine changes in pitch intervals²⁰. More specifically, it seems that the posterior supratemporal regions are more involved in encoding the height of pitch, whereas the anterior regions are representing the chroma, that is the pitch category irrespectively of the octave²¹. Moreover, these areas participate in auditory imagery of melodies²² and in the processing of the contour and Gestalt patterns of melodies, allowing for recognition and mistakes discrimination²³.

Beyond music-related functions, superior temporal regions are recruited for phonological processing and multimodal integration of sensory information. In particular, accumulating evidence has shown that STS and posterior STG, together with early auditory regions (HG), are involved for the processing of speech sounds, abstract representation of speech sounds and phonemes, and audiovisual integration

mechanisms. Therefore, temporal regions seem to represent fundamental structures for both language and music processing²⁴.

The inferior frontal gyrus has been described as an important hub of both the dorsal and ventral auditory streams. The dorsal auditory stream connects the auditory cortex with the parietal lobe, which projects in turn to the inferior frontal gyrus pars opercularis (Brodmann area 44). This area has been related to the articulatory network, dedicated to specific functions of speech comprehension and production, and highly connected to premotor and insular cortices²⁵. The ventral auditory stream connects the auditory cortex with the middle temporal gyrus and temporal pole, which in turn connects to the inferior frontal gyrus pars triangularis (Brodmann area 45). This area has been associated with semantic processing²⁶. These two regions within the inferior frontal gyrus constitute Broca's area.

The parietal lobe.

The parietal lobe has been also described as an integration area of sensory inputs. The superior parietal lobule includes Brodmann areas 5 and 7, which are involved in somatosensory processing and visuomotor coordination, respectively. The inferior parietal lobule includes Brodmann areas 39 and 40, the angular gyrus and supramarginal gyrus, respectively. The angular gyrus has been related to projection of visual information to Wernicke's area, memory retrieval and theory of mind²⁷. The supramarginal gyrus is an integration hub of somatosensory input, and has been involved in language perception and processing²⁸. Notably, these areas are also important relays in the dorsal auditory stream and the posterior part of the default mode network (DMN).

The salience network.

It has been proposed that the insula and the anterior cingulate cortex (ACC) form the salience network (SN), that coordinates between the DMN and the central executive network (CEN)²⁹. The ACC has been related to cognitive and emotional processing. The cognitive component projects to prefrontal, motor and parietal areas to process top-down and bottom-up stimuli. The emotional component projects to amygdala, nucleus accumbens, hypothalamus, hippocampus, and insula, assessing the salience of emotional and motivational information³⁰. The insula integrates information from the internal physiological state, and projects to the ACC, ventral striatum and prefrontal cortex to initiate adaptive responses³¹. Thus, enhanced function of these areas after musical training appears to result in a more efficient coordination between networks.

The sensorimotor system.

The precentral and postcentral gyrus represent the primary motor and somatosensory cortex, respectively. These two areas are divided by the central sulcus, whose extension represent the sensation and motion of segregated body parts. Our findings show both convergent and divergent effect of musical training in these areas, suggesting a more complex picture than previously thought. For example, neuroadaptations in the sensorimotor system may vary depending on the musical instrument of use³².

The basal ganglia are nuclei of neurons important for the initiation and suppression of movements. In the motor loop of the basal ganglia, inputs from motor cortices project to the dorsal striatum, composed by the putamen and caudate. In the presence of adequate dopaminergic signaling, the direct pathway works to facilitate movement, while the indirect pathway suppresses it. By effective disinhibition processes, the striatum transiently inhibits the pallidum, and in turn, the motor area of the thalamus is disinhibited and free to project back to the motor cortex and initiate a motor program down the corticospinal tract. Similarly, the subthalamic nucleus in the indirect pathway is transiently inhibited when suppressing movement, increasing the inhibition of the pallidum over the thalamus, therefore

blocking the motor cortex activity³³. Our findings show neuroadaptive processes in the putamen and caudate of musicians, presumably resulting in enhancement of such disinhibition mechanisms reflected in fine movement control.

The cerebellum has been shown to play a crucial role in multiple cognitive processes such as sensory discrimination, rhythmic perception and production, working memory, language, and cognition³⁴. Previous fMRI studies in humans suggest that the cerebellum shows segregated activations for motor and cognitive tasks. Motor tasks seem to activate lobules IV-VI in the superior parts of the anterior cerebellum. In contrast, attentional or working memory tasks activate posterior cerebellar hemispheres, namely lobule VIIA, which is divided to crus I and crus II, as well as lobule VIIB³⁵. Musicians and non-musicians show GM volume differences in the cerebellum, specifically in area Crus I, as reported in our results. It has been demonstrated that the activity in crus I/II has a specific relationship with cognitive performance and is linked with lateral prefrontal areas activated by cognitive load increase³⁶. In other words, the crus I/II seems to optimize the response time when the cognitive load increases. Additionally, it has been suggested that crus I/II is associated with beat discrimination thresholds. Thus, there is a positive correlation between GM volume in crus I and beat discrimination performance, evidenced by enhanced ability in musicians³⁷.

White matter.

With regards to WM, we found significant clusters comparing M vs NM that were located in the WM tracts of internal capsule, corpus callosum, longitudinal fasciculus, and thalamic radiations. The internal capsule is a WM structure which connects basal ganglia regions and carries information from and to surrounding cerebral cortex. Connecting fibers in basal ganglia might be thickened by musical expertise because of their involvement in motor control, rhythmic processing, sequence learning, reinforcement learning and memory processes³⁸. In general, basal ganglia structures are recruited during working memory processing for musical motifs³⁹ and the most ventral regions are a core structure of the reward circuit. Interestingly, they are found to be more active in musicians as compared to non-musicians while listening to expressive music⁴⁰.

Lastly, the observed increases in the CC volume may be related to the facilitation of necessary communications between hemispheres and functional networks underlying the coordination of complex sequential bimanual motor sequences and the recall of stored motor programs⁴¹.

3.3. Limitations and future perspectives.

This comprehensive review had the scope to provide a unified synthesis of the effects of musical training on brain structure and functions. Our qualitative review further highlights that previous studies in this field are characterized by heterogeneity of methods, paradigms, and sample backgrounds, as well as relevant missing information. While arguing that the field will benefit from more clarity (e.g. thorough description of methods) and consistency across future studies, we also delineate limitations for our meta-analysis. For example, we set a contrast based on the comparison M vs NM with the aim to narrow down the heterogeneity of the sample and methods used. However, by doing so we relied on two assumptions: (1) the data we pool is based on best research practices; (2) the validity of the GingerALE method. Indeed, to conduct the ALE meta-analysis, we pooled peak coordinates derived from the included studies, rather than using original raw structural MRI images. Thus, the accuracy of our findings relies on the result of a statistical estimation of coordinate-based anatomic foci (input), treated as spatial probability distributions centered at the given coordinates. The heterogeneity of the methods in use in previous studies (ranging from preprocessing software, smoothing, statistical thresholds and participants' characteristics) are not under our control and represent potential

confounders for the results. Perhaps a regression-based assessment of the influence of those heterogeneous factors on the findings would sharpen the results. However, meta-regression analysis is not compatible with GingerALE. When assessing publication bias, all clusters resulting from the primary outcome showed an FSN greater than the minimum imposed of 17, indicating a robust convergence of foci in these regions (further information is reported in **Supplementary Table 9**).

Lastly, on a more theoretical perspective, our results contribute but do not solve the long-standing “nature vs nurture” debate. Indeed, based on evidence that musical training stimulates higher-cognitive functions, auditory-motor integration, attention, memory and engages reward networks, some have suggested that it may be particularly effective in driving neuroplastic mechanisms⁴². However, we are indeed blind to whether the highlighted differences emerging when comparing M Vs NM are training-dependent or due to innate predispositions. In line with the above-mentioned view there are multiple studies demonstrating a strong correlational link between the length of musical training and neuro-functional-anatomical changes. The idea of use-dependent structural changes is further supported by the study conducted by Gaser & Schlaug⁴³ which reported that amateur musicians showed an intermediate increase in gray matter volume when compared to NM and M. Interestingly, the same pattern was found when comparing cognitive abilities, with amateurs showing higher cognitive abilities than NM, but lower than M⁴⁴. Lastly, beyond the statistical power of common inferential studies, a longitudinal neuroanatomical study conducted by Hyde et al.⁴⁵ in children, provided the most compelling evidence in support of the association between musical training and neuroplastic effects. The study showed that, despite the absence of behavioral or brain differences at baseline (prior to any training) in children assigned to musical training and control groups, structural brain changes were visible after only 15 months of musical training. In particular, they found increases in the volume of corpus callosum, auditory and motor cortices and these group differences correlated with improvements in musically relevant motor and auditory skills.

Altogether, the most reasonable conclusion is that the observed neuro-anatomical and -functional changes may be attributed to the interaction between brain maturation processes and the environmental demands of musical training^{13,43,50}. Only when a consistent amount of longitudinal studies will be published, future systematic reviews and meta-analyses will be able to determine the causal link between musical training and brain changes.

4. Conclusions.

The musician's brain has been repeatedly suggested as an ideal example of neuroplasticity. Yet, decades of cognitive neuroscience have provided a scattered and partially controversial series of findings. Thus, with the present comprehensive coordinate-based meta-analysis, we have made a first necessary attempt to provide a unified quantitative and qualitative synthesis of the link between musical training and neuro-anatomical and -functional changes. Results highlight that music experts possess highly developed auditory, sensorimotor and subcortical regions. Indeed, we show that music experts exhibit bilateral widespread neuro-anatomical and functional differences as compared to laypersons, challenging the previous simplistic perspective focused on the left-lateralization of the brain for music-related functions. Conversely, these findings contribute to the notion that musical training is a complex multisensory experience and cannot be anchored to a few numbers of specialized brain regions only.

In conclusion, our findings rediscover the musician's brain and refine the knowledge related to its neuroplasticity. On top of this, we propose that long-lasting professional trainings represent remarkable resources to gain insights on the neuroplastic mechanisms that regulate the human brain.

5. Methods.

5.1. Literature search, screening and extraction.

This systematic review and meta-analysis followed procedures from the Cochrane Handbook for Systematic Reviews⁴⁸ and from the Centre for Reviews and Dissemination (Centre for Reviews and Dissemination, 2014). The review protocol was registered with PROSPERO No. [CRD42017060365]. This review was carried in accordance with the PRISMA statement⁴⁹.

5.1.1. Search strategy.

Systematic search was performed using PubMed, PsycInfo and Scopus, of publications that reported brain structural or functional differences between M and NM. The search (August, 2020) included MeSH terms (“music”, “education”, “brain”, “motor skills”, “magnetic resonance imaging”) and key words (“musical training”, “musician”). No years or places of publication were imposed.

5.1.2. Study eligibility.

For qualitative synthesis, studies were included if they met the following criteria: (1) studies comparing brain structure and function between musicians and non-musicians, (2) in adult population, (3) by means of magnetic resonance imaging, in either structural modality (e.g., voxel-based morphometry [VBM]) or functional modality (e.g., functional magnetic resonance imaging[fMRI]). For quantitative synthesis (meta-analysis), studies were included if results were reported in stereotactic coordinates either Talairach or Montreal Neurological Institute (MNI) three-dimensional-coordinate system.

Studies were excluded using the following criteria: (1) review articles with no original experimental data, (2) neuroimaging data from non-MRI studies (e.g. PET), (3) pathological population, (4) longitudinal designs, (5) functional connectivity analyses, and (6) analyses were region-of-interest (ROI)-based and not whole-brain based (only quantitative synthesis).

Two reviewers (AC and VP) independently screened by title and abstract and selected articles for full-text review, and also performed full-text reviews. Screening and data extraction were performed using the Covidence tool⁵⁰. Any disagreements that arose between the reviewers were resolved through discussion or by a third and/or fourth reviewer (LB, EB).

5.1.3. Data extraction.

From each study, the following variables were extracted: first author, year of publication, population of interest, number of participants, age, sex, absolute pitch, musical feature, years of education, years of musical training, age of musical training onset, weekly training, musical instrument, MRI-system, MRI-model, head-coil, image acquisition parameters of T1, T2* and DWI sequences, repetition time (TR), echo time (TE), voxel size, analysis method and software. The main outcome to extract was any difference in structure or function, in stereotactic coordinates, comparing a musician group and a non-musician group. If any of these points were not reported in the original article, authors were contacted to retrieve this information. Six authors were contacted, with 2 positive answers.

5.2. Quality assessment of MRI studies.

Criteria for MRI quality reporting was selected from a set of guidelines for the standardized reporting of MRI studies^{51,52}. Such guidelines dictate a more consistent and coherent policy for the reporting of MRI methods to ensure that methods can be understood and replicated.

5.3. Activation likelihood estimation (ALE) and meta-analytic connectivity modelling (MACM).

To test the convergence of findings from the neuroimaging studies, we used the anatomic/activation likelihood estimation (ALE) method implemented in the GingerALE software v3.0.2¹⁸, a widely used technique for coordinate-based meta-analysis of neuroimaging data. Statistically significant foci from between-group contrasts were extracted and recorded for each study. If necessary, coordinates were converted from Talairach coordinates to MNI space using the Lancaster transform (icbm2tal) incorporated in GingerALE^{53,54}. The ALE method treats activation foci (input) not as single points, but as spatial probability distributions centered at the given coordinates. Therefore, the algorithm tests to what extent the spatial locations of the foci correlate across independently conducted MRI studies investigating the same construct and assesses them against a null-distribution of random spatial association between experiments⁴⁶. Statistical significance of the ALE scores was determined by a permutation test using cluster-level inference at $p < 0.05$ (FWE). As we did not impose any minimum cluster size of supra-threshold voxels, small volume clusters should be interpreted with caution.

First, the primary outcome was to identify brain structural and functional differences measured by MRI between musicians (M) and non-musicians (NM), pooling all coordinates (foci) together, to examine comprehensively the neural signatures of musical expertise. To test the directionality of the primary outcome, foci were pooled reporting higher volume/activity in musicians ($M > NM$) and lower volume/activity in musicians ($NM > M$). Second, we conducted a contrast analysis to test for similarity (conjunction) and difference (subtraction), between the results of functional and structural foci, independently. Finally, a series of *post hoc* subgroup analyses were conducted to identify brain regions related to specific musical features such as: improvisation, language, melody, memory, motor, pitch, rhythm and timbre.

Meta-analytic connectivity modelling (MACM) was performed to analyze co-activation patterns of regions-of-interest (ROI) resulting from the primary outcome, aiming to functionally segregate each region's putative contribution to behavioral domains^{55,56}. Co-activation analyses were performed using Sleuth⁵⁷ and GingerALE from the BrainMap platform. To identify regions of significant convergence, an ALE meta-analysis was performed over all foci retrieved after searching Sleuth by each music-related ROI independently, and included the experiment level search criteria of “context: normal mapping” and “activations: activation only”. Music-related ROIs were created in Mango⁵⁸ with a 5mm-radius sphere. The functional characterization of music-related ROIs was based on the “Behavioral Domain” (BD) meta-data categories available for each neuroimaging study in the BrainMap database which include action, perception, emotion, cognition and interoception.

All meta-analytic results (ALE maps) were visualized using Mango on the MNI152 1mm standard brain, and resulting coordinates were cross-referenced to the Harvard-Oxford Cortical and Subcortical Atlas and the Juelich Histological Atlas via NeuroVault⁵⁹ and FSLeyes⁶⁰, respectively.

5.4. Fail-Safe N analysis (FSN).

Finally, as traditional meta-analyses, coordinate-based meta-analyses such as ALE can be subject to different forms of publication bias which may impact results and invalidate findings (e.g., the “file drawer problem”). Thus, the Fail-Safe N analysis (FSN)⁶¹ was performed as a measure of robustness against potential publication bias. It refers to the amount of contra-evidence that can be added to a meta-analysis before the results change, and can be obtained for each cluster that survives thresholding in an ALE meta-analysis. For normal human brain mapping, it is estimated that a 95% confidence interval for the number of studies that report no local maxima varies from 5 to 30 per 100 published studies.

Using the upper bound and the fact that the present ALE meta-analysis consists of 58 studies, an estimate for the number of unpublished studies is 17. Therefore, the minimum FSN was defined as 17. A higher FSN indicates more stable results and hence a higher robustness. Such analyses were conducted only on the primary outcome.

Acknowledgments.

The Center for Music in the Brain (MIB) is supported by the Danish National Research Foundation (grant number DNRF 117). The authors wish to thank Hella Kastbjerg for assistance with language check and proof reading.

Authors' contributions.

AC, VPN and EB designed the meta-analysis. VPN guided AC and LB in the initial steps of the meta-analysis after which AC and LB conducted the screening of the studies, and VPN controlled the final selection. AC performed the sensitivity analysis, wrote the first draft of the manuscript and prepared the initial versions of the tables. EB and LB edited and wrote paragraphs in the Introduction and Discussion, whereas VPN wrote paragraphs in the Methods, Results and Discussion sections. VPN prepared the figures and tables. PV contributed to financially supporting the study and commented the initial versions of the manuscripts. EB finalized the paper for submission and all authors edited the manuscript and approved its final version.

Conflict of interest.

There is no actual or potential financial and other conflict of interest related to this manuscript.

References

1. Wolf, C. & Linden, D. E. J. Biological pathways to adaptability-interactions between genome, epigenome, nervous system and environment for adaptive behavior On the origin of adaptability. *Genes, Brain Behav.* **11**, 3–28 (2012).
2. Jäncke, L. The plastic human brain. *Restorative Neurology and Neuroscience* **27**, 521–538 (2009).
3. Lüscher, C. & Malenka, R. C. NMDA receptor-dependent long-term potentiation and long-term depression (LTP/LTD). *Cold Spring Harb. Perspect. Biol.* **4**, 1–15 (2012).
4. Münte, T. F., Altenmüller, E. & Jäncke, L. The musician's brain as a model of neuroplasticity. *Nat. Rev. Neurosci.* **3**, 473–478 (2002).
5. Bermudez, P. & Zatorre, R. Differences in Gray Matter between Musicians and Nonmusicians. *Ann. N. Y. Acad. Sci.* **1060**, 395–399 (2005).
6. Wollman, I., Penhune, V., Segado, M., Carpentier, T. & Zatorre, R. J. Neural network retuning and neural predictors of learning success associated with cello training. *PNAS Proc. Natl. Acad. Sci. United States Am.* **115**, E6056–E6064 (2018).
7. Reybrouck, M. *et al.* Music and and Brain Brain Plasticity : Music Plasticity : How How Sounds Sounds Trigger Trigger Neurogenerative Adaptations Neurogenerative Adaptations. doi:10.5772/intechopen.74318
8. Pantev, C. & Herholz, S. C. Plasticity of the human auditory cortex related to musical training. *Neuroscience and Biobehavioral Reviews* (2011). doi:10.1016/j.neubiorev.2011.06.010
9. Barrett, K. C., Ashley, R., Strait, D. L. & Kraus, N. Art and science: how musical training shapes the brain. *Front. Psychol.* (2013). doi:10.3389/fpsyg.2013.00713
10. Schlaug, G. The brain of musicians. A model for functional and structural adaptation. *Ann. N. Y. Acad. Sci.* **930**, 281–299 (2001).
11. Ashburner, J. & Friston, K. J. Voxel-Based Morphometry—The Methods. (2000). doi:10.1006/nimg.2000.0582
12. Alexander, A. L., Lee, J. E., Lazar, M. & Field, A. S. Diffusion Tensor Imaging of the Brain. (2007).
13. Glover, G. H. Overview of functional magnetic resonance imaging. *Neurosurgery Clinics of North America* **22**, 133–139 (2011).
14. Vaquero, L. *et al.* Structural neuroplasticity in expert pianists depends on the age of musical training onset. *Neuroimage* **126**, 106–119 (2016).
15. James, C. E. *et al.* Musical training intensity yields opposite effects on grey matter density in cognitive versus sensorimotor networks. *Brain Struct. Funct.* **219**, 353–366 (2014).
16. Baer, L. H. *et al.* Regional cerebellar volumes are related to early musical training and finger tapping performance. *Neuroimage* **109**, 130–139 (2015).
17. Clayton, K. K. *et al.* Executive function, visual attention and the cocktail party problem in musicians and non-musicians. *PLoS One* **11**, 17 (2016).
18. Eickhoff, S. B. *et al.* Coordinate-Based Activation Likelihood Estimation Meta-Analysis of Neuroimaging Data: A Random-Effects Approach Based on Empirical Estimates of Spatial Uncertainty. (2009). doi:10.1002/hbm.20718
19. Schneider, P. *et al.* Structural and functional asymmetry of lateral Heschl's gyrus reflects pitch perception preference. *Nat. Neurosci.* **8**, 1241–1247 (2005).

20. Brattico, E., Tervaniemi, M., Näätänen, R. & Peretz, I. Musical scale properties are automatically processed in the human auditory cortex. *Brain Res.* **1117**, 162–174 (2006).
21. Warren, J. D., Uppenkamp, S., Patterson, R. D. & Griffiths, T. D. Analyzing Pitch Chroma and Pitch Height in the Human Brain. in *Annals of the New York Academy of Sciences* **999**, 212–214 (2003).
22. Zatorre, R. J. There's more to auditory cortex than meets the ear. *Hear. Res.* **229**, 24–30 (2007).
23. Hyde, K. L., Peretz, I. & Zatorre, R. J. Evidence for the role of the right auditory cortex in fine pitch resolution. *Neuropsychologia* **46**, 632–639 (2008).
24. Peretz, I., Vuvan, D., Lagrois, M.-É. & Armony, J. L. Neural overlap in processing music and speech. *Philos. Trans. R. Soc. B* **370**, 20140090 (2015).
25. Brown, S., Martinez, M. J. & Parsons, L. M. Music and language side by side in the brain: a PET study of the generation of melodies and sentences. *Eur. J. Neurosci.* **23**, 2791–2803 (2006).
26. DeWitt, I. & Rauschecker, J. P. Phoneme and word recognition in the auditory ventral stream. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 2709 (2012).
27. Seghier, M. L. The angular gyrus: Multiple functions and multiple subdivisions. *Neuroscientist* **19**, 43–61 (2013).
28. Hartwigsen, G. *et al.* Phonological decisions require both the left and right supramarginal gyri. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 16494–16499 (2010).
29. Menon, V. & Uddin, L. Q. Saliency, switching, attention and control: a network model of insula function. *Brain structure & function* **214**, 655–667 (2010).
30. Bush, G., Luu, P. & Posner, M. I. Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences* **4**, 215–222 (2000).
31. Brown, S., Martinez, M. J. & Parsons, L. M. Passive music listening spontaneously engages limbic and paralimbic systems. *Neuroreport* **15**, 2033–2037 (2004).
32. Schneider, P., Sluming, V., Roberts, N., Bleeck, S. & Rupp, A. Structural, functional, and perceptual differences in Heschl's gyrus and musical instrument preference. *Ann. N. Y. Acad. Sci.* **1060**, 387–394 (2005).
33. Purves, D. *et al.* *Neuroscience 5th. Edition.* (Sinauer Associates, Inc., Sunderland., 2012).
34. Kim, S., Ugurbil, K. & Strick, P. Activation of a cerebellar output nucleus during cognitive processing. *Science (80-.)*. **265**, 949–951 (1994).
35. Desmond, J. E., Chen, S. H. A. & Shieh, P. B. Cerebellar Transcranial Magnetic Stimulation Impairs Verbal Working Memory. (2005). doi:10.1002/ana.20604
36. Salmi, J. *et al.* Cognitive and Motor Loops of the Human Cerebro-cerebellar System. *J. Cogn. Neurosci.* **22**, 2663–2676 (2010).
37. Grahn, J. A. & Brett, M. Rhythm and Beat Perception in Motor Areas of the Brain. *J. Cogn. Neurosci.* **19**, 893–906 (2007).
38. Lehericy, S. *et al.* Distinct basal ganglia territories are engaged in early and advanced motor sequence learning. *Proc. Natl. Acad. Sci.* **102**, 12566–12571 (2005).
39. Koelsch, S. *et al.* Functional architecture of verbal and tonal working memory: An fMRI study. *Hum. Brain Mapp.* **30**, 859–873 (2009).

40. Brattico, E. *et al.* It's sad but I like it: The neural dissociation between musical emotions and liking in experts and laypersons. *Front. Hum. Neurosci.* **9**, 21 (2016).
41. Burunat, I. *et al.* Action in perception: Prominent visuo-motor functional symmetry in musicians during music listening. *PLoS One* **10**, (2015).
42. Steele, C. J., Bailey, J. A., Zatorre, R. J. & Penhune, V. B. Early musical training and white-matter plasticity in the corpus callosum: evidence for a sensitive period. *J Neurosci* **33**, 1282–1290 (2013).
43. Gaser, C. & Schlaug, G. *Brain Structures Differ between Musicians and Non-Musicians*. (2003).
44. Criscuolo, A., Bonetti, L., Särkämö, T., Kliuchko, M. & Brattico, E. On the Association Between Musical Training, Intelligence and Executive Functions in Adulthood. *Front. Psychol.* **10**, 1704 (2019).
45. Hyde, K. L. *et al.* The effects of musical training on structural brain development: A longitudinal study. *Ann. N. Y. Acad. Sci.* **1169**, 182–186 (2009).
46. Hudziak, J. J. *et al.* Cortical thickness maturation and duration of music training: Health-promoting activities shape brain development. *J. Am. Acad. Child Adolesc. Psychiatry* **53**, 1153–1161.e2 (2014).
47. Bailey, J. A., Zatorre, R. J. & Penhune, V. B. Early Musical Training is linked to gray matter Structure in the Ventral Premotor Cortex and Auditory-motor Rhythm Synchronization performance. *J. Neurosci.* **26**, 755–767 (2014).
48. Higgins, J. P. & Green, S. *Cochrane Handbook for Systematic Reviews of Interventions*. (2011).
49. Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G. & Group, T. P. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. **6**, 1–5 (2009).
50. Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia. Available at www.covidence.org. Available at: <https://support.covidence.org/help/how-can-i-cite-covidence>.
51. Poldrack, R. A. *et al.* Guidelines for reporting an fMRI study. *Neuroimage* **40**, 409–414 (2008).
52. Nichols, T. E. *et al.* Best practices in data analysis and sharing in neuroimaging using MRI. *Nat. Neurosci.* **20**, 299–303 (2017).
53. Eickhoff, S. B., Bzdok, D., Laird, A. R., Kurth, F. & Fox, P. T. Activation likelihood estimation meta-analysis revisited. *Neuroimage* **59**, 2349–2361 (2012).
54. Laird, A. R. *et al.* Comparison of the disparity between Talairach and MNI coordinates in functional neuroimaging data: Validation of the Lancaster transform. (2010). doi:10.1016/j.neuroimage.2010.02.048
55. Laird, A. R. *et al.* Networks of task co-activations. *Neuroimage* **80**, 505–514 (2013).
56. Robinson, J. L., Laird, A. R., Glahn, D. C., Lovallo, W. R. & Fox, P. T. Metaanalytic Connectivity Modeling: Delineating the Functional Connectivity of the Human Amygdala. *Hum. Brain Mapp.* **31**, 173–184 (2010).
57. Laird, A. R., Lancaster, J. L. & Fox, P. T. *BrainMap: The Social Evolution of a Human Brain Mapping Database*. *Neuroinformatics* (2005).
58. Lancaster, J. L. *et al.* Automated regional behavioral analysis for human brain images. *Front.*

Neuroinform. **6**, 1–12 (2012).

59. Gorgolewski, K. J. *et al.* NeuroVault.org: a web-based repository for collecting and sharing unthresholded statistical maps of the human brain. *Front. Neuroinform.* **9**, 1–9 (2015).
60. Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W. & Smith, S. M. FSL. *Neuroimage* **62**, 782–790 (2012).
61. Acar, F., Seurinck, R., Eickhoff, S. B. & Moerkerke, B. Assessing robustness against potential publication bias in Activation Likelihood Estimation (ALE) meta-analyses for fMRI. *PLoS One* **13**, e0208177 (2018).

Figure legends.

Figure 1. Anatomic likelihood estimation meta-analytic results for studies comparing brain structure and function between M and NM at cluster level inference $p < 0.05$ (FWE). The primary outcome included ALE meta-analysis of all included foci, and the contrast M vs NM for structural and functional modalities, independently. M > NM = higher volume/activity in musicians; NM < NM = lower volume/activity in musicians. Z, peak Z-value.

Figure 2. Conjunction analyses comparing foci from structural (CT, DTI, VBM) and functional (fMRI) studies, at cluster level inference $p < 0.05$ (FWE). Contrast analyses were performed for structural and functional studies, and were tested for similarity (conjunction) and difference (subtraction), to illustrate common and/or distinct areas between the neuroimaging modalities. ALE, anatomic likelihood estimation value; Z, peak Z-value.

Figure 3. Musical feature subgroup meta-analyses, at cluster level inference $p < 0.05$ (FWE). Z, peak Z-value.

Table 1. Characteristics of studies included in qualitative synthesis.											M (age)		NM (age)		M (education)		NM (education)		Training (years)		Onset (years)		Training (hr/week)		M vs NM	
Author	Year	Method	Feature	M	NM	AP	Males	Females	Total	Instrument	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD	Expertise	Foci		
1 Abdul-K	2011a	VBM	structure	26	26	-	52	0	52	b,c,d	43.2	9.3	42.8	10.8	-	-	-	-	20.4	9.4	9.6	2.4	36	5	Pro	N
2 Abdul-K	2011b	DTI	structure	10	10	-	14	6	20	a,b	38	14.6	34.5	11.7	13.8	1.2	13.5	2.5	29.9	14.1	6.6	2.05	-	-	Pro	N
3 Acer	2018	VBM+DTI	structure	7	7	-	14	0	14	b	22.8	2.1	23.4	1.9	-	-	-	-	6.3	1.5	10	-	10	2.5	Pro	Y
4 Amunts	1997	VBM	structure	21	30	-	51	0	51	a	26.9	4.6	26.4	3.9	-	-	-	-	-	-	-	-	-	-	Pro	Y
5 Angulo-P	2014	fMRI	timbre	28	25	-	29	24	53	a,b	28	7	29	9	-	-	-	-	-	-	-	-	-	-	Pro	Y
6 Bailey	2014	VBM+DBM	structure	30	20	-	-	-	50	-	23.5	3.9	26.2	4.3	-	-	-	-	11.7	4	5.8	1.2	15.2	9.97	Pro	Y
7 Bangert	2006	fMRI	melody+motor	7	7	-	6	8	14	a	28.5	7.3	28.4	5.8	-	-	-	-	20	8.7	8.5	4.8	-	-	Pro	Y
8 Baumann	2007	fMRI	melody+motor	7	7	-	5	9	14	a	25.7	3.2	30	5	-	-	-	-	10	-	-	-	15	-	Pro	Y
9 Bengtsson	2005	DTI	structure	8	8	-	16	0	16	a	32.6	5.7	33.1	9.8	-	-	-	-	-	5.8	1.4	-	-	-	Pro	Y
10 Berkowitz	2010	fMRI	improvisation+rhythm+motor	12	12	-	-	-	24	a	21.9	-	22.9	-	-	-	-	-	13	-	-	-	-	-	Pro	Y
11 Bermudez	2005	VBM	structure	43	51	22	-	-	94	-	-	-	-	-	-	-	-	-	> 10	-	-	-	-	-	Pro	Y
12 Bermudez	2009	VBM+CT	structure	71	64	27	49	86	135	-	23.3	3.2	24.4	4.9	-	-	-	-	16.9	3.4	6.5	2.5	13.9	9.7	Pro	Y
13 Bianchi	2017	fMRI	pitch	16	15	1	14	17	31	a,b,c,d	26	-	25	-	-	-	-	-	12.4	-	7	-	-	-	Pro	Y
14 Chen	2008	fMRI	rhythm	12	12	-	-	-	24	-	23.2	-	23.8	-	-	-	-	-	-	-	-	-	-	-	Pro	Y
15 Choi	2015	CT	structure	14	14	-	0	28	28	c	20.3	1.2	20.1	1.2	-	-	-	-	7.9	1.2	-	-	-	-	Pro/Ama	Y
16 De Manzano	2018	CT	structure	9	9	-	8	10	18	a	37	6	37	6	-	-	-	-	-	-	-	-	-	-	Ama	N
17 Du	2017	fMRI	language/speech	15	15	-	16	14	30	a,b,c,d	21.4	2.7	22.1	4.4	10.5	1.7	2.5	1.7	16.3	3.8	5.1	1.6	-	-	Pro	Y
18 Elmer	2012	fMRI	language/speech	12	13	-	9	16	25	a,b,c,d	24.9	5.5	25.3	3.5	-	-	-	-	-	5.9	1.3	-	-	-	Pro	Y
19 Elmer	2013	CT	structure+language/speech	13	13	-	13	13	26	a,b,c,d	25.2	5.74	25.3	3.52	-	-	-	-	-	6.22	1.06	-	-	-	Pro	Y
20 Elmer	2016	DTI	structure+language/speech	13	13	-	13	13	26	a,b,c,d	25.2	5.7	25.3	3.5	-	-	-	-	-	6.22	1.06	-	-	-	Pro	N
21 Gaab	2003	fMRI	pitch+memory/familiarity	10	10	-	10	10	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Pro	N
22 Gaab	2006	fMRI	language/speech	20	20	-	20	20	40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Pro	N
23 Gagnepain	2017	fMRI	memory/familiarity+melody	20	19	-	20	19	39	a,b,c,d	22.9	3.05	24.6	3.91	15.15	0.99	16.32	2.08	15.3	3.7	7.55	1.87	-	-	Pro/Ama	Y
25 Gaser	2003	VBM	structure	20	40	-	80	0	80	a,b	23.1	3.83	26.9	4.9	-	-	-	-	17	-	6	1.8	15.6	-	Pro/Ama	Y
24 Gärtner	2013	DBM	structure	20	19	-	39	0	30	a,b	43.3	3.8	43.5	3.8	-	-	-	-	36.2	4.5	7.8	2.6	19.6	-	Pro	N
26 Giacosa	2016	DTI	structure	18	19	-	13	24	37	a,b,c,d	22.9	3.4	25.4	5.1	2.37	0.98	2.58	1.12	15	3.6	-	-	-	-	Pro	Y
27 Groussard	2010	VBM	structure	20	20	-	20	20	40	a,b,c	22.9	3.05	24.6	3.8	15.15	0.99	16.35	2.03	15.3	3.6	7.55	1.87	-	-	Pro	Y
28 Groussard	2014	VBM	structure	33	11	-	26	18	44	a,b,c	24.2	3.5	25	3.4	15.45	1.13	16.9	2.17	15.5	2	6.6	1.4	7.5	-	Pro/Ama	Y
29 Halwani	2011	DTI	structure	22	11	-	-	-	33	a,b,c,d	26.5	5.2	27.5	10.3	-	-	-	-	14.7	7	7	2.3	-	-	Pro	N
30 Han	2009	VBM+DTI	structure	18	21	-	16	23	39	a	22.6	2.6	22.4	2.6	-	-	-	-	10.4	4.2	12.2	3.3	-	-	Pro	Y
31 Harris	2015	fMRI	improvisation+memory/familiarity	24	12	-	35	1	36	a	40.2	13	43.7	9.6	-	-	-	-	-	-	-	-	-	-	Pro	Y
32 Haslinger	2004	fMRI	motor	12	12	-	12	12	24	a	23	2.2	25	1.2	-	-	-	-	15.9	2	-	-	-	-	Pro	Y
33 Haslinger	2005	fMRI	motor	12	12	-	12	12	24	a	23	2.4	25	1.2	-	-	-	-	16.9	2.7	-	-	-	-	Pro	Y
34 Herdener	2010	fMRI	pitch+memory/familiarity	7	7	-	12	2	14	-	42.6	10.3	42.6	11.5	-	-	-	-	-	6.1	2	20.8	23.1	-	Pro	Y
35 Herdener	2014	fMRI	rhythm+language/speech+pitch	11	11	-	22	0	22	d	31.7	7.29	27.6	5.44	-	-	-	-	-	8.42	1.89	21.5	14.55	-	Pro	Y
36 Hernández	2019	VBM	structure	32	26	-	38	20	58	a,b,c,d,e	20.1	2.1	20.7	2.22	-	-	-	-	10.3	1.32	-	-	-	-	Pro	N
37 Huang	2010	fMRI	memory/familiarity	10	10	-	0	20	20	a	21.3	1.1	21.6	1.3	-	-	-	-	-	-	-	-	-	-	Ama	Y
38 Hutchinson	2003	VBM	structure	60	60	-	60	60	120	a,b	25.1	4.8	24.8	4.7	-	-	-	-	19.6	4.6	5.6	1.9	-	-	Pro	Y
39 Imfeld	2009	DTI	structure	26	13	13	16	23	39	a,b,c	24.6	2.9	25.6	5.3	-	-	-	-	17.8	2.9	6.8	2.1	-	-	Pro	Y
40 James	2014	VBM	structure	20	19	-	-	-	59	a	24.5	4.5	24	4.5	-	-	-	-	18.3	2.6	6.2	1.9	-	-	Pro/Ama	Y
41 Karpati	2017	VBM+CT	structure	19	20	-	14	25	39	a,b,c,d	22.9	3.4	25.4	5.1	-	-	-	-	15.4	3.4	-	-	-	-	Pro	Y
42 Kleber	2010	fMRI	singing+motor	10	18	-	22	27	49	e	38.1	-	25.2	-	-	-	-	-	21.3	-	-	-	27.6	-	Pro	Y
43 Kleber	2016	VBM	singing+motor	27	28	-	15	40	55	e	26.6	3.5	24.9	3.9	-	-	-	-	10.4	4.2	15.7	3.7	17.1	6.6	Pro	Y
44 Koelsch	2005	fMRI	pitch	10	10	-	10	10	20	a	26.8	-	25.6	-	-	-	-	-	9.4	-	-	-	-	-	Pro/Ama	N
45 Koeneke	2004	fMRI	motor	7	7	-	8	6	14	a	24	3.4	24.3	5.6	-	-	-	-	16.4	-	-	-	-	-	Pro	Y
46 Krings	2000	fMRI	motor	4	4	-	4	4	8	a	36.5	-	-	-	-	-	-	-	30	-	-	-	20	-	Pro	N
47 Krishnan	2018	fMRI	timbre	40	20	-	53	7	60	b,e	28.4	6.8	27.8	8.9	-	-	-	-	5.9	3.9	11.2	2.3	-	-	Pro	Y
48 Lee	2011	fMRI	audiovisual+language/speech	18	19	-	-	-	37	a	24.3	5.3	27.1	3.4	-	-	-	-	16.4	5.5	7.9	1.9	3.33	1.64	Ama	Y
49 Limb	2006	fMRI	rhythm	12	12	-	18	6	24	-	31	6.52	34	14.9	-	-	-	-	-	-	-	-	-	-	Pro	Y
50 Liu	2018	fMRI	tempo+emotion	21	27	-	23	25	48	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Pro	Y
51 Matsui	2013	fMRI	musical structure	12	15	-	0	27	27	a	29.1	-	23.4	-	-	-	-	-	-	-	-	-	-	-	Pro	Y
52 Mathews	2020	fMRI	rhythm+harmony+motor	29	25	-	30	24	54	-	23.8	2.84	23.2	2.5	-	-	-	-	11.5	3.27	-	-	11.67	10		

64 Rüber	2015	DTI	structure	20	10	-	16	14	30	a,b	24.7	5.1	26.5	2.5	-	-	-	-	-	6.5	2.3	3.3	2.5	Pro/Ama	N	
65 Sakreida	2018	fMRI	motor+rhythm	15	16	-	15	16	31	a,b,c	20.8	2.3	20.4	1.5	-	-	-	11.6	3.4	-	-	10.9	-	Pro	Y	
66 Sato	2015	VBM	structure	23	15	-	0	38	38	a,b,c	21.2	-	21.6	-	-	-	-	-	4	-	-	-	-	Pro	Y	
67 Schlaffke	2020	fMRI+DTI	motor+structure	20	24	-	44	0	44	d	-	-	-	-	-	-	-	17	6	-	-	10.5	8	Pro	N	
68 Schlaug	1995a	VBM	structure	30	30	-	44	16	60	a,b	26.1	3.8	26.5	4.6	-	-	-	-	-	-	-	-	-	Pro	N	
69 Schlaug	1995b	VBM	structure	30	30	11	44	16	60	a,b	26.1	3.8	26.5	4.6	-	-	-	-	-	-	-	-	-	Pro	N	
70 Schlaug	2005	VBM	structure	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Pro	N	
71 Schmithorst	2002	DTI	structure	5	6	-	-	-	11	-	31.2	11.2	42.2	20.1	-	-	-	-	-	-	-	-	-	Ama	N	
72 Schmithorst	2003	fMRI	melody+harmony	7	8	-	11	4	15	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Ama	N	
73 Schmithorst	2004	fMRI	math	7	8	-	11	4	15	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Ama	Y	
75 Schneider	2002	VBM	structure	16	8	-	16	8	24	a	-	-	-	-	-	-	-	-	-	-	-	-	-	Pro/Ama	N	
74 Seung	2005	fMRI	pitch	5	5	-	0	10	10	a	-	-	-	-	-	-	-	-	-	-	-	-	-	Pro	Y	
76 Sluming	2002	VBM	structure	26	26	-	52	0	52	b,c	43.2	9.3	42.8	10.8	-	-	-	-	20.4	9.4	9.6	2.4	36	5	Pro	Y
77 Sluming	2007	fMRI	motor	10	10	-	20	0	20	b,c	42.2	10.1	40.1	11.8	-	-	-	-	-	-	-	-	-	-	Pro	Y
78 Steele	2013	DTI	structure	36	17	-	34	19	53	-	22.7	4.14	26.4	4.71	-	-	-	-	11.5	3.22	5.72	1.13	15	10.2	Pro	N
79 Vaquero	2016	VBM	structure	36	17	-	27	26	53	a	24.9	4.86	24.1	4.39	-	-	-	-	-	5.19	0.69	14.26	9.31	Pro	Y	
80 Vaquero	2020	DTI	structure	31	31	-	28	34	62	-	23.1	4.63	25	4.25	14.03	3.08	14.06	2.83	15.7	4.26	7.19	2.39	-	-	Pro	N
81 Wang	2019	VBM	structure	21	21	-	0	42	42	e	24.1	2.19	23.2	2.65	-	-	-	-	12	5	12	5	-	-	Pro	Y
82 Zarate	2005	fMRI	pitch+singing	12	12	-	12	12	24	e	-	-	-	-	-	-	-	-	9.5	-	-	-	-	-	Ama	N
83 Zarate	2008	fMRI	pitch+singing	12	12	-	12	12	24	e	-	-	-	-	-	-	-	-	10	3.4	-	-	-	-	Ama	N
84 Zuk	2014	fMRI	memory/familiarity	15	15	-	18	12	30	a,b,c	-	-	-	-	-	-	-	-	5.2	1.33	5.73	1.62	21.87	11.49	Pro	N

M, musicians; NM, non-musicians; CT, cortical thickness; DBM, deformation-based morphometry; DTI, diffusion tensor imaging; fMRI, functional magnetic resonance imaging; VBM, voxel-based morphometry; Pro, professional musician; Ama, amateur; a, piano; b, strings; c, wind; d, percussion; e, voice; Y, yes; N, no.

Table 2. Anatomic likelihood estimation meta-analytic results for studies comparing brain structure and function between M and NM at cluster level inference $p < 0.05$ (FWE).

Cluster number	Volume (mm ³)	MNI coordinates			ALE	P	Z	Label (Side region BA)		
		x	y	z						
1. All foci: 675 foci, 79 experiments, 2780 subjects										
1	42464	50	10	22	3E-02	3E-07	5.0	R Inferior Frontal Gyrus BA9		
		54	-10	4	3E-02	1E-06	4.7	R Superior Temporal Gyrus BA22		
		54	24	10	3E-02	2E-06	4.6	R Inferior Frontal Gyrus BA45		
		50	-32	18	3E-02	5E-06	4.4	R Superior Temporal Gyrus BA41		
		64	-14	0	2E-02	1E-05	4.2	R Superior Temporal Gyrus BA22		
		54	12	0	2E-02	2E-04	3.6	R Precentral Gyrus BA44		
		60	-44	-6	2E-02	3E-04	3.4	R Middle Temporal Gyrus BA21		
		62	-8	-8	2E-02	3E-04	3.4	R Middle Temporal Gyrus BA21		
		28	2	-4	2E-02	8E-04	3.1	R Putamen		
		58	-36	24	2E-02	9E-04	3.1	R Insula BA13		
		58	-2	-6	2E-02	1E-03	3.1	R Superior Temporal Gyrus BA22		
		50	-28	-4	2E-02	1E-03	3.1	R Insula BA22		
		32	22	-10	2E-02	1E-03	3.1	R Insula BA47		
		58	2	34	2E-02	1E-03	3.0	R Precentral Gyrus BA6		
		52	-44	36	2E-02	2E-03	2.9	R Supramarginal Gyrus BA40		
		52	-62	-4	1E-02	2E-03	2.9	R Inferior Temporal Gyrus BA19		
		54	-60	-8	1E-02	3E-03	2.7	R Middle Occipital Gyrus BA19		
		36	22	8	1E-02	3E-03	2.7	R Insula BA13		
		66	-40	8	1E-02	7E-03	2.4	R Middle Temporal Gyrus BA22		
		56	-10	22	1E-02	8E-03	2.4	R Postcentral Gyrus BA43		
		56	0	22	1E-02	1E-02	2.3	R Precentral Gyrus BA6		
		54	-44	16	1E-02	1E-02	2.3	R Superior Temporal Gyrus BA13		
		36	30	-18	1E-02	1E-02	2.3	R Inferior Frontal Gyrus BA47		
		56	-4	24	1E-02	1E-02	2.3	R Precentral Gyrus BA6		
		40	20	-2	1E-02	1E-02	2.3	R Insula		
		54	36	14	1E-02	1E-02	2.3	R Middle Frontal Gyrus BA46		
		30	14	12	1E-02	1E-02	2.3	R Claustrum		
		18	6	8	1E-02	1E-02	2.3	R Putamen		
		38	-2	12	1E-02	1E-02	2.2	R Insula BA13		
		64	-30	-4	1E-02	1E-02	2.2	R Middle Temporal Gyrus BA21		
		42	2	-2	1E-02	1E-02	2.2	R Claustrum		
		40	2	6	1E-02	1E-02	2.2	R Claustrum		
		52	-22	-14	1E-02	2E-02	2.2	R Superior Temporal Gyrus BA22		
		40	32	-16	1E-02	2E-02	2.2	R Inferior Frontal Gyrus BA47		
		66	-42	-24	1E-02	2E-02	2.2	R Inferior Temporal Gyrus BA20		
		18	12	10	1E-02	2E-02	2.1	R Caudate Body		
		12	10	-8	1E-02	2E-02	2.1	R Caudate Head		
		36	-8	12	1E-02	2E-02	2.1	R Claustrum		
		50	-12	26	1E-02	2E-02	2.1	R Precentral Gyrus BA6		
		18	14	-4	1E-02	2E-02	2.1	R Putamen		
		36	22	-16	1E-02	2E-02	2.0	R Inferior Frontal Gyrus BA47		
		68	-44	-14	1E-02	2E-02	2.0	R Middle Temporal Gyrus BA20		
		46	38	14	1E-02	2E-02	2.0	R Middle Frontal Gyrus BA46		
		50	46	18	9E-03	2E-02	2.0	R Middle Frontal Gyrus BA46		
		42	-48	-12	9E-03	2E-02	2.0	R Fusiform Gyrus BA37		
		68	-38	-12	9E-03	3E-02	1.9	R Middle Temporal Gyrus BA21		
		60	-8	-18	9E-03	3E-02	1.9	R Middle Temporal Gyrus BA21		
		2	27056	-58	-18	4	3E-02	4E-08	5.4	L Superior Temporal Gyrus BA41
				-52	-6	2	3E-02	2E-06	4.6	L Superior Temporal Gyrus BA22
				-44	-40	42	2E-02	3E-05	4.0	L Supramarginal Gyrus BA40
-58	-46			16	2E-02	4E-05	3.9	L Superior Temporal Gyrus BA22		
-54	-26			38	2E-02	5E-05	3.9	L Postcentral Gyrus BA2		
-58	-30			14	2E-02	5E-05	3.9	L Superior Temporal Gyrus BA42		
-54	4			-12	2E-02	4E-04	3.3	L Superior Temporal Gyrus BA38		
-48	-2			-10	2E-02	7E-04	3.2	L Superior Temporal Gyrus BA22		
-56	-36			22	2E-02	8E-04	3.2	L Insula BA13		
-66	-24			30	2E-02	9E-04	3.1	L Inferior Parietal Lobule BA40		
-30	-14			-14	2E-02	9E-04	3.1	L Parahippocampal Gyrus		
-48	-52			30	2E-02	1E-03	3.1	L Superior Temporal Gyrus BA39		
-42	-26			2	2E-02	2E-03	3.0	L Superior Temporal Gyrus BA22		
-50	-36			56	2E-02	2E-03	2.9	L Inferior Parietal Lobule BA40		
-40	-42			12	1E-02	2E-03	2.8	L Superior Temporal Gyrus BA41		
-60	-14			34	1E-02	2E-03	2.8	L Postcentral Gyrus BA3		
-20	-16			-16	1E-02	1E-02	2.3	L Parahippocampal Gyrus BA28		
-48	-42			18	1E-02	1E-02	2.2	L Superior Temporal Gyrus BA13		

3	17744	-18	-4	-14	1E-02	2E-02	2.2	L Parahippocampal Gyrus
		-14	-14	-14	1E-02	2E-02	2.2	L Parahippocampal Gyrus BA28
		-38	-20	-2	1E-02	2E-02	2.1	L Claustrum
		-48	6	-24	1E-02	2E-02	2.1	L Superior Temporal Gyrus BA38
		-42	-26	52	1E-02	2E-02	2.1	L Postcentral Gyrus BA40
		-48	-50	10	9E-03	2E-02	2.0	L Superior Temporal Gyrus BA39
		-72	-28	4	9E-03	2E-02	2.0	L Superior Temporal Gyrus BA22
		-50	-50	54	9E-03	3E-02	1.9	L Inferior Parietal Lobule BA40
		-50	6	36	3E-02	1E-07	5.2	L Precentral Gyrus BA6
		-50	8	16	3E-02	3E-07	5.0	L Inferior Frontal Gyrus BA44
		-50	10	22	3E-02	8E-07	4.8	L Inferior Frontal Gyrus BA9
		-22	-2	54	3E-02	1E-06	4.7	L Middle Frontal Gyrus BA6
		-48	4	26	3E-02	5E-06	4.4	L Precentral Gyrus BA6
		-52	16	44	2E-02	7E-04	3.2	L Middle Frontal Gyrus BA6
		-48	2	50	2E-02	2E-03	2.9	L Precentral Gyrus BA6
		-34	-4	44	1E-02	3E-03	2.7	L Middle Frontal Gyrus BA6
		-40	-18	56	1E-02	4E-03	2.7	L Postcentral Gyrus BA3
		-28	-14	68	1E-02	5E-03	2.5	L Precentral Gyrus BA6
		-46	-4	54	1E-02	6E-03	2.5	L Precentral Gyrus BA6
		-40	-6	56	1E-02	8E-03	2.4	L Precentral Gyrus BA6
		-36	-26	66	1E-02	1E-02	2.2	L Postcentral Gyrus BA3
		-36	-18	48	1E-02	1E-02	2.2	L Postcentral Gyrus BA3
		-34	8	60	9E-03	3E-02	1.9	L Middle Frontal Gyrus BA6
4	17400	46	-32	44	2E-02	5E-05	3.9	R Inferior Parietal Lobule BA40
		44	-50	48	2E-02	7E-05	3.8	R Inferior Parietal Lobule BA40
		62	-14	40	2E-02	2E-04	3.6	R Precentral Gyrus BA4
		32	-16	64	2E-02	2E-04	3.5	R Precentral Gyrus BA4
		54	-22	42	2E-02	4E-04	3.4	R Postcentral Gyrus BA2
		40	-42	46	2E-02	5E-04	3.3	R Inferior Parietal Lobule BA40
		42	-48	56	2E-02	7E-04	3.2	R Inferior Parietal Lobule BA40
		38	-20	40	2E-02	1E-03	3.1	R Postcentral Gyrus BA2
		42	-54	42	2E-02	1E-03	3.1	R Inferior Parietal Lobule BA40
		18	6	52	2E-02	1E-03	3.0	R Cingulate Gyrus BA24
		24	6	60	1E-02	2E-03	2.8	R Sub-Gyral BA6
		52	0	48	1E-02	3E-03	2.8	R Precentral Gyrus BA6
		30	-2	66	1E-02	3E-03	2.8	R Middle Frontal Gyrus BA6
		34	-34	46	1E-02	3E-03	2.7	No Gray Matter found
		42	-58	52	1E-02	5E-03	2.6	R Inferior Parietal Lobule BA7
		48	-6	48	1E-02	5E-03	2.6	R Precentral Gyrus BA6
		28	-62	56	1E-02	6E-03	2.5	R Superior Parietal Lobule BA7
		40	2	50	1E-02	1E-02	2.3	R Middle Frontal Gyrus BA6
		36	4	52	1E-02	1E-02	2.2	R Middle Frontal Gyrus BA6
		28	4	50	1E-02	2E-02	2.2	R Middle Frontal Gyrus BA6
		60	-54	44	1E-02	2E-02	2.2	R Inferior Parietal Lobule BA40
		38	-18	52	9E-03	2E-02	2.0	R Precentral Gyrus BA4
		<hr/>						
2. $M > NM$: 509 foci, 59 experiments, 2189 subjects								
1	36264	-58	-18	4	3E-02	8E-09	5.6	L Superior Temporal Gyrus BA41
		-52	-6	2	3E-02	5E-07	4.9	L Superior Temporal Gyrus BA22
		-50	8	18	3E-02	3E-06	4.5	L Inferior Frontal Gyrus BA9
		-54	8	12	2E-02	5E-06	4.4	L Inferior Frontal Gyrus BA44
		-48	4	26	2E-02	5E-06	4.4	L Precentral Gyrus BA6
		-58	-46	16	2E-02	1E-05	4.2	L Superior Temporal Gyrus BA22
		-58	-30	14	2E-02	2E-05	4.1	L Superior Temporal Gyrus BA42
		-44	-40	40	2E-02	3E-05	4.0	L Supramarginal Gyrus BA40
		-54	-26	36	2E-02	7E-05	3.8	L Inferior Parietal Lobule BA40
		-52	4	36	2E-02	9E-05	3.7	L Precentral Gyrus BA6
		-54	4	-12	2E-02	2E-04	3.6	L Superior Temporal Gyrus BA38
		-48	-2	-10	2E-02	3E-04	3.5	L Superior Temporal Gyrus BA22
		-66	-24	30	2E-02	4E-04	3.4	L Inferior Parietal Lobule BA40
		-58	-36	20	2E-02	4E-04	3.4	L Insula BA13
		-52	16	44	2E-02	4E-04	3.4	L Middle Frontal Gyrus BA6
		-52	-34	-6	2E-02	5E-04	3.3	L Middle Temporal Gyrus BA21
		-48	2	50	2E-02	9E-04	3.1	L Precentral Gyrus BA6
		-48	-52	30	1E-02	1E-03	3.0	L Superior Temporal Gyrus BA39
		-42	-28	4	1E-02	2E-03	2.8	L Superior Temporal Gyrus BA13
		-46	-4	54	1E-02	3E-03	2.7	L Precentral Gyrus BA6
		-40	-6	56	1E-02	4E-03	2.6	L Precentral Gyrus BA6
		-36	-24	60	1E-02	5E-03	2.6	L Precentral Gyrus BA4
		-36	-26	66	1E-02	7E-03	2.4	L Postcentral Gyrus BA3
-36	-18	48	1E-02	8E-03	2.4	L Postcentral Gyrus BA3		

2	35744	-48	6	-24	1E-02	9E-03	2.4	L Superior Temporal Gyrus BA38
		-66	-40	-2	1E-02	1E-02	2.2	L Middle Temporal Gyrus BA21
		-72	-28	4	9E-03	1E-02	2.2	L Superior Temporal Gyrus BA22
		-60	-42	0	9E-03	2E-02	2.1	L Middle Temporal Gyrus BA21
		-42	-26	50	9E-03	2E-02	2.1	L Postcentral Gyrus BA40
		-36	-28	12	9E-03	2E-02	2.0	L Transverse Temporal Gyrus BA41
		-32	-38	36	8E-03	2E-02	2.0	L Sub-Gyral BA40
		54	-10	4	3E-02	4E-07	4.9	R Superior Temporal Gyrus BA22
		54	24	10	3E-02	5E-07	4.9	R Inferior Frontal Gyrus BA45
		50	10	22	3E-02	1E-06	4.7	R Inferior Frontal Gyrus BA9
		66	-14	0	2E-02	4E-05	3.9	R Superior Temporal Gyrus BA22
		54	12	0	2E-02	6E-05	3.9	R Precentral Gyrus BA44
		60	-42	-4	2E-02	1E-04	3.7	R Middle Temporal Gyrus BA21
		52	-32	18	2E-02	1E-04	3.7	R Superior Temporal Gyrus BA41
		42	-32	16	2E-02	1E-04	3.6	R Superior Temporal Gyrus BA41
		58	-36	24	2E-02	4E-04	3.3	R Insula BA13
		50	-28	-4	2E-02	5E-04	3.3	R Insula BA22
		58	0	-6	2E-02	6E-04	3.3	R Superior Temporal Gyrus BA22
		52	-62	-4	1E-02	1E-03	3.1	R Inferior Temporal Gyrus BA19
		52	-44	36	1E-02	1E-03	3.1	R Supramarginal Gyrus BA40
		54	-60	-8	1E-02	2E-03	3.0	R Middle Occipital Gyrus BA19
		36	22	8	1E-02	2E-03	2.9	R Insula BA13
		54	6	32	1E-02	3E-03	2.7	R Precentral Gyrus BA6
		66	-40	8	1E-02	4E-03	2.7	R Middle Temporal Gyrus BA22
		54	-44	16	1E-02	5E-03	2.6	R Superior Temporal Gyrus BA13
		60	-30	8	1E-02	5E-03	2.6	R Superior Temporal Gyrus BA41
		40	20	-2	1E-02	6E-03	2.5	R Insula
		52	-22	-14	1E-02	8E-03	2.4	R Superior Temporal Gyrus BA22
		50	-64	-18	1E-02	8E-03	2.4	R Fusiform Gyrus BA37
		64	-30	-6	1E-02	9E-03	2.4	R Middle Temporal Gyrus BA21
		66	-42	-24	1E-02	9E-03	2.4	R Inferior Temporal Gyrus BA20
		68	-44	-14	1E-02	1E-02	2.3	R Middle Temporal Gyrus BA20
		34	22	-8	1E-02	1E-02	2.2	R Claustrum
		40	-40	24	9E-03	2E-02	2.1	R Insula BA13
		60	-8	-18	9E-03	2E-02	2.1	R Middle Temporal Gyrus BA21
3. NM > M: 166 foci, 20 experiments, 591 subjects								
1	768	-50	-36	56	1E-02	3E-05	4	L Inferior Parietal Lobule BA40
		-46	-40	52	1E-02	7E-05	3.8	L Inferior Parietal Lobule BA40
2	624	-48	8	36	2E-02	6E-06	4.4	L Precentral Gyrus BA6
3	232	48	-34	66	2E-02	3E-05	4	R Postcentral Gyrus BA2
4	200	64	-14	38	1E-02	7E-05	3.8	R Precentral Gyrus BA4
5	184	22	-66	-24	1E-02	2E-04	3.6	R Cerebellum, Declive
6	168	58	12	22	1E-02	2E-04	3.5	R Inferior Frontal Gyrus BA9
7	168	-6	18	48	1E-02	3E-04	3.4	L Superior Frontal Gyrus BA6
8	160	18	6	10	1E-02	5E-04	3.3	R Caudate Body
		18	12	10	1E-02	7E-04	3.2	R Caudate Body
9	120	62	-22	2	1E-02	3E-04	3.4	R Superior Temporal Gyrus BA41
10	104	38	-60	50	1E-02	3E-04	3.4	R Superior Parietal Lobule BA7

M, musicians; NM, non-musicians; BA, Brodmann area; ALE, anatomic likelihood estimation; P, p-value; Z, peak z-value; R, right; L, left.

ALL FOCI

M vs NM

M vs NM (STRUCTURAL)

M vs NM (FUNCTIONAL)

ALL FOCI

M > NM

NM > M

STRUCTURAL

FUNCTIONAL

STRUCTURAL + FUNCTIONAL

STRUCTURAL - FUNCTIONAL

FUNCTIONAL - STRUCTURAL