Spontaneous *Escherichia coli* persists with week-long survival dynamics and lasting memory of a short starvation pulse

Mikkel Skjoldan Svenningsen¹, Sine Lo Svenningsen², Michael Askvad Sørensen^{2,*}, Namiko Mitarai^{1,**}

Abstract

The vast majority of a bacterial population is killed within a time scale comparable to their generation time when treated with a lethal concentration of antibiotics. However, a small subpopulation typically survives for an extended period. To investigate the long-term killing dynamics of bacterial cells we constructed a week-long killing assay and followed the survival fraction of *Escherichia coli* K12 exposed to a high concentration of ciprofloxacin. We found that long-term survivors were formed during exponential growth, with some cells surviving at least 7 days. In contrast to the biphasic killing dynamics typically used to describe persistence, the long-term dynamics contained at least three timescales, which greatly enhances predictions of the population survival time. Furthermore, we observed a surprisingly long memory effect of a brief starvation pulse, which was dependent on the (p)ppGpp synthase *relA*. Specifically, one hour of carbon starvation prior to antibiotics exposure increased the surviving fraction by nearly 100-fold even after 4 days of ciprofloxacin treatment.

Importance: Bacterial infections pose an increasing worldwide health risk due to the rise of antibiotics resistance. This motivates the question of how to administer antibiotics while minimizing the time the bacteria are exposed to the antibiotic to reduce the development of resistance, and poses a need for understanding bacterial killing dynamics. We investigate the long-term time-dependent survival of *E. coli* during one week of exposure to ciprofloxacin. We demonstrate that a subpopulation of bacteria survived for several days even when ciprofloxacin was added to exponentially growing cultures. Furthermore, a short starvation pulse resulted in a long-lasting effect on survival. The revealed killing dynamics show that long-term persistence is not caused by simply one subpopulation of antibiotic-tolerant cells, but is a complex, time-dependent phenomenon.

Keywords: Bacterial persistence, Stress response, Growth physiology, Antimicrobials, (p)ppGpp

1. Introduction

Bacterial populations are quickly decimated during a typical antibiotics assault. Within a few generation times, the far majority of cells will be dead. However, it is typically recommended to use extended durations of treatment, ranging from several days to months, prolonging the exposure time of bacterial pathogens to the antibiotic [1]. The WHO is now considering the benefits of shortening the duration of antibiotics administration while still keeping the treatment effective, due to concerns of increasing antibiotic resistance occurring as a

consequence of increased exposure [1]. To find the optimal treatment duration, one needs to understand the killing dynamics of bacteria when exposed to antibiotics, especially the bacterial cells surviving for longer times. The long-term survivors are typically referred to as persister cells, a subgroup of cells that survive antibiotics for an extended period compared to the average of the population, but have not acquired mutations that make them resistant to the antibiotic [2, 3]. Most research on persister cells is done within the well-defined conditions of the laboratory, but despite these strongly simplified conditions, and more than seventy years of research, laboratory persisters are still far from understood [4, 5].

One pending question is whether and how much persisters form spontaneously during the exponential growth phase. Such persisters are called type-II [3] or

*correspondence: mas@bio.ku.dk

**correspondence: mitarai@nbi.ku.dk

¹The Niels Bohr Institute, University of Copenhagen

²The Department of Biology, University of Copenhagen

spontaneous [4] persisters. It was repeatedly shown that stress-triggered (or type I) persisters are formed in high numbers during the stationary phase, but research on spontaneous persister formation during the exponential phase is sparse [4]. The research has mostly been confounded by a lack of careful attention to the presence of stationary phase cells carried over from the starter cultures, which artificially elevated the persister fraction of exponential cultures [5, 6, 4]. One carefully executed study showed that no *E. coli* persister cells were formed during fast exponential growth in rich medium [7], whereas other studies merely showed reduced levels during exponential growth [3, 5]. A benefit of analysing the exponential growth phase is the well-defined physiology of this state [8, 9]. This makes it possible to vary the growth physiology in a controlled manner, especially by varying the growth rate through culturing bacteria in media of different nutrient quality. It was previously shown that the bacterial growth rate strongly correlates with the death rate during the initial period of killing with beta-lactams [10, 11, 12]. This poses the additional questions of how the growth rate at the time of antibiotics exposure affects the short- and long-term killing dynamics.

The current standard for persister identification at the population level is a biphasic killing curve, where two time-scales are identified in the time-kill curves [4]. Persisters are identified as the subpopulation with a second, slower killing rate than the rapid death rate of the primary population. If only two timescales are present in the killing dynamics, the population survival time can be extrapolated from the second slow killing rate. Notably, a few studies have reported deviations from the biphasic model. In one case, a heavier tail of long-term survivors was observed, possibly due to low level contamination of the persister assay [13]. In another case, a powerlaw tail in the short-term killing dynamics was observed [14]. Altogether, the long-term survival of the antibiotics-tolerant subpopulation is gravely understudied. Most *in vitro* lab research on persisters of fast growing bacteria as *E. coli* is carried out for three to five hours [15, 7, 5], though some studies increase the exposure time to 24–50 hours [2, 16, 17, 3]. Investigation of long-term survival beyond the typical five-hour persister assay might reveal new insights into bacterial killing dynamics that are relevant for the week-long antibiotics treatment of bacterial infections recommended by the WHO [1].

Lastly, the molecular mechanism(s) of persister formation is still unknown [18, 12]. Many intracellular components have been proposed to play a role [19, 15, 20, 21, 22, 23], but so far no single mechanism

convincingly explains persister formation. In fact, bacterial persistence presents as a very complex and diverse problem, where the survival fraction could be composed of different subpopulations.

Despite the complexity of persistence, it has been established that stationary phase cultures contain a greater persister fraction than exponentially growing cultures [5]. Stationary phase bacteria may refer to bacteria in a multitude of different physiological states, but is typically associated with starvation stress [24]. Furthermore, the second messenger (p)ppGpp, which accumulates during starvation responses was frequently shown to correlate positively with persistence formation [17, 25, 5, 21]. Hence, it is critical for persistence research to understand the degree to which (p)ppGpp levels affect persistence, and under which circumstances.

The present study investigates persistence in the balanced exponential growth phase where (p)ppGpp levels are relatively low and correlate inversely with the growth rate. It deals with whether *E. coli* forms spontaneous persisters in the exponential phase, their dependence on the growth rate, how long they survive and how their formation relates to (p)ppGpp levels. We followed the long-term survival of *E. coli* K12 populations exposed to a lethal concentration of ciprofloxacin for one week. The growth rate of the *E. coli* population at the time of antibiotics exposure was varied using growth medium with either of two different carbon sources. In addition, a knockout strain was constructed in the wild-type background, removing the gene *relA* and, thus, introducing a (p)ppGpp synthesis deficiency. Furthermore, we compared the killing dynamics with and without a short carbon-starvation period immediately prior to the killing assay. The starvation pulse had a considerable influence on persister formation and had a very long memory effect on the survival of the population.

2. Results

Long-term persister assay of exponentially growing cells

First, we investigated whether long-term persister cells form during exponential growth in glucose minimal medium. We treated balanced cultures of *E. coli* K-12 with ciprofloxacin and monitored the killing dynamics for one week of antibiotics treatment. Balanced growth was obtained by culturing the cells for more than twenty doubling times in the target medium at 37 °C, keeping the cell density of the culture below an OD₄₃₆ of 0.3 by repeated back-dilutions. Cultures were then treated for a week with 10 µg/mL ciprofloxacin

Figure 1: Killing dynamics of exponential phase *E. coli* persisters. Killing dynamics for exponential phase persisters. The bacteria were grown in glucose minimal medium. All three biological replicates are shown for each data point. The black line is the geometrical mean. Each separate phase of a triphasic fit are shown respectively as cyan, green and blue dashed lines. The second phase of a biphasic fit to the first seven hours is shown as a brown dotted line.

and their killing dynamics were monitored by repeated platings of culture aliquots on antibiotics-free growth medium (See methods for detail).

E. coli persisters were formed during exponential balanced growth in glucose minimal medium, as seen in Fig. 1. There was a fast initial killing at a rate of about $1/0.3 \text{ (h}^{-1}\text{)}$, with a slower killing rate already after two hours. When we fit a biphasic curve (summation of two exponential functions) to the data up to 7 hours, the second phase of killing is at a rate of about $1/2.3 \text{ (h}^{-1}\text{)}$, shown by a brown line in the inset in Fig. 1. However, for longer times, this fit significantly underestimates the survival time of the bacterial population (Fig. 1). In other words, there is an even slower phase of killing, extending from seven hours to four days. Lastly, from day five to day seven, the remaining cells were killed at a very slow rate, however, this part of the data is less reliable due to the small numbers of recovered colonies.

To identify the various phases in a quantitative manner, the data was fitted with a sum of exponentials, and the appropriate number of exponentials was chosen with a χ^2 test (see SI Appendix section 2). The test rejects a double exponential as a good fit, but accepts the hypothesis of a triple exponential. The best fit obtained was the first phase of killing at a rate $1/0.3 \text{ (h}^{-1}\text{)}$ (a cyan dashed line), the second phase of killing at a rate of $1/8 \text{ (h}^{-1}\text{)}$ (a green dashed line), and the third phase of killing at a rate of $1/57 \text{ (h}^{-1}\text{)}$ (a blue dashed line). The long-term killing dynamics thus had more than two time-scales,

Figure 2: Killing dynamics in different medium with and without the *relA* gene. Each line represents three biological replicates. (A) The wildtype killing dynamics in either glycerol or glucose minimal medium. (B) The $\Delta relA$ killing dynamics in either glycerol or glucose minimal medium. (C) Killing dynamics in glucose minimal medium for comparison between the wildtype and the $\Delta relA$ strain. The $\Delta relA$ mutant had more survivors than the wildtype from day 4 and later, but there was no statistically significant difference at any of the timepoints (SI Appendix section 2). (D) Killing dynamics in glycerol minimal medium for comparison between the wildtype and the $\Delta relA$ strain. (E) Survival fraction after two hours of ciprofloxacin treatment compared to the growth rate prior to antibiotics treatment.

and the dominant phase of killing is only apparent after several days of measurement.

The exponential phase growth rate determines many aspects of bacterial physiology, including the macromolecular composition [8, 26]. The growth rate at the time of antibiotics exposure has previously been linked to short-term survival of antibiotics [10, 11], and could also affect long-term survival. For that reason, the long-term killing assay was repeated with glycerol as the carbon source, which strongly affected the wildtype growth rate. In glucose minimal medium, the wildtype doubling time was $50 \pm 1.4 \text{ min}$, while it was $106 \pm 3.0 \text{ min}$ in glycerol minimal medium. This difference had an impact on the initial phase of killing for up to seven hours, showing slower killing and a higher survivor fraction in glycerol (Fig. 2A inset). However, the long-term killing curve in glycerol minimal medium was not merely a decelerated version of the killing curve in glucose minimal medium. Though more cells survive in glycerol than in

glucose for up to two days (Fig. 2A), the survival curve in glycerol medium at 1 to 2 days had a steeper drop than in glucose medium, resulting in comparable surviving fractions after 3 days. In fact, in two of the three biological replicates of wildtype cultures grown in glycerol, almost no survivors were observed after 3 days of killing (see SI Appendix section 1). Overall, the wildtype killing dynamics in glycerol had more than two time scales, with a best fit of four separate phases of killing (SI Appendix section 2).

Deletion of *relA* affects the killing dynamics

In glycerol minimal medium, the steady state (p)ppGpp level of the wildtype strain is higher than in glucose minimal medium [27]. Because the (p)ppGpp level has been frequently associated with persister formation, we next aimed at investigating the effect of the initial (p)ppGpp level on long-term survival. *E. coli* encodes the primary (p)ppGpp synthetase RelA and the secondary (p)ppGpp synthetase SpoT, the latter of which is bifunctional as a (p)ppGpp hydrolase. The nutrient-dependent steady-state growth rates are inversely related to the concentrations of (p)ppGpp, both for *relA*⁺ and *relA*⁻ strains [27], and for many carbon sources, the growth rates and (p)ppGpp levels of *relA*⁺/*relA*⁻ strain pairs are indistinguishable due to (p)ppGpp synthesis by SpoT. However, in low energy carbon sources like glycerol or acetate, SpoT produces insufficient (p)ppGpp to suppress the growth rate when RelA is missing, leading to an enhanced growth rate of the RelA mutant strain relative to the wildtype [27]. We constructed a *ΔrelA* mutant to clarify the role of (p)ppGpp in the killing dynamics.

As expected, the difference between the growth rate in glucose and in glycerol minimal medium was smaller for the *ΔrelA* strain, with a doubling time of 47 ± 1.5 min in glucose minimal medium and 74 ± 1.6 min in glycerol minimal medium. The short term (up to 4 hours) survival under antibiotics exposure was also correlated with the growth rate in the *ΔrelA* strain (Fig. 1B inset).

Like wildtype cells, the *ΔrelA* mutant formed long-term survivors in both glucose and glycerol minimal medium with more than two phases of killing (Fig. 2B, see SI Appendix section 2 for statistical analysis). In glucose, the two strains grew at similar rates, and the early killing dynamics of the *ΔrelA* mutant were very similar to that of the wildtype (Fig. 2C). In glycerol, the faster-growing *ΔrelA* mutant showed a significantly lower level of persisters than the wildtype in the initial phase of killing up to one day (Fig. 2D, see SI Appendix section 2), indicating the importance of growth rate, or

ppGpp level, for persister formation in this phase. Interestingly, the long-term survival of the *ΔrelA* mutant and wildtype were comparable at later times (after 3 days) in glycerol medium, and somewhat higher than the wildtype strain in glucose medium. Thus, survival in the long term is not simply dependent on the population growth rate at the time of antibiotics exposure.

A starvation pulse prior to the antibiotic application affects the long-term persistence of wildtype cells in glucose minimal medium

A sudden downshift of the carbon source is known to give a spike of the (p)ppGpp level in the wildtype strain just after the downshift, while the spike is significantly lower in a RelA strain [28]. We then wondered if a short pulse of carbon source starvation to the exponentially growing cells prior to the killing assay will give a quantifiable difference in the long-term persistence between the wildtype strain and the *ΔrelA* strain.

In order to test this, part of the cultures in balanced growth were filtered into growth medium without a carbon source and starved for 1 hour, before the carbon source was replenished (Figure 3A). Figure 3B shows that the 1-hour starvation pulse resulted in a quick rise of the ppGpp level peaking at about 15 min after the downshift for the wildtype strain in glucose medium, while only a mild increase of the ppGpp level was seen in the *ΔrelA* strain. Antibiotics were added simultaneously with the carbon source replenishment (Fig. 3A, see methods for details). Remarkably, the short carbon starvation prior to the addition of antibiotics had long-term effects on the killing dynamics. This was especially visible for the wildtype strain grown in glucose minimal medium, where the brief starvation period reproducibly resulted in almost 100-fold more persisters for up to four days. The difference was abolished by removing *relA*, as seen in Fig. 4B; the *ΔrelA* strain only exhibited increased survival for the first six hours following starvation. As such, the long-term memory of the starvation pulse is seemingly a *relA*-dependent effect. However, the *ΔrelA* strain in the steady state growth in glucose had more long-term survivors than the wild-type strain, and the downshift brought the wildtype strain survival fraction to a level similar to the *ΔrelA* strain.

The effect of the downshift was smaller in the glycerol medium (Fig. 4CD). In the wildtype strain, the average persister level with downshift was higher up to 4 days, but the statistical significance of the difference was confirmed only up to 7 hours due to the larger data scatter for later time points (Fig. 4C). The effect of

Figure 3: The experimental setup for the long-term persistence assay. (A) Cells were in balanced exponential growth prior to the killing assay. A part of the culture was filtered and then resuspended in medium with or without the carbon source, and incubated for 1 hour. After 1 hour, the ciprofloxacin was added to both of the samples, and at the same time the carbon source was added to the culture that has been starved for 1 hour. The first sample was taken just before the additions. After that, samples were taken at 2, 4, 6, 7, 21+24-n hours after the antibiotic addition for $n \in [0; 6]$. The samples were washed and plated on agar plates containing the target medium. (B) The 1-hour time course of ppGpp level for the culture grown in the glucose medium, filtered, and resuspended in fresh medium without carbon source. The time zero is the time of the resuspension. The wild type (blue circles) shows clear peak around 15 minutes after the resuspension. $\Delta relA$ strain shows only mild increase of the ppGpp level.

downshift disappeared faster in the $\Delta relA$ strain already after 2 days (Fig. 4D).

3. Discussion

We expanded the understanding of bacterial killing dynamics with a new long-term persister assay. The use of minimal medium facilitated the formation of long-term persisters during exponential growth, in contrast to growth and killing in rich medium [7]. Spontaneous persisters were observed during the exponential growth phase, both in glucose and glycerol minimal medium, and in some cases they survived at least one week. This long-term survival does not require *relA*, although the residual (p)ppGpp synthesized by SpoT is likely necessary. In fact, there is an increase in long-term survival of the $\Delta relA$ mutant in glucose.

We have shown that a one-hour starvation pulse prior to the addition of the antibiotic affects long-term survival. This finding is consistent with a previous study of a temporal nitrogen downshift prior to antibiotics treatment, which was shown to elevate the persister level at 24 hours in a *relA* dependent manner [29]. Our study demonstrated that the memory can be remarkably long-lasting, as one-hour carbon starvation gave an increase in survival for at least 4 days in the wildtype strain grown in glucose medium. The molecular mechanism underlying this long-term memory is yet to be investigated.

Figure 4: The effect of a short starvation-pulse on the survival dynamics. All strains and conditions are shown with and without the downshift. The black line is with the downshift, the colored line is without the downshift. The magenta dots represent statistically different datapoints as tested by an unequal variances two-sided t-test. The cyan interval illustrates the downshift period prior to the killing assay. (A) The wildtype in glucose. A downshift in glucose significantly enhances wildtype survival for four days and seems to be increased for up to all seven days. (B) The $\Delta relA$ mutant in glucose. (C) The wildtype in glycerol. (D) The $\Delta relA$ mutant in glycerol.

304 tigated, but in all likelihood it is linked to the abrupt
305 RelA-mediated rise in (p)ppGpp upon starvation, since
306 the starvation-pulse effect on long-term survival was
307 abolished in the $\Delta relA$ mutant. In further support of this
308 hypothesis, there was no long-term effect when glyc-
309 erol was used as the carbon source, which could be due
310 to the high basal level of (p)ppGpp in glycerol relative
311 to glucose minimal medium [8]. The sensitivity of the
312 survival fraction to the rather short perturbation may be
313 consistent with the idea that there is a threshold in some
314 molecule concentration to determine if the cell becomes
315 persister or not [30, 31], since a small perturbation can
316 have a major impact on the occurrence of rare expres-
317 sion patterns that exceed a extreme threshold [32]. This
318 observation also alerts us that small perturbations in the
319 experimental procedure may strongly affect the result of
320 persister assays.

321 This study shows that the long-term killing of *E. coli*
322 in ciprofloxacin is not adequately described by biphasic
323 dynamics. At least three phases of killing were present
324 in the data. Thus, despite the emphasis on a biphasic
325 behaviour to define persistence [4], we propose a third,
326 or even fourth, phase of killing that may even be more
327 clinically relevant. The presence of additional phases
328 also means that the population survival time will be un-
329 derestimated by predictions from the biphasic killing as-
330 sumption. Indeed, it is not sufficient to measure killing
331 dynamics for only five hours and then extrapolate the
332 population survival time from there.

333 The population growth rate was found to be posi-
334 tively correlated with the killing rate in the initial phase.
335 However, the correlation was diminished in the longer
336 term survival, and lost in the third phase of killing. In
337 shorter persister assays, a difference in growth rate, such
338 as between different mutants, might strongly confound
339 results when analyzing persistence fractions. These dif-
340 ferences seem smaller and less relevant in later phases
341 of killing.

342 This investigation of long-term killing dynamics has
343 added to the concept of bacterial persistence as a com-
344 plex phenomenon. During long-term antibiotics treat-
345 ment, different mechanisms could account for bacterial
346 survival on different timescales (an hour, a day, a week),
347 although the (p)ppGpp level at the time of initial expo-
348 sure to the antibiotic seems to be important in all cases.
349 As such, persistence seems to be a time-dependent phe-
350 nomenon, where different survival mechanisms account
351 for different bacterial life spans. The presence of sev-
352 eral phases in the killing dynamics begs the question if
353 a more extended concept should replace the simplified
354 concept of bacterial persistence.

4. Methods and Materials

Strains

357 The strain MAS1081 (MG1655 *rph*⁺ *gatC*⁺ *glpR*⁺)
358 was used as the wildtype [33]. The $\Delta relA$, MAS1191,
359 is MAS1081 made $\Delta relA251 :: Kan$ by P1 transduction
360 from CF1651 [34] followed by selection on kanamycin.

Long-term killing assay

361 A single colony of the *E. coli* strain grown on an
362 agar plate was incubated overnight in the target medium
363 (MOPS minimal medium with either glucose or glyc-
364 erol as the carbon source [35]. See the SI Appendix sec-
365 tion 1 for the recipe). The overnight culture was diluted
366 1 : 10⁷ in 10 mL target medium in a 100 mL Erlenmeyer
367 flask. The flask was continuously shaken at 160 RPM
368 in a 37°C room. Hours later, the culture was diluted
369 further, at least 1 : 10², in 100 mL preheated medium
370 in a 1L Erlenmeyer flask, reaching a total dilution of at
371 least 1 : 10⁹. Once the OD₄₃₆ reached a detectable level
372 of 0.01, at least 5 samples were measured at different
373 timepoints to establish a growth rate. The OD₄₃₆ was
374 consistently kept below 0.3. A part of the culture, typ-
375 ically around 20 mL, was filtered, and the cells on the
376 filter were resuspended in 40 mL of the target medium
377 (w/o carbon source) in a 300 mL Erlenmeyer flask. As
378 such, both the control and the starved culture were ex-
379 posed to filtration and resuspension. The medium vol-
380 ume never exceeded 14 percent of the flask volume. The
381 starvation was verified by measuring OD₄₃₆, to confirm
382 either increase in biomass (control) or no growth (starv-
383 ing culture). See the SI Appendix section 1 for confir-
384 mation of the downshift. After one hour of starvation, a
385 sample was taken immediately before ciprofloxacin (10
386 µg/mL) was added along with the carbon source, that
387 was replenished to end the downshift. Samples were
388 taken at times 2,4,6/7, 21+24-*n* hours for *n* ∈ [0; 6]. The
389 samples were put on ice for a few minutes and then cen-
390 trifuged for ten minutes at 4°C at 10,000g. The super-
391 natant was removed and the cell pellet was resuspended
392 in room temperature MOPS buffer with no supplements.
393 The sample was diluted appropriately, never more than
394 1 : 100 per step, corresponding to 10 µL in 990 µL.
395 The sample was plated with 200 µL per plate on mini-
396 mal medium plates containing the target medium. The
397 plates were kept at 37°C for at least one week and all
398 colonies were counted. The whole experiment is illus-
399 trated in figure 3.

Controls

401 After each completed experiment (7 days), 200 µL
402 of culture, still containing antibiotics, was spread on a
403

plate with the target medium. This was left at 37°C for at least 7 days, and no growth confirmed the absence of a growing resistant culture in the flask. In addition, the activity of the antibiotics in the culture was tested after seven days by dropping 20 μ L on a lawn of growing *E. coli*.

Analysis of time-scales in the killing dynamics

Time-scales in the killing dynamics were statistically identified by fitting a sum of exponential functions to the data. The model with the least number of parameters, that could not significantly be rejected, was then chosen [36]. The functional form of the models was a sum of exponential functions. The number of exponential functions in the sum corresponded to the number of time-scales. A biphasic killing curve would for example be well fitted by the sum of two exponentials. The χ^2 -test for goodness of fit is used for identifying the appropriate model using the Minuit minimization software [37, 38] (see the SI Appendix section 2 for specifics of the analysis).

Statistical analysis

All killing curves are based on three biological replicates. The mean is determined as the mean of the logarithmic values, which corresponds to the geometric mean. This is done to get a more adequate mean-value representation in log-space. The uncertainties are also calculated as the standard deviations of the log transformed values. Whenever a datapoint had the value zero, which happened frequently, that value was replaced with the detection limit, to get a sensible value in log-space. An unequal variance two-sided t-test was used to determine significant differences between two datapoints at the same timepoint ($P < 0.05$).

ppGpp measurements

Cultures were grown for two generations in the presence of 75 μ Ci/mL 32P-phosphate at a total phosphate concentration of 0.33 mM. At the time of starvation, cultures were filtered, washed in medium without glucose and phosphate and resuspended in medium without glucose but containing 32P at the same specific activity as during growth. These steps were performed at 37°C and lasted less than 2 min. For determination of the nucleotide pools, 100 μ L of culture was harvested into 20 μ L 2M formic acid at 0°C. After centrifugation the nucleotides in the supernatant were separated by chromatography on polyethyleneimine-cellulose plates. The activities of the individual spots were quantified by PhosphoImager scans (Typhoon Phosphor Imager

FLA7000 (GE Healthcare)) of the plates. The specific activity of the signal was determined from a medium sample from the individual cultures spotted onto the same brand of plates.

Acknowledgments

MSS and NM thank S. Semsey for fruitful discussions. This work was supported by the Danish National Research Foundation (DNRF120) and the Independent Research Fund Denmark (8049-00071B and 8021-00280A).

References

- [1] M. D. Pezzani, G. Be, P. Cattaneo, A. Zaffagnini, F. Gobbi, P. Rodari, Z. Bisoffi, E. Tacconelli, Evidence based review on optimal duration of antibiotic therapy for bacterial infections to support antimicrobial stewardship recommendations WHO Secretariat Nicola Magrini, Secretary of the Expert Committee on Selection and Use of Essential Medicines; In (2019) 1–28.
- [2] J. W. Bigger, Treatment of Staphylococcal Infections With Penicillin By Intermittent Sterilisation, *The Lancet* 244 (1944) 497–500. doi:10.1016/S0140-6736(00)74210-3.
- [3] N. Q. Balaban, J. Merrin, R. Chait, L. Kowalik, S. Leibler, Bacterial Persistence as a Phenotypic Switch, *Science* 305 (2004) 1622–1625. doi:10.1126/science.1099390.
- [4] N. Q. Balaban, S. Helaine, A. Camilli, J. J. Collins, J.-M. Ghigo, W.-D. Hardt, A. Harms, M. Heinemann, Definitions and guidelines for research on antibiotic persistence, *Nature Reviews Microbiology* 17 (2019) 441–448. URL: <http://dx.doi.org/10.1038/s41579-019-0196-3>. doi:10.1038/s41579-019-0196-3.
- [5] A. Harms, C. Fino, M. A. Sørensen, S. Semsey, K. Gerdes, Prophages and Growth Dynamics Confound Experimental Results with Antibiotic-Tolerant Persister Cells, *mBio* 8 (2017) 1–18. doi:10.1128/mBio.01964-17.
- [6] A. Brauner, O. Fridman, O. Gefen, N. Q. Balaban, Distinguishing between resistance, tolerance and persistence to antibiotic treatment, *Nature Reviews Microbiology* 14 (2016) 320–330. URL: <http://dx.doi.org/10.1038/nrmicro.2016.34>. doi:10.1038/nrmicro.2016.34.
- [7] I. Keren, N. Kaldalu, A. Spoering, Y. Wang, K. Lewis, Persister cells and tolerance to antimicrobials, *FEMS Microbiology Letters* 230 (2004) 13–18. doi:10.1016/S0378-1097(03)00856-5.
- [8] H. Bremer, P. P. Dennis, Modulation of Chemical Composition and Other Parameters of the Cell by Growth Rate, *Escherichia coli and Salmonella: cellular and molecular biology* 2 (1987) 1527–1542. doi:10.1046/j.1365-2486.2000.06017.x.
- [9] M. Scott, C. W. Gunderson, E. M. Mateescu, Z. Zhang, T. Hwa, American Association for the Advancement of Science, *Science* 330 (2010) 1099–1102. doi:https://doi.org/10.1126/science.1192588.
- [10] E. Tuomanen, R. Cozens, W. Tosch, O. Zak, A. Tomasz, The rate of killing of *Escherichia coli* by beta-lactam antibiotics is strictly proportional to the rate of bacterial growth., *Journal of general microbiology* 132 (1986) 1297–304. doi:10.1099/00221287-132-5-1297.
- [11] A. J. Lee, S. Wang, H. R. Meredith, B. Zhuang, Z. Dai, L. You, Robust, linear correlations between growth rates and β -lactam-mediated lysis rates (2018) 1–6. doi:10.1073/pnas.1719504115.

- [12] M. H. Pontes, E. A. Groisman, Slow growth determines non-heritable antibiotic resistance in *Salmonella enterica*, *Science Signaling* 12 (2019) 1–11. doi:10.1126/scisignal.aax3938.
- [13] E. M. Windels, Z. B. Meriem, T. Zahir, K. J. Verstrepen, P. Hersen, B. V. den Bergh, J. Michiels, Isolation of persisters enabled by β -lactam-induced filamentation reveals their single-cell awakening characteristics, *Communications Biology* (2019) 600700. URL: <https://www.biorxiv.org/content/10.1101/600700v1>. doi:10.1101/600700.
- [14] E. Simsek, M. Kim, Power-law tail in lag time distribution underlies bacterial persistence 116 (2019). doi:10.1073/pnas.1903836116.
- [15] F. Goormaghtigh, N. Fraikin, M. Putrins, T. Hallaert, V. Hauyryliuk, A. Garcia-Pino, A. Sjödin, S. Kasvandik, K. Udekku, T. Tenson, N. Kaldalu, L. Van Melderen, Reassessing the Role of Type II Toxin-Antitoxin Systems in Formation of *Escherichia coli* Type II Persister Cells, *mBio* 9 (2018) 1–14.
- [16] F. Goormaghtigh, L. V. Melderen, Single-cell imaging and characterization of *Escherichia coli* persister cells to ofloxacin in exponential cultures, *Science Advances* 5 (2019).
- [17] S. Helaine, A. M. Cheverton, K. G. Watson, L. M. Faure, S. A. Matthews, D. W. Holden, Internalization of *Salmonella* by Macrophages Induces Formation of Nonreplicating Persisters, *Science* 343 (2014) 204–209.
- [18] A. Harms, E. Maisonneuve, K. Gerdes, Mechanisms of bacterial persistence during stress and antibiotic exposure, *Science* 354 (2016). doi:10.1126/science.aaf4268.
- [19] E. A. Zalis, A. S. Nuxoll, S. Manuse, G. Clair, L. C. Radlinski, B. P. Conlon, J. Adkins, K. Lewis, Stochastic Variation in Expression of the Tricarboxylic Acid Cycle Produces Persister Cells, *mBio* 10 (2019) 1–10.
- [20] Y. Shan, A. B. Gandt, S. E. Rowe, J. P. Deisinger, B. P. Conlon, K. Lewis, ATP-Dependent Persister Formation in *Escherichia coli*, *mBio* 8 (2017).
- [21] M. S. Svenningsen, A. Veress, A. Harms, N. Mitarai, S. Semsey, Birth and Resuscitation of (p) ppGpp Induced Antibiotic Tolerant Persister Cells, *Scientific Reports* 9 (2019). doi:10.1038/s41598-019-42403-7.
- [22] R. Yamasaki, S. Song, M. Benedik, T. Wood, Persister Cells Resuscitate Using Membrane Sensors that Activate Chemotaxis, Lower cAMP Levels, and Revive Ribosomes, *SSRN Electronic Journal* 23 (2019) 100792. URL: <https://doi.org/10.1016/j.isci.2019.100792>. doi:10.2139/ssrn.3399578.
- [23] R. C. Molina-Quiroz, C. Silva-Valanzuela, J. Brewster, E. Castro-Nallar, S. B. Levy, A. Camilli, Cyclic AMP Regulates Bacterial Persistence through Repression of the Oxidative Stress Response and SOS-Dependent DNA Repair in Uropathogenic *Escherichia coli*, *mBio* 9 (2018) e02144–17. doi:10.1128/mBio.02144-17.
- [24] T. Nyström, Stationary-Phase Physiology, *Annual Review of Microbiology* 58 (2004) 161–181. doi:10.1146/annurev.micro.58.030603.123818.
- [25] J. L. Radzikowski, H. Schramke, M. Heinemann, Bacterial persistence from a system-level perspective, *Current Opinion in Biotechnology* 46 (2017) 98–105. URL: <http://dx.doi.org/10.1016/j.copbio.2017.02.012>. doi:10.1016/j.copbio.2017.02.012.
- [26] S. Hui, J. M. Silverman, S. S. Chen, D. W. Erickson, M. Basan, J. Wang, T. Hwa, J. R. Williamson, Quantitative proteomic analysis reveals a simple strategy of global resource allocation in bacteria, *Molecular Systems Biology* 11 (2015) e784–e784. doi:10.15252/msb.20145697.
- [27] J. Ryals, R. Little, H. Bremer, Control of rRNA and tRNA syntheses in *Escherichia coli* by guanosine tetraphosphate, *Journal of Bacteriology* 151 (1982) 1261–1268. doi:10.1128/jb.151.3.1261-1268.1982.
- [28] R. A. Lazzarini, M. Cashel, J. Gallant, On the regulation of guanosine tetraphosphate levels in stringent and relaxed strains of *Escherichia coli*, *Journal of Biological Chemistry* 246 (1971) 4381–4385.
- [29] D. R. Brown, Nitrogen starvation induces persister cell formation in *Escherichia coli*, *Journal of Bacteriology* 201 (2019). doi:10.1128/JB.00622-18.
- [30] E. Rotem, A. Loinger, I. Ronin, I. Levin-Reisman, C. Gabay, N. Shoshitaishvili, O. Biham, N. Q. Balaban, Regulation of phenotypic variability by a threshold-based mechanism underlies bacterial persistence, *Proceedings of the National Academy of Sciences of the United States of America* 107 (2010) 12541–12546. doi:10.1073/pnas.1004333107.
- [31] J. E. Michiels, B. Van den Bergh, N. Verstraeten, J. Michiels, Molecular mechanisms and clinical implications of bacterial persistence, *Drug Resistance Updates* 29 (2016) 76–89. doi:10.1016/j.drug.2016.10.002.
- [32] M. S. Svenningsen, S. Semsey, N. Mitarai, Gene Expression Changes with Minor Effects on the Population Average Have Major Effects on the Occurrence of Cells with Extreme Protein Concentrations, *mSphere* 4 (2019) 1–6. doi:10.1128/msphere.00575-18.
- [33] B. Gummeson, S. A. Shah, A. S. Borum, M. Fessler, N. Mitarai, M. A. Sørensen, S. L. Svenningsen, Valine-induced isoleucine starvation in *Escherichia coli* k-12 studied by spike-in normalized rna sequencing, *Frontiers in Genetics* 11 (2020) 144.
- [34] H. Xiao, M. Kalman, K. Ikehara, S. Zemel, G. Glaser, M. Cashel, Residual guanosine 3',5'-bispyrophosphate synthetic activity of relA null mutants can be eliminated by spoT null mutations, *Journal of Biological Chemistry* 266 (1991) 5980–5990.
- [35] F. C. Neidhardt, P. L. Bloch, D. F. Smith, Culture medium for enterobacteria, *Journal of Bacteriology* 119 (1974) 736–747. doi:10.1128/jb.119.3.736-747.1974.
- [36] R. J. Barlow, STATISTICS: A GUIDE AND REFERENCE TO THE USE OF STATISTICAL METHODS IN THE PHYSICAL SCIENCES, Wiley, 1993.
- [37] F. James, M. Roos, Minuit – A System for Function Minimization and Analysis of the Parameter Errors and Correlations, *Computer Physics Communications* 10 (1975) 343–367.
- [38] iminuit – A Python interface to Minuit, 2020.