Crosstalk in oxygen homeostasis networks: SKN-1/NRF inhibits the HIF-1 hypoxia-inducible factor in *Caenorhabditis elegans*

Dingxia Feng^{*¶}, Zhiwei Zhai[¶], Zhiyong Shao, Yi Zhang, and Jo Anne Powell-Coffman*

Department of Genetics, Development and Cell Biology, Iowa State University, Ames, Iowa,
United States of America

*corresponding author:

E-mail: japc@iastate.edu (JP)

E-mail: fengdi@indiana.edu (DF)

[¶]co-first authors contributed equally

Abstract

During development, homeostasis, and disease, organisms must balance responses that allow adaptation to low oxygen (hypoxia) with those that protect cells from oxidative stress. The evolutionarily conserved hypoxia-inducible factors are central to these processes, as they orchestrate transcriptional responses to oxygen deprivation. Here, we employ genetic strategies in *C. elegans* to identify stress-responsive genes and pathways that modulate the HIF-1 hypoxia-inducible factor and facilitate oxygen homeostasis. Through a genome-wide RNAi screen, we show that RNAi-mediated mitochondrial or proteasomal dysfunction increases the expression of hypoxia-responsive reporter *Pnhr-57:GFP* in *C. elegans*. Interestingly, only a subset of these effects requires *hif-1*. Of particular importance, we found that *skn-1* RNAi increases the expression of hypoxia-responsive reporter *Pnhr-57:GFP* and elevates HIF-1 protein levels. The SKN-1/NRF transcription factor has been shown to promote oxidative stress resistance. We present evidence that the crosstalk between HIF-1 and SKN-1 is mediated by EGL-9, the prolyl hydroxylase that targets HIF-1 for oxygen-dependent degradation. Treatment that induces SKN-1, such as heat, increases expression of a *Pegl-9:GFP* reporter, and this effect requires *skn-1* function and a putative SKN-1 binding site in *egl-9* regulatory sequences. Collectively, these data support a model in which SKN-1 promotes *egl-9* transcription, thereby inhibiting HIF-1. We propose that this interaction enables animals to adapt quickly to changes in cellular oxygenation and to better survive accompanying oxidative stress.

Introduction

Oxygen homeostasis has profound effects on health and fitness. Oxygen serves as the terminal electron acceptor in the oxidative phosphorylation processes that generate energy for life. When

oxygen levels are low (hypoxia), cells and tissues must adapt quickly by increasing oxygen delivery, adjusting the levels of key metabolic enzymes, and limiting the accumulation of misfolded proteins. While oxygen is essential, it is also highly reactive. The reactive oxygen species (ROS) generated by cellular metabolism and signaling processes can damage macromolecules, and excess ROS are thought to contribute to cellular aging and deterioration [1-3]. One of the central challenges of aerobic life is to coordinate the biological networks that control disparate aspects of oxygen homeostasis.

This balance between surviving hypoxic stress and mitigating the potential damage caused by reactive oxygen species is especially important in cardiovascular development and disease. When ischemia blocks circulation to a mammalian tissue, oxygen levels drop, and cells induce hypoxia-inducible transcription factors (HIFs). Upon reperfusion and reoxygenation of the tissue, mammalian cells respond by rapidly degrading HIF and inducing the NRF2 transcription factor [4, 5]. However, intermittent hypoxia has been shown to induce both HIF-1 α and NRF2 [6, 7]. NRF2 activates phase II detoxification genes to mitigate the effects of oxidative insults [8, 9].

Although mammalian HIF and NRF2 share some common target genes such as aldehyde dehydrogenase 1A1 or heme oxygenase-1 HO-1, the genes induced by re-oxygenation are largely distinct from those that respond to oxygen deprivation [5, 10, 11]. Crosstalk between these two pathways is complex and context specific in mammals [12], as these important transcription factors facilitate the rapid changes in gene expression needed to limit reperfusion injury and regulate oxygen-dependent developmental processes.

C. elegans has been proven to be an excellent model system for studying the regulatory networks that govern oxygen homeostasis. The *C. elegans* genome encodes a single hypoxia-inducible factor alpha subunit (HIF-1), and the *hif-1* gene has been shown to have important roles in stress

response and in aging [13-17]. HIF protein levels and HIF activity are tightly regulated. When oxygen is abundant, the HIF alpha subunit is hydroxylated by the PHD/EGL-9 enzymes. Once modified, HIF α protein interacts with the Von Hippel-Lindau tumor suppressor (VHL) and is targeted for ubiquitination and proteasomal degradation [18-23]. Thus, in hypoxic conditions, HIF-1 protein is stable, and the transcription factor complex can activate the expression of a battery of genes that enable adaptation to low oxygen. This pathway for oxygen-dependent degradation of HIF protein is evolutionarily conserved. The *C. elegans* *hif-1*, *aha-1*, *egl-9*, and *vhl-1* genes are orthologous to mammalian *HIF α* , *HIF β* , *PHD*, and *VHL*, respectively [24-26]. The targets of *C. elegans* HIF-1 include *egl-9* and *rhy-1*, genes that inhibit HIF-1 expression and activity [27-30]. In wild-type animals, these negative feedback loops keep HIF-1 activity in check and limit the potentially adverse effects of HIF-1 over-activation. The *C. elegans* *skn-1* gene is homologous to mammalian *NRF1/2/3* [31]. SKN-1 regulates the expression of a battery of genes with cytoprotective functions, including phase II detoxification genes [32, 33]. SKN-1 is activated by a range of stresses or toxicants that cause oxidative stress, and SKN-1 promotes resistance to these insults [32, 34-37]. Here, we investigate the cellular processes and transcriptional networks that regulate HIF-1 function. We describe an unbiased genetic screen to identify genes that inhibit *C. elegans* HIF-1. We discover that SKN-1/NRF represses HIF-1 protein levels. Hence, SKN-1-mediated repression of HIF-1 may provide a mechanism by which cells can rapidly respond to specific environmental stresses and optimize gene expression to achieve oxygen homeostasis. We investigate the hypothesis that this cross talk is mediated by EGL-9, the oxygen-sensing prolyl hydroxylase that modulates HIF-1 stability and activity.

Results

To identify genes and cellular processes that attenuated HIF-1-mediated gene expression, we conducted a genome-wide RNAi screen. This experimental strategy relied on the *Pnhr-57:GFP* reporter gene, which had been shown to be responsive to HIF-1 and hypoxia [28, 29]. Through chromatin immunoprecipitation experiments, we confirmed that *nhr-57* was a direct target of HIF-1 (S1 Fig). We screened a bacterial RNAi library representing ~80% of *C. elegans* genes [38], and we identified 179 genes for which RNAi increased *Pnhr-57:GFP* expression, as assayed by inspection under a fluorescent stereomicroscope (screen design illustrated in Fig 1A). These genes are listed in S1 Table and their functions are summarized in Fig 1B. Remarkably, 89 genes were predicted to have mitochondrial or metabolic functions, and the majority of these were electron transport chain components (42 genes) or subunits of mitochondrial ribosomes (24 genes). The second largest category of the genes was protein folding or turnover (33 genes), and the majority of these were proteasomal components. The remaining major categories include vesicular transport, transporters and channels, signaling and cytoskeleton, transcription and DNA or RNA processing.

Fig 1. Genome-wide RNAi screen to identify negative regulators of HIF-1-mediated gene expression. (A) Illustration of screen design. The screen identified RNAi treatments that increased the expression of *Pnhr-57:GFP*, a reporter that has been shown to be induced by hypoxia and positively regulated by HIF-1. (B) Functional summary of the 179 genes identified from the RNAi screen.

Recognizing that most eukaryotic genes are coordinately regulated by multiple transcription

factors, we did a secondary screen to identify those RNAi treatments that had a clear *hif-1*-dependent effect on *Pnhr-57:GFP* expression. To do this, we compared the *Pnhr-57:GFP* induction of these 179 RNAi treatments in wild-type animals and *hif-1*-deficient animals. Most of the 179 RNAi treatments increased the expression of *Pnhr-57:GFP* independent of *hif-1*. However, the *Pnhr-57:GFP* induction by 13 RNAi treatments showed a strong *hif-1*-dependent effect (S2 Table). Among these 13 genes, as expected, RNAi for *egl-9*, *rhy-1*, and *vhl-1*, previously characterized negative regulators of *C. elegans* HIF-1 [26, 29], increased *Pnhr-57:GFP* expression in wild-type animals, but not in *hif-1* mutants. These results validated the efficacy of our screen approach, and gave us the confidence to continue investigating the potentially new negative regulators of HIF-1 among these 13 genes.

***skn-1* attenuates HIF-1 protein levels and HIF-1 function**

We were especially intrigued by the finding that *skn-1* RNAi increased the expression of HIF-1-responsive reporter. Transcription factor SKN-1 has been shown to have critical roles in enabling *C. elegans* to respond to oxidative stress [32-34, 39]. Our finding suggested a potential crosstalk between hypoxia response and oxidative stress response. To quantify the effect of *skn-1* RNAi on this HIF-1-responsive reporter, we examined *Pnhr-57:GFP* levels using protein blots. In the normal room air culture conditions, *Pnhr-57:GFP* was 40% higher in *skn-1* RNAi compared to control RNAi (Fig 2A) (** $p < 0.01$, from three independent experiments). To gain insight to the effects of this interaction in hypoxic conditions, we moved the animals to 0.5% oxygen. After 4 hours of hypoxia treatment, *Pnhr-57:GFP* was 50% higher in *skn-1* RNAi compared to control RNAi (Fig 2A) (** $p < 0.01$, from three independent experiments). Thus, *skn-1* RNAi increased *Pnhr-57:GFP* levels under normoxic and hypoxic conditions.

We next asked whether *skn-1* RNAi increased HIF-1 protein levels. We tested two *skn-1* RNAi

constructs, and each resulted in an increase of HIF-1 protein levels by 2 to 3-fold as shown in Fig 2B (** $p < 0.01$, from three independent experiments). In sum, these results showed that *skn-1* RNAi increased HIF-1 protein level and HIF-1 reporter expression.

Fig 2. Identification of SKN-1 as a regulator of HIF-1. (A) *skn-1* RNAi increased expression of the *Pnhr-57::GFP* reporter. Reporter gene expression was quantitated in L4-stage animals in normal culture conditions or after 2 or 4 hours of hypoxia treatment (0.5% oxygen). The protein levels were calculated from three independent experiments and normalized to 0 hour hypoxia control RNAi. A representative western blot is shown. In each lane, lysates from 80 L4-stage worms were loaded. Asterisks indicate significant differences between control RNAi and *skn-1* RNAi at any given time point. *: $p < 0.05$; **: $p < 0.01$. (B) *skn-1* RNAi increased HIF-1 protein levels. These animals expressed an epitope-tagged HIF-1 protein [40]. The protein levels were calculated from three independent experiments and normalized to control RNAi. Two different RNAi clones were assayed, and they are designated as *skn-1(1)* and *skn-1(2)* here. A representative western blot is shown. In each lane, lysates from 100 L4-stage worms were loaded. Asterisks indicate significant differences between control RNAi and *skn-1* RNAi at any given time point. **: $p < 0.01$.

Differential requirements for *skn-1* and *hif-1*

The finding that SKN-1 repressed HIF-1 protein levels suggested that there might be conditions in which it would be beneficial for the animal to express one of these two stress-responsive transcription factors, but not the other. To address this, we examined the relative requirements for *skn-1* and *hif-1* more closely.

In previous studies, we and others had shown that *hif-1* was required for survival in moderate hypoxia [25, 41]. As validated in the experiments described in Table 1, loss of *hif-1* impaired animal development and survival in 0.5% oxygen: after 24 hours of hypoxia treatment, only 75.8% of *hif-1*-deficient eggs hatched, and only 25.6% developed to adulthood within 72 hours. In contrast, *skn-1* RNAi had no effect on *C. elegans* development and survival in hypoxic conditions: after 24 hours of hypoxia treatment, 99.4% of *skn-1* RNAi treated eggs hatched and completed normal development to adulthood within 72 hours.

Table 1. Relative requirements for *skn-1* and *hif-1*: survival in 0.5% oxygen.

Genotype	% hatched $\pm$ SEM	% survive to adult $\pm$ SEM	n ^a
N2 (wild type)	99.1 $\pm$ 0.25	99.1 $\pm$ 0.3	672
<i>hif-1(ia04)</i>	75.8 $\pm$ 2.5	25.6 $\pm$ 0.9	616
N2;control(RNAi)	99.6 $\pm$ 0.19	99.6 $\pm$ 0.2	558
N2; <i>skn-1</i> (RNAi)	99.4 $\pm$ 0.42	99.4 $\pm$ 0.4	640

^a n is the total number of animals assayed in three independent experiments.

Prior studies also suggested that there were differential requirements for HIF-1 and SKN-1 in oxidative stress conditions. Mutants carrying loss-of-function mutations in *skn-1* have been shown to decrease the ability of *C. elegans* to survive exposure to agents that cause oxidative stress [34, 42-44]. In contrast, *C. elegans* carrying loss-of-function mutations in *hif-1* have been reported to be relatively resistant to peroxide [40]. We compared these phenotypes directly, and the data are provided in Table 2. These experiments confirmed that, while *skn-1*-deficient animals were sensitive to t-butyl peroxide, mutants lacking *hif-1* were remarkably resistant to this oxidative stress: while none of the *skn-1*-deficient mutants survived 6 hours of t-butyl

peroxide treatment, 97.5% of *hif-1*-deficient mutants survived 10 hours of t-butyl peroxide treatment.

Table 2. Relative requirements for *skn-1* and *hif-1*: survival on t-butyl-peroxide.

Exposure time	Genotype	Mean Survival $\pm$ SEM	N ^a
6 hours	N2 (wild type)	95.2 $\pm$ 2.6	106
	<i>skn-1(zu67)</i>	0.0 $\pm$ 0	117
	<i>hif-1(ia04)</i>	100.0 $\pm$ 0	120
8 hours	N2 (wild type)	29.08 $\pm$ 7.7	106
	<i>skn-1(zu67)</i>	0.0 $\pm$ 0	117
	<i>hif-1(ia04)</i>	99.17 $\pm$ 0.8	120
10 hours	N2 (wild type)	2.78 $\pm$ 2.8	106
	<i>skn-1(zu67)</i>	0.0 $\pm$ 0	117
	<i>hif-1(ia04)</i>	97.50 $\pm$ 2.5	120

^a n is the total number of animals assayed in three independent experiments.

SKN-1/NRF promotes *egl-9* expression

We next sought to discover the mechanism by which SKN-1 inhibits HIF-1 protein levels. In silico analyses identified a potential SKN-1 binding site in the *egl-9* promoter region (Fig 3A). EGL-9 is a central inhibitor of HIF-1 protein levels [26, 27] and of HIF-1 transcriptional activity [29, 30]. This suggested a model in which the SKN-1 DNA-binding complex bound directly to the *egl-9* regulatory sequences to promote *egl-9* expression, which, in turn, would ultimately decrease HIF-1 protein levels. To test this, we employed real-time quantitative RT-PCR to compare *egl-9* mRNA levels in worms fed with *skn-1* RNAi versus control RNAi. To produce

reliable and reproducible results, *egl-9* mRNA levels were quantitated in three independent real-time quantitative RT-PCR experiments in L4-stage animals in room air or hypoxic conditions (0.5% oxygen). Each sample was performed with three technical replicates, and they produced similar C_t values. There are seven isoforms of *egl-9* mRNA transcripts (https://wormbase.org/species/c_elegans/gene/WBGene00001178#0-9f-10). The real-time quantitative PCR primer set used in this study can detect six *egl-9* mRNA isoforms. In room air, *skn-1* RNAi decreased *egl-9* mRNA levels by 30% compared to control RNAi (** $p < 0.01$, from three independent experiments) (Fig 3B). HIF-1 has been shown to activate *egl-9* mRNA expression under hypoxia, creating a negative feedback loop [27, 28]. In accordance with this, the inhibition effects of *skn-1* RNAi on *egl-9* mRNA levels were minimized by placing the animals in hypoxic conditions (Fig 3B).

Fig 3. Identification of *egl-9* as a potential transcriptional target of SKN-1. (A) Sequence from the *egl-9* promoter was aligned with established SKN-1 binding sites in *gcs-1*, *med-1*, and *med-2*. Asterisks identify sequence identities shared by all four promoter regions in this interval, and predicted SKN-1 binding sites are in red. (B) *skn-1* RNAi decreased *egl-9* mRNA levels. *egl-9* mRNA levels were quantitated from three independent real-time quantitative RT-PCR experiments. The values at each time point were normalized to the 0 hour hypoxia control RNAi. Asterisks indicate significant differences between control RNAi and *skn-1* RNAi at any given time point. *: $p < 0.05$; **: $p < 0.01$.

To test the hypothesis that conditions that activate SKN-1 can promote *egl-9* promoter activity, we generated a reporter construct in which 1.6 kb of *egl-9* regulatory sequence directed the

expression of GFP (Fig 4A). To distinguish the effects of SKN-1 on *egl-9* expression from those of HIF-1, we conducted these experiments in a *hif-1* mutant background. In agreement with prior studies [45], *Pegl-9::GFP* was visible in several tissues, including the body muscle, vulva, pharynx, anterior intestine, rectal cells and additional cells in the tail in standard culture conditions (20 °C) (Figs 4B and 4C). When the animals were treated with heat shock conditions that had been shown to activate SKN-1 (29 °C for 20 hours) [34], we observed dramatic induction of *Pegl-9::GFP* in the intestine. (Figs 4D and 4E).

Fig 4. Heat shock alters *Pegl-9::GFP* expression. (A) The *Pegl-9::GFP* construct includes 1.6 kb of sequence 5' to the *egl-9* translational start. GFP coding sequence is diagramed as a green box. The red oval indicates the position of the putative SKN-1 binding site. (B-E) *Pegl-9::GFP* expression in L4-stage animals under normal culture conditions and heat shock. Animals are shown as DIC images (B and D) and corresponding images of GFP fluorescence (C and E). In all images, the head is to the right. (B and C) Under normal conditions, *Pegl-9::GFP* was expressed in the body muscle, vulva, pharynx, anterior intestine, rectal cells and additional cells in the tail. (D and E) After heat shock treatment (29°C for 20 hours), *Pegl-9::GFP* was strongly induced in the intestine. The *Pegl-9::GFP* expression patterns in the L1, L2, L3 and adults were similar to that in the L4 worms, under both normal and heat shock conditions (data not shown).

We next asked whether heat shock induction of *Pegl-9::GFP* required *skn-1* function. We found that heat shock increased *Pegl-9::GFP* by 2.5-fold in animals carrying the wild-type *skn-1* allele. However, the heat shock induction of *Pegl-9::GFP* was abolished in *skn-1(zu67)* loss-of-function mutants (Fig 5A). Analyses of another independent *Pegl-9::GFP* transgenic line yielded similar

results (data not shown).

To test the hypothesis that the putative SKN-1 binding site in the *egl-9* promoter was required for heat shock induction of *Pegl-9:GFP*, we generated the *P(m)egl-9:GFP* construct, which contained mutations in the putative SKN-1 binding site (in red type in Fig 3A). Heat shock increased *Pegl-9:GFP* by 2.1-fold. However, heat shock failed to induce the expression of *P(m)egl-9:GFP* (Fig 5B). Experiments with a second *P(m)egl-9:GFP* transgenic line gave similar results (data not shown). Collectively, these data demonstrated that heat shock induction of *Pegl-9:GFP* required *skn-1* function and the putative SKN-1 binding site in the *egl-9* promoter.

Fig 5. SKN-1 acts through the putative SKN-1 binding site in the *egl-9* promoter to

activate *egl-9* expression. (A) Heat shock induced *Pegl-9:GFP* in animals carrying the wild-type *skn-1* allele, but did not induce the reporter in animals carrying the *skn-1(zu67)* loss-of-function mutation. The vertical axis shows the log2 fold changes of GFP caused by heat shock for each strain with standard errors, as determined by four biological replicates. A representative western blot is shown. For each sample, 20 L4-satge worms were boiled and lysates corresponding to 10 worms were loaded to each lane. (B) Heat shock increased the expression of *Pegl-9:GFP*, but did not increase the expression of the reporter in which the putative SKN-1 binding site was mutated (*P(m)egl-9:GFP*). The vertical axis shows the log2 fold changes of GFP caused by heat shock for each strain with standard errors, as determined by five biological replicates. A representative western blot is shown. For each sample, 20 L4-satge worms were boiled and lysates corresponding to 10 worms were loaded to each lane.

Discussion

This study provides new insights to the mechanisms that allow animals to respond appropriately to diverse stresses. While HIF-1 and SKN-1 are both stress responsive transcription factors, they have distinct functions. For example, *skn-1*-deficient animals are less able to survive exposure to peroxide, while *hif-1*-deficient mutants are relatively resistant to this oxidizing agent [40]. Conversely, while a deletion mutation in *hif-1* dramatically impairs survival in 0.5% oxygen [25], *skn-1* RNAi has little effect (Table 1). Thus, animals may benefit from cross-talk between these two transcription factors as they are challenged by oxygen deprivation and oxidative stress. Here, we provide evidence that SKN-1 promotes *egl-9* expression, thereby attenuating HIF-1 function.

SKN-1 promotes *egl-9* expression, thereby inhibiting HIF-1

Our data support a model in which SKN-1 binds directly to the *egl-9* promoter to increase *egl-9* expression. This conclusion is further substantiated by the genome wide chromatin immunoprecipitation experiments that determined that SKN-1:GFP associated with DNA sequences in the *egl-9* 5' regulatory region [46]. EGL-9 functions as a cellular oxygen sensor, and it mediates oxygen-dependent degradation of HIF-1 [26]. Hence, *skn-1* RNAi results in increased HIF-1 protein levels (Fig 2B). We expect that this regulatory interaction, in which SKN-1 can quickly down-regulate HIF-1, would allow animals to adapt quickly to changes in cellular conditions.

Prior studies have identified genes that are positively regulated by either SKN-1 or HIF-1, and these gene lists are largely non-overlapping [28, 32, 47-49]. Genes that are commonly regulated by both SKN-1 and HIF-1 include K10H10.2/*cysl-2*, F57B9.1, M05D6.5, and *rhy-1*. Two lines of evidence suggest that *rhy-1* may be a direct target of SKN-1. First, Oliveira et al. (2009)

identified a potential SKN-1 binding site in the *rhy-1* promoter region. Second, the modENCODE project found that *rhy-1* 5' regulatory sequences associated with SKN-1 *in vivo* [46]. Like *egl-9*, *rhy-1* is also a negative regulator of HIF-1 [28, 29]. Collectively, these data suggest that SKN-1 may act through both *egl-9* and *rhy-1* to reduce HIF-1 function. Interactions between HIF-1 and SKN-1 are likely to be different in select cells, developmental stages, or environmental contexts. HIF-1, EGL-9 and SKN-1 each have other developmental functions, and some of these are specific to certain cell types [34, 50-54]. The downstream effects of SKN-1 or HIF-1 activation are further influenced by cellular or environmental contexts. For example, *C. elegans* SKN-1 can be activated by arsenite or by t-butyl hydroperoxide, but only a subset of SKN-1 targets are activated by either toxicant [32]. Also, while *C. elegans* SKN-1 and HIF-1 have distinct roles in peroxide and hypoxia stress responses, their functions overlap in selenium and hydrogen sulfide stress responses [13, 15, 35-37]. Similarly, the mammalian NRF and HIF transcription factors are very sensitive to environmental and physiological cues or stresses, and their regulatory relationships are context specific. Ischemia causes tissue hypoxia, which stabilizes HIF transcription factors. Once the ischemic tissue is reperfused, HIF transcription factors are degraded quickly and NRF2 is up-regulated, presumably to limit the oxidative damage [4, 5]. However, intermittent hypoxia has been shown to induce both HIF-1 α and NRF2 [6, 7]. While NRF2 signaling activates HIF-1 in several cancer types [12], studies of the anti-inflammatory drug andrographolide in endothelial cells revealed interactions between NRF2 and the PHDs that modulate HIF-1 [55]. NRF2 and the HIF transcription factors have key roles in angiogenesis and iron regulation, and their functions can converge on developmental processes or feedback loops that modulate their activities [12, 56-58].

***Pnhr-57:GFP* as a marker for hypoxia-induced gene expression:**

discoveries and insights from a genome-wide screen

Prior studies have demonstrated that *nhr-57* was induced by hypoxia in a *hif-1*-dependent manner and that over-expression of *Pnhr-57:GFP* in *egl-9* mutants required *hif-1* [13, 27-30]. Moreover, Bellier *et al.* (2009) found that HIF-1-mediated induction of *nhr-57* helped to protect *C. elegans* from the lethal effects of pore-forming toxins [59]. While these studies showed that *nhr-57* is a direct target of HIF-1, other transcription factors must also contribute to its expression. The studies presented here show that, while genes such as *egl-9* or *rhy-1* clearly regulate HIF-1 to control *Pnhr-57:GFP* expression, many other RNAi treatments can activate *Pnhr-57:GFP* through *hif-1*-independent pathways. The *Pnhr-57:GFP* reporter will continue to be a valuable marker, but these data inform our interpretations of studies that employ this reporter. While *Pnhr-57:GFP* is clearly regulated by HIF-1, it is important to compare expression of the reporter in wild-type and *hif-1*-deficient animals before drawing conclusions about HIF-1 activity.

We found that diverse RNAi treatments that compromise metabolic function or protein homeostasis increased *Pnhr-57:GFP* expression, and this effect did not require a functional *hif-1* gene. Interestingly, many of the genes integral to these processes have been shown to have roles in stress response and aging [60-65]. RNAi-mediated depletion of proteasomal components has also been shown to impact resistance to polyglutamine toxicity and to induce expression of *Pgpdh-1:GFP*, a marker for osmotic stress and glycerol production [66, 67].

Further characterization of the 179 RNAi treatments that increased *Pnhr-57:GFP* identified 13 genes that had much stronger effects in animals carrying a wild-type *hif-1* gene. These genes included *vhl-1*, *egl-9*, and *rhy-1*. These three genes had all been identified in prior studies as

negative regulators of HIF-1 [26, 29]. The succinate dehydrogenase subunit *sdhb-1* was also found to have *hif-1*-dependent effects. This is especially interesting, since studies in cancer cell lines have shown that succinate can inhibit the enzymatic activities of HIF prolyl hydroxylases [68, 69]. *sams-1* and *sbp-1* encode the *C. elegans* S-adenosyl methionine synthetase and the SREBP homologs, respectively. The RNAi treatments of *sams-1* and *sbp-1* have a lesser impact on *Pnhr-57::GFP* levels in *hif-1* mutants, suggesting that the effects of *sams-1* and *sbp-1* RNAi on the reporter are mediated by HIF-1 (S2 Fig). Both of these genes have key roles in methionine metabolism and fatty acid biosynthesis [70], and it will be interesting to investigate the ways in which these important processes intersect with hypoxia response.

Materials and methods

Strains

The following strains were used in this study: wild-type N2 Bristol; ZG430: *Pnhr-57::GFP(iaIs07)IV*; *egl-9(sa307)V*; *hif-1(ia04)V*; *Phif-1:hif-1a:Myc:HA (iaIs28)*; ZG120: *Pnhr-57::GFP(iaIs07)IV*; ZG509: *rrf-3(pk1426)II*; *Pnhr-57::GFP(iaIs07)IV*; ZG508: *rrf-3(pk1426)II*; ZG429: *hif-1(ia04)V*; *Phif-1:hif-1a:Myc:HA(iaIs28)*; ZG472: *hif-1(ia04)V*; *Pegl-9::GFP(iaEx84)*; ZG487: *hif-1(ia04)V*; *P(m)egl-9::GFP(iaEx96)*; ZG488: *skn-1(zu67)IV*; *hif-1(ia04)V*; *Pegl-9::GFP(iaEx84)*. The transgenes expressing epitope-tagged HIF-1 protein were described and characterized previously [40].

RNAi experiments

The RNAi screen was conducted as previously described [71], with few modifications. Each bacterial clone (expressing double-stranded RNA for one gene) was cultured in L-broth with 50 ug/mL ampicillin and 12.5 ug/mL tetracycline overnight at 37°C. The following morning, the

bacteria were inoculated into new L-broth with 100 ug/mL ampicillin for 6 hours at 37°C before seeding on 24-well NGM agar plates with 25 ug/mL carbenicillin and 2 mM IPTG. Each RNAi clone was plated in duplicate. The following day, 15-25 L1-stage worms were added to each well. The plates were incubated at 15°C for 5-6 days, and then the worms were screened for positive *Pnhr-57::GFP* green fluorescence by stereomicroscopy. Bacterial RNAi clones that increased the reporter were rescreened in two independent replicates, and the plasmid inserts were validated by sequencing.

skn-1 RNAi causes maternal-effect lethality, in this study we examined the effects of first generation *skn-1* RNAi. Dead egg percentages given by the first generation *skn-1* RNAi adults were measured to check the *skn-1* RNAi efficiency. We routinely achieved as high as 90% dead egg percentages from the first generation *skn-1* RNAi adults, indicating high *skn-1* RNAi efficiency.

Hypoxia stress and oxidative stress assays

To assess the relative effects of t-butyl-peroxide exposure, animals in the first day of adulthood were placed on NGM plates containing 7.5 mM t-butyl-peroxide, in the presence of bacterial food. The survival was scored after treating the animals for 6, 8 or 10 hours.

For hypoxia experiments, adults were allowed to lay eggs on standard NGM plates with OP50 bacterial food for 2 hours. The adults were then removed, and the plates with embryos were placed in a hypoxia chamber (0.5% oxygen; 21°C) for 24 hours. After 24 hours, the plates were removed from the hypoxia chamber, and the un-hatched eggs were counted immediately. The plates were then maintained in room air (21°C). The adult worms were counted at 72 hour since the eggs had been laid. Wild-type control animals hatched within 24 hours and reached adulthood within 72 hours.

***Pegl-9:GFP* expression constructs**

To generate the *Pegl-9:GFP* construct, a fragment that contained 1.6 kb of sequence upstream of the initiation ATG of *egl-9* gene was amplified by PCR using the forward primer 5'-CGCGCATGCGTGTATGTGTGTGAAAGAG-3' and the reverse primer 5'-GCGGTCGACGCAACTTTTTTCTGTCACATTCAG-3'. The PCR product was cloned into the green fluorescence protein (GFP) vector pPD95.75 (gift from Andrew Fire). To create the *P(m)egl-9:GFP* point mutation construct, the predicted SKN-1 binding site TTTGTCAT [34, 72] was altered to CGACGGGC. Transgenic animals were generated by injection of DNA into the gonadal syncytium, using standard methods with *rol-6* (pRF4) as the co-injection marker [73]. For each construct, two independent transgenic lines were generated and assayed.

Protein Blots

To assay the expression of GFP or HIF-1 proteins, 20-100 L4-stage worms were collected and boiled for 5 min in 1X SDS sample buffer, and the lysates were size fractionated on polyacrylamide gels and analyzed by Western blots. The GFP-specific mouse monoclonal antibody (from Roche) was used at 1:500. The HA-specific mouse monoclonal antibody (from Cell Signaling) was used at 1:250. The secondary antibody (goat anti-mouse IgG+IgM from Biorad) was used at 1:2000 dilutions. The western blot images were analyzed by the Image J software. For each assay, three to five independent biological replicates were included.

RNA extraction and real-time quantitative RT-PCR

Total RNA was isolated from synchronized L4-stage animals using Trizol (Invitrogen) and RNeasy Mini Kit (Qiagen). After being treated by RNase free DNase (Promega), total RNA was reverse transcribed to complementary DNA using Oligo dT₁₈ primer and AffinityScript reverse

transcriptase (Stratagene). Real-time quantitative PCR was performed using the iQ SYBR GREEN supermix (Bio-Rad) and Stratagene Mx4000 multiplex PCR system. In the assay, three biological replicates were included. And for each sample, three technical replicates were performed and they gave similar C_t values. And cDNA from 62.5 ng of total RNA was added to each PCR reaction. Relative mRNA quantification was performed using the efficiency-corrected comparative quantification method [74]. *inf-1*, a gene not regulated by hypoxia, was used as the reference gene [28]. The primer sequences for *inf-1* real-time quantitative PCR are included in S1 Fig. The primers for *egl-9* real-time quantitative PCR are the forward primer 5'-GCCGACTTTCAATCCACTTC-3' and reverse primer 5'-AATGATCGGAGATCGACTGG-3'. There are seven isoforms of *egl-9* mRNA transcripts (https://wormbase.org/species/c_elegans/gene/WBGene00001178#0-9f-10). This primer set can detect six out of seven *egl-9* mRNA isoforms. The isoform d.1 will not be detected by this primer set, because the forward primer is located within the eighth intron of the unspliced isoform d.1.

References

1. Beckman KB, Ames BN. The free radical theory of aging matures. *Physiol Rev.* 1998;78(2):547-81. Epub 1998/04/30. PubMed PMID: 9562038.
2. Finkel T. Oxidant signals and oxidative stress. *Current opinion in cell biology.* 2003;15(2):247-54. Epub 2003/03/22. PubMed PMID: 12648682.
3. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. *Nature.* 2000;408(6809):239-47. Epub 2000/11/23. doi: 10.1038/35041687. PubMed PMID: 11089981.
4. Kim YJ, Ahn JY, Liang P, Ip C, Zhang Y, Park YM. Human prx1 gene is a target of Nrf2 and is up-regulated by hypoxia/reoxygenation: implication to tumor biology. *Cancer Res.* 2007;67(2):546-54. Epub 2007/01/20. doi: 67/2/546 [pii] 10.1158/0008-5472.CAN-06-2401. PubMed PMID: 17234762.
5. Leonard MO, Kieran NE, Howell K, Burne MJ, Varadarajan R, Dhakshinamoorthy S, et al. Reoxygenation-specific activation of the antioxidant transcription factor Nrf2 mediates cytoprotective gene expression in ischemia-reperfusion injury. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology.* 2006;20(14):2624-6. Epub 2006/12/05. doi: 10.1096/fj.06-5097fje. PubMed PMID: 17142801.
6. Malec V, Gottschald OR, Li S, Rose F, Seeger W, Hanze J. HIF-1 alpha signaling is augmented during intermittent hypoxia by induction of the Nrf2 pathway in NOX1-expressing adenocarcinoma A549

- cells. Free radical biology & medicine. 2010;48(12):1626-35. Epub 2010/03/30. doi: 10.1016/j.freeradbiomed.2010.03.008. PubMed PMID: 20347035.
7. Martinive P, Defresne F, Bouzin C, Saliez J, Lair F, Gregoire V, et al. Preconditioning of the tumor vasculature and tumor cells by intermittent hypoxia: implications for anticancer therapies. Cancer Res. 2006;66(24):11736-44. doi: 10.1158/0008-5472.CAN-06-2056. PubMed PMID: 17178869.
8. Thimmulappa RK, Mai KH, Srisuma S, Kensler TW, Yamamoto M, Biswal S. Identification of Nrf2-regulated genes induced by the chemopreventive agent sulforaphane by oligonucleotide microarray. Cancer Res. 2002;62(18):5196-203. Epub 2002/09/18. PubMed PMID: 12234984.
9. Hayes JD, McMahon M. Molecular basis for the contribution of the antioxidant responsive element to cancer chemoprevention. Cancer Lett. 2001;174(2):103-13. Epub 2001/11/02. doi: S0304383501006954 [pii]. PubMed PMID: 11689285.
10. Hough RB, Piatigorsky J. Preferential transcription of rabbit Aldh1a1 in the cornea: implication of hypoxia-related pathways. Molecular and cellular biology. 2004;24(3):1324-40. Epub 2004/01/20. PubMed PMID: 14729976; PubMed Central PMCID: PMC321433.
11. Yeligar SM, Machida K, Kalra VK. Ethanol-induced HO-1 and NQO1 are differentially regulated by HIF-1alpha and Nrf2 to attenuate inflammatory cytokine expression. The Journal of biological chemistry. 2010;285(46):35359-73. Epub 2010/09/14. doi: 10.1074/jbc.M110.138636. PubMed PMID: 20833713; PubMed Central PMCID: PMC2975160.
12. Toth RK, Warfel NA. Strange Bedfellows: Nuclear Factor, Erythroid 2-Like 2 (Nrf2) and Hypoxia-Inducible Factor 1 (HIF-1) in Tumor Hypoxia. Antioxidants (Basel). 2017;6(2). doi: 10.3390/antiox6020027. PubMed PMID: 28383481; PubMed Central PMCID: PMCPMC5488007.
13. Budde MW, Roth MB. Hydrogen sulfide increases hypoxia-inducible factor-1 activity independently of von Hippel-Lindau tumor suppressor-1 in C. elegans. Mol Biol Cell. 2010;21(1):212-7. Epub 2009/11/06. doi: E09-03-0199 [pii] 10.1091/mbc.E09-03-0199. PubMed PMID: 19889840; PubMed Central PMCID: PMC2801715.
14. Powell-Coffman JA. Hypoxia signaling and resistance in C. elegans. Trends Endocrinol Metab. 2010;21(7):435-40. Epub 2010/03/26. doi: S1043-2760(10)00041-X [pii] 10.1016/j.tem.2010.02.006. PubMed PMID: 20335046.
15. Romanelli-Credrez L, Doitsidou M, Alkema MJ, Salinas G. HIF-1 Has a Central Role in Caenorhabditis elegans Organismal Response to Selenium. Front Genet. 2020;11:63. doi: 10.3389/fgene.2020.00063. PubMed PMID: 32161616; PubMed Central PMCID: PMCPMC7052493.
16. Saldanha JN, Parashar A, Pandey S, Powell-Coffman JA. Multiparameter behavioral analyses provide insights to mechanisms of cyanide resistance in Caenorhabditis elegans. Toxicol Sci. 2013;135(1):156-68. doi: 10.1093/toxsci/kft138. PubMed PMID: 23805000; PubMed Central PMCID: PMCPMC3748764.
17. Shao Z, Zhang Y, Ye Q, Saldanha JN, Powell-Coffman JA. C. elegans SWAN-1 Binds to EGL-9 and regulates HIF-1-mediated resistance to the bacterial pathogen Pseudomonas aeruginosa PAO1. PLoS Pathog. 2010;6(8):e1001075. doi: 10.1371/journal.ppat.1001075. PubMed PMID: 20865124; PubMed Central PMCID: PMCPMC2928816.
18. Kaelin WG, Jr., Ratcliffe PJ. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. Mol Cell. 2008;30(4):393-402. Epub 2008/05/24. doi: S1097-2765(08)00292-X [pii] 10.1016/j.molcel.2008.04.009. PubMed PMID: 18498744.
19. Semenza GL. Oxygen sensing, homeostasis, and disease. N Engl J Med. 2011;365(6):537-47. doi: 10.1056/NEJMr1011165. PubMed PMID: 21830968.
20. Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O2 tension. Proc Natl Acad Sci U S A. 1995;92(12):5510-4. doi: 10.1073/pnas.92.12.5510. PubMed PMID: 7539918; PubMed Central PMCID: PMCPMC41725.
21. Maxwell PH, Wiesener MS, Chang GW, Clifford SC, Vaux EC, Cockman ME, et al. The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. Nature. 1999;399(6733):271-5. doi: 10.1038/20459. PubMed PMID: 10353251.

22. Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, et al. Targeting of HIF- α to the von Hippel-Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science*. 2001;292(5516):468-72. doi: 10.1126/science.1059796. PubMed PMID: 11292861.
23. Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, et al. HIF α targeted for VHL-mediated destruction by proline hydroxylation: implications for O₂ sensing. *Science*. 2001;292(5516):464-8. doi: 10.1126/science.1059817. PubMed PMID: 11292862.
24. Powell-Coffman JA, Bradfield CA, Wood WB. *Caenorhabditis elegans* orthologs of the aryl hydrocarbon receptor and its heterodimerization partner the aryl hydrocarbon receptor nuclear translocator. *Proc Natl Acad Sci U S A*. 1998;95(6):2844-9. Epub 1998/04/18. PubMed PMID: 9501178.
25. Jiang H, Guo R, Powell-Coffman JA. The *Caenorhabditis elegans* hif-1 gene encodes a bHLH-PAS protein that is required for adaptation to hypoxia. *Proc Natl Acad Sci U S A*. 2001;98(14):7916-21. Epub 2001/06/28. doi: 10.1073/pnas.141234698. PubMed PMID: 11427734; PubMed Central PMCID: PMC35443.
26. Epstein AC, Gleadle JM, McNeill LA, Hewitson KS, O'Rourke J, Mole DR, et al. *C. elegans* EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. *Cell*. 2001;107(1):43-54. Epub 2001/10/12. doi: S0092-8674(01)00507-4 [pii]. PubMed PMID: 11595184.
27. Bishop T, Lau KW, Epstein AC, Kim SK, Jiang M, O'Rourke D, et al. Genetic analysis of pathways regulated by the von Hippel-Lindau tumor suppressor in *Caenorhabditis elegans*. *PLoS Biol*. 2004;2(10):e289. Epub 2004/09/14. doi: 10.1371/journal.pbio.0020289. PubMed PMID: 15361934.
28. Shen C, Nettleton D, Jiang M, Kim SK, Powell-Coffman JA. Roles of the HIF-1 hypoxia-inducible factor during hypoxia response in *Caenorhabditis elegans*. *The Journal of biological chemistry*. 2005;280(21):20580-8. Epub 2005/03/23. doi: M501894200 [pii]. PubMed PMID: 15781453.
29. Shen C, Shao Z, Powell-Coffman JA. The *Caenorhabditis elegans* rhy-1 gene inhibits HIF-1 hypoxia-inducible factor activity in a negative feedback loop that does not include vhl-1. *Genetics*. 2006;174(3):1205-14. Epub 2006/09/19. doi: genetics.106.063594 [pii]. PubMed PMID: 16980385.
30. Shao Z, Zhang Y, Powell-Coffman JA. Two distinct roles for EGL-9 in the regulation of HIF-1-mediated gene expression in *Caenorhabditis elegans*. *Genetics*. 2009;183(3):821-9. Epub 2009/09/10. doi: genetics.109.107284 [pii]. PubMed PMID: 19737748; PubMed Central PMCID: PMC2778979.
31. Walker AK, See R, Batchelder C, Kophengnavong T, Gronniger JT, Shi Y, et al. A conserved transcription motif suggesting functional parallels between *Caenorhabditis elegans* SKN-1 and Cap'n'Collar-related basic leucine zipper proteins. *The Journal of biological chemistry*. 2000;275(29):22166-71. Epub 2000/04/15. doi: 10.1074/jbc.M001746200. PubMed PMID: 10764775.
32. Oliveira RP, Abate JP, Dilks K, Landis J, Ashraf J, Murphy CT, et al. Condition-adapted stress and longevity gene regulation by *Caenorhabditis elegans* SKN-1/Nrf. *Aging Cell*. 2009;8(5):524-41. Epub 2009/07/07. doi: ACE501 [pii]. PubMed PMID: 19575768.
33. Park SK, Tedesco PM, Johnson TE. Oxidative stress and longevity in *Caenorhabditis elegans* as mediated by SKN-1. *Aging Cell*. 2009;8(3):258-69. Epub 2009/07/25. doi: 10.1111/j.1474-9726.2009.00473.x. PubMed PMID: 19627265; PubMed Central PMCID: PMC2762118.
34. An JH, Blackwell TK. SKN-1 links *C. elegans* mesendodermal specification to a conserved oxidative stress response. *Genes Dev*. 2003;17(15):1882-93. Epub 2003/07/19. doi: 10.1101/gad.1107803. PubMed PMID: 12869585.
35. Horsman JW, Heinis FI, Miller DL. A Novel Mechanism To Prevent H₂S Toxicity in *Caenorhabditis elegans*. *Genetics*. 2019;213(2):481-90. doi: 10.1534/genetics.119.302326. PubMed PMID: 31371406; PubMed Central PMCID: PMC6781907.
36. Li WH, Chang CH, Huang CW, Wei CC, Liao VH. Selenite enhances immune response against

- Pseudomonas aeruginosa PA14 via SKN-1 in Caenorhabditis elegans. PLoS One. 2014;9(8):e105810. doi: 10.1371/journal.pone.0105810. PubMed PMID: 25147937; PubMed Central PMCID: PMC4141825.
37. Miller DL, Budde MW, Roth MB. HIF-1 and SKN-1 coordinate the transcriptional response to hydrogen sulfide in Caenorhabditis elegans. PLoS One. 2011;6(9):e25476. Epub 2011/10/08. doi: 10.1371/journal.pone.0025476 PONE-D-11-09953 [pii]. PubMed PMID: 21980473; PubMed Central PMCID: PMC3183046.
38. Kamath RS, Ahringer J. Genome-wide RNAi screening in Caenorhabditis elegans. Methods. 2003;30(4):313-21. Epub 2003/06/28. doi: S1046202303000501 [pii]. PubMed PMID: 12828945.
39. Inoue H, Hisamoto N, An JH, Oliveira RP, Nishida E, Blackwell TK, et al. The C. elegans p38 MAPK pathway regulates nuclear localization of the transcription factor SKN-1 in oxidative stress response. Genes Dev. 2005;19(19):2278-83. Epub 2005/09/17. doi: gad.1324805 [pii] 10.1101/gad.1324805. PubMed PMID: 16166371.
40. Zhang Y, Shao Z, Zhai Z, Shen C, Powell-Coffman JA. The HIF-1 hypoxia-inducible factor modulates lifespan in C. elegans. PLoS One. 2009;4(7):e6348. Epub 2009/07/28. doi: 10.1371/journal.pone.0006348. PubMed PMID: 19633713.
41. Padilla PA, Nystul TG, Zager RA, Johnson AC, Roth MB. Dephosphorylation of cell cycle-regulated proteins correlates with anoxia-induced suspended animation in Caenorhabditis elegans. Mol Biol Cell. 2002;13(5):1473-83. Epub 2002/05/15. doi: 10.1091/mbc.01-12-0594. PubMed PMID: 12006646.
42. Li X, Matilainen O, Jin C, Glover-Cutter KM, Holmberg CI, Blackwell TK. Specific SKN-1/Nrf stress responses to perturbations in translation elongation and proteasome activity. PLoS Genet. 2011;7(6):e1002119. Epub 2011/06/23. doi: 10.1371/journal.pgen.1002119. PubMed PMID: 21695230; PubMed Central PMCID: PMC3111486.
43. Tullet JM, Hertweck M, An JH, Baker J, Hwang JY, Liu S, et al. Direct inhibition of the longevity-promoting factor SKN-1 by insulin-like signaling in C. elegans. Cell. 2008;132(6):1025-38. Epub 2008/03/25. doi: S0092-8674(08)00130-X [pii] 10.1016/j.cell.2008.01.030. PubMed PMID: 18358814.
44. Wang J, Robida-Stubbs S, Tullet JM, Rual JF, Vidal M, Blackwell TK. RNAi screening implicates a SKN-1-dependent transcriptional response in stress resistance and longevity deriving from translation inhibition. PLoS Genet. 2010;6(8). Epub 2010/08/12. doi: 10.1371/journal.pgen.1001048. PubMed PMID: 20700440; PubMed Central PMCID: PMC2916858.
45. Darby C, Cosma CL, Thomas JH, Manoil C. Lethal paralysis of Caenorhabditis elegans by Pseudomonas aeruginosa. Proc Natl Acad Sci U S A. 1999;96(26):15202-7. Epub 1999/12/28. PubMed PMID: 10611362.
46. Niu W, Lu ZJ, Zhong M, Sarov M, Murray JI, Brdlik CM, et al. Diverse transcription factor binding features revealed by genome-wide ChIP-seq in C. elegans. Genome research. 2011;21(2):245-54. Epub 2010/12/24. doi: 10.1101/gr.114587.110. PubMed PMID: 21177963; PubMed Central PMCID: PMC3032928.
47. Glover-Cutter KM, Lin S, Blackwell TK. Integration of the unfolded protein and oxidative stress responses through SKN-1/Nrf. PLoS Genet. 2013;9(9):e1003701. doi: 10.1371/journal.pgen.1003701. PubMed PMID: 24068940; PubMed Central PMCID: PMC3772064.
48. Paek J, Lo JY, Narasimhan SD, Nguyen TN, Glover-Cutter K, Robida-Stubbs S, et al. Mitochondrial SKN-1/Nrf mediates a conserved starvation response. Cell Metab. 2012;16(4):526-37. doi: 10.1016/j.cmet.2012.09.007. PubMed PMID: 23040073; PubMed Central PMCID: PMC3774140.
49. Robida-Stubbs S, Glover-Cutter K, Lamming DW, Mizunuma M, Narasimhan SD, Neumann-Haefelin E, et al. TOR signaling and rapamycin influence longevity by regulating SKN-1/Nrf and DAF-16/FoxO. Cell Metab. 2012;15(5):713-24. doi: 10.1016/j.cmet.2012.04.007. PubMed PMID: 22560223; PubMed Central PMCID: PMC3348514.
50. Bishop NA, Guarente L. Two neurons mediate diet-restriction-induced longevity in C. elegans. Nature. 2007;447(7144):545-9. Epub 2007/06/01. doi: nature05904 [pii]

- 10.1038/nature05904. PubMed PMID: 17538612.
51. Chen D, Thomas EL, Kapahi P. HIF-1 modulates dietary restriction-mediated lifespan extension via IRE-1 in *Caenorhabditis elegans*. *PLoS Genet*. 2009;5(5):e1000486. Epub 2009/05/23. doi: 10.1371/journal.pgen.1000486. PubMed PMID: 19461873.
52. Park EC, Ghose P, Shao Z, Ye Q, Kang L, Xu XZ, et al. Hypoxia regulates glutamate receptor trafficking through an HIF-independent mechanism. *The EMBO journal*. 2012;31(6):1379-93. Epub 2012/01/19. doi: 10.1038/emboj.2011.499. PubMed PMID: 22252129; PubMed Central PMCID: PMC3321172.
53. Pocock R, Hobert O. Oxygen levels affect axon guidance and neuronal migration in *Caenorhabditis elegans*. *Nat Neurosci*. 2008;11(8):894-900. Epub 2008/07/01. doi: nn.2152 [pii] 10.1038/nn.2152. PubMed PMID: 18587389.
54. Chang AJ, Bargmann CI. Hypoxia and the HIF-1 transcriptional pathway reorganize a neuronal circuit for oxygen-dependent behavior in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*. 2008;105(20):7321-6. doi: 10.1073/pnas.0802164105. PubMed PMID: 18477695; PubMed Central PMCID: PMCPMC2438248.
55. Lin HC, Su SL, Lu CY, Lin AH, Lin WC, Liu CS, et al. Andrographolide inhibits hypoxia-induced HIF-1 α -driven endothelin 1 secretion by activating Nrf2/HO-1 and promoting the expression of prolyl hydroxylases 2/3 in human endothelial cells. *Environ Toxicol*. 2017;32(3):918-30. doi: 10.1002/tox.22293. PubMed PMID: 27297870.
56. Duarte TL, Talbot NP, Drakesmith H. NRF2 and Hypoxia-Inducible Factors: Key Players in the Redox Control of Systemic Iron Homeostasis. *Antioxid Redox Signal*. 2020. doi: 10.1089/ars.2020.8148. PubMed PMID: 32791852.
57. Li L, Pan H, Wang H, Li X, Bu X, Wang Q, et al. Interplay between VEGF and Nrf2 regulates angiogenesis due to intracranial venous hypertension. *Sci Rep*. 2016;6:37338. doi: 10.1038/srep37338. PubMed PMID: 27869147; PubMed Central PMCID: PMCPMC5116754.
58. Wong BW, Kuchnio A, Bruning U, Carmeliet P. Emerging novel functions of the oxygen-sensing prolyl hydroxylase domain enzymes. *Trends Biochem Sci*. 2013;38(1):3-11. doi: 10.1016/j.tibs.2012.10.004. PubMed PMID: 23200187.
59. Bellier A, Chen CS, Kao CY, Cinar HN, Aroian RV. Hypoxia and the hypoxic response pathway protect against pore-forming toxins in *C. elegans*. *PLoS Pathog*. 2009;5(12):e1000689. Epub 2009/12/17. doi: 10.1371/journal.ppat.1000689. PubMed PMID: 20011506.
60. Lee SS, Lee RY, Fraser AG, Kamath RS, Ahringer J, Ruvkun G. A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat Genet*. 2003;33(1):40-8. Epub 2002/11/26. doi: 10.1038/ng1056 ng1056 [pii]. PubMed PMID: 12447374.
61. Hansen M, Hsu AL, Dillin A, Kenyon C. New genes tied to endocrine, metabolic, and dietary regulation of lifespan from a *Caenorhabditis elegans* genomic RNAi screen. *PLoS Genet*. 2005;1(1):119-28. Epub 2005/08/17. doi: 10.1371/journal.pgen.0010017. PubMed PMID: 16103914; PubMed Central PMCID: PMC1183531.
62. Hamilton B, Dong Y, Shindo M, Liu W, Odell I, Ruvkun G, et al. A systematic RNAi screen for longevity genes in *C. elegans*. *Genes Dev*. 2005;19(13):1544-55. Epub 2005/07/07. doi: 19/13/1544 [pii] 10.1101/gad.1308205. PubMed PMID: 15998808.
63. Curran SP, Ruvkun G. Lifespan regulation by evolutionarily conserved genes essential for viability. *PLoS Genet*. 2007;3(4):e56. Epub 2007/04/07. doi: 07-PLGE-RA-0112R1 [pii] 10.1371/journal.pgen.0030056. PubMed PMID: 17411345.
64. Kim Y, Sun H. Functional genomic approach to identify novel genes involved in the regulation of oxidative stress resistance and animal lifespan. *Aging Cell*. 2007;6(4):489-503. Epub 2007/07/05. doi: ACE302 [pii] 10.1111/j.1474-9726.2007.00302.x. PubMed PMID: 17608836.
65. Sugimoto T, Mori C, Takanami T, Sasagawa Y, Saito R, Ichiishi E, et al. *Caenorhabditis elegans* par2.1/mtssb-1 is essential for mitochondrial DNA replication and its defect causes comprehensive

transcriptional alterations including a hypoxia response. *Experimental cell research*. 2008;314(1):103-14. Epub 2007/09/29. doi: 10.1016/j.yexcr.2007.08.015. PubMed PMID: 17900564.

66. Lamitina T, Huang CG, Strange K. Genome-wide RNAi screening identifies protein damage as a regulator of osmoprotective gene expression. *Proc Natl Acad Sci U S A*. 2006;103(32):12173-8. Epub 2006/08/02. doi: 0602987103 [pii] 10.1073/pnas.0602987103. PubMed PMID: 16880390.

67. Nollen EA, Garcia SM, van Haaften G, Kim S, Chavez A, Morimoto RI, et al. Genome-wide RNA interference screen identifies previously undescribed regulators of polyglutamine aggregation. *Proc Natl Acad Sci U S A*. 2004;101(17):6403-8. Epub 2004/04/16. doi: 10.1073/pnas.0307697101 0307697101 [pii]. PubMed PMID: 15084750.

68. Selak MA, Armour SM, MacKenzie ED, Boulahbel H, Watson DG, Mansfield KD, et al. Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF- α prolyl hydroxylase. *Cancer Cell*. 2005;7(1):77-85. Epub 2005/01/18. doi: S153561080400368X [pii] 10.1016/j.ccr.2004.11.022. PubMed PMID: 15652751.

69. Koivunen P, Hirsila M, Remes AM, Hassinen IE, Kivirikko KI, Myllyharju J. Inhibition of hypoxia-inducible factor (HIF) hydroxylases by citric acid cycle intermediates: possible links between cell metabolism and stabilization of HIF. *The Journal of biological chemistry*. 2007;282(7):4524-32. Epub 2006/12/22. doi: M610415200 [pii] 10.1074/jbc.M610415200. PubMed PMID: 17182618.

70. Walker AK, Jacobs RL, Watts JL, Rottiers V, Jiang K, Finnegan DM, et al. A conserved SREBP-1/phosphatidylcholine feedback circuit regulates lipogenesis in metazoans. *Cell*. 2011;147(4):840-52. Epub 2011/11/01. doi: 10.1016/j.cell.2011.09.045. PubMed PMID: 22035958; PubMed Central PMCID: PMC3384509.

71. Simmer F, Moorman C, van der Linden AM, Kuijk E, van den Berghe PV, Kamath RS, et al. Genome-wide RNAi of *C. elegans* using the hypersensitive rrf-3 strain reveals novel gene functions. *PLoS Biol*. 2003;1(1):E12. Epub 2003/10/14. doi: 10.1371/journal.pbio.0000012. PubMed PMID: 14551910; PubMed Central PMCID: PMC212692.

72. Blackwell TK, Bowerman B, Priess JR, Weintraub H. Formation of a monomeric DNA binding domain by Skn-1 bZIP and homeodomain elements. *Science*. 1994;266(5185):621-8. Epub 1994/10/28. PubMed PMID: 7939715.

73. Mello CC, Kramer JM, Stinchcomb D, Ambros V. Efficient gene transfer in *C.elegans*: extrachromosomal maintenance and integration of transforming sequences. *The EMBO journal*. 1991;10(12):3959-70. Epub 1991/12/01. PubMed PMID: 1935914.

74. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res*. 2001;29(9):e45. Epub 2001/05/09. PubMed PMID: 11328886; PubMed Central PMCID: PMC55695.

Supporting information

S1 Fig. *nhr-57* promoter HIF-1 chromatin immunoprecipitation experiments. (A) *nhr-57* promoter sequence and positions of primers used for real-time quantitative PCR assays in HIF-1 chromatin immunoprecipitation experiments. (B) Primer sequences for real-time quantitative PCR assays in *nhr-57* promoter HIF-1 chromatin immunoprecipitation experiments. (C) Chromatin co-

immunoprecipitation data. In these experiments the endogenous *hif-1* locus was disrupted by the *ia04* large deletion and HIF-1 function was restored by the *Phif-1:hif-1a:Myc:HA* transgene [40]. The relative amounts of *nhr-57* promoter regions that co-immunoprecipitated with HIF-1:Myc:HA was determined by real-time quantitative PCR. The bars show the average enriched fold from at least three independent replicates. *inf-1*, a gene not regulated by HIF-1, was used as the reference gene.

S2 Fig. RNAi inactivation of *sams-1* or *sbp-1* increased *Pnhr-57:GFP* expression. (A) RNAi for *sams-1* (S-adenosyl methionine synthetase) increased *Pnhr-57:GFP* expression more than 7-fold in animals carrying the wild-type *hif-1* allele relative to control RNAi, and increased the reporter 3-fold in animals carrying the *hif-1(ia04)* deletion. The difference in RNAi effect between *hif-1(+)* and *hif-1(ia04)* strains is statistically significant ($*p < 0.05$, from six independent experiments, by student *t*-test). (B) RNAi for the SREBP homolog *sbp-1* increased expression of the reporter more than 3-fold in animals carrying the wild-type *hif-1* allele, but had no effect on *Pnhr-57:GFP* expression in *hif-1(ia04)* mutants. The difference in RNAi effect between *hif-1(+)* and *hif-1(ia04)* strains is statistically significant ($*p < 0.05$, from five independent experiments, by student *t*-test). GFP levels were determined by protein blots, and the control animals were fed on bacteria carrying the empty RNAi vector (L4440). The experiments were conducted in RNAi-sensitive strains (*rrf-3(pk1426)*).

S1 Table. Genes increased *Pnhr-57:GFP* expression when knocked-down by RNAi.

S2 Table. Genes for which RNAi caused *hif-1*-dependent increase of *Pnhr-57:GFP* expression.

A

RNAi inactivation of **negative** regulator
of HIF-1 activity

B

■ Metabolism: Electron transport chain

■ Metabolism: Mitochondrial translation

■ Metabolism: Others

■ Protein folding and turnover

■ Vesicular transport

■ Transporters and channels

■ Signal transduction and cytoskeleton

■ Transcription/translation/DNA/RNA processing

■ Others

A

<i>egl-9</i>	-365	C	TTT	GTC	ATC	GTG	TCT	CTT	TG	AT	CCG	-340
<i>gcs-1</i>	-122	C	TTT	AT	CAT	CAT	GA	-	GAT	TTA	ATGTT	-98
<i>med-1</i>	-124	C	TCT	GTC	AT	CAT	GAT	GAT	TTTT	TG	GAG	-99
<i>med-2</i>	-124	C	TCT	GTC	AT	CAT	GAT	GAT	TTTT	TA	GAG	-99

* * * * * * * * * * * *