Embryo Lethality Assay for Evaluating Virulence of Isolates from Bacterial Chondronecrosis with Osteomyelitis in Broilers.

Nnamdi S. Ekesi,^{a,*} Amer Hasan,^{a,b} Alia Parveen,^{a,†} Abdulkarim Shwani,^a Douglas D. Rhoads^{a,#}

^a-Program in Cell and Molecular Biology, University of Arkansas, Fayetteville, Arkansas, USA

^b-Department of Veterinary Public Health, College of Veterinary Medicine, University of Baghdad, Baghdad, Iraq

[#]Corresponding author: 601 Science Engineering, Department of Biological Sciences, University of Arkansas, Fayetteville, AR USA 72701. drhoads@uark.edu

Running Title: ELA for BCO Isolates

Word Count: Abstract: 206; Importance: 72; Text: 2307

*Present address: Nnamdi S. Ekesi, Division of Microbiology, National Center for Toxicological Research, Redfield, AR, USA

†Present address: Alia Parveen, University College Dublin, Belfield, Dublin, Ireland

Abstract

We used an embryo lethality assay (ELA) to assess virulence for different isolates from cases of bacterial chondronecrosis with osteomyelitis (BCO) in broilers. ELA has been used to measure virulence and lethal dosage of *Enterococcus faecalis* and *Enterococcus cecorum*. We hypothesized that ELA could substitute for more laborious and costly assessments of BCO isolate pathogenicity using live birds. We evaluated two different levels of bacteria injected into eggs from layer and commercial broiler embryos. Significant findings include a) *Escherichia coli* from neighboring farms operated by the same integrator had very different embryo lethality, b) isolate *Staphylococcus agnetis* 908 had low virulence in ELA, even though this isolate can induce more than 50% BCO lameness, c) *Enterococcus cecorum* 1415 also had low pathogenicity; even though it was recovered from severe bilateral tibial dyschondroplasia, d) human and chicken isolates of *S. aureus* had significant pathogenicity, e) virulence for some isolates was highly variable possibly corresponding with quality of the embryos/fertile eggs used, and f) ELA pathogenicity was much lower for our BCO isolates than previous reports which may reflect maternal environment. Overall, ELA virulence and BCO virulence are not always concordant indicating that that ELA may not be an effective measure for assessing virulence with respect to BCO.

Importance

Lameness is among the most significant animal welfare issues in the poultry industry. Bacterial infections are a major cause of lameness and different bacterial species have been obtained from lame broilers. Reliable lab-based assays are required to assess relative virulence of

bacteria obtained from lame broilers. Embryo Lethality Assays have been used to compare virulence. Our results suggest that this assay may not be an effective measure of virulence related to lameness.

Keywords: Embryo lethality; Lameness; Virulence; Bacterial chondronecrosis with osteomyelitis

Introduction

Bacterial chondronecrosis with osteomyelitis (BCO) is the leading cause of lameness in rapidly growing broilers (1-4). Lameness in broilers is significant as an animal welfare issue, and as a financial cost, in the poultry industry (2). Our research group isolated and characterized an isolate of *Staphylococcus agnetis*, designated 908, from lame broilers on our research farm (1). *S. agnetis* 908 is closely related to isolates from sub-clinical mastitis in dairy cows (5). *S. agnetis* 908 can induce greater than 50% BCO lameness by 56 days of age when administered in a single dose in drinking water at 10^4 to 10^5 CFU/mL on day 20 (1, 3, 6, 7). Our current model for lameness etiology is that environmental stress can lead to increased leakage (translocation) of bacteria across the gut and pulmonary epithelia into the blood system (1-3, 8-10). Particular species that are able to survive in the blood stream may colonize the growth plate, a vulnerable niche in the blood system of the rapidly growing leg bones of fast-growing broilers (2, 3, 11). Distinct bacterial species have been isolated from lame birds including *Staphylococcus aureus*, *Enterococcus cecorum*, and *Escherichia coli* (12-32). However, there are few comparisons of different BCO-associated species, or isolates, for pathogenicity (18, 33). In this study, we investigated the pathogenicity of BCO isolates of *S. agnetis*, *Staphylococcus chromogenes*, *E. coli*, *E. cecorum*, and *S. aureus* using an embryo lethality assay (ELA). The isolates were obtained

from BCO lesions on our research farm or commercial broiler farms in Arkansas. ELA has been used to correlate the expression frequency of nine virulence-associated *E. coli* genes with embryo mortality (34). Borst et al. (33) used this technique to compare the virulence of *E. cecorum* isolated from broiler spinal lesions (kinky back) to non-pathogenic *E. cecorum* strains isolated from ceca of unaffected birds. Blanco et al. (35) used ELA to determine the virulence and the lethal dose of *Enterococcus faecalis*.

Methods

Microbiology

Isolation and handling of the isolates has been described (1, 36). Media included: CHROMagar Orientation (CO; DRG International, Springfield Township, NJ), tryptic soy broth (Difco brand, Becton, Dickinson and Company, Franklin Lakes, NJ); and Luria broth (LB; per liter 10 g tryptone, 5 g yeast extract, 5 g NaCl).

Embryo Lethality Assay

Fertilized eggs were obtained from leghorns (LCL) and Cobb700 commercial broilers (BCL) on the University of Arkansas research farm. The eggs were washed with warm soapy water containing about 1% household bleach. Eggs were incubated (NatureForm™ Hatchery Systems, Jacksonville, FL, USA) at 37.5°C, a relative humidity of 56%, on autorotate. On day 12, stationary-phase bacterial cultures were diluted 1:200 in 1x phosphate buffer saline (1xPBS; 150mM NaCl, 10 mM potassium phosphate pH 7.2). CFU concentration was computed from Absorbance at 650 nm using pre-calibrated standard curves for each isolate, and then diluted in 1xPBS to the required concentration. Eggs were candled, and fertile eggs were injected using a

tuberculin syringe and 20G needle (Becton, Dickinson and Company) with 100μL of the appropriate bacterial suspension, or vehicle control, into the allantois cavity as described (33). The opening was sealed with transparent box tape. Inoculated embryos were scored for mortality every day for 4 days after bacteria administration (33, 35).

Statistical Analysis

The results of the ELA were analyzed with either Pearson's Chi-squared (χ^2) or Fisher's Exact (FE) analysis using SAS and R software (SAS Institute. 2011; RStudio Team. 2016). Significant differences were accepted at $P < 0.05$.

Results

Embryo Lethality Assay with BCO isolates

To establish a suitable assay for comparing different isolates, we first injected *E. coli* 1413 a dilution series ($10^3 - 10^8$ CFUs) in sterile 1xPBS to estimate the lethal dosage for Leghorn Chicken Line (LCL) embryos (Figure 1). Injections of 1413 above 10^5 CFUs had embryo lethality of 80 to 100%. We therefore assessed different BCO isolates at 10^5 and 10^6 CFUs (Table 1). We included *S. agnetis* 908 recovered from a femoral BCO lesion on our research facility as this isolate can induce lameness $\geq 50\%$ by day 56 when administered in drinking water for two days to 20-day old broilers (1, 3, 6, 7). Surprisingly, 908 injections of even 10^6 CFUs resulted in only 14% embryo lethality, a level not statistically different from 1xPBS control treatment (Figure 2A). For the methicillin-sensitive human *S. aureus* isolate 1302, originally retrieved from a wound (Table 1), injections of 10^5 or 10^6 CFUs resulted in 80% embryo lethality (Figure 2B). *Staphylococcus chromogenes* 1401 was recovered from an infected T4 vertebra of a chicken

with “kinky back” (Table 1). Injections of 10^6 CFUs resulted in only 7% embryo death, less than the 1xPBS control for that experiment (Figure 2C). *E. coli* 1409 was recovered from a tibial head necrosis lesion (Table 1). Injection of 10^5 to 10^6 CFUs resulted in no lethality through day 4 (Figure 2D). *E. coli* 1413 was isolated from the blood of a lame bird with bilateral BCO of the tibiae and femorae, where *E. coli* was also recovered from multiple lesions (Table 1). As before, injections of 10^5 or 10^6 CFUs into LCL resulted in approximately 80% embryo lethality (Figure 2E). *E. cecorum* 1415 was isolated from a tibial head abscess in a case of bilateral tibial dyschondroplasia (Table 1). ELA results for 10^5 CFU showed slightly more lethality than 10^6 CFUs but neither was statistically different from the PBS control (Figure 2F). We used two isolates (1510 & 1514) of *S. aureus* obtained from BCO lesions from two different birds in a commercial broiler house lameness outbreak (Table 1; Ekesi 2020). Draft genome assemblies for these isolates were highly related. The isolates showed different ELA results with 1510 lethality of 60% for 10^6 CFU, while 1514 produced 47% lethality. Only the 1510 results were statistically different from the 1xPBS control (Figure 2G & 2I). *E. coli* 1512 and 1527 were recovered from the left and right femoral lesions of the same bird (Table 1; Ekesi, 2020). Draft genome assemblies for both 1512 and 1527 were determined to be virtually identical. ELA results for 1512 yielded 87% lethality for 10^6 CFUs and 52% lethality for 10^5 , but only the 10^6 results were statistically different from the 1xPBS control (Figure 2H). Therefore, the only isolates that showed significant lethality using LCL embryos were the human isolate *S. aureus* 1302, and chicken isolates *E. coli* 1413, *S. aureus* 1510, and *E. coli* 1512.

We then extend the analyses to ELA using Broiler Chicken Line (BCL) embryos. As shown in Figure 3, significant embryo lethality was obtained with *S. aureus* 1302 and 1514, and *E. coli*

1413 and 1512. *S. agnetis* 908, *S. chromogenes* 1401, *E. coli* 1409, and *E. cecorum* 1415, showed no virulence for either 10^5 or 10^6 CFU. We did note that for all four isolates that showed lethality for BCL, both the 10^5 and 10^6 CFU injections showed significant embryo mortality (Figure 3 panels B, E, G, and H). For LCL, the 10^5 injections were only different from the 1xPBS control for *S. aureus* 1302 and *E. coli* 1413. However, for the other two isolates, we might reach significance for the 10^5 CFU injections with more embryos. We also note that lethality was more rapid in the BCL than with LCL embryos (Figures 2 and 3).

1413 *Repeatability of ELA as a Measure of Virulence*

We noted that there was occasional variation in the ELA results for some isolates including *S. aureus* 1510 and 1514, which should be nearly identical, and *E. coli* 1409. We therefore compiled results from 11 experiments using LCL or BCL injected with 1xPBS or *E. coli* 1409 spanning nearly two years (Table 2). Embryo survival at day 5 for 1409 ranged from 100 to 11% in LCL and 73 to 44% in BCL. Close inspection of the data across experiments shows that embryo survival for 1x PBS was also varied with different batches of embryos. For the experiment of 2/20/020 the 1xPBS control was actually lower than that for 1409. We did not see this level of variability for *E. coli* 1413 as it was always highly virulent. We suspect that the variable ELA results for some isolates is likely due to differences in the particular set of eggs available for experiments. Factors affecting embryo quality might include age of hens, environmental factors in the housing (feed, heat, cold, air-quality, lighting, etc.), or post-lay conditions for specific groups of eggs (days post lay, fresh collection, etc.).

1413 **Discussion**

We performed ELA with different bacterial isolates isolated from lame broilers to estimate relative pathogenicity. We observed that *S. agnetis* 908 is not pathogenic in the ELA even though we have shown that this isolate readily infects young broilers when administered at 10^4 or 10^5 CFU/ml in drinking water at 20 and 21 days of age (1, 3, 6, 7). Broilers challenged with *S. agnetis* 908 begin to develop lameness by 41 days of age and by 56 days of age 50% of the birds will be clinically lame with BCO lesions in proximal femoral and tibial heads. Many of the birds develop bacteremia with hundreds to thousands of CFU/ml of blood. Additionally, the treated birds spread the infection to birds within the same room, with 30-40% of those birds lame by 56 days of age (3, 6). ELA in broiler embryos has been used to compare virulence of *E. cecorum* from BCO birds (primarily kinky back) and feces (33). Kinky back isolates showed lethality of >70% while cecal isolates showed lethality of <40%. We used *E. cecorum* 1415 collected from an infected vertebrae in a kinky back bird (36) but it showed no significant lethality (<40%) in the ELA. We compared three *E. coli* isolates from BCO lame birds (36) and found they had very different apparent virulence in the ELA. Even though these three isolates were from three different commercial broiler farms in Arkansas that were experiencing BCO outbreaks, we have shown that all 3 are very different based on whole-genome comparisons (36). This is surprising given that 1409 and 1413 were isolated on the same day from two different farms within 5 km of each other that were operated by the same integrator and supplied from the same hatchery. *S. aureus* isolates showed virulence in the ELA, including an isolate from human infection, and isolates from a BCO outbreak on a different farm operated by a different integrator. We sampled 11 lame birds from that farm and determined that 7 of the birds were infected with *S. aureus*. Genome analyses showed that the BCO *S. aureus* isolates were highly related and very

closely related to *S. aureus* isolates obtained from diseased birds or broiler meat dating from 2010 in Oklahoma all the way back to the 1970s in Europe (36, 37). The clade has been identified multiple times in Arkansas and Oklahoma for at least a decade. The clade appears to be exclusively associated with poultry, so virulence in the ELA is not surprising. The genomic comparisons of *E. coli* and *S. aureus* chicken BCO isolates led us to propose that *E. coli* association with BCO is not exclusively poultry specific and that this species appears to be more of a generalist, whereas *S. aureus* and *S. agnetis* appear to be specialists and do not readily jump back and forth, infecting different host species (5, 36).

Comparison of our results with those of Borst *et al.* (33) identified several differences. They used BCO isolates and cecal isolates of *E. cecorum* in specific pathogen free (SPF) and broiler embryos. They reported that injections of 10^2 CFU of the BCO isolates resulted in viabilities of 33% for broiler and 42% in SPF, embryos. With higher inocula the SPF embryos appeared more susceptible than the broiler embryos. Further, broiler embryos injected with 10^2 of cecal isolates produced higher viability, circa 60%. In our experiments significant loss of viability required 10^5 CFU. Although there was some loss of viability at lower injection levels we did not reach statistical significance for 10^3 or 10^4 CFU, when we used our most virulent isolate, *E. coli* 1413. Lack of significance at the lower injection quantities reflects that we used only 5 embryos per injection level. When we used more (n=15) embryos for the screens of different isolates we still had many isolates that failed to show significance when the viabilities were above 60% even with 10^5 or 10^6 CFU. In our hands, with our broiler embryo sources, far higher numbers of bacteria were required to reach significant levels of lethality. This difference may be because our embryos were from broilers (and layers) raised under standard conditions, whereas Borst *et*

al. used embryos from SPF stocks and a broiler “flock intensively managed and has no history of *E. cecorum*-associated disease” (33). Thus, in our breeder stocks there may be higher maternal exposure to bacterial pathogens which would result in significant deposition of maternal antibodies in the eggs, which could provide enhanced immunity to the embryos. Considering that we saw marked differences in the ELA results for *E. coli* 1409 with different sets of embryos across different flocks, the quality and sources of embryos appear to be critical. Indeed, in our experiments, broiler embryos generally were more susceptible than our layer embryos. Our data suggest caution in drawing broad conclusions when using ELA for evaluation of relative virulence of BCO isolates. Concerns include: a) the lack of significant virulence by *S. agnetis* 908 which we have shown is hypervirulent in inducing lameness and spreading through a broiler facility (3, 6, 7), b) the lack of virulence of *E. cecorum* 1415 which should have been virulent based on earlier reports using ELA (33), and c) the variability of ELA results with different sets of embryos for hypo- or moderately virulent isolates (e.g., *E. coli* 1409). There remains a need to develop new, convenient, and reliable, laboratory assays for assessing pathogenicity for chicken bacterial isolates related to BCO.

Acknowledgements

Partial support for this work was from Zinpro LLC, Chr Hansen, and the Arkansas Biosciences Institute. The funders had not input on the design or interpretations of these experiments.

AH was supported by a scholarship from the Iraqi Ministry of Higher Education and Scientific Research. AP was supported in part by a Fulbright Fellowship.

References

- 208 1. Al-Rubaye AAK, Couger MB, Ojha S, Pummill JF, Koon JA, II, Wideman RF, Jr., Rhoads DD. 2015.
209 Genome analysis of *Staphylococcus agnetis*, an agent of lameness in broiler chickens. PLoS One
210 10:e0143336.
- 211 2. Wideman RF. 2016. Bacterial chondronecrosis with osteomyelitis and lameness in broilers: a
212 review. Poult Sci 95 325-344.
- 213 3. Al-Rubaye AAK, Ekesi NS, Zaki S, Emami NK, Wideman RF, Rhoads DD. 2017. Chondronecrosis
214 with osteomyelitis in broilers: Further defining a bacterial challenge model using the wire
215 flooring model. Poult Sci 96:332-340.
- 216 4. Weimer SL, Wideman RF, Scanes CG, Mauromoustakos A, Christensen KD, Vizzier-Thaxton Y.
217 2019. The utility of infrared thermography for evaluating lameness attributable to bacterial
218 chondronecrosis with osteomyelitis. Poult Sci 98:1575-1588.
- 219 5. Shwani A, Adkins PRF, Ekesi NS, Alrubaye A, Calcutt MJ, Middleton JR, Rhoads DD. 2020. Whole
220 genome comparisons of *Staphylococcus agnetis* isolates from cattle and chickens. Appl Environ
221 Microbiol 86:e00484-20.
- 222 6. Alrubaye A, Ekesi NS, Hasan A, Koltes DA, Wideman Jr R, Rhoads D. 2020. Chondronecrosis with
223 osteomyelitis in broilers: Further defining a bacterial challenge model using standard litter
224 flooring and protection with probiotics. Poult Sci 99:6474-6480.
- 225 7. Alrubaye AAK, Ekesi NS, Hasan A, Elkins E, Ojha S, Zaki S, Dridi S, Wideman RF, Rebollo MA,
226 Rhoads DD. 2020. Chondronecrosis with Osteomyelitis in Broilers: Further Defining Lameness-
227 Inducing Models with Wire or Litter Flooring, to Evaluate Protection with Organic Trace
228 Minerals. Poult Sci 99:5422-5429.
- 229 8. Wideman RF, Prisby RD. 2013. Bone circulatory disturbances in the development of
230 spontaneous bacterial chondronecrosis with osteomyelitis: A translational model for the
231 pathogenesis of femoral head necrosis. Front Endocrinol (Lausanne) 3:183.

- 232 9. Jiang T, Mandal RK, Jr. RFW, Khatiwara A, Pevzner I, Kwon YM. 2015. Molecular survey of
233 bacterial communities associated with bacterial chondronecrosis with osteomyelitis (BCO) in
234 broilers. PLoS One 10:e0124403.
- 235 10. Mandal RK, Jiang T, Al-Rubaye AA, Rhoads DD, Wideman RF, Zhao J, Pevzner I, Kwon YM. 2016.
236 An investigation into blood microbiota and its potential association with bacterial
237 chondronecrosis with osteomyelitis (BCO) in broilers. Sci Rep 6:25882.
- 238 11. Wideman RF, Al-Rubaye A, Gilley A, Reynolds D, Lester H, Yoho D, Hughes JM, Pevzner I. 2013.
239 Susceptibility of 4 commercial broiler crosses to lameness attributable to bacterial
240 chondronecrosis with osteomyelitis. Poult Sci 92:2311-2325.
- 241 12. Carnaghan RBA. 1966. Spinal cord compression due to spondylitis caused by *Staphylococcus*
242 *pyogenes*. J Comp Pathol 76:9-14.
- 243 13. Duff SR. 1984. Dyschondroplasia of the caput femoris in skeletally immature broilers. Res Vet Sci
244 37:293-302.
- 245 14. Emslie KR, Fenner LM, Nade S. 1984. Acute haematogenous osteomyelitis: II. The effect of a
246 metaphyseal abscess on the surrounding blood supply. J Pathol 142:129-134.
- 247 15. Duff S, Randall C. 1987. Observations on femoral head abnormalities in broilers. Res Vet Sci
248 42:17-23.
- 249 16. Duff S. 1989. Disturbed endochondral ossification in the axial skeleton of young fowls. J Comp
250 Pathol 101:399-409.
- 251 17. Duff SRI. 1989. Physeal clefts and disturbed endochondral ossification in broiler fowls. J Comp
252 Pathol 101:75-86.
- 253 18. Braga JFV, Chanteloup NK, Trotureau A, Baucheron S, Guabiraba R, Ecco R, Schouler C. 2016.
254 Diversity of *Escherichia coli* strains involved in vertebral osteomyelitis and arthritis in broilers in
255 Brazil. BMC Vet Res 12:140.

- 256 19. Nairn ME, Watson ARA. 1972. Leg weakness of poultry—a clinical and pathological
257 characterisation. Aust Vet J 48:645-656.
- 258 20. McCaskey PC, Rowland GN, Page RK, Minear LR. 1982. Focal Failures of Endochondral
259 Ossification in the Broiler. Avian Dis 26:701-717.
- 260 21. Mutalib A, Riddell C, Osborne AD. 1983. Studies on the pathogenesis of Staphylococcal
261 osteomyelitis in chickens. II. Role of the respiratory tract as a route of infection. Avian Dis
262 27:157-160.
- 263 22. Riddell C. 1983. Pathology of the skeleton and tendons of broiler chickens reared to roaster
264 weights. I. crippled chickens. Avian Dis 27:950-962.
- 265 23. Griffiths G, Hopkinson W, Lloyd J. 1984. Staphylococcal necrosis of the head of the femur in
266 broiler chickens. Aust Vet J 61:293-293.
- 267 24. Duff SRI. 1990. Diseases of the musculoskeletal system. In Jordan F (ed), Poultry Diseases 3rd ed.
268 Bailliere, Tindall. UK.
- 269 25. Hocking PM. 1992. Musculo-skeletal disease in heavy breeding birds., p 297-309. In Whitehead
270 CC (ed), Bone Biology and Skeletal Disorders in Poultry. Carfax Publishing Company, Abingdon,
271 United Kingdom.
- 272 26. Tate CR, Mitchell WC, Miller RG. 1993. *Staphylococcus hyicus* associated with turkey stifle joint
273 osteomyelitis. Avian Dis 37:905-907.
- 274 27. Thorp BH. 1994. Skeletal disorders in the fowl: a review. Avian Pathol 23:203-236.
- 275 28. Skeeles KJ. 1997. Staphylococcosis, p 247-253. In B.W. Calnek, Barnes HJ, Beard CW, McDougald
276 LR, Saif YM (ed), Diseases of Poultry 10 ed. Iowa State University Press, Ames, IA, USA.
- 277 29. Thorp B, Waddington D. 1997. Relationships between the bone pathologies, ash and mineral
278 content of long bones in 35-day-old broiler chickens. Res Vet Sci 62:67-73.

30. McNamee PT, Smyth JA. 2000. Bacterial chondronecrosis with osteomyelitis ('femoral head necrosis') of broiler chickens: a review. *Avian Pathol* 29:477-495.
31. Joiner KS, Hoerr FJ, van Santen E, Ewald SJ. 2005. The avian major histocompatibility complex influences bacterial skeletal disease in broiler breeder chickens. *Vet Pathol* 42:275-281.
32. Gaußmann B, Hess C, Grafl B, Kovacs M, Troxler S, Stessl B, Hess M, Paudel S. 2018. *Escherichia coli* isolates from femoral bone marrow of broilers exhibit diverse pheno- and genotypic characteristics that do not correlate with macroscopic lesions of bacterial chondronecrosis with osteomyelitis. *Avian Pathol* 47:271-280.
33. Borst LB, Suyemoto MM, Keelara S, Dunningan SE, Guy JS, Barnes HJ. 2014. A Chicken Embryo Lethality Assay for Pathogenic *Enterococcus cecorum*. *Avian Dis* 58:244-248.
34. Oh JY, Kang MS, Yoon H, Choi HW, An BK, Shin EG, Kim YJ, Kim MJ, Kwon JH, Kwon YK. 2012. The embryo lethality of *Escherichia coli* isolates and its relationship to the presence of virulence-associated genes. *Poult Sci* 91:370-375.
35. Blanco AE, Caverio D, Icken W, Voss M, Schmutz M, Preisinger R, Sharifi AR. 2018. Genetic approach to select against embryo mortality caused by *Enterococcus faecalis* infection in laying hens. *Poult Sci* 97:4177-4186.
36. Ekesi NS, Hasan A, Alrubaye A, Rhoads D. 2021. Analysis of Genomes of Bacterial Isolates from Lameness Outbreaks in Broilers. *Poult Sci* (in review).
37. Lowder BV, Guinane CM, Ben Zakour NL, Weinert LA, Conway-Morris A, Cartwright RA, Simpson AJ, Rambaut A, Nübel U, Fitzgerald JR. 2009. Recent human-to-poultry host jump, adaptation, and pandemic spread of *Staphylococcus aureus*. *PNAS USA* 106:19545-19550.

301 Table 1: Details of bacterial isolates utilized for ELA, including species, strain designation, host,
302 and primary citation. In Isolate Source Lane indicates a bird number from BCO sampling.
303 Abbreviations include: LT: Left Tibia, RT: Right tibia, LF: Left femur, and RF: Right femur.

Species	Designation	Host	Isolate Source	Citation
<i>S. agnetis</i>	908	Broiler	Femoral BCO; UA Research Farm	(1)
<i>S. chromogenes</i>	1401	Broiler	Thoracic Vertebrae; Lane3	(36)
<i>E. cecorum</i>	1415	Broiler	LT/RT; Lane5	(36)
<i>E. coli</i>	1409	Broiler	RT; Lane3	(36)
	1413	Broiler	Blood; Lane12	(36)
	1512	Broiler	LF; Lane18	(36)
	1527	Broiler	RF; Lane18	(36)
<i>S. aureus</i>	1510	Broiler	LT Lane14	(36)
	1514	Broiler	RF Lane15	(36)
	1302	Human	Wound	ATCC 29213

304

Table 2. Variability of ELA results for *E. coli* 1409. On the given dates for the indicated count (n) of embryos from layers (LCL) or broilers (BCL), were injected with 100 μ l of 1x PBS or 10⁶ CFU of *E. coli* 1409. Percent embryo viability is presented on the given days post injection. Averages and SEM are provided at the bottom across all experiments.

Date	n	Embryos	Inject	Day			
				1	2	3	4
11/21/2018	5	LCL	1x PBS	100	100	100	100
11/21/2018	5	LCL	1409	100	80	80	80
12/14/2018	15	LCL	1x PBS	100	100	100	100
12/14/2018	15	LCL	1409	100	100	100	100
12/31/2019	15	LCL	1x PBS	100	87	87	87
12/31/2019	15	LCL	1409	73	40	33	33
1/20/2020	15	LCL	1x PBS	100	100	100	100
1/20/2020	15	LCL	1409	87	67	60	47
2/20/2020	15	LCL	1x PBS	87	67	47	47
2/20/2020	15	LCL	1409	87	67	67	67
3/20/2020	9	LCL	1x PBS	100	100	89	89
3/20/2020	9	LCL	1409	33	33	22	11
3/31/2020	15	LCL	1x PBS	100	100	100	100
3/31/2020	15	LCL	1409	60	60	60	47
9/6/2020	10	LCL	1409	70	60	60	60
9/25/2020	14	LCL	1x PBS	100	100	100	100
9/25/2020	14	LCL	1409	50	29	29	14
1/5/2019	9	BCL	1x PBS	100	100	100	100
1/5/2019	9	BCL	1409	44	44	44	44
1/22/2019	15	BCL	1x PBS	100	100	100	100
1/22/2019	15	BCL	1409	87	80	73	73
Average		All	1x PBS	99	95	92	92
		LCL	1409	73	60	57	51
		BCL	1409	66	62	59	59
		All	1x PBS	1	3	5	5
SEM		LCL	1409	8	8	9	10
		BCL	1409	22	18	15	15

Figure 1. Layer chicken line embryo lethality for day post injection (DPI) with 1xPBS, different numbers of colony forming units (CFU) of *E. coli* 1413. The percentage (n=5) of live embryos (Y-axis) for different doses is graphed for four days post-injection (X-axis). Isolate source is described in Table 1. Asterisk (*) indicates where treatment was significantly different from 1xPBS (P < 0.05) on day 4.

Figure 2. Layer chicken line embryo lethality for day post injection (DPI) with 1xPBS, 10^5 , or 10^6 CFU of different bacterial isolates. Isolates were: A: 908 (n=14), B: 1302 (n=15), C: 1401 (n=14), D: 1409 (n=15), E: 1413 (n=15), F: 1415 (n=9), G: 1510 (n=15), H: 1512 (n=15), and I: 1514 (n=15). Details are as in Figure 1. Asterisk (*) indicates treatments significantly different from 1xPBS ($P < 0.05$) on day 4.

Figure 3. Broiler chicken line embryo lethality for injections of 10^5 or 10^6 CFU of bacterial isolates. Isolates were: A. 908, B. 1302, C. 1401, D. 1409, E. 1413, F. 1415, G. 1514, and H.1527. Details are as in Figures 1 and 2. For each treatment n=15.