Thermal biology of two sympatric Lacertids lizards (*Lacerta diplochondrodes* and *Parvilacerta parva*) from Western Anatolia

Mehmet Kürşat Şahin¹, Arda Cem Kuyucu²

¹Karamanoğlu Mehmet Bey University, Department of Biology,

²Turkey, Hacettepe University, Department of Biology, Faculty of Science, Ankara, Turkey

Abstract

Sympatric lizard species differing in morphology present convenient models for studying the differentiation in thermal behavior and the role of morphological differences in thermal biology. Here we studied the thermal biology of two sympatric lizard species which occur together sympatrically in western Anatolia, Frig Valley. These two species differ in body size, with the larger *Lacerta diplochondrodes* and smaller *Parvilacerta parva*. Field body temperatures of the individuals belonging to both species were recorded in the activity period. Additionally, several environmental parameters including solar radiation, substrate temperature, air temperature and wind speed were also monitored to investigate the relative effect of these abiotic parameters on thermal biology of the two species. The field body temperature and temperature excess (difference between body and substrate temperature) of two species while being relatively close to each other, showed seasonal differences. Solar radiation, substrate temperature and air temperature were the main effective factors on thermal biology in the field. Additionally, although body size did not have a direct significant effect on body temperature or temperature excess, the interaction between body size and wind were effective on temperature excess. In conclusion, our study partially supports the conservation of thermal biology of related lizard species.

Keywords: *Lacerta diplochondrodes*, *Parvilacerta parva*, thermal biology, body size, body temperature.

1. Introduction

Temperature is one of the most important environmental parameters that strongly influence almost all biochemical reactions and thus most physiological processes linked to fitness of organisms (Angilletta, 2009; Hochachka and Somero, 2002; Ortega et al., 2016a; Pörtner, 2002; Somero, 2004). Eventually, body temperature is one of the most effective factors on the performance of ectothermic animals (Gaitán-Espitia et al., 2014; Huey and Kingsolver, 1989; Huey and Pianka, 2018; Huey and Stevenson, 1979). Thermal performance of ectotherms generally conforms to an asymmetric reaction-norm curve where optimal performance peaks between maximal and minimal thresholds (Angilletta, 2006; Chown et al., 2010; Huey and Stevenson, 1979; Kingsolver and Huey, 1998; Rezende and Bozinovic, 2019; Schulte et al., 2011). Organisms strive to keep their body temperature close to this optimum by thermoregulation to maximize performance in life history including growth, reproduction, foraging, locomotion (Angilletta, 2009; Aparicio Ramirez et al., 2020; Lailvaux and Irschick, 2007; Rutschmann et al., 2020; Seebacher and Franklin, 2005).

Body temperatures of ectotherms, including reptiles, strongly depend on thermal properties of their microhabitats (Goller et al., 2014; Gómez Alés et al., 2017; Huey and Slatkin, 1976; Ortega and Pérez-Mellado, 2016). However, terrestrial environments show thermal variation to varying degrees in several spatial and temporal levels, including altitudinal (Kuyucu et al., 2018; Lymburner and Blouin-Demers, 2020; Van Damme et al., 1989; Zamora-Camacho et al., 2016), seasonal (Díaz et al., 2006; Huey and Pianka, 1977; Liz et al., 2019; Van Damme et al., 1987) and microhabitat level (Adolph, 1990; Goller et al., 2014; Ortega et al., 2016b; Scheers and Van

Damme, 2002). Because of this heterogeneity of the thermal environment, behavioral thermoregulation holds crucial importance for ectothermic organisms including reptiles (Cowles and Bogert, 1944; Huey and Slatkin, 1976; Kearney et al., 2009; McMaster and Downs, 2013; Rangel-Patiño et al., 2020). The thermoregulatory strategies vary both among and between species, which depends on the costs and benefits of thermoregulation which also depend on several biotic and abiotic factors in the environment (Huey and Slatkin 1976, Lymburner and Blouin Demers 2020). Species that rely on behavioral thermoregulation can utilize solar radiation by basking (heliothermy) or regulate their heat exchange with the substrate (thigmothermy) (Belluere and Carrascal, 2002; Garrick, 2008; Penniket and Cree, 2015; Rutschmann et al., 2020) to keep their body temperature in set point limits. Most Lacertids are classified as heliothermic, since they effectively shuttle between sun and shade in order to keep their body temperature (T_b) in certain optimum intervals (Basson et al., 2017; Díaz et al., 1996; Van Damme et al., 1987).

It is shown that even sympatric squamate species that live in the same environments might differ in their usage of thermal micro-habitats (Angilletta, 2009; Gomes et al., 2016; Gómez Alés et al., 2017). Therefore, differentiations in thermoregulatory strategies can also result in alterations for their habitat use in spatial and temporal scales (Buckley et al., 2015; Lymburner and Blouin-Demers, 2020; Neel and McBrayer, 2018; Paterson and Blouin-Demers, 2018). Microhabitat selection (Castilla and Bauwens, 1991; Ortega and Pérez-Mellado, 2016), reproduction activities (Ibargüengoytia et al., 2008; Luo et al., 2010; Meiri et al., 2012; Wang et al., 2016), foraging rates (Kearney et al., 2020; Verwaijen and Van Damme, 2007) and growth patterns (Angilletta et al., 2002; Have and Jong, 1996; Refsnider et al., 2019; Sears et al., 2016) are partly determined by thermal factors in ectotherms' environment. The heterogeneity of the spatial thermal

environment has an important role in thermoregulation (Logan et al., 2019, 2015; Sears et al., 2011). This heterogeneity provides more opportunities for activities and regulations for lizards by enabling lizards to behaviorally regulate their energetic thermoregulation costs (Ortega and Martín-Vallejo, 2019).

Air temperature, substrate temperature, humidity, solar radiation, wind speed, availability of shade can be assessed as some of the most important environmental factors that influence thermal activities of lizards (Fei et al., 2012b; Kiefer et al., 2007; Logan et al., 2015). In addition to abiotic factors in the environment morphological and physical characteristics can also influence the thermal biology of ectotherms, such as coloration and body size (Clusella-Trullas and Chown, 2014; Sagonas et al., 2013a; Tanaka, 2005). For instance, darker or melanic coloration might increase the amount of solar radiation absorbed by integument, providing potential advantage to darker colored ectotherms in colder environments (Castella et al., 2013; Clusella-Trullas et al., 2008; Matthews et al., 2016; Stuart-fox et al., 2017). Another important factor on heat exchange is body size, case studies on ectotherm species have shown that generally there is a negative correlation between body size and heat exchange rate (Belluere and Carrascal, 2002; Herczeg et al., 2007; Lactin and Johnson, 1997; Rutschmann et al., 2020), i.e. larger individuals heat up more slowly compared to smaller ones by absorbing solar radiation and conduction with hot substrates (Belluere and Carrascal 2002), however this might also lead to losing heat in a slower rate in colder environments (Garrrick, 2008). Small size also can be both an advantage and disadvantage, for example it might enable smaller ectotherms to heat up more rapidly to optimal temperatures, which is particularly useful for lizards under predator pressure and should move quickly when predation danger is present, on the other hand this higher heating rate carries with it the danger of overheating and individuals have to retreat

quickly to shelters before body temperatures reaches harmful levels under very high temperatures (Gardner et al., 2011). Additionally, individuals with larger body size are able to maintain their body temperature more efficiently in shelter after basking, needing less shuttling between different thermal spaces, reducing the exposure time to predators (Dzialowski and O'Connor, 2001; O'Connor et al., 2000; Sagonas et al., 2013a), on the other hand large ectotherms are also sensitive to overheating in hot and open environments as they also cool down slower (Maia-Carneiro and Rocha, 2013; Porter and Gates, 1969). Wind speed is another important parameter that might be influential on body temperature of ectotherms due to heat loss by convection, and high wind speed might significantly reduce the effectiveness of thermoregulation (Maia-Carnero et al. 2012, Logan et al. 2015, Ortega et al. 2017).

Here, we study and compare the thermal biology of two sympatric lizards in western Anatolia with emphasis on important biotic and abiotic variables in the environment, including body size, sex, air temperature, substrate temperature, solar radiation and wind speed. We monitored *Lacerta diplochondrodes* and *Parvilacerta parva* in Phrygian valley during their active season by several *in situ* observations. Thermoregulation dynamics of various lacertid species were studied in mainland and islands of Greece in the last decade (Sagonas et al. 2013 a,b). However, there is a serious lack of field-based thermal ecology studies on not only our model species but also for all terrestrial reptiles in the Anatolian Peninsula. Therefore, three questions were set for this work: i) Do these two species differ in their thermal biology? ii) To what degree body size is effective on this potential difference. iii) Which microclimatic variables are effective on the thermal biology of these two species and to what extent?

2. Materials and methods

2.1 Model organisms

Lacerta diplochondrodes Wettstein, 1952 (formerly known as *Lacerta trilineata* in Western Anatolia) a large-sized lizard of Lacertidae, is evaluated to be endemic to the Anatolian Peninsula after recent phylogeographic queries (Kornilios et al., 2020). This large-sized Lacertid can reach up to 50 cm as a body length. The dorsal colour is brownish and there are 5 light coloured longitudinal lines present in juveniles. There may also be a dashed line on both sides of the ventralia. As the lizard grows the color of dorsalia turns to green, light lines disappear and small and dark spots occur densely. These spots are generally more common in females. Light blue coloring is observed on both sides of the head in males. The ventralia is yellowish white in males and pinkish yellow in females. It lives in steppe habitat, where shrubs and rocks are abundant. Despite its large size, its movements are very fast and it tends to hide between the roots of shrubs.

The smaller sized *Parvilacerta parva* (Boulenger, 1887) inhabits steppe vegetation along inner parts of Anatolian Peninsula and Armenian mountainous steppes. It is a dwarf lizard with a body length of up to 14 cm. The dorsal plates are keeled, and the coloration of this part is grayish or light brown with black and white spots. Similarly, there are also spots on lateral sides of the body. While the coloration of ventralia is yellowish or white in males, it is generally white in females. Therefore, the highlighted dimorphic morphological character is the presence of blue spots on ventral laterals in only males.

2.2 Study area

Although the entire Phrygian region covers parts of Ankara, Afyonkarahisar and Eskişehir provinces, plus most of Kütahya province and northern slopes of Konya, Isparta and Burdur in Western Anatolia, the study area range is only northern slopes of this region in Afyonkarahisar (Lat: 39.046638, Lon: 30.522929, Elevation: 1145 m) (Fig. 1). There are many deep valleys in this slope and elevation varies between 1100 and 1500 meters. The rock formations are shaped from soft volcanic tuffs, which are dated to Tertiary Era (Gürsoy et al., 2003). This also allows metamorphism of the greywackes and their sedimentation by corrosion process causes alluvial soil type. The microclimate provided by this soil type has its own characteristics: Hot and dry in summers and cool and less humid in winters. Vegetation in the Phrygian Valley is highly dominated by floristic steppe elements. The study area, which is the northern slopes of valley, includes various annual *Poaceae* species, that covers almost the entire area. Apart from that, shrubs are abundant in several areas, where *L. diplochondrodes* specimens were observed. However, due to Valley's geological status tuff based bedrock landscape is surrounding the study area (Gürsoy et al., 2003).

The mean annual temperature in the study area is 10.2° C and as a result of terrestrial climate, the hottest months are July (22.1° C) and August (20.6° C) (TSMS, 2020). The precipitation is high in spring but mean precipitation is still low due to two factors: i) distance from the sea side, ii) low humidity. The mean annual precipitation is 406.1 mm and the cloudiness rate is 4.7/10 (TSMS, 2020). Therefore, according to Köppen-Geiger climate classification, the climatological conditions of the study area displays “Csa” pattern (Warm climate with hot dry summer) (Beck et al., 2018; Turkes, 2020).

2.3. *Field Sampling*

Lizards from both species were simultaneously studied between May and September 2018 for 45 non-rainy days (9 days in May, 7 days in June, 5 days in July, 15 days in August and 9 days in September). In order to evaluate the activity and variation in body temperature of lizards, they were captured from 07.00 to 18.00 during their diurnal activity times by visual encounter survey (VES) method (Maia-Carneiro et al., 2012), an effective method for capturing the lizards which live in open habitats and are locally abundant.

The body temperatures (T_b) were measured with an infrared thermometer (Fluke 62 MAX). Due to their low invasiveness, non-contact methods such as infrared thermometers and infrared thermography are now widely used in thermal biology studies on reptiles (Barroso et al., 2016; Carretero, 2012; Tattersall, 2016). The measurements were taken from lizards' dorsal part between their head and neck immediately 2-5 seconds after capture to avoid heat transfer to animal (0.1°C sensitivity). The hemipenis presence/absence is also an easy way for the sexual identification of specimens. Total body length (BL) was measured from snout to the tip of the tail by a digital caliper (Ecotone ®, to the nearest 0.1 mm). After these examinations, the lizards were released to the same point where they were originally captured.

Air temperatures (T_a) were taken with a digital thermometer (HTC 288-ATH ®). Substrate temperatures (T_s) were taken with the same digital infrared laser thermometer (Fluke Model 62 ®) that was used for body temperature measurements from the lizard capture point. Substrate type was recorded from the lizard capture point as grass, rock or soil. Wind speed (m/s) was measured with a Windtronic 2 ® anemometer and solar radiation (sol) (w/m^2) was measured with an Apogee ® pyranometer.

All field works were done under permission of the Ministry of Agriculture and Forest (27.02.2018, 72784983-488.04-49994) and Hacettepe University Local Ethical Committee (31.01.2018/52338575-4).

2.4. Statistical Analysis

We built two ordinary least square linear models for T_b and T_{ex} . Compliance to normal distribution of parameters were checked with Shapiro-Wilks test. The distribution of T_b did not conform to a normal distribution, approaching a more leptokurtic distribution so we used a bijective inverse transformation using the `lambertW` package for R (Goerg, 2020) to transform the data to a normal distribution. T_b was taken as the dependent variable and species, sampling month (May to September), sampled substrate (soil, grass and rock) and sex (juvenile, male and female) were determined as categorical predictors while body size (measured as body length in mm), air temperature (T_a), substrate temperature (T_s), solar radiation (w/m^2) and wind speed (m/s) were continuous predictors. Additionally, we included the following interactions, body length (BL): solar radiation, body size: wind speed, species: month and species: substrate. As the distribution of T_{ex} conformed to a normal distribution, we built the linear model without data transformation to compare the effects of same environmental and morphological parameters on temperature excess including interactions, except T_a which was left out which since it was directly related with T_{ex} . After building the initial models we used AIC (Akaike Information Criteria) (Aho et al., 2014; Burnham and Anderson, 2004; Symonds and Moussalli, 2011) backward stepwise elimination to gather the most informative model. Variance inflation was also checked and variables with $VIF > 4$ were excluded from the final models to minimize collinearity. The substrate use between species were compared by building cross tables for

proportion of observations in three substrate types and comparisons were made by chi-square test. All statistical analyses and transformations were carried out in R statistical software (R. Core Team 2018).

3. Results

On general both mean T_b (33.7 ± 0.13 °C) and T_{ex} (9.38 ± 0.14 °C) of the smaller (BL = 114.59 ± 0.84 mm) *P. parva* was higher compared to larger *L. diplochondrodes* (T_b : 32 ± 0.12 °C, T_{ex} : 8.99 ± 0.18 °C, BL = 326.2 ± 2.55 mm) (Table 1). As expected, body temperature reached its highest points during the middle of summer for both species. It can be seen from Table 2 and Fig 2 that temperature excess parameters show a monthly fluctuating pattern, as for both species T_{ex} decreased between May and June, increasing again between June and July and decreasing after, on the other hand although T_{ex} continued to decrease for *Parvilacerta*, for *Lacerta diplochondrodes*, it increased again in September (Table 2). According to the final model for body temperature, sampling month (September), type of substrate, T_a , T_s and solar radiation were significant factors. Additionally, although species was not a significant factor the interaction between species (*P. parva*) and substrate (rock) had significant effect (Table 3). For the temperature excess model, significant factors were sampling month, substrate (rock) and solar radiation. Additionally, we found that interactions species: month, species: substrate and body size: wind speed had significant effect on temperature excess (Table 4). The interaction between species and months i.e. the differential pattern of temperature excess is apparent in Fig. 3. We also detected that the two species differed in habitat use where *P. parva* had higher preference for grassy habitat (54% grass, 42% soil, 3% rock) whereas *L. diplochondrodes* had higher preference for open soil (29% grass, 69% soil, 1% rock) ($\chi^2 = 77.25$, df = 2, $p < 0.001$).

4. Discussion

Here we studied the thermal biology of two sympatric Lacertid species (*L. diplochondrodes* and *P. parva*) in Phrygian Valley, Turkey. As expected for heliothermic lizards that depend on basking for thermoregulation; air temperature (T_a), substrate temperature (T_s), seasonality (months) and solar radiation were the most effective parameters on both body temperature and temperature excess (Clusella-Trullas et al., 2008; Garrick, 2008; Matthews et al., 2016). On the other hand, body size was not a statistically significant factor in T_b and T_{ex} by itself. Although numerous studies (especially those which successfully implement detailed biophysical models into experimental study frames) documented the significant effect of body size on heating and cooling rate of lizards (Fei et al., 2012a; Garrick, 2008; Herczeg et al., 2007; Shine and Kearney, 2001), this relation might not be detected directly in field conditions where several other important environmental, morphological and behavioral factors can conceal this interaction (Christian et al., 2006; Corkery et al., 2018; Sagonas et al., 2013a; Stevenson, 1985). First, the effect of body size is a two way process as increased thermal inertia slows both the heat gain and heat loss from ectotherm bodies (Garrick, 2008; Herczeg et al., 2007; Maia-Carneiro and Rocha, 2013; Zamora-Camacho et al., 2014). On the other hand, despite not observing a substantial relation between either body size or wind speed and T_{ex} , we detected a significant negative effect of the interaction between body size and wind speed on T_{ex} . This finding points that heat loss by convection of a larger area of exposed body surface might be an important factor on thermal biology in the field. Previously, it was shown that increased wind speed can significantly decrease the effectiveness of thermoregulation and impend activity in exposed open high elevation habitats (Huang et al., 2020; Logan et al., 2015; Maia-Carneiro et al., 2012; Ortega et al., 2017). Our study area is an open steppe that is relatively more exposed to wind effects with

low bushes and grass as vegetation, additionally the wind speed in the area was relatively high during the measurements (mean wind speed: 2-5 m/s). Added together these components might be increasing the effect of wind speed on body temperatures in the field.

Although observation of gender differences in thermal biology of reptiles are relatively rare, gender as a factor should be taken into consideration due to the biological importance of these rare cases (Huey and Pianka, 2007). A proportion of cases where significant difference observed for thermal biology this variation can be connected with sexual size dimorphism (Lailvaux, 2007; Ortega et al., 2016b). On the other hand thermal biology differences between genders might also be connected with other gender specific differences in performance and behavior such as locomotion, stamina, assimilation efficiency and micro-habitat preference (Beal et al., 2014; Lailvaux et al., 2003). In our study there was not any visible sexual size dimorphism for both species and we did not detect any significant effect of gender or ontogeny (between juveniles and adults) on body temperature or temperature excess. These results suggest that regardless of gender or ontogeny the individuals belonging to two species might have similar thermoregulatory behaviour in the field.

Although we measured body temperatures from the neck region by infrared thermography our measurements ($32 \pm 0.12^{\circ}\text{C}$) for *L. diplochondrodes* were close to body temperature measurements taken from *L. trilineata*, a very closely related species in mainland Greece [Stymphalia: $31.94 \pm 1.65^{\circ}\text{C}$, Karditsa: $32.30 \pm 1.54^{\circ}\text{C}$ st.dev, (Sagonas et al., 2013c)]. Additionally, the relatively close pattern of body temperatures across different months suggest that these two species maintain similar body temperatures to each other through variable environmental temperatures. These results are in compliance with results from other studies and reviews, which suggest that thermal biology of lizards is a relatively conserved trait across

different species (Clusella-Trullas and Chown, 2014; Gómez Alés et al., 2017; Grigg et al., 2013; Rowe et al., 2020).

It was shown that sympatrically coexisting lizards might differentiate their niches, and this also might affect their thermoregulatory behavior (Akashi et al., 2016; Gomes et al., 2016; Gómez Alés et al., 2017; Sagonas et al., 2017). In our study study we have found that the two species differed in their preferences for different habitat types. On the other hand, we did not detect a significant relation between thermal biology and preference of grass and soil substrates, which are the main habitat types in the study area. The smaller *P. parva* might be preferring grassy habitats for the suitability to hide smaller lizards from predators. The main difference for T_b and T_{ex} between species were seasonal as shown by the effect of the interaction between month and species, thus our results confirm that seasonality is the main factor on the field body temperature of lacertids, considering that other abiotic parameters including temperature and solar radiation are also highly dependent on seasonality (Ortega and Pérez-Mellado, 2016). Additionally, aside from the direct effects of seasonality on body temperature other important life history activities such as reproduction, foraging and feeding depend on both thermal biology and seasonality patterns (Adolph and Porter, 1996; Christian et al., 2003; Van Damme et al., 1987). This seasonal difference was much more pronounced for T_{ex} as *P. parva* shows higher T_{ex} for both July and August with high significance (Table 1) while for T_b this effect was marginally significant for August (Table 2). The relatively stable state of T_b compared to the difference in T_{ex} between months indicates that the role of behaviour in maintenance of optimal body temperature might be important, as the higher amount of difference between T_b and T_a in warmer periods of year might indicate a higher frequency of overheating avoidance (Vicente Liz et al., 2019). Given that both of these two species are diurnal with most active times being spring and

summer, efficient thermoregulatory behaviour is crucial for heat avoidance while maintaining an optimal body temperature for foraging and metabolism (Huey and Pianka, 2018; Sepúlveda et al., 2014). It is also striking that while smaller *P. parva* has higher T_{ex} than *L. diplochondrodes* in July when ground temperature is at its highest, this picture is reversed in September when temperatures are lower. Since we did not detect a direct effect of body size on temperature excess this pattern might have been caused by the temporal behavioral differences between the two species.

Regarding the usage of substrate in the study area, the effect of rock type of substrate was significant and negative on T_b and T_{ex} . However, the interaction between *P. parva* and rock substrate was significant and positive. Which might be due to the effect of rocky substrate being smaller on *P. parva*, on the other hand both of these species rarely prefer rock, as we observed the individuals from two species on rocky substrates only in 22 times out of 1059 total observations. In addition rocky substrates in the study area were relatively rare with most of the area being either shrubs or naked soil.

Although thermal ecology of various lizard species was investigated in Aegean Islands and mainland Greece in the east Mediterranean region (Sagonas et al., 2017, 2013c, 2013b) to our knowledge this study is the first to investigate the thermal biology of lizard species occurring in Anatolia. On the other hand, additional studies on the operative temperatures (Hertz et al., 1993; Ortega et al., 2014; Sagonas et al., 2013a), voluntary and critical thermal limits (Aparicio Ramirez et al., 2020; Camacho and Rusch, 2017; Qu et al., 2011) and also thermal preferences (Goller et al., 2014; Van Berkel and Clusella-Trullas, 2018; Wang et al., 2013) are necessary for a more realized thermal ecology framework for these species. We believe that this study will also be a first step in constructing a framework for predicting the responses of reptile populations

inhabiting western Anatolia to climate change. As thermal traits, particularly thermal behavior and tolerance plays a significant role in determining the responses of ectothermic species to global environmental change (Buckley et al., 2015; Enriquez-Urzelai et al., 2020; Kearney et al., 2009), knowledge of the thermal ecology and physiology differences and microhabitat preferences would provide useful information to infer and prevent the effects of climate change on reptile species under threat (Ceia-Hasse et al., 2014; Kearney et al., 2009).

Acknowledgments

M.K. Sahin contributed to the study design, collected the field data and contributed to the manuscript draft. A.C. Kuyucu designed and supervised the study, carried out the statistical analysis and drafted the manuscript. This work was supported by Hacettepe University Scientific Research Coordination Unit. (Project Number FHD -2018-16903).

Data accessibility

Data supporting this study are provided in the link, doi:10.17632/g868g5m438.1

References

- Adolph, S.C., 1990. Influence of behavioral thermoregulation on microhabitat use by two *Sceloporus* lizards. *Ecology* 71, 315–327.
- Adolph, S.C., Porter, W.P., 1996. Growth , seasonality , and lizard life histories : age and size at maturity. *Oikos* 77, 267–278.
- Aho, K., Derryberry, D., Peterson, T., 2014. Model selection for ecologists: the worldview of AIC and BIC. *Ecology* 95, 631–636.
- Akashi, H.D., Cádiz Díaz, A., Shigenobu, S., Makino, T., Kawata, M., 2016. Differentially expressed genes associated with adaptation to different thermal environments in three sympatric Cuban Anolis lizards. *Mol. Ecol.* 25, 2273–2285.
- Angilletta, M.J., 2009. Thermal adaptation: a theoretical and empirical synthesis. Oxford University

Press, USA.

- Angilletta, M.J., 2006. Estimating and comparing thermal performance curves. *J. Therm. Biol.* 31, 541–545.
- Angilletta, M.J., Niewiarowski, P.H., Navas, C.A., 2002. The evolution of thermal physiology in ectotherms. *J. Therm. Biol.* 27, 249–268.
- Aparicio Ramirez, A., Perez, K., Telemeco, R.S., 2020. Thermoregulation and thermal performance of crested geckos (*Correlophus ciliatus*) suggest an extended optimality hypothesis for the evolution of thermoregulatory set-points. *J. Exp. Zool. Part A Ecol. Integr. Physiol.* 1–10.
- Barroso, F.M., Carretero, M.A., Silva, F., Sannolo, M., 2016. Assessing the reliability of thermography to infer internal body temperatures of lizards. *J. Therm. Biol.* 62, 90–96.
- Basson, C.H., Levy, O., Angilletta, M.J., Clusella-Trullas, S., 2017. Lizards paid a greater opportunity cost to thermoregulate in a less heterogeneous environment. *Funct. Ecol.* 31, 856–865.
- Beal, M.S., Lattanzio, M.S., Miles, D.B., 2014. Differences in the thermal physiology of adult Yarrow’s spiny lizards (*Sceloporus jarrovi*) in relation to sex and body size. *Ecol. Evol.* 4, 4220–4229.
- Beck, H.E., Zimmermann, N.E., McVicar, T.R., Vergopolan, N., Berg, A., Wood, E.F., 2018. Present and future köppen-geiger climate classification maps at 1-km resolution. *Sci. Data* 5, 1–12.
- Belliure, J., Carrascal, L.M., 2002. Influence of heat transmission mode on heating rates and on the selection of patches for heating in a mediterranean lizard. *Physiol. Biochem. Zool.* 75, 369–376.
- Buckley, L.B., Ehrenberger, J.C., Angilletta, M.J., 2015. Thermoregulatory behaviour limits local adaptation of thermal niches and confers sensitivity to climate change. *Funct. Ecol.* 29, 1038–1047.
- Burnham, K.P., Anderson, D.R., 2004. Multimodel inference: Understanding AIC and BIC in model selection. *Sociol. Methods Res.* 33, 261–304.
- Camacho, A., Rusch, T.W., 2017. Methods and pitfalls of measuring thermal preference and tolerance in lizards. *J. Therm. Biol.* 68, 63–72.
- Carretero, M.A., 2012. Measuring body temperatures in small lacertids: Infrared vs. contact thermometers. *Basic Appl. Herpetol.* 26, 99–105.
- Castella, B., Golay, J., Monney, J.C., Golay, P., Mebert, K., Dubey, S., 2013. Melanism, body condition and elevational distribution in the asp viper. *J. Zool.* 290, 273–280.
- Castilla, A.M., Bauwens, D., 1991. Thermal biology, microhabitat selection, and conservation of the insular lizard *Podarcis hispanica atrata*. *Oecologia* 85, 366–374.

- Ceia-Hasse, A., Sinervo, B., Vicente, L., Pereira, H.M., 2014. Integrating ecophysiological models into species distribution projections of European reptile range shifts in response to climate change. *Ecography (Cop.)*. 37, 679–688.
- Chown, S.L., Hoffmann, A.A., Kristensen, T.N., Angilletta, M.J., Stenseth, N.C., Pertoldi, C., 2010. Adapting to climate change: A perspective from evolutionary physiology. *Clim. Res.* 43, 3–15.
- Christian, K.A., Tracy, C.R., Tracy, C.R., 2006. Evaluating thermoregulation in reptiles: an appropriate null model. *am. nat.* 168, 421–430.
- Christian, K.A., Webb, J.K., Schultz, T.J., 2003. Energetics of bluetongue lizards (*Tiliqua scincoides*) in a seasonal tropical environment. *Oecologia* 136, 515–523.
- Clusella-Trullas, S., Chown, S.L., 2014. Lizard thermal trait variation at multiple scales: A review. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.*
- Clusella-Trullas, S., Terblanche, J.S., Blackburn, T.M., Chown, S.L., 2008. Testing the thermal melanism hypothesis: A macrophysiological approach. *Funct. Ecol.* 22, 232–238.
- Corkery, I., Bell, B.D., Nelson, N.J., 2018. Thermoregulation of a temperate reptile in a forested habitat. *Zoology* 127, 63–69.
- Cowles, R.B., Bogert, C.M., 1944. A preliminary study of the thermal requirements of desert reptiles. *Iguana* 83, 53.
- Díaz, J.A., Bauwens, D., Asensio, B., 1996. A comparative study of the relation between heating rates and ambient temperatures in Lacertid lizards. *Physiol. Zool.* 69, 1359–1383.
- Díaz, J.A., Iraeta, P., Monasterio, C., 2006. Seasonality provokes a shift of thermal preferences in a temperate lizard, but altitude does not. *J. Therm. Biol.* 31, 237–242.
- Dzialowski, E.M., O'Connor, M.P., 2001. Physiological control of warming and cooling during simulated shuttling and basking in lizards. *Physiol. Biochem. Zool.* 74, 679–693.
- Enriquez-Urzelai, U., Tingley, R., Kearney, M.R., Sacco, M., Palacio, A.S., Tejedo, M., Nicleza, A.G., 2020. The roles of acclimation and behaviour in buffering climate change impacts along elevational gradients. *J. Anim. Ecol.* 89, 1722–1734.
- Fei, T., Skidmore, A.K., Venus, V., Wang, T., Schlerf, M., Toxopeus, B., van Overjijk, S., Bian, M., Liu, Y., 2012a. A body temperature model for lizards as estimated from the thermal environment. *J. Therm. Biol.* 37, 56–64.
- Fei, T., Skidmore, A.K., Venus, V., Wang, T., Toxopeus, B., Bian, M., Liu, Y., 2012b. Predicting micro

- thermal habitat of lizards in a dynamic thermal environment. *Ecol. Modell.* 231, 126–133.
- Gaitán-Espitia, J.D., Bacigalupe, L.D., Opitz, T., Lagos, N.A., Timmermann, T., Lardies, M.A., 2014. Geographic variation in thermal physiological performance of the intertidal crab *Petrolisthes violaceus* along a latitudinal gradient. *J. Exp. Biol.* 217, 4379–4386.
- Gardner, J.L., Peters, A., Kearney, M.R., Joseph, L., Heinsohn, R., 2011. Declining body size: A third universal response to warming? *Trends Ecol. Evol.* 26, 285–291.
- Garrick, D., 2008. Body surface temperature and length in relation to the thermal biology of lizards. *Biosci. Horizons Int. J. Student Res.* 1, 136–142.
- Goerg GM (2020). LambertW: Probabilistic Models to Analyze and Gaussianize Heavy-Tailed, Skewed Data. R package: version 0.6.6.
- Goller, M., Goller, F., French, S.S., 2014. A heterogeneous thermal environment enables remarkable behavioral thermoregulation in *Uta stansburiana*. *Ecol. Evol.* 4, 3319–3329.
- Gomes, V., Carretero, M.A., Kaliontzopoulou, A., 2016. The relevance of morphology for habitat use and locomotion in two species of wall lizards. *Acta Oecologica* 70, 87–95.
- Gómez Alés, R., Acosta, J.C., Laspiur, A., 2017. Thermal biology in two syntopic lizards, *Phymaturus extrilidus* and *Liolaemus parvus*, in the Puna region of Argentina. *J. Therm. Biol.* 68, 73–82.
- Grigg, J.W., Buckley, L.B., Grigg, J.W., Buckley, L.B., 2013. Conservatism of lizard thermal tolerances and body temperatures across evolutionary history and geography Conservatism of lizard thermal tolerances and body temperatures across evolutionary history and geography.
- Gürsoy, H., Piper, J.D.A., Tatar, O., 2003. Neotectonic deformation in the western sector of tectonic escape in Anatolia: Palaeomagnetic study of the Afyon region, central Turkey. *Tectonophysics* 374, 57–79.
- Have, V. Der, Jong, D., 1996. Adult size in ectotherms: temperature effects on growth and differentiation. *J. Theor. Biol.* 183, 329–340.
- Herczeg, G., Török, J., Korsós, Z., 2007. Size-dependent heating rates determine the spatial and temporal distribution of small-bodied lizards. *Amphib. Reptil.* 28, 347–356.
- Hertz, P.E., Huey, R.B., Stevenson, R.D., 1993. Evaluating temperature regulation by field-active ectotherms: The fallacy of the inappropriate question. *Am. Nat.* 142, 796–818. <https://doi.org/10.1086/285573>
- Hochachka, P.W., Somero, G.N., 2002. Biochemical adaptation: mechanism and process in physiological

- evolution. Oxford university press, 2002.
- Huang, S.P., Lin, Y.C., Lin, T.E., Richard, R., 2020. Thermal physiology explains the elevational range for a lizard, *Eutropis longicaudata*, in Taiwan. J. Therm. Biol. 93.
- Huey, R.B., Kingsolver, J.G., 1989. Evolution of thermal sensitivity of ectotherm performance. Trends Ecol. Evol. 4, 131–135.
- Huey, R.B., Pianka, E.R., 2018. Body temperature distributions of active diurnal lizards in three deserts: Skewed up or skewed down? Funct. Ecol. 32, 334–344.
- Huey, R.B., Pianka, E.R., 2007. Lizard thermal biology: Do genders differ? Am. Nat. 170, 473–478.
- Huey, R.B., Pianka, E.R., 1977. Seasonal variation in thermoregulatory behavior and body temperature of diurnal Kalahari lizards. Ecology 58, 1066–1075.
- Huey, R.B., Slatkin, M., 1976. Cost and benefits of lizard thermoregulation. Q. Rev. Biol. 51, 363–384.
- Huey, R.B., Stevenson, R., 1979. Integrating thermal physiology and ecology of ectotherms: A discussion of approaches. Am. Zool. 19, 357– 366.
- Ibargüengoytia, N.R., Acosta, J.C., Boretto, J.M., Villavicencio, H.J., Marinero, J.A., Krenz, J.D., 2008. Field thermal biology in *Phymaturus* lizards: Comparisons from the Andes to the Patagonian steppe in Argentina. J. Arid Environ. 72, 1620–1630.
- Kearney, M., Shine, R., Porter, W.P., 2009. The potential for behavioral thermoregulation to buffer “cold-blooded” animals against climate warming. Proc. Natl. Acad. Sci. U. S. A. 106, 3835–3840. <https://doi.org/10.1073/pnas.0808913106>
- Kearney, M.R., Porter, W.P., Huey, R.B., 2020. Modelling the joint effects of body size and microclimate on heat budgets and foraging opportunities of ectotherms. Methods Ecol. Evol. 0–2.
- Kiefer, M.C., Van Sluys, M., Rocha, C.F.D., 2007. Thermoregulatory behaviour in *Tropidurus torquatus* (Squamata, Tropiduridae) from Brazilian coastal populations: An estimate of passive and active thermoregulation in lizards. Acta Zool. 88, 81–87.
- Kingsolver, J.G., Huey, R.B., 1998. Evolutionary analyses of morphological and physiological plasticity in thermally variable environments. Am. Zool. 38, 1–15.
- Kornilios, P., Thanou, E., Lymberakis, P., Ilgaz, Ç., Kumlutaş, Y., Leaché, A., 2020. A phylogenomic resolution for the taxonomy of Aegean green lizards. Zool. Scr. 49, 14–27.
- Kuyucu, A.C., Sahin, M.K., Caglar, S.S., 2018. The relation between melanism and thermal biology in a colour polymorphic bush cricket, *Isophya rizeensis*. J. Therm. Biol. 71, 212–220.

- Lactin, D.J., Johnson, D.L., 1997. Response of body temperature to solar radiation in restrained nymphal migratory grasshoppers (Orthoptera: Acrididae): Influences of orientation and body size. *Physiol. Entomol.* 22, 131–139.
- Lailvaux, S.P., 2007. Interactive effects of sex and temperature on locomotion in reptiles. *Integr. Comp. Biol.* 47, 189–199.
- Lailvaux, S.P., Alexander, G.J., Whiting, M.J., 2003. Sex-based differences and similarities in locomotor performance, thermal preferences, and escape behaviour in the lizard *Platysaurus intermedius wilhelmi*. *Physiol. Biochem. Zool.* 76, 511–21.
- Lailvaux, S.P., Irschick, D.J., 2007. Effects of temperature and sex on jump performance and biomechanics in the lizard *Anolis carolinensis*. *Funct. Ecol.* 21, 534–543.
- Liz, A.V., Santos, V., Ribeiro, T., Guimarães, M., Verrastro, L., 2019. Are lizards sensitive to anomalous seasonal temperatures? Long-term thermobiological variability in a subtropical species. *PLoS One* 14, 1–16.
- Logan, M.L., Fernandez, S.G., Calsbeek, R., 2015. Abiotic constraints on the activity of tropical lizards. *Funct. Ecol.* 29, 694–700.
- Logan, M.L., van Berkel, J., Clusella-Trullas, S., 2019. The Bogert Effect and environmental heterogeneity. *Oecologia* 191, 817–827.
- Luo, L.G., Ding, G.H., Ji, X., 2010. Income breeding and temperature-induced plasticity in reproductive traits in lizards. *J. Exp. Biol.* 213, 2073–2078.
- Lymburner, A.H., Blouin-Demers, G., 2020. Changes in thermal quality of the environment along an elevational gradient affect investment in thermoregulation by Yarrow’s spiny lizards. *J. Zool.* 312, 133–143.
- Maia-Carneiro, T., Dorigo, T.A., Rocha, C.F.D., 2012. Influences of seasonality, thermal environment and wind intensity on the thermal ecology of Brazilian sand lizards in a Restinga remnant. *South Am. J. Herpetol.* 7, 241–251.
- Maia-Carneiro, T., Rocha, C.F.D., 2013. Influences of sex, ontogeny and body size on the thermal ecology of *Liolaemus lutzae* (Squamata, Liolaemidae) in a restinga remnant in southeastern Brazil. *J. Therm. Biol.* 38, 41–46.
- Matthews, G., Goulet, C.T., Delhey, K., Chapple, D.G., 2016. The effect of skin reflectance on thermal traits in a small heliothermic ectotherm. *J. Therm. Biol.* 60, 109–124.

- McMaster, M.K., Downs, C.T., 2013. Thermoregulation in leopard tortoises in the Nama-Karoo: The importance of behaviour and core body temperatures. *J. Therm. Biol.* 38, 178–185.
- Meiri, S., Brown, J.H., Sibly, R.M., 2012. The ecology of lizard reproductive output. *Glob. Ecol. Biogeogr.* 21, 592–602.
- Neel, L.K., McBrayer, L.D., 2018. Habitat management alters thermal opportunity. *Funct. Ecol.* 32, 2029–2039.
- O'Connor, M.P., Zimmerman, L.C., Dzialowski, E.M., Spotila, J.R., 2000. Thick-walled physical models improve estimates of operative temperatures for moderate to large-sized reptiles. *J. Therm. Biol.* 25, 293–304.
- Ortega, Z., Martín-Vallejo, F.J., 2019. Main factors affecting lacertid lizard thermal ecology. *Integr. Zool.* 14, 293–305.
- Ortega, Z., Mencía, A., Pérez-Mellado, V., 2017. Wind constraints on the thermoregulation of high mountain lizards. *Int. J. Biometeorol.* 61, 565–573.
- Ortega, Z., Mencía, A., Pérez-Mellado, V., 2016a. The peak of thermoregulation effectiveness: Thermal biology of the Pyrenean rock lizard, *Iberolacerta bonnali* (Squamata, Lacertidae). *J. Therm. Biol.* 56, 77–83.
- Ortega, Z., Mencía, A., Pérez-Mellado, V., 2016b. Sexual differences in behavioral thermoregulation of the lizard *Scelarcis perspicillata*. *J. Therm. Biol.* 61, 44–49.
- Ortega, Z., Pérez-Mellado, V., 2016. Seasonal patterns of body temperature and microhabitat selection in a lacertid lizard. *Acta Oecologica* 77, 201–206.
- Ortega, Z., Pérez-Mellado, V., Garrido, M., Guerra, C., Villa-García, A., Alonso-Fernández, T., 2014. Seasonal changes in thermal biology of *Podarcis lilfordi* (Squamata, Lacertidae) consistently depend on habitat traits. *J. Therm. Biol.* 39, 32–39.
- Paterson, J.E., Blouin-Demers, G., 2018. Density-dependent habitat selection predicts fitness and abundance in a small lizard. *Oikos* 127, 448–459.
- Penniket, S., Cree, A., 2015. Adherence to Bergmann's rule by lizards may depend on thermoregulatory mode: support from a nocturnal gecko. *Oecologia* 178, 427–440.
- Porter, W.P., Gates, D.M., 1969. thermodynamic equilibria of animals with environment. *Ecol. Monogr.* 39, 227–244.
- Pörtner, H.O., 2002. Climate variations and the physiological basis of temperature dependent

- biogeography: Systemic to molecular hierarchy of thermal tolerance in animals. *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* 132, 739–761.
- Qu, Y., Li, H., Gao, J., Xu, X., Ji, X., 2011. Thermal preference, thermal tolerance and the thermal dependence of digestive performance in two *Phrynocephalus* lizards (Agamidae), with a review of species studied. *Curr. Zool.* 57, 684–700.
- Rangel-Patiño, C.A., Mastachi-Loza, C.A., Eifler, D., García-Morales, C., Ruiz-Gómez, M. de L., 2020. When things get hot: Thermoregulation behavior in the lizard *Sceloporus aeneus* at different thermal conditions. *J. Therm. Biol.* 89.
- Refsnider, J.M., Clifton, I.T., Vazquez, T.K., 2019. Developmental plasticity of thermal ecology traits in reptiles: Trends, potential benefits, and research needs. *J. Therm. Biol.* 84, 74–82.
- Rezende, E.L., Bozinovic, F., 2019. Thermal performance across levels of biological organization. *Philos. Trans. R. Soc. B Biol. Sci.* 374.
- Rowe, J.W., Clark, D.L., Martin, C.E., Valle, C., 2020. Diel and seasonal variations in the thermal biology of San Cristobal Lava Lizards (*Microlophus bivittatus*). *J. Therm. Biol.* 88, 102518.
- Rutschmann, A., Rozen-rechels, D., Dupoué, A., Blaimont, P., de Villemereuil, P., Miles, D.B., Richard, M., Clobert, J., 2020. Climate dependent heating and cooling efficiency in the common lizard. 1–26.
- Sagonas, K., Kapsalas, G., Valakos, E., Pafilis, P., 2017. Living in sympatry: The effect of habitat partitioning on the thermoregulation of three Mediterranean lizards. *J. Therm. Biol.* 65, 130–137.
- Sagonas, K., Meiri, S., Valakos, E.D., Pafilis, P., 2013a. The effect of body size on the thermoregulation of lizards on hot, dry Mediterranean islands. *J. Therm. Biol.* 38, 92–97.
- Sagonas, K., Meiri, S., Valakos, E.D., Pafilis, P., 2013b. The effect of body size on the thermoregulation of lizards on hot, dry Mediterranean islands. *J. Therm. Biol.* 38, 92–97.
- Sagonas, K., Valakos, E.D., Pafilis, P., 2013c. The impact of insularity on the thermoregulation of a Mediterranean lizard. *J. Therm. Biol.* 38, 480–486.
- Scheers, H., Van Damme, R., 2002. Micro-scale differences in thermal habitat quality and a possible case of evolutionary flexibility in the thermal physiology of lacertid lizards. *Oecologia* 132, 323–331.
- Schulte, P.M., Healy, T.M., Fanguie, N.A., 2011. Thermal performance curves, phenotypic plasticity, and the time scales of temperature exposure. *Integr. Comp. Biol.* 51, 691–702.
- Sears, M.W., Angilletta, M.J., Schuler, M.S., Borchert, J., Dilliplane, K.F., Stegman, M., Rusch, T.W., Mitchell, W.A., 2016. Configuration of the thermal landscape determines thermoregulatory

- performance of ectotherms. Proc. Natl. Acad. Sci. 113, 10595 LP – 10600.
- Sears, M.W., Raskin, E., Angilletta Jr, M.J., 2011. The world is not flat: defining relevant thermal landscapes in the context of climate change. Integr. Comp. Biol. 51, 666–675.
- Seebacher, F., Franklin, C.E., 2005. Physiological mechanisms of thermoregulation in reptiles: A review. J. Comp. Physiol. B Biochem. Syst. Environ. Physiol. 175, 533–541.
- Sepúlveda, M., Sabat, P., Porter, W.P., Fariña, J.M., 2014. One solution for two challenges: The lizard *Microlophus atacamensis* avoids overheating by foraging in intertidal shores. PLoS One 9, 1–11.
- Shine, R., Kearney, M., 2001. Field Studies of Reptile Thermoregulation : How Well Do Physical Models Predict Operative Temperatures ? Func. Ecol. 15(2), 282-288.
- Somero, G.N., 2004. Adaptation of enzymes to temperature: Searching for basic “strategies.” Comp. Biochem. Physiol. - B Biochem. Mol. Biol. 139, 321–333.
- Stevenson, R.D., 1985. Body size and limits to the daily range of body temperature in terrestrial ectotherms. Am. Nat. 125, 102–117.
- Stuart-fox, D., Newton, E., Clusella-trullas, S., Stuart-fox, D., 2017. Thermal consequences of colour and near-infrared reflectance. Phil. Trans. R. Soc. B 372, 20160345.
- Symonds, M.R.E., Moussalli, A., 2011. A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike’s information criterion. Behav. Ecol. Sociobiol. 65, 13–21.
- Tanaka, K., 2005. Thermal Aspects of Melanistic and Striped Morphs of the Snake *Elaphe quadrivirgata*. Zoolog. Sci. 22, 1173–1179.
- Tattersall, G.J., 2016. Infrared thermography: A non-invasive window into thermal physiology. Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol. 202, 78–98.
- Turkes, M., 2020. Climate and Drought in Turkey, in: Water Resources of Turkey. Springer, pp. 85–125.
- TSMS, Turkish State of Meteorological Service, 2020. Link: <https://mgm.gov.tr/eng/forecast-cities.aspx>.
- Van Berkel, J., Clusella-Trullas, S., 2018. Behavioral thermoregulation is highly repeatable and unaffected by digestive status in *Agama atra*. Integr. Zool. 13, 482–493.
- Van Damme, R., Bauwens, D., Castilla, A.M., Verheyen, R.F., 1989. Altitudinal variation of the thermal biology and running performance in the lizard *Podarcis tiliguerta*. Oecologia 80, 516–524.
- Van Damme, R., Bauwens, D., Verheyen, R.F., 1987. Thermoregulatory responses to environmental

- seasonality by the lizard *Lacerta vivipara*. *Herpetologica* 43, 405–415.
- Verwajen, D., Van Damme, R., 2007. Correlated evolution of thermal characteristics and foraging strategy in lacertid lizards. *J. Therm. Biol.* 32, 388–395.
- Vicente Liz, A., Santos, V., Ribeiro, T., Guimarães, M., Verrastro, L., 2019. Are lizards sensitive to anomalous seasonal temperatures? Long-term thermobiological variability in a subtropical species. *PLoS One* 14, e0226399.
- Wang, Y., Zeng, Z.G., Li, S.R., Bi, J.H., Du, W.G., 2016. Low precipitation aggravates the impact of extreme high temperatures on lizard reproduction. *Oecologia* 182, 961–971.
- Wang, Z., Lu, H., Ma, L., Ji, X., 2013. Differences in thermal preference and tolerance among three *Phrynocephalus* lizards (Agamidae) with different body sizes and habitat use. *Asian Herpetol. Res.* 4, 214–220.
- Zamora-Camacho, F.J., Reguera, S., Moreno-Rueda, G., 2016. Thermoregulation in the lizard *Psammodromus algirus* along a 2200-m elevational gradient in Sierra Nevada (Spain). *Int. J. Biometeorol.* 60, 687–697.
- Zamora-Camacho, F.J., Reguera, S., Moreno-Rueda, G., 2014. Bergmann’s Rule rules body size in an ectotherm: heat conservation in a lizard along a 2200-metre elevational gradient. *J. Evol. Biol.* 27, 2820–2828.

Figures

Figure 1. Study area: Phrygian Valley, Afyon, Turkey

A)

B)

Figure 2. A) Monthly boxplots of body temperature (T_b) values for *L. diplochondrodes* and *P. parva*. B) Monthly boxplots of temperature excess (T_{ex}) values for *L. diplochondrodes* and *P. parva*.

Figure 3. **A)** Boxplots of temperature excess (T_{ex}) values for *L. diplochondrodes* and *P. parva*. **B)** Boxplot values of body temperature (T_b) values for *L. diplochondrodes* and *P. parva*. **C)** Scatterplot of body temperature (T_b) with substrate temperature (T_s). **D)** Scatterplot of temperature excess (T_{ex}) with solar radiation (w/m^2).

Tables

Table 1 Mean body temperature (T_b), mean temperature excess (T_{ex}) and mean body length of two lizard species.

Species	Mean $T_b \pm$ St. error	$T_{ex} \pm$ St. error	BL $\pm$ St. error
<i>Lacerta diplochondrodes</i>	32 $\pm$ 0.12 °C	8.99 $\pm$ 0.18 °C	326.2 $\pm$ 2.55 mm
<i>Parvilacerta parva</i>	33.7 $\pm$ 0.13 °C	9.38 $\pm$ 0.13 °C	114.59 $\pm$ 0.84 mm

Table 2 Monthly Mean body temperature (T_b) and mean temperature excess (T_{ex}) of two lizard species.

Species	Month	$T_b \pm$ St. error	$T_{ex} \pm$ St. error	N
<i>Lacerta diplochondrodes</i>	May	30.6 $\pm$ 0.25 °C	11.1 $\pm$ 0.37 °C	91
	June	32.4 $\pm$ 0.14 °C	9.1 $\pm$ 0.47 °C	70
	July	33.1 $\pm$ 0.13 °C	10.4 $\pm$ 0.36 °C	49
	August	32.9 $\pm$ 0.21 °C	6.83 $\pm$ 0.27 °C	150
	September	30.9 $\pm$ 0.28 °C	9.62 $\pm$ 0.41 °C	90
<i>Parvilacerta parva</i>	May	32.6 $\pm$ 0.19 °C	11.3 $\pm$ 0.24 °C	119
	June	34.7 $\pm$ 0.25 °C	9.35 $\pm$ 0.25 °C	89
	July	34.9 $\pm$ 0.34 °C	12.4 $\pm$ 0.34 °C	58
	August	34.9 $\pm$ 0.18 °C	8.34 $\pm$ 0.22 °C	223
	September	31.4 $\pm$ 0.39 °C	7.97 $\pm$ 0.3 °C	120

Table 3 Results of the best model for parameters effective on body temperature (T_b).

	Coefficient	Std.error	t value	P
Intercept	24.07	0.89	27.10	<0.001***
Species: <i>P. parva</i>	0.58	0.48	1.21	0.23
June	-0.001	0.36	-0.002	~1
July	0.39	0.40	0.98	0.33
August	-0.18	0.34	-0.54	0.59
September	-0.72	0.34	-2.11	0.034*
Substrate: Rock	-1.31	0.60	-2.18	0.03*
Substrate: Soil	0.081	0.20	0.41	0.68
Body length	-0.0008	0.0018	-0.44	0.66
Solar rad	0.002	0.00048	4.60	<0.001***
T_a	0.18	0.024	7.57	<0.001***
T_s	0.093	0.021	4.43	<0.001***
Species: <i>P. parva</i> June	0.41	0.44	0.92	0.36
Species: <i>P. parva</i> July	0.83	0.5	1.63	0.1
Species: <i>P. parva</i> August	0.92	0.39	2.37	0.018*
Species: <i>P. parva</i> September	-0.51	0.44	-1.16	0.25
Species: <i>P. parva</i> Substrate: Rock	2.73	0.99	2.77	<0.01**
Species: <i>P. parva</i> Substrate: Soil	-0.12	0.27	-0.43	0.67
*: 0.01 < p < 0.05 **: 0.001 < p < 0.01 ***: p < 0.001				

Table 4 Results of the best model for parameters effective on body temperature (T_{ex}).

	Coefficient	Std.error	t value	P
Intercept	11.45	1.34	8.56	<0.001***
Species: <i>P. parva</i>	-0.87	0.75	-1.28	0.20
Month:June	-3.44	0.58	-5.94	<0.001***
Month:July	-1.77	0.62	-2.84	<0.01**
Month:August	-4.9	0.49	-10.0	<0.001***
Month:September	-2.10	0.53	-3.98	<0.001***
Body length	0.0019	0.0034	0.55	0.58
T_s	-0.06	0.032	-1.87	0.062
Substrate:Rock	-3.29	0.94	-3.5	<0.001***
Substrate:Soil	0.28	0.31	0.93	0.36
Solar rad.	0.0034	0.00075	4.62	<0.001***
Wind Speed	0.061	0.083	0.73	0.46
Species: <i>P. parva</i> June	1.32	0.73	1.8	0.073
Species: <i>P. parva</i> July	2.83	0.79	3.57	<0.001***
Species: <i>P. parva</i> August	2.0	0.61	3.3	<0.001***
Species: <i>P. parva</i> September	-0.91	0.69	-1.31	0.19
Species: <i>P. parva</i> Substrate: Rock	4.68	1.55	3.02	<0.01**
Species: <i>P. parva</i> Substrate: Soil	-0.17	0.43	-0.39	0.70
Body length:Wind speed	-0.00082	0.00036	-2.27	<0.023*
*: 0.01 < p < 0.05 **: 0.001 < p < 0.01 ***: p < 0.001				