Original article

A Mendelian randomization study of glycemic and anthropometric traits and Parkinson's disease

Running head: Glucose, body weight and Parkinson's disease

Sandeep Grover^{1,2} PhD, Ricarda Graf² BSc, Christine Klein^{3, 4} MD, Norbert Brüggemann^{4,5,6} MD, Inke R. König² PhD, Fabiola Del Greco M^{*7} PhD, Manu Sharma^{*1} PhD

¹Centre for Genetic Epidemiology, Institute for Clinical Epidemiology and Applied Biometry, University of Tübingen, Tübingen, Germany

²Institut für Medizinische Biometrie und Statistik, Universität zu Lübeck, Universitätsklinikum Schleswig-Holstein, Campus Lübeck, Germany

³Department of Psychiatry and Psychotherapy, University of Lübeck, Lübeck, Germany

⁴Institute of Neurogenetics, University of Lübeck, Lübeck, Germany

⁵Department of Neurology, University Medical Center Schleswig-Holstein, Campus Lübeck, Lübeck, Germany

Institute of Neurogenetics, University of Lübeck, Lübeck, Germany

⁶Center for Brain, Behavior, and Metabolism, University of Lübeck, Lübeck, Germany

⁷Institute for Biomedicine, Eurac Research, Bolzano, Italy

* FDG and MS should be considered joint senior authors

Corresponding Author:

Dr Manu Sharma

E-Mail: manu.sharma@uni-tuebingen.de

Phone: 0049-7071-2978259

Conflict of Interest

All authors declare no conflicts of interests.

Financial disclosure

Character count:

Title: 93, Running head: 44

Number of words:

Abstract: 250, Body of the Manuscript: 3664 (Excluding abstract and references)

Number of Figures: 2; Number of Tables: 3

Abstract

Background: Impaired glucose and obesity are frequently observed in patients with Parkinson's disease (PD), although it is unclear whether the impairment precedes or results from the neurodegeneration.

Objective: We aimed to assess whether glycemic and anthropometric traits can influence the risk of PD in 33,674 cases and 449,056 healthy controls using the Mendelian randomization (MR) framework.

Methods: We investigated causality with a two-sample MR approach in the European population to compute effect estimates with summary statistics from available discovery meta-analyses of genome-wide association studies (GWAS) on glycemic and anthropometric traits.

Results: We considered a threshold of $p\text{-value}=0.0038$ as significant after accounting for multiple testing, and $p\text{-value}<0.05$ was considered to be a suggestive evidence for a potential association. We observed a protective effect of waist-hip ratio (WHR) on PD (Inverse variance-weighted (IVW): $OR_{IVW}=0.735$; 95%CI= 0.622–0.868; $p\text{-value}=0.0003$; I^2 index=22.0%; MR-Egger intercept $p\text{-value}=0.1508$; Cochran Q test $p\text{-value}=0.0003$). The association was further retained after the exclusion of overlapping UK biobank (UKB) samples between the WHR and PD datasets ($OR_{IVW}=0.791$; 95%CI=0.659–0.950; $p\text{-value}=0.012$; I^2 index=13.0%; MR-Egger intercept $p\text{-value}=0.733$; Cochran Q test $p\text{-value}=0.035$). The sensitivity analysis provided suggestive evidence of an increased risk of PD on fasting glucose (FG) ($\beta_{IVW}=0.0188$; 95%CI=0.0062–0.0313, $p\text{-value}=0.0055$; I^2 index=0.0%; MR-Egger intercept $p\text{-value}=0.0957$; Cochran Q test $p\text{-value}=0.4555$) and protective effect of PD on T2D (Weighted median effect: $OR_{WME}=0.946$; 95%CI=0.9290–0.983; $p\text{-value}=0.0051$; Weighted mode effect: $OR_{MBE}=0.943$; 95%CI=0.904–0.983; $p\text{-value}=0.0116$).

Conclusions: Our results showed that central or abdominal obesity may be protective against PD development, independent of glucose levels.

Keywords: Mendelian randomization, Causal inference, Neurodegenerative disorders, Parkinson's disease, Glycemic traits, Type 2 diabetes, body weight, anthropometric traits

Introduction

The lack of neuroprotective or disease-modifying therapy has considerably hampered the management of Parkinson's disease (PD). However, several recent preclinical and clinical studies have shown the potential beneficial effects of type 2 diabetes (T2D) specific treatment in exerting neuroprotection against PD, possibly by modulating glucose homeostasis and body weight¹⁻⁵. Traditionally, insulin has been implicated in the general hormonal regulation of glucose metabolism, as insulin crosses the blood-brain barrier to modulate brain energy homeostasis, with a minor contribution from internal neuronal secretion⁶. A recent study also reported significantly higher blood glucose in non-diabetic PD patients compared to healthy controls during an oral glucose tolerance test, with no significant increase in insulin levels⁷. The study further reported an association of higher blood glucose levels with a higher BMI. Recently, type 2 diabetes (T2D) – characterized by high blood sugar, insulin resistance, and low insulin sensitivities – was shown to be associated with higher motor scores in patients with PD⁸. Change in body weight is also long known to occur during the clinical course of PD and with its treatment. A handful of observational studies with highly heterogeneous epidemiological study designs have investigated the association of body weight with PD, showing conflicting results although weight loss appears to be a consistent finding in more advanced PD⁹⁻¹². In summary, inconclusive evidence from observational studies suggest that several highly correlated glycemic and anthropometric traits could alter the risk associated with development of PD. This could be attributed to limited sample sizes, presence of inherent confounding and reporting bias in observational studies.

Genome-wide association (GWAS) or meta-analyses of GWAS often have larger sample sizes with adequate coverage of the human genome, making GWAS-based Mendelian randomization (MR) an

attractive approach. MR has recently evolved as an alternative statistical approach that can, against potential confounding, judge potentially causal relationships between risk factors (e.g. altered glucose metabolism or body mass index) and an outcome (e.g. PD)¹³. In principle, MR allows the use of genetic variants as proxy representatives of exposure from one population to test an association with an outcome in a completely independent population. The approach mimics the randomization of exposure in randomized controlled trials (RCTs) and, thereby, addresses hidden confounding factors. To date, MR studies exploring the causal role of altered glucose or insulin homeostasis in PD are lacking. However, a previously published study explored the role of body mass index (BMI) on PD and showed a protective role of body mass index (BMI) (OR = 0.82, 95% CI = 0.69-0.98)^{14, 15}. Most recently, the availability of GWAS datasets from the UK Biobank has further made it possible to take advantage of an increased power associated with a higher sample size by meta-analyzing it with previously existing large scale consortium datasets on various phenotypes of interest¹⁶⁻¹⁸.

In the present study, we expand the spectrum by assessing the impact and influence of several glycemic traits including 2-hour post-challenge glucose (2hrGlu), fasting glucose (FG), fasting insulin (FI), fasting insulin (FPI), homeostasis model assessment of β -cell function (HOMA-B), homeostasis model assessment of insulin resistance (HOMA-IR), glycated hemoglobin (HbA1c), Modified Stumvoll Insulin Sensitivity Index (ISI), and Type II diabetes (T2D) as well as anthropometric traits including body mass index (BMI), waist-hip ratio (WHR), waist circumference (WC), hip circumference (HC), adult height (AH) and birth weight (BW) on PD.

Methods

Study design and identification of datasets

We conducted a two-sample MR study using summary estimates to examine the lifelong effect of glycemic and anthropometric traits on the risk of PD in the European population. We reviewed the most recent meta-analyses of discovery GWAS datasets in the literature and identified genetic instruments that influence glycemic traits including 2hGlu, FG, FI, FPI, HOMA-B, HOMA-IR, HbA1c, ISI, T2D and anthropometric traits including BMI, WHR, WC, HC, AH, and BW^{16, 18-28} (**Table 1**). For the outcome

dataset, we used the discovery cohort of a recent meta-analysis of GWAS on 33,674 PD cases and 449,056 controls¹⁷.

Prioritization of genetic variants

We extracted significant SNPs from each GWAS dataset by employing a cutoff of 5×10^{-8} . All SNPs with F-statistics < 10 were further excluded for a possible violation of MR assumption I^{13,29}. A clumping window of 10,000 kb and linkage disequilibrium (LD; i.e. r^2) cutoff of 0.001 was applied in the European population in the 1000Genome Phase 3v5 dataset to identify the leading SNP that represents each significantly associated locus³⁰. If a specific leading SNP was not available in the PD dataset, a proxy SNP ($r^2 > 0.8$) was identified by using the European population in the 1000Genome Phase 3v5 dataset, when possible. The statistical power to detect a causal association was estimated by the method described by Brion et al.³¹. Specifically, we set the sample size of the outcome dataset to 482,750 with 7.498% as proportion of PD patients in the dataset, a continuous exposure with a variance $\geq 1\%$ and a threshold p-value of 3.8×10^{-3} (see the section below).

Effect estimation using MR and test of pleiotropy

We used the inverse variance-weighted (IVW) method with first order weights as primary method to investigate the direct causal role of glycemic traits and anthropometric traits on PD¹⁶⁻²⁸. If a genetic instrument consisted of a single SNP, we used the Wald ratio along with the delta method to estimate the causal effect and standard error, respectively. We applied a conservative Bonferroni correction to account for the number of 13 independent tests (local significance level = $0.05/\text{number of tests}$) for a global significance of 0.05. All other results from statistical tests are interpreted descriptively. We used the intercept deviation test with MR-Egger's and MR-Egger intercept and the I^2 index to evaluate the heterogeneity of Wald estimates. Additionally, we also estimated the Cochran Q statistic for the IVW method as well as Rucker's Q` statistics for the MR-Egger's method to evaluated the heterogeneity³²⁻³⁵.

Sensitivity analysis

We employed MR-Egger, weighted median (WME), and weighted mode methods (MBE) methods to check the reliability of estimates by relaxing some of the MR assumption, thereby allowing instruments with varying proportions of pleiotropic variants, as previously explained^{33, 36-39}. To avoid the overlapping of samples from UK Biobank, which has been included in recently published GWAS, we computed casual effect estimates by using PD datasets without UK biobank samples, as used in the previous study (9,581 PD cases and 33,245 controls)^{35, 40, 41}. For glycemic traits, a number of studies reported GWAS results from replication and/or pooled data (**Supplementary Table 1a**). We therefore compared our primary results with those using genetic instruments from these additional studies to explore the issue of the bias due to the winner's curse, led by the selection of the instruments done from the same dataset (discovery meta-analysis) used for the causal effect estimation.

Given that BMI has been shown to influence the role of glycemic and anthropometric traits on several diseases, including PD⁷, we also estimated the effect of genetic instruments adjusted for BMI for 2hrGlu, FG, FI, ISI, T2D, and WHR to identify their overall influence on the causal effect estimates for PD. A summary of GWAS datasets used to study the influence of GWAS study design and BMI adjusted datasets is provided in **Supplementary Tables 1a and 1b** respectively.

We employed a leave-one and leave-one-group-out cross-validation approach to check the influence of outlier variants as well as that of variants known to be associated with confounders of the relationship between glycemic, and anthropometric traits and PD. We used the PhenoScanner database to identify potential pleiotropic genetic instruments that are known to be associated with potential confounders⁴². Finally, we conducted a reverse directional MR by identifying genetic variants representing proxy markers of PD using the same study.

Results

Prioritization of genetic instruments and power analysis

The depth of genomic coverage and number of individuals in different discovery GWAS datasets is provided in **Table 1**. The table further shows the variance explained by genetic instruments for different

exposure datasets and availability of genetic instruments in the PD dataset, estimated by the formula $2 \times \beta^2 \times \text{EAF} \times (1 - \text{EAF})$, where β is the estimated genetic effect of the exposure and EAF is the corresponding allele frequency⁴³.

Two glyceic exposures (HOMA-IR and ISI) were not further analyzed given the absence of significant variants; however, we analyzed them as outcome with the PD as exposure in the reverse causation investigation. Thus, the number of primary statistical tests was reduced to the investigation of the causal effect of seven glyceic traits and six anthropometric traits on PD, leading to 13 tests. The significance level was accordingly set to $0.05/13 = 3.8 \times 10^{-3}$.

Our power analysis suggests that our study has $\approx 80\%$ power to detect a true OR of 1.208 or 0.799 for PD per SD of the continuous phenotype assuming that the proportion of the continuous phenotype explained by the genetic instrument is $\geq 1\%$ at a type 1 error rate of 3.8×10^{-3} .

Effect estimation and sensitivity analysis

The causal effect estimates of glyceic traits and anthropometric traits on PD are shown in **Table 2**, which also provides various measures to evaluate the robustness of the effect estimates. The summary data used to compute effect estimates and sensitivity analysis is further presented in **Supplementary Table 2**. Among all the glyceic and anthropometric traits, we found that a 1-standard deviation (SD) increase in waist-hip ratio (WHR) was associated with a 26.5% lower risk of PD in the European population ($\text{OR}_{\text{IVW}}=0.735$; 95%CI=0.622–0.868 per 1-SD of WHR; p-value=0.0003; $I^2=25.9\%$; MR-Egger intercept p-value=0.1508; Cochran Q test p-value=0.0003) (**Table 2a**). We further observed a similar effect using the WME method ($\text{OR}_{\text{WME}}=0.810$; 95%CI=0.721–0.911). The distribution of individual SNP-level effect estimates along with the effect estimates computed through different MR methods for the effect of WHR on PD are shown as scatter and funnel plots in **Figure 1**. After ruling out the effect of weak instrument bias on account of overlapping UKB samples, the protective effect of WHR on PD was retained suggesting the reliability of the observed findings ($\text{OR}_{\text{IVW}}=0.791$; 95% CI=0.659–0.950; p-value=0.012; $I^2=13.0\%$; MR-Egger intercept p-value=0.733; Cochran Q test p-value=0.035). Using the PhenoScanner database, out of 357 SNPs WHR associated SNPs employed in causal effect

analysis; we identified a total of 127 pleiotropic SNPs that have been previously shown to be associated with non-anthropometric traits such as blood cell count, glycemic traits, lipid levels, and respiratory capacity (data not shown). In our sensitivity analysis that excluded these pleiotropic SNPs our instrument was weaker thus explaining the diminished protective effect estimate ($OR_{IVW}=0.801$; 95% CI=0.640–1.000; p-value=0.052). The leave-one out sensitivity analysis also failed to show influence of any single SNP, suggesting reliability of our findings (data not shown here). We further observed a loss of association when using genetic instruments for WHR which were adjusted for BMI, suggesting a potential role of BMI in influencing the causal association with PD.

The observed findings of the potential causal role of anthropometric traits was further confirmed by the absence of the causal effect of any of the glycemic traits, including FG and T2D on PD. This lack of association of glycemic traits further persisted when we used genetic instruments that were prioritized from a small proportion of moderately associated SNPs which were followed up in a pooled cohort for 2hGlu, FG, FI, and FPI (**Supplementary Table 3a**). Similarly, no association was observed for HOMA-B and HOMA-IR where genetic instruments were available for the replication cohorts. In addition, we did not observe any influence of the BMI-adjusted instruments that were available for 2hrGlu, FG, FI, and T2D, regardless of the GWAS study cohort that was used to extract the instrument (**Supplementary Table 3b**). With respect to other glycemic traits, we did not find genetic instruments for ISI. However, we were able to explore the causality using the single genetic instrument for BMI adjusted ISI phenotype which hinted at an association using genetic instruments prioritized from the discovery cohort only ($OR_{wald} = 0.532$, 95% CI=0.286-0.990; p-value=0.0464).

Lastly, we checked reliability of the observed relationships between various glycemic and anthropometric traits, and PD by conducting MR analyses in reverse direction, as shown in **Table 3**. We did not observe any causal effect of PD on WHR ($\beta_{IVW}=-0.0077$; 95% CI=-0.0277–0.0122; p-value=0.4288; $I^2=84.1\%$; MR-Egger intercept p-value=0.8462; Cochran Q test p-value<0.0001). On the contrary, in our sensitivity analysis we observed the strongest effect on FG with a 1-log odds increase in genetic predisposition to PD being associated with 0.0188 mmol/l increase in FG concentration ($\beta_{IVW} =$

0.0188 per log-odds of PD; 95% CI=0.0062–0.0313; p-value=0.0055; $I^2=0.0\%$; MR-Egger intercept p-value=0.0957; Cochran Q test p-value=0.4555). Additionally, although IVW method failed to detect influence of PD on T2D ($OR_{IVW}=0.973$; 95% CI=0.921–1.028; p-value=0.3258; $I^2=76.9\%$; MR-Egger intercept p-value=0.4711; Cochran Q test p-value= 5.28×10^{-11}), the genetic predisposition to PD was associated with approximately 5.0% lower risk of T2D using other methods ($OR_{WME}=0.946$; 95% CI=0.930–0.973; p-value=0.0051; $OR_{MBE}=0.943$; 95% CI=0.940–0.983; p-value=0.0116). Our findings further motivated us to explore the triangulation relationship between the traits shown to be related to PD using the MR approach. Our data suggested a bidirectional causal relationship between T2D and FG as well as T2D and WHR (**Figure 2**) (data not shown). We further observed WHR as a potential risk factor for a higher FG with the absence of any effect of FG on WHR (**Figure 2**).

Discussion

The present study using a two-sample MR approach aimed to understand the role of glycemic and anthropometric on PD and observed that an increase in WHR is a strong protective factor for PD. Furthermore, sensitivity analyses provided suggestive evidence, though not conclusive, of higher glucose tolerance and protection against T2D in PD patients.

Dopamine neurotransmission in the human brain is known to modulate the rewarding properties of food. Previous studies have further shown that dopamine receptors are under expressed in obese individuals, thereby initiating a feedback loop to compensate for lower dopamine secretion⁴⁴. It is, however, not known whether an altered dopaminergic metabolism in overweight individuals could influence the PD susceptibility. Several indicators have been used to measure overweight and obesity. While WHR and WC are predominantly used as measurements of central obesity, BMI is used as measurement of overall obesity. It has also been shown that WHR and WC may be regarded as better alternatives to BMI to measure obesity, especially in individuals with cardiovascular risk factors including T2D^{45, 46}. Numerous observational studies have previously explored the association between both measures of obesity and PD with mixed results^{9, 10}. A recent meta-analysis of ten cohort studies with 2706 PD cases showed an absence of association of BMI with PD¹⁰. In contrast, a recent nationwide

health check-up data for the whole South Korean population comprising 44,205 incident cases identified
risky association of abdominal obesity with PD (HR=1.13, 95% CI=1.10–1.16)⁹.
Using an MR approach, we observed a significant risk reduction of 26.5% with every one unit of SD
increase in WHR, while not observing any role of BMI. Our results are in contrast with a previously
reported protective causal association of BMI with PD, which observed a significant risk reduction of
18%¹⁴. The previous study however assessed 77 loci for exploring causal effect of BMI comprising
13,708 PD cases and 95,282 controls compared to 548 loci assessed in a pooled dataset of 33,764 PD
cases 449,056 controls in the present study. The discrepancy in the number of prioritized loci between the
two studies is attributed to the use of GIANT dataset on BMI in the previous study compared to pooled
dataset of GIANT and UKB on BMI used in the present study. Another recent MR study prioritizing
genetic instruments using the UKB dataset only and exploring the casual role of 401 exposures did not
detect a casual role of BMI and WHR¹⁵. Nevertheless, they observed a consistent protective causal
association of nine adiposity related traits with the strongest effect observed for arm fat percentage. In
line with our findings, these studies collectively argue that the assessment of fat mass vs. fat-free mass
e.g. by using body plethysmography reflects the underlying causative or protective factors more closely
than body weight or BMI. Interestingly, a recent study reported a slight reduction of 1.12% in PD risk for
every 1kg/m² increase in BMI without any clear evidence of heterogeneity⁴⁷. The study employed
23andMe dataset for both BMI and PD datasets and the absence of heterogeneity in the observed findings
possibly suggest that a considerable heterogeneity observed in previous MR studies could be attributed to
the clinical heterogeneity among the commonly employed IPDGC PD cases. Our study provided strong
evidence regarding the role of WHR in PD, nonetheless, further studies are highly warranted to
disentangle the protective role of WHR in PD.

The results observed in our present study provided further suggest a potential role of glucose
metabolism in PD, and data obtained herein agree with the previously published epidemiological studies.
For example, a recent study reported a significantly higher area under time curve (AUC) for the blood
glucose levels in 50 non-diabetic PD patients compared to 50 healthy controls during a 75g oral glucose

tolerance test (1187 ± 229 vs 1101 ± 201 mmol/min.; $p=0.05$), with no significant difference in AUC for blood insulin levels (6681 ± 3495 vs 7271 ± 6070 mmol/min.; $p=0.57$)⁷. The study also reported that higher blood glucose levels were associated with higher BMI ($p\text{-value}<0.0001$). Another recent longitudinal study identified high blood glucose as a risk marker for PD progression⁴⁸. The 48-month follow-up study exploring the role of 44 clinical variables in 135 patients with early PD, identified high FG levels ($p\text{-value}=0.013$) and T2D ($p\text{-value}=0.033$), among several other factors as significant predictors of annual cognitive decline in PD. The study further observed significant differences in the baseline levels of glucose when compared to 109 healthy controls. Our results are henceforth in consent with these results suggesting that PD promotes dysregulation of glucose metabolism.

The relationship between PD and T2D as observed in our study is intriguing. We observed association only in sensitivity analyses. Thus, the results should be interpreted with caution. Nevertheless, several longitudinal studies have previously explored the influence of pre-existing T2D on the predisposition to PD with contradictory results. A prospective follow-up of 147,096 predominantly Caucasian participants in the Cancer Prevention Study II Nutrition Cohort from the United States found no association of the history of diabetes with PD risk ($RR=0.88$; 95% $CI=0.62\text{--}1.25$)⁴⁹. Another study that comprised two large US cohorts – the Nurses’ Health Study (121,046 women) and the Health Professionals Follow-up Study (50,833 men) – observed similar results ($RR=1.04$; 95% $CI=0.74\text{--}1.46$)⁵⁰. In contrast, a follow-up study in 51,552 Finnish individuals demonstrated an increased incidence of PD among patients with T2D ($HR=1.85$; 95% $CI=1.23\text{--}2.80$)⁵¹. Most recently, a meta-analysis of four cohort studies (3284 PD cases and 32,695 diabetes cases) confirmed the finding that the onset of diabetes was a risk factor for PD ($RR=1.37$; 95% $CI=1.21\text{--}1.55$)⁵². However, the same study reported the absence of an association in pooled populations of five case-control studies (6487 PD cases and 1387 diabetes cases; $OR=0.75$; 95% $CI=0.50\text{--}1.11$). In summary, findings of the association between T2D and PD have been highly heterogeneous and could be attributed to the varying age of onset of T2D and PD. Indeed previously published meta-analysis showed that the earlier onset of T2D before the onset of PD found to be a major risk factor for future PD patients⁵³. Our close examination of GWAS datasets used in our study

showed a remarkable variability in age of onset in PD and T2D subjects. For example, the majority of PD patients had an age of onset ranging from 48.9 to 71.2 years as compared to the average 52.5 years as an age of onset in T2D patients. Thus, it is conceivable that the earlier prodromal phase of PD, as compared to diabetes onset, could be protective against T2D in PD patients. Nevertheless, an in-depth clinical evaluation of PD subjects is highly warranted to further discern the role of T2D in PD etiology.

Despite our inability to stratify patients by the age of onset, our study has several strengths. We adopted a comprehensive approach that included several known markers of insulin metabolism. However, we observed that the genetic instruments for FI, HOMA-B, and HOMA-IR explained a very low amount of variance and, therefore, potential causation with PD might not be completely ruled out. An important limitation of this study could be the unavailability of individual-level data, which could have enabled us to confirm the absence of pleiotropic variants by using various potential confounding variables between WHR and PD. For instance, it is known that different markers of obesity show gender specific cut-offs and possibility of specific gender in influencing the observed causal estimates cannot be ruled out^{54, 55}. Although we observed a suggestive reverse causal association of T2D with PD using different MR methods, we did not observe similar results with HbA1C, which is a known biomarker for prediabetes or diabetes. One of the reasons for this could be that the GWAS on HbA1c with 123,491 individuals from the general population was underpowered when compared to the GWAS on T2D that included 898,129 individuals^{20, 21}. Another potential limitation could be the existence of overlapping UKB samples between WHR and PD datasets, which could have led to possible bias in our findings. Nevertheless, our sensitivity analysis demonstrates retention of the association even after the exclusion of UKB samples from the PD dataset, highlighting the robustness of our novel finding⁵⁶. Lastly, we could not conduct a causal association analysis among different glycemic traits within PD patients.

Despite these limitations, our study represents one of the most comprehensive studies to date that has explored the potential causal role of glycemic and anthropometric traits on PD. Our analyses suggest that central obesity may play a role in conferring protection against PD. An extensive sensitivity analysis further suggested a possible role of PD in altered glucose metabolism independent of insulin activity.

Furthermore, we showed that, despite high fasting glucose levels, PD patients may be protected against T2D. We further suggest the adoption of a cautionary approach when drawing clinical interpretations from the results of the current study, because additional lines of evidence may be generated, including the potential complex relationship of anthropometric, glycemic and PD with other unexplored traits.

Author contributions

1. Research project: A. Conception, B. Organization, C. Execution; 2. Statistical Analysis: A. Design, B. Execution, C. Review and Critique; 3. Manuscript Preparation: A. Writing of the first draft, B. Review and Critique;

S.G.: 1A, 1B, 1C, 2A, 2B, 3A;

R.G.: 2A, 2B;

F.D.G.: 2C;

N.B.: 3B

C.K.: 3B

I.R.K.: 2C, 3B;

M.S.: 1A, 1B, 2C, 3B

Acknowledgments

This study is, in-part, supported by the EU Joint Programme - Neurodegenerative Diseases Research (JPND) project under the aegis of JPND (www.jpnd.eu) through Germany, BMBF, funding code 01ED1406. Dr. Sharma is further funded by the Michael J Fox Foundation, USA Genetic Diversity in PD Program: GAP-India Grant ID: 17473. and also supported by the grants from the German Research Council (DFG/SH 599/6-1 to M.S.), MSA Coalition, and Michael J Fox Foundation. This study was also supported by grants from the German Research Foundation (Research Unit ProtectMove, FOR 2488) to I.R.K., F.D.G., N.B., C.K).

Data on PD GWAS was contributed by IPDGC team and downloaded from <https://pdgenetics.org/>. Data on glycemic traits were contributed by MAGIC investigators and downloaded from

www.magicinvestigators.org. Data on T2D were contributed by DIAGRAM investigators and downloaded from www.diagram-consortium.org. Data on BMI, WHR, WC, HC and height were provided by GIANT consortium and downloaded from <https://portals.broadinstitute.org/collaboration/giant/>. Data on birth weight was provided by EEG consortium and downloaded from <https://egg-consortium.org/>.

References

1. Kim DS, Choi HI, Wang Y, Luo Y, Hoffer BJ, Greig NH. A New Treatment Strategy for Parkinson's Disease through the Gut-Brain Axis: The Glucagon-Like Peptide-1 Receptor Pathway. *Cell Transplant* 2017;26(9):1560-1571.
2. Katout M, Zhu H, Rutsky J, et al. Effect of GLP-1 mimetics on blood pressure and relationship to weight loss and glycemia lowering: results of a systematic meta-analysis and meta-regression. *Am J Hypertens* 2014;27(1):130-139.
3. Athauda D, MacLagan K, Skene SS, et al. Exenatide once weekly versus placebo in Parkinson's disease: a randomised, double-blind, placebo-controlled trial. *Lancet* 2017;390(10103):1664-1675.
4. Athauda D, Gulyani S, Karnati HK, et al. Utility of Neuronal-Derived Exosomes to Examine Molecular Mechanisms That Affect Motor Function in Patients With Parkinson Disease: A Secondary Analysis of the Exenatide-PD Trial. *JAMA Neurol* 2019;76(4):420-429.
5. Li Y, Perry T, Kindy MS, et al. GLP-1 receptor stimulation preserves primary cortical and dopaminergic neurons in cellular and rodent models of stroke and Parkinsonism. *Proc Natl Acad Sci U S A* 2009;106(4):1285-1290.
6. Kuwabara T, Kagalwala MN, Onuma Y, et al. Insulin biosynthesis in neuronal progenitors derived from adult hippocampus and the olfactory bulb. *EMBO Mol Med* 2011;3(12):742-754.
7. Marques A, Dutheil F, Durand E, et al. Glucose dysregulation in Parkinson's disease: Too much glucose or not enough insulin? *Parkinsonism Relat Disord* 2018;55:122-127.
8. Pagano G, Polychronis S, Wilson H, et al. Diabetes mellitus and Parkinson disease. *Neurology* 2018;90(19):e1654-e1662.
9. Nam GE, Kim SM, Han K, et al. Metabolic syndrome and risk of Parkinson disease: A nationwide cohort study. *PLoS Med* 2018;15(8):e1002640.
10. Wang YL, Wang YT, Li JF, Zhang YZ, Yin HL, Han B. Body Mass Index and Risk of Parkinson's Disease: A Dose-Response Meta-Analysis of Prospective Studies. *PLoS One* 2015;10(6):e0131778.

11. Wills AM, Perez A, Wang J, et al. Association Between Change in Body Mass Index, Unified Parkinson's Disease Rating Scale Scores, and Survival Among Persons With Parkinson Disease: Secondary Analysis of Longitudinal Data From NINDS Exploratory Trials in Parkinson Disease Long-term Study 1. *JAMA Neurol* 2016;73(3):321-328.
12. van der Marck MA, Dicke HC, Uc EY, et al. Body mass index in Parkinson's disease: a meta-analysis. *Parkinsonism Relat Disord* 2012;18(3):263-267.
13. Grover S, Del Greco MF, Stein CM, Ziegler A. Mendelian Randomization. *Methods Mol Biol* 2017;1666:581-628.
14. Noyce AJ, Kia DA, Hemani G, et al. Estimating the causal influence of body mass index on risk of Parkinson disease: A Mendelian randomisation study. *PLoS Med* 2017;14(6):e1002314.
15. Noyce AJ, Bandres-Ciga S, Kim J, et al. The Parkinson's Disease Mendelian Randomization Research Portal. *Mov Disord* 2019;34(12):1864-1872.
16. Pulit SL, Stoneman C, Morris AP, et al. Meta-analysis of genome-wide association studies for body fat distribution in 694 649 individuals of European ancestry. *Hum Mol Genet* 2019;28(1):166-174.
17. Nalls MA, Blauwendraat C, Vallerga CL, et al. Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol* 2019;18(12):1091-1102.
18. Yengo L, Sidorenko J, Kemper KE, et al. Meta-analysis of genome-wide association studies for height and body mass index in approximately 700000 individuals of European ancestry. *Hum Mol Genet* 2018;27(20):3641-3649.
19. Scott RA, Lagou V, Welch RP, et al. Large-scale association analyses identify new loci influencing glycemic traits and provide insight into the underlying biological pathways. *Nat Genet* 2012;44(9):991-1005.
20. Wheeler E, Leong A, Liu CT, et al. Impact of common genetic determinants of Hemoglobin A1c on type 2 diabetes risk and diagnosis in ancestrally diverse populations: A transethnic genome-wide meta-analysis. *PLoS Med* 2017;14(9):e1002383.
21. Mahajan A, Taliun D, Thurner M, et al. Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. *Nat Genet* 2018;50(11):1505-1513.
22. Manning AK, Hivert MF, Scott RA, et al. A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. *Nat Genet* 2012;44(6):659-669.
23. Dupuis J, Langenberg C, Prokopenko I, et al. New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet* 2010;42(2):105-116.

24. Strawbridge RJ, Dupuis J, Prokopenko I, et al. Genome-wide association identifies nine common variants associated with fasting proinsulin levels and provides new insights into the pathophysiology of type 2 diabetes. *Diabetes* 2011;60(10):2624-2634.
25. Saxena R, Hivert MF, Langenberg C, et al. Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. *Nat Genet* 2010;42(2):142-148.
26. Walford GA, Gustafsson S, Rybin D, et al. Genome-Wide Association Study of the Modified Stumvoll Insulin Sensitivity Index Identifies BCL2 and FAM19A2 as Novel Insulin Sensitivity Loci. *Diabetes* 2016;65(10):3200-3211.
27. Shungin D, Winkler TW, Croteau-Chonka DC, et al. New genetic loci link adipose and insulin biology to body fat distribution. *Nature* 2015;518(7538):187-196.
28. Horikoshi M, Beaumont RN, Day FR, et al. Genome-wide associations for birth weight and correlations with adult disease. *Nature* 2016;538(7624):248-252.
29. Pichler I, Del Greco MF, Gogele M, et al. Serum iron levels and the risk of Parkinson disease: a Mendelian randomization study. *PLoS Med* 2013;10(6):e1001462.
30. Consortium GP, Auton A, Brooks LD, et al. A global reference for human genetic variation. *Nature* 2015;526(7571):68-74.
31. Brion MJ, Shakhbazov K, Visscher PM. Calculating statistical power in Mendelian randomization studies. *Int J Epidemiol* 2013;42(5):1497-1501.
32. Greco MF, Minelli C, Sheehan NA, Thompson JR. Detecting pleiotropy in Mendelian randomisation studies with summary data and a continuous outcome. *Stat Med* 2015;34(21):2926-2940.
33. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 2015;44(2):512-525.
34. Rucker G, Schwarzer G, Carpenter JR, Binder H, Schumacher M. Treatment-effect estimates adjusted for small-study effects via a limit meta-analysis. *Biostatistics* 2011;12(1):122-142.
35. Grover S, Lill CM, Kasten M, Klein C, Del Greco MF, König IR. Risky behaviors and Parkinson disease: A mendelian randomization study. *Neurology* 2019;93(15):e1412-e1424.
36. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol* 2016;40(4):304-314.
37. Burgess S BJ. Integrating summarized data from multiple genetic variants in Mendelian randomization: bias and coverage properties of inverse-variance weighted methods. *arXiv:151204486* 2015.
38. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data Mendelian randomization via the zero modal pleiotropy assumption. *Int J Epidemiol* 2017;46(6):1985-1998.

39. Grover S, Del Greco M F, Kasten M, Klein C, Lill CM, Koenig IR. Risky behaviors and Parkinson's disease: A Mendelian randomization study in up to 1 million study participants. *bioRxiv* 2018.
40. Nalls MA, Pankratz N, Lill CM, et al. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat Genet* 2014;46(9):989-993.
41. Lill CM, Roehr JT, McQueen MB, et al. Comprehensive research synopsis and systematic meta-analyses in Parkinson's disease genetics: The PDGene database. *PLoS Genet* 2012;8(3):e1002548.
42. Staley JR, Blackshaw J, Kamat MA, et al. PhenoScanner: a database of human genotype-phenotype associations. *Bioinformatics* 2016;32(20):3207-3209.
43. Falconer DS, Mackay TFC. *Introduction to Quantitative Genetics*. 4 ed. Harlow, Essex, UK: Longmans Green, 1996.
44. Wang GJ, Volkow ND, Logan J, et al. Brain dopamine and obesity. *Lancet* 2001;357(9253):354-357.
45. Dalton M, Cameron AJ, Zimmet PZ, et al. Waist circumference, waist-hip ratio and body mass index and their correlation with cardiovascular disease risk factors in Australian adults. *J Intern Med* 2003;254(6):555-563.
46. Janssen I, Katzmarzyk PT, Ross R. Body mass index, waist circumference, and health risk: evidence in support of current National Institutes of Health guidelines. *Arch Intern Med* 2002;162(18):2074-2079.
47. Heilbron K, Jensen MP, Bandres-Ciga S, et al. Unhealthy Behaviours and Parkinsons Disease: A Mendelian Randomisation Study. *medRxiv* 2020:2020.2003.2025.20039230.
48. Mollenhauer B, Zimmermann J, Sixel-Doring F, et al. Baseline predictors for progression 4 years after Parkinson's disease diagnosis in the De Novo Parkinson Cohort (DeNoPa). *Mov Disord* 2019;34(1):67-77.
49. Palacios N, Gao X, McCullough ML, et al. Obesity, diabetes, and risk of Parkinson's disease. *Mov Disord* 2011;26(12):2253-2259.
50. Simon KC, Chen H, Schwarzschild M, Ascherio A. Hypertension, hypercholesterolemia, diabetes, and risk of Parkinson disease. *Neurology* 2007;69(17):1688-1695.
51. Hu G, Jousilahti P, Bidel S, Antikainen R, Tuomilehto J. Type 2 diabetes and the risk of Parkinson's disease. *Diabetes Care* 2007;30(4):842-847.
52. Cereda E, Barichella M, Pedrolli C, et al. Diabetes and risk of Parkinson's disease: a systematic review and meta-analysis. *Diabetes Care* 2011;34(12):2614-2623.
53. Xu Q, Park Y, Huang X, et al. Diabetes and risk of Parkinson's disease. *Diabetes Care* 2011;34(4):910-915.

54. Schorr M, Dichtel LE, Gerweck AV, et al. Sex differences in body composition and association with cardiometabolic risk. *Biol Sex Differ* 2018;9(1):28.
55. Macek P, Biskup M, Terek-Derszniak M, et al. Optimal cut-off values for anthropometric measures of obesity in screening for cardiometabolic disorders in adults. *Sci Rep* 2020;10(1):11253.
56. Burgess S, Davies NM, Thompson SG. Bias due to participant overlap in two-sample Mendelian randomization. *Genetic epidemiology* 2016;40(7):597-608.

List of figures and tables

Figure 1. Graphical representation of causal association analysis and assessment of pleiotropy

A. Scatterplot showing causal effect estimates computed using various MR methods for the association of Parkinson disease (PD) as exposure with Waist Hip Ratio (WHR) as an outcome.

B. Funnel plot showing the extent of heterogeneity among the individual Wald ratio estimates for the association of Parkinson disease (PD) as exposure with Waist Hip Ratio (WHR) as an outcome.

Figure 2. Graphical representation of causal relationship between Parkinson's disease, glycemic and anthropometric traits. *Effect estimate computed using weighted median effect method (WME).

Table 1. Details of the discovery GWAS datasets that explored and prioritized genetic instruments used for direct and reverse casual analysis in the present study. The direct analysis was done using Parkinson's disease (PD) as an outcome, and the reverse was done using glycemic traits and modifiable anthropometric traits as an outcome.

Table 2. Causal effect estimates using different Mendelian randomization (MR) methods and heterogeneity analysis of causal effect estimates for Parkinson's disease (PD) by using various (a) glycemic traits as exposures. (b) anthropometric traits as exposures.

Table 3. Causal effect estimates using different Mendelian randomization methods and heterogeneity analysis of causal effect estimates for various (a) glycemic traits and (b) modifiable anthropometric traits by using Parkinson's disease as an exposure.

Supplementary Table 1a. Details of follow-up genetic variants in the replication and pooled exposure GWAS datasets and prioritized genetic instruments used for the secondary analysis

Supplementary Table 1b. Details of exposure GWAS datasets adjusted for BMI and prioritized genetic instruments used for the sensitivity analysis

Supplementary Table 2. Harmonized summary effect estimates from exposure and outcome datasets used for the conduct of Mendelian randomization (MR)

Supplementary Table 3. Sensitivity analysis exploring the (a) influence of GWAS study design (b) BMI adjusted traits on causal effect estimates

Financial disclosures

Author	Sandeep Grover
Stock Ownership in medically-related fields	None
Consultancies	None
Advisory Boards	None
Partnerships	None
Honoraria	None
Grants	None
Intellectual Property Rights	None
Expert Testimony	None
Employment	Centre for Genetic Epidemiology, Institute for Clinical Epidemiology and Applied Biometry, University of Tübingen, Tübingen, Germany

Contracts	None
Royalties	None
Other	None

508

Author	Ricarda Graf
Stock Ownership in medically-related fields	None
Consultancies	None
Advisory Boards	None
Partnerships	None
Honoraria	None
Grants	None
Intellectual Property Rights	None
Expert Testimony	None
Employment	Institut für medizinische Statistik, Universität Göttingen, Göttingen, Germany
Contracts	None
Royalties	None
Other	None

509

Author	Christine Klein
Stock Ownership in medically-related fields	None
Consultancies	Medical Consultant for Centogene and Retromer Therapeutics
Advisory Boards	Else Kroener Fresenius Foundation
Partnerships	None

Honoraria	Speaking Honorarium (Desitin)
Grants	DFG, EU, BMBF, MJFF, Centogene
Intellectual Property Rights	None
Expert Testimony	None
Employment	University of Luebeck and University Hospital of Schleswig Holstein
Contracts	None
Royalties	Oxford University Press
Other	None

510

Author	Norbert Brüggemann
Stock Ownership in medically-related fields	None
Consultancies	Centogene, Bridgebio
Advisory Boards	Bridgebio
Partnerships	None
Honoraria	None
Grants	DFG (BR4328.2-2, GRK1957)
Intellectual Property Rights	None
Expert Testimony	None
Employment	University Medical Center Schleswig-Holstein Lübeck, Department of Neurology; University of Lübeck
Contracts	None
Royalties	None
Other	None

511

Author	Inke R. König
Stock Ownership in medically-related fields	None
Consultancies	None
Advisory Boards	None
Partnerships	None
Honoraria	None
Grants	German Research Foundation, BMBF, German Cancer Aid
Intellectual Property Rights	None
Expert Testimony	None
Employment	University of Lübeck, Germany
Contracts	None
Royalties	None
Other	None

512

Author	Fabiola Del Greco M
Stock Ownership in medically-related fields	None
Consultancies	None
Advisory Boards	None
Partnerships	None
Honoraria	None
Grants	None
Intellectual Property Rights	None
Expert Testimony	None
Employment	Eurac research, Bolzano, Italy

Contracts	None
Royalties	None
Other	None

513

Author	Manu Sharma
Stock Ownership in medically-related fields	None
Consultancies	None
Advisory Boards	None
Partnerships	None
Honoraria	None
Grants	Michael J Fox Foundation, DFG, MSA Coalition
Intellectual Property Rights	None
Expert Testimony	None
Employment	Centre for Genetic Epidemiology, Institute for Clinical Epidemiology and Applied Biometry, University of Tübingen, Tübingen, Germany
Contracts	None
Royalties	None
Other	None

514

A. Scatterplot showing causal effect estimates

B. Funnel plot showing individual SNP level causal effect estimates

Type 2 Diabetes (T2D)

Parkinsons Disease (PD)

Fasting Glucose (FG)

Waist Hip Ratio (WHR)

OR = 4.19 (1.84-9.56); P = 0.0016

$\beta = 0.11$ (0.09-0.13); P = <0.0001

OR = 1.18 (0.86-1.62); P = 0.2910

$\beta = 0.02$ (0.006-0.03); P = 0.0055

OR = 0.95 (0.93-0.96); P = 0.0051*

OR = 1.03 (0.97-1.09); P = 0.2780

OR = 2.96 (2.60-3.36); P < 0.001

$\beta = 0.07$ (0.05-0.09); P < 0.0001

OR = 0.73 (0.62-0.87); P = 0.0003

$\beta = -0.008$ (-0.03-0.01); P = 0.4288

OR = 0.98 (0.94-1.03); P = 0.5572

$\beta = 0.09$ (0.06-0.12); P = < 0.0001