The island rule explains consistent patterns of body size evolution in terrestrial vertebrates

Ana Benítez-López^{1,2*}, Luca Santini^{1,3}, Juan Gallego-Zamorano¹, Borja Milá⁴, Patrick Walkden⁵, Mark A.J. Huijbregts^{1,†}, Joseph A. Tobias^{5,†}

¹Department of Environmental Science, Institute for Wetland and Water Research, Radboud University, P.O. Box 9010, NL-6500 GL, Nijmegen, the Netherlands.

²Integrative Ecology Group, Estación Biológica de Doñana, CSIC, 41092, Sevilla, Spain

³National Research Council, Institute of Research on Terrestrial Ecosystems (CNR-IRET), Via Salaria km 29.300, 00015, Monterotondo (Rome), Italy

⁴National Museum of Natural Sciences, Spanish National Research Council (CSIC), Madrid 28006, Spain

⁵Department of Life Sciences, Imperial College London, Silwood Park, Buckhurst Road, Ascot, Berkshire SL5 7PY, United Kingdom

*Correspondence to: ana.benitez@ebd.csic.es; abenitez81@gmail.com

†These two authors contributed equally

Abstract

Island faunas can be characterized by gigantism in small animals and dwarfism in large animals, but the extent to which this so-called ‘island rule’ provides a general explanation for evolutionary trajectories on islands remains contentious. Here we use a phylogenetic meta-analysis to assess patterns and drivers of body size evolution across a global sample of paired island-mainland populations of terrestrial vertebrates. We show that ‘island rule’ effects are widespread in mammals, birds and reptiles, but less evident in amphibians, which mostly tend towards gigantism. We also found that the magnitude of insular dwarfism and gigantism is mediated by climate as well as island size and isolation, with more pronounced effects in smaller, more remote islands for mammals and reptiles. We conclude that the island rule is pervasive across vertebrates, but that the implications for body size evolution are nuanced and depend on an array of context-dependent ecological pressures and environmental conditions.

Introduction

From giant pigeons to dwarf elephants, islands have long been known to generate evolutionary oddities¹. Understanding the processes by which island lineages evolve remains a prominent theme in evolutionary biology, not least because they include many of the world's most bizarre and highly threatened organisms². The classic insular pattern of both small-animal gigantism and large-animal dwarfism in relation to mainland relatives has been described as a macro-evolutionary or biogeographical rule – the ‘island rule’³⁻⁵ (Fig. 1). However, previous research has cast doubt on the generality of this pattern⁶, suggesting that body size shifts are asymmetrical, with reduced size in some clades (e.g. carnivores, heteromyid rodents, and artiodactyls) or increased size in others (e.g. murid rodents)^{7,8}. Even in these cases, the underlying mechanisms driving patterns of insular gigantism and dwarfism remain unclear.

Several mechanisms have been proposed to explain the island rule, including reduced predation, relaxed competition and resource limitation in island environments⁹. In theory, each of these factors may be accentuated in smaller, more isolated islands, where lower levels of interspecific competition and predation could lead to ‘ecological release’, allowing small-bodied species to increase in body size^{5,9}. Conversely, among large-bodied species, limited resource availability could select for smaller body sizes with reduced energy requirements, leading to insular dwarfism. Climatic conditions may also influence body size evolution on islands since primary productivity and associated resource availability are strongly influenced by climate^{9,10}. Although previous studies of body size evolution on islands have tested the effects of these different mechanisms, many have focused on relatively restricted geographic and taxonomic scales and did not directly address the island rule in its broad sense across multiple species within a taxon¹⁰⁻¹³, with notable exceptions^{9,14-16}.

Most work on the island rule has been restricted to mammals (e.g.^{4,7,14,17}), although the hypothesis has also been tested in amphibians¹⁸, reptiles¹⁹⁻²¹, birds^{15,22}, fish²³, insects²⁴, molluscs²⁵, and plants²⁶. The highly inconsistent results of these studies (e.g.^{5,6,27}) are perhaps unsurprising because they typically deal with single species or pool together data on different traits from numerous sources without controlling for variation in study design or accounting for sampling variance. Accordingly, a recent systematic review based on a simplified scoring system²⁷ concluded that empirical support for the island rule is not only potentially biased but

also generally low, particularly for non-mammalian taxa. However, scoring approaches provide only limited information as they do not account for heterogeneity between studies, taxonomic representativeness, sample size, or precision in the estimates.

These limitations are best addressed with formal meta-analyses^{28,29}, hence we tested the island rule hypothesis by applying phylogenetic meta-regressions to a global dataset of 2,479 island-mainland comparisons for 1,166 insular and 886 mainland species of terrestrial vertebrates (Supplementary Dataset 1, Fig. 2). Our analytical framework allows us to control for multiple types of variation, including data source, sample size imbalance, intraspecific and intra-population variability, and phylogenetic relatedness (see Methods). For each island-mainland comparison, we calculated the log response ratio ($\ln RR$) as the natural logarithm of the ratio between the mean body size of individuals from an insular population M_i and that of mainland relatives M_m ($\ln RR = \log[M_i/M_m]$)³⁰. Then, we regressed $\ln RR$ against the body mass of the mainland population (M_m) (Fig. 1).

This framework provides a clear set of predictions in the context of evolutionary trajectories on islands^{4,6,14}. Specifically, since negative values of $\ln RR$ indicate dwarfism and positive values indicate gigantism, a positive intercept and negative slope of the $\ln RR$ -mainland mass relationship supports the island rule (Fig. 1). Given the contentiousness of the generality in the island rule, we assessed the robustness of our results against potential biases derived of regressing ratios^{31,32}, using small samples, imputing missing data, or the influence of using data from the island rule literature or derived from other studies focused on unrelated questions (i.e. publication bias; see Methods). Finally, we use our framework to assess how body size shifts are related to island size, island isolation, island productivity and climate, as well as species diet. The extent to which these different factors explain insular body size shifts allows us to re-evaluate a range of hypotheses for the mechanisms underlying “island rule” effects on body size, including ecological release⁹, immigrant selection⁹, resource limitation^{9,33,34}, thermoregulation^{9,15,35}, water availability^{36,37} and starvation resistance^{9,33} (Supplementary Table 1, Extended Data Fig. 1).

Results

The generality of the island rule

We found that $\ln RR$ and mainland body mass were negatively related for mammals, birds and reptiles, with small species tending to gigantism and large species to dwarfism (Fig. 3). The relationship was weakly negative but statistically non-significant for amphibians, with a tendency towards gigantism across all body sizes (Fig. 3, Table 1). We obtained similar results using size ratios corrected for small sample size ($\ln RR^d$), or by regressing island mass against mainland mass, with support for the island rule across all groups except for amphibians (Supplementary Table 3-4). This indicates that our analyses are robust to small sample size bias³⁸ or any potential spurious correlation associated to ratio regression models^{31,32} (Extended Data Fig. 2). Further, neither imputation nor publication bias influenced our results (Supplementary Table 5-6), with no apparent differences between island-mainland comparisons sampled from studies formally testing the island rule or compiled from unrelated data sets.

Mainland body mass explained 11.4, 7.0 and 17.6% of the variance in mammals, birds and reptiles, respectively. The amount of further variance accounted for by phylogeny (0.0–29.8%), data source (1.8–25.1%), and species (25.9–53.2%) fluctuated widely among taxa (Extended Data Fig. 3). Phylogeny accounted for a relatively large amount of variance in mammals (20.1%) and reptiles (29.8%), but even in these cases the overall patterns were not driven by large effects in particular clades. Some groups tended towards gigantism and others towards dwarfism, while others contained both dwarfs or giants depending on body size (e.g., Primata, Rodentia, and Carnivora in mammals, and Viperidae, Scincidae and Iguanidae in reptiles; Extended Data Fig. 4).

Ecological mechanisms underlying body size evolution on islands

The pattern of body size evolution in our island-mainland comparisons provides some insight into the likely mechanisms driving “island rule” effects (Extended Data Fig. 5-8, Supplementary Table 7, Supplementary Dataset 2). Overall, insular size shifts arise through some combination of ecological release from predation and competition, resource limitation, biased colonization (i.e. immigrant selection), and starvation resistance. The fact that no single factor explained island effects on body size is not surprising because some hypotheses shared overlapping predictions, making them difficult to disentangle.

Shifts in body mass of mammals were mostly explained by island size and spatial isolation ($Q_m = 12.20$, $P = 0.002$, Fig. 4a), resulting in more pronounced gigantism or dwarfism in small and remote islands. Birds showed similar size shifts in relation to spatial isolation and island area, but these were not statistically significant (Supplementary Table 7). In both mammals and birds, temperature had similar effects across the size range, with body size consistently larger in cool islands and smaller in warm islands (Extended Data Figs. 5e, 6e). Hence, in these groups, even large species that had undergone dwarfism were significantly larger in cool insular environments than in warm ones. Contrary to the starvation resistance hypothesis, small-sized birds did not become larger in highly seasonal islands, but large-sized birds had reduced dwarfism on islands with high seasonality in temperatures ($Q_m = 12.33$, $P < 0.001$, Extended Data Fig. 6).

In reptiles, the combination of island area and spatial isolation were the most important factors explaining variation in body size (Fig. 4c), with productivity and seasonality also supported but with weaker effects (Extended Data Fig. 7). Similar to mammals, the tendency towards dwarfism or gigantism in large-bodied or small-bodied reptiles was more apparent in isolated small-sized islands, with stronger effects of area than isolation (Supplementary Table 7). The effects of productivity and seasonality were only partially in line with predictions, as small-sized species were larger on islands with high seasonality, but smaller on islands with high productivity (Extended Data Fig. 7). In turn, large-bodied reptiles were smaller on islands with low productivity and high seasonality.

Finally, the relationship between size ratio and mainland mass in amphibians was slightly steeper in small and remote islands (Fig. 4d), with island area being marginally more important than spatial isolation (Extended Data Fig. 8). The effect of seasonality was clearer, with amphibian species inhabiting islands with high seasonality (unpredictable environments) tending toward gigantism, whereas those from islands with low seasonality (predictable environments) being similar in size to mainland counterparts (Extended Data Fig. 8). We found no effects of diet for any of the four taxa, or precipitation for amphibians, contrary to the water availability hypothesis.

Discussion

Based on comprehensive morphometric data from a worldwide sample of island fauna, we show consistent patterns of body size evolution across terrestrial vertebrates in accordance with predictions of the island rule. This finding was robust to alternative modelling approaches (island mass vs mainland mass regressions), small sample bias, data imputation, and publication bias. Moreover, we have demonstrated that insular size shifts are contextual and depend not only on the body size of mainland relatives (island rule *sensu stricto*) but also on the physiographic and climatic characteristics of particular island environments⁹.

Repeated evolutionary trajectories on islands

We found a clear negative relationship between insular body size variation and the body mass of mainland individuals in mammals, birds, and reptiles. Mainland body mass explains between 7.0 and 17.6% of the variation in insular size divergence in these three taxonomic groups, which is similar to that reported in smaller-scale studies of bats (15%), birds (13%), snakes (42%), non-volant and terrestrial mammals (11–21%), and turtles (8%)^{5,14,15,39,40}. Contrary to these earlier studies, our analyses are corrected not only for phylogenetic relatedness, but also for variability between species and intrapopulation variability, thereby strengthening the evidence for predictable evolutionary trajectories on islands. Nevertheless, the island rule provides only a partial explanation for these trajectories because substantial variation around the trend line remains unexplained. We also conducted the first multispecies test of island rule effects in amphibians, showing that the relationship goes in the expected direction but with a weak effect (1.4 %), possibly because the body mass range in amphibians is narrower and limited to small sizes (~ 0.5-50 g) and thus most amphibians tend to gigantism on islands with reduced predation risk.

Our findings are in contrast with a number of studies rejecting the island rule, including a recent review of evidence from across mammals, birds and reptiles²⁷, as well as other taxon-specific studies focused on lizards^{20,41} and turtles²¹. On the other hand, the patterns we detect are consistent with analyses supporting the island rule in snakes¹⁹, mammals^{4,9} and birds^{5,15}. We conclude that the contradictory results of previous studies may have been related to sampling bias, heterogeneity between sources and species, and phylogenetic relatedness (i.e. statistical non-independence). By accounting for these effects in our global models we are able to

demonstrate that vertebrate animals evolve in largely consistent ways on islands. Further, we have shown that the island rule is not clade-specific and instead applies to numerous clades within major taxonomic groups, particularly in mammals and birds.

A corollary that emerges from the island rule is that body size converges on islands. Specifically, if insular environments select for intermediate body sizes, closer to the optimal size of the focal clade, then the size spectrum of organisms found on islands should be narrower compared to the mainland^{42,43}. Theoretically, the optimal body size towards which small and large species may converge in low-diversity systems such as islands should correspond to the point where the trend intersects the horizontal dashed line in the relationship between size ratio and mainland mass, at which point fitness is maximized⁴² (but see⁴⁴). Interestingly, the shift between dwarfism and gigantism in our models occurred at approximately 100-250 g in endotherms, slightly larger than the 100 g adult body mass proposed for mammals⁴² (but see⁴³), and the mode of the global body size distribution of birds that separate between small- and large-bodied species (60 g)^{22,45,46}. Additionally, our analyses suggest that the optimal body size for island reptiles should be ca. 20 g, which is marginally higher than the modal body size of Lepidosaur (14.1 g)⁴⁷. Whether there is an optimal body size in island biotas has been the subject of much debate⁴⁴, but overall we expect that phenotypic variability in morphometric traits will be substantially narrowed if directional selection is operating in island assemblages, a feature that warrants further investigation. Additionally, optimal phenotypes should vary with the environmental characteristics of islands, in particular their area and isolation, climate, productivity and seasonality. For example, in mammals, our results suggest that the optimal body size would be ca. 100 g and ca. 900 g in warm and cold islands, respectively.

Ecological mechanisms influencing body size variation

Because body size is intimately linked to many physiological and ecological characteristics of vertebrates, it may be associated with a variety of environmental factors. As a consequence, the body size of colonizing species may predictably evolve as the result of selective pressures associated with insular environments (e.g., low food resources, few competitors, no predators) and others that act across larger geographic scales (e.g., climate). For mammals and reptiles, our results suggest that insular body size shifts are indeed governed by spatial isolation and island size, with individuals becoming dwarfs or giants in remote islands of limited size. Furthermore,

the slope of the relationship between size ratio and mainland mass was slightly steeper for birds and amphibians in small remote islands than in large islands near continental land masses (Fig. 4). This points to a combination of resource limitation (with small islands having fewer resources to maintain large-sized organisms^{48,49}) along with release from interspecific competition and predation pressure in small, species-poor islands. The pattern is also consistent with biased colonization favoring larger individuals with higher dispersal abilities (immigration selection⁵⁰). Conversely, our results showed that body size divergence on islands close to the mainland was minimal, reflecting two non-mutually exclusive processes. First, many of these islands were connected to the continent by land bridges so recently that phenotypic differences have not had time to accumulate. Second, regular dispersal between mainland and island populations promotes gene flow, with introgression counteracting divergent selection^{51,52}.

Besides island physiography (area and isolation), other relevant factors were temperature conditions in endotherms and resource availability and seasonality in ectothermic organisms. Mammals and birds both responded to island temperature in line with the heat conservation hypothesis, with small- and large-sized species exhibiting exacerbated gigantism and diminished dwarfism, presumably to conserve heat in colder, harsher insular environments. Additionally, temperature seasonality was an important determinant of the size of large-bodied birds, with populations on highly seasonal islands being similar in size to mainland populations. One possibility is that larger size in these cases may help maintain energy reserves during periods of low food availability, allowing them to thrive in otherwise hostile environments. Another possibility is that bird populations on highly seasonal islands – which tend to be situated at relatively high latitudes – are more often seasonally mobile or even migratory, potentially increasing gene flow with mainland populations or weakening adaptation to the local environment⁵³. These findings add new insights to previous results regarding the role of thermal and feeding ecology on morphological divergence in island birds^{54,55}. Traditionally, changes in feeding ecology were thought to be the prime force in driving morphological divergence in island birds^{54,55}. Yet, our results imply that physiological mechanisms related to heat conservation (‘thermoregulation hypothesis’) and energy constraints (‘starvation resistance hypothesis’) may also shape body size evolution in island birds.

In reptiles, we find some evidence that resource availability and seasonality are important factors explaining body size evolution, with some deviations from the patterns predicted. As hypothesized, large species are much smaller on islands with low resource availability, and small species are larger on islands with high seasonality. Yet, unexpectedly, small species are larger on islands with low productivity, perhaps because increased intraspecific competition favors large individuals under the high population densities that reptiles often attain on islands^{56,57}.

Overall, most amphibians tended to gigantism, presumably as a result of increased growth rate or lower mortality due to reduced predation pressure on islands⁵⁸. Additionally, we found that body size of amphibians consistently increased on islands where resources were highly seasonal and unpredictable, perhaps to maximize energy reserves and withstand long periods without food, for example during aestivation or hibernation⁵⁹. We did not find a clear relationship between precipitation and body size, suggesting that water availability is not a key factor. It appears that gigantism in island amphibians is mostly driven by physiological mechanisms that maximize growth rate, particularly in smaller, more isolated islands. These findings should be further explored when more data on island-mainland pairwise populations of amphibians become available.

Body size evolution in extinct species

Our analyses focused on extant species for which we could gather information on the variation around the morphometric estimates, along with sample size (essential for meta-analyses). The widespread extinction of large species on islands, including dwarf morphotypes of large species such as insular elephants in Sicily and the Aegean islands^{60,61}, may have masked the historical pattern of phenotypic variation on islands⁶². Giant insular birds^{54,63}, primates^{64,65}, and lizards⁶⁶, along with large insular turtle species, went extinct during the Holocene and late Pleistocene⁶⁷, most likely because of overhunting and the introduction of invasive species^{68,69}. Overall, it is estimated that human colonization of oceanic islands was followed by the extinction of 27% of insular endemic mammals⁷⁰, as well as over 2000 bird species in the Pacific region alone⁷¹, with these losses biased towards large-bodied, flightless, ground-nesting species⁶⁸. Extinct species may shed new light on size evolution in insular vertebrates because species extinctions have substantially altered the biogeography of body size in island faunas, potentially leading to downsized insular communities^{72,73}. For example, the predominance in our dataset of smaller-

bodied organisms could reflect the extinction of large species on islands⁶⁸, or simply the fact that few islands support large species. Either way, further studies should include data from extinct species as this may alter or strengthen the signal that we report for extant species³⁹.

We foresee that, under global change, the extinction of insular species and the introduction of novel (invasive) species may trigger new equilibria, with concomitant shifts in the composition of insular communities and the opening of novel niches to which species may respond via genetic adaptations and phenotypic plasticity. Recent evidence indicates that even introduced species on islands, which were not included in our analysis, predictably evolve towards dwarfism or gigantism⁷⁴⁻⁷⁶. In theory, as the Anthropocene gathers pace, further extinctions will drive a decline in mean body size of the overall island community, pushing optimal body sizes towards the lower end of body size ranges in the different vertebrate groups.

Conclusions

Of the many evolutionary implications of living on islands – together known as the ‘island syndrome’² – the effects on body size are the most widely known and controversial. We have shown that these ‘island rule’ effects are widespread in vertebrate animals, although the evidence for amphibians is inconclusive. Morphological changes were directional for species at the extremes of the body size range in mammals, birds and reptiles, following the predicted pattern of convergence towards intermediate “optimum” body sizes, in line with optimal body size theory^{42,43,45}. Although this convergence towards morphological optima may result from natural selection or phenotypic plasticity, the exact mechanism producing these changes on islands is still not well understood. Nonetheless, we found that consistent transitions towards intermediate body sizes were associated with a combination of factors, indicating a range of different ecological mechanisms. Our results highlight the contextual nature of insular size shifts, where island physiographic, climatic and ecological characteristics play a fundamental role in shaping body size evolution, reinforcing the idea that large-scale macroevolutionary patterns do not arise from single mechanisms but are often the result of multiple processes acting together^{77,78}.

Methods

Data collection

We collected baseline morphometric data from articles included in a recent assessment of the island rule²⁷, as well as other compilations assembled to test the hypothesis in reptiles²⁰, mammals⁶, and birds¹⁵. To expand this sample, we then performed a literature search (February 2020) in Web Of Science Core Collection (WOS) using the following search string: (“island rule” OR “island effect” OR “island syndrome” OR island*) AND (gigantism OR dwarfism OR “body size” OR weight OR SVL OR snout-vent length OR length OR size) AND (mammal* OR bird* OR avian OR amphibia* OR reptile*) (Appendix 1). Because this search was complementary to the data we have gathered from previous compilations^{6,15,20,27}, we only downloaded the first 500 hits out of a total of 33,431 hits ordered by relevance, and removed duplicates already included in our dataset. We reviewed every island-mainland comparison reported in published studies and traced primary source data when possible to extract original measurements. We also extracted data from all studies containing morphometric measurements for insular populations when these could be matched with equivalent data published elsewhere for relevant mainland taxa. We excluded problematic data, such as comparisons that were not supported by taxonomic or phylogenetic evidence, or which reported morphometric data restricted to single specimens or without sample size. In addition, we excluded comparisons based on extinct taxa since they are often known from very few or incomplete specimens (Supplementary Dataset 3).

It has been argued that research on the island rule might be prone to ascertainment bias, where researchers are more likely to notice and measure animals of extreme body size when conducting research on islands⁴¹. To help overcome this problem, we collected body size data not only from studies testing the island rule, or reporting dwarfism and gigantism in island fauna, but also from studies that did not specifically test hypotheses related to the island rule. We matched unpaired insular populations with independent data from mainland populations by performing species-specific searches in WOS and Google Scholar. We also compiled morphometric data for 442 insular and 407 mainland bird species from an independent global dataset of avian functional traits⁷⁹.

Large islands may be more ‘mainland like’ in relation to factors that are thought to affect body size (i.e. competition, resource availability and predation⁵). Thus, when major islands were at least 10 times larger than a nearby island, we treated the large island as the mainland comparison, following previous studies testing the island rule^{4,5,20}. Consequently, a single mid-sized island can simultaneously be treated as the continent in comparisons with smaller islands, and the island in comparisons with larger continents. When authors reported data referring to an entire archipelago instead of a specific island (3.2% of cases), we used the size of the largest island as island area. Removing these cases from our analyses did not qualitatively affect our results (Supplementary Table 8).

Our final dataset contained 529 data sources and 2,479 island-mainland comparisons^{7,10,36,58,79-602}. In total, we collated morphometric measurements for 63,561 insular and 154,875 mainland specimens representing mammals (1,058 island-mainland comparisons), birds (695 comparisons), reptiles (547 comparisons) and amphibians (179 comparisons) from across the globe (Fig. 2). 2,068 island-mainland comparisons (83.4%) were within species (e.g. subspecies) comparisons, and 411 (16.6%) were between-species comparisons. Insular populations were sampled from an array of islands varying widely in size (0.0009–785,753 km²), climate, and level of spatial isolation (0.03–3,835 km from mainland). To explore the drivers of body-size shifts in insular populations, we also sampled species with a wide range of average body masses (0.18–234,335 g). We collated data on body size indices (body mass, body length, cranial and dental measurements) of different taxa in island and mainland populations following strict morphological, phylogenetic and biogeographic criteria. Specifically, we always compared the same body size index for island and mainland populations. For within-species comparisons, we compared island and mainland populations based on the information given by the authors of the relevant study (e.g. taking note of which mainland source populations are likely to inhabit a particular island because of colonization history or isolation via rising sea levels^{89,101,240,386,548}). When we matched comparisons independently, we used information published in the study reporting the insular form, selecting the geographically closest mainland population whenever possible. In addition, we prioritized latitudinal alignment of mainland and island populations to avoid confounding effects of latitudinal variation in body size. In the case of island endemics, we compared island populations to their closest mainland relative whenever these were identifiable by phylogenetic data or other information reported in each particular study. This usually meant

selecting their sister species or the geographically closest representative of a sister clade or polytomy (Supplementary Dataset 1). If we could not reliably establish the closest mainland relative, we discarded the data (see Supplementary Dataset 3).

When more than one body size index was reported in published studies, we prioritized those indices related most closely to body mass (Supplementary Table 2). For mammals, we selected indices in this order of preference: body mass, body length, cranial length (greatest skull length or condylobasal length), and dentition (e.g. canine length)⁵. For birds, preferred indices were body mass, wing length, tarsus length and bill length. Finally, for amphibians and reptiles, size was reported as body mass, snout-vent-length (SVL), carapace length (CL, for turtles) and total length (TL, including SVL and tail length). In all cases, we included measurements for adults only. To avoid size biases attributable to sexual size dimorphism, we calculated the pooled mean for both sexes and the combined SD using standard formulae for combining groups⁶⁰³. When information was only available for one sex (male or female), we restricted our size comparisons to the sex for which we had morphometric data in both mainland and island populations. Data from zoos or studies that could not be georeferenced were discarded.

To overcome the problem that different authors report size using different indices, we used allometric relationships to convert island and mainland size to body mass equivalents, thereby enabling cross-taxa and cross-study comparisons. Although this conversion is imprecise, morphological indices and body mass are nonetheless highly correlated across the global scale and wide range of body sizes within our samples (providing more accurate predictions than simply assuming an exponent ~ 3 , as in previous studies testing the island rule^{5,9}). We used published allometric relationships where available (see Supplementary Table 2), or derived them based on published datasets^{47,179,604-609} and other data sources (Supplementary Dataset 4). To calculate allometric relationships, we used OLS (Ordinary Least Square) models of the $\log^{10}$ transformed body mass against the $\log^{10}$ transformed body size index (e.g. condylobasal length, Supplementary Table 2, Supplementary Dataset 4).

For birds, we complemented published data with standardized morphometric measurements from 3,618 museum specimens and live individuals of 436 insular and 404 mainland bird species (see⁷⁹). We used wing length in the main analyses instead of tarsus length because the former is a better predictor of body mass in our dataset ($R^2_{wing} = 0.89$ vs $R^2_{tarsus} = 0.69$, Supplementary Table

2) (see also⁶¹⁰). Although wing length may change during moult or thereafter because of wear, these effects are negligible in relation to interspecific differences⁷⁹ and minimized by calculating averages across multiple individuals. Further, interobserver differences between measurements may explain some variation in wing length estimates, but again this bias was shown to have negligible effects in our dataset by comparing repeated measures from different observers (see⁷⁹). To assess the consistency in our results, we repeated analyses using tarsus length, another popular proxy of overall body size in birds⁶¹¹. Our results were unchanged (Fig. S2).

To select suitable comparisons for museum specimens, we first classified species as either insular or continental by overlapping IUCN range polygons with a GIS land layer including continental land masses. For each insular species we then identified continental sister species from avian phylogenies⁶¹², using the method described above. We excluded bird species that are highly pelagic or aerial (e.g. swifts) and fully migratory species because in these groups it is unclear whether insular and mainland forms experience different environments¹⁵. Further, we also excluded flightless bird species, because morphological changes may be due to flightlessness rather than island dwelling per se¹⁵.

We calculated the response ratio (*lnRR*, eq. 1) as effect size in our meta-regressions, where we divided the mean body mass of individuals from an insular population $\bar{M}_i$ by that of the nearest mainland relative, $\bar{M}_m$, and then applied the natural logarithm. Unlike unlogged ratios, the sampling distribution of *lnRR* is normal, particularly for small samples³⁰, and thus less prone to statistical artefacts associated with ratio-based regressions.

$$\ln RR = \ln \left(\frac{\bar{M}_i}{\bar{M}_m} \right) \quad (\text{Eq. 1})$$

Response ratios greater than zero indicate a shift towards larger sizes (gigantism) whereas ratios less than zero indicate shifts towards smaller sizes (dwarfism). Besides mean measurements, we recorded measures of variation, i.e. standard deviation (SD), standard error (SE) or coefficient of variation (CV), and sample sizes of the body size indices in island and mainland organisms. SD and sample sizes were used to calculate sampling variances (Eq. 2), which were then used to weight each response ratio (coupled with the amount of heterogeneity, i.e. the variance in the underlying effects)³⁰.

$$\hat{\sigma}^2(\ln RR) = \frac{SD_i^2}{N_i \bar{x}_i^2} + \frac{SD_m^2}{N_m \bar{x}_m^2} ; \quad (\text{Eq. 2})$$

SDs were extracted from raw data when possible. If ranges were provided instead of SD (or SE or CV), we calculated SD following⁶¹³. If neither ranges nor measures of variation were reported, but the reported sample size was > 1, we imputed SD based on the coefficient of variation from all complete cases (“Bracken approach”⁶¹⁴). Imputation was done for 22% of all cases in mammals, 1.1% in birds, 11% in reptiles and 7.3% in amphibians, all within the upper limit of imputations (<30% of all cases per group) advised in previous studies⁹⁰.

For each study and island-mainland comparison, we compiled the mainland and island names, the study reference, the body size index used, the geographic coordinates, the distance to the closest mainland (spatial isolation, km) and the island area (km²). We completed missing data on island characteristics using the UNEP island database (<http://islands.unep.ch/>) and the Threatened Island Biodiversity Database (TIB, <http://tib.islandconservation.org/>). Missing information was calculated using Google Earth. Additionally, we extracted the Normalized Difference Vegetation Index (NDVI) as a proxy for resource availability on islands⁶¹⁵. We also calculated the standard deviation of NDVI to assess seasonality in leaf or vegetation cover, as an index of seasonality in available resources. NDVI was downloaded from NASA Ames Ecological Forecasting Lab (<https://lpdaacsvc.cr.usgs.gov/appears/task/area>).

Because climate influences both resource requirements and primary productivity, body size evolution should also be influenced by climatic conditions on islands. We thus extracted island climatic conditions from WorldClim v. 2.0 (<http://worldclim.org>⁶¹⁶). Specifically, we used variables that are more closely associated with the proposed underlying mechanisms of Bergmann’s rule (i.e. thermoregulation and starvation resistance): mean annual temperature, annual precipitation, and seasonality of temperature and precipitation⁶¹⁷. We assumed that the time period for these bioclimatic variables (1970–2000), although not necessarily matching the actual time period of body size evolution in the insular populations, roughly represents the climatic conditions in the Holocene, a period relatively climatically stable where most of our populations became isolated (i.e., after the last glacial maximum; see also⁹). Because climatic variability across cells substantially exceeds variation within cells in the Holocene, current layers are considered adequate for geographic comparisons. All spatial variables were downloaded at

0.1-degree resolution, and we averaged all cells per island to obtain a mean value of each environmental variable (e.g., temperature, NDVI, precipitation, etc.). Finally, for each species included in our dataset, we collated diet information from EltonTraits for birds and mammals⁶¹⁸, and from other sources for reptiles^{608,619}, and classified species as carnivores (> 50% diet consisting of vertebrates) or non-carnivores (< 50% diet consisting of vertebrates), following previous studies^{79,620}. As all amphibians in our dataset are carnivores⁶²¹, we did not record their diet.

Data analyses

To test the island rule hypothesis, we used phylogenetic meta-regressions between $\ln RR$ and body mass of mainland relatives, following most previous studies of the island rule (e.g.^{4,5,7,6,622,623}). A negative slope for this relationship would support the island rule (Fig. 1).

The use of multiple populations of the same species can overestimate the actual number of degrees of freedom, generating type-1 errors. We controlled for this by adding ‘Species’ as a random effect intercept in our analyses. Additionally, body size evolution in insular vertebrates is heavily influenced by phylogenetic effects, with species within entire clades seemingly showing either dwarfism or gigantism⁶. Thus, we accounted for phylogeny by including the phylogenetic relatedness correlation matrix as a random effect. The species term captures the similarities of effect sizes within the same species, while the phylogenetic term represents the similarity due to relatedness⁶²⁴. We also added ‘Source’ as a random effect intercept to account for between-source variability and the fact that we had multiple response ratios per study. In some cases, ‘Source’ represented the combination of two sources of data, one for the island size and one for the mainland size. Finally, we included an observation level random effect, which represents the residual variance that needs to be explicitly modelled in a meta-analysis²⁹. Total heterogeneity, and heterogeneity due to phylogeny, source and species identity were computed following Nakagawa & Santos (2012)²⁹.

We tested the robustness of our results against several potential limitations. Because multiple island populations were often compared with a single mainland population, we accounted for these repeated measures in a variance-covariance matrix where the diagonal includes the sampling variances, and the off-diagonals of the matrix represent the shared variance (covariance) among the response ratios due to the common mainland population⁶²⁵. Further, we

compared our main results to models fitted with $\ln RR$ and sampling variances corrected for small sample size³⁸. Another potential problem is that regressions using ratios may lead to spurious correlations^{31,32}. Thus, we conducted an additional analysis testing the statistical significance of body size trends by regressing island mass against mainland mass, following previous studies^{4,5,20,41}. Phylogenetic meta-regressions were run using island mass as the response variable, and mainland mass as the predictor (both transformed with natural-log), with random effects as specified above, and sampling variance $sd_i^2/mass_i^2 \cdot N_i$. This approach has some limitations in being harder to visualize and less effective in considering the sampling variance of measurements (representing intrapopulation variability), yet nonetheless provides an alternative approach for assessing the robustness of our results, in line with previous studies^{4,5,20,41}. Finally, we assessed publication bias by testing the influence of data source on the relationship between size ratio and mainland mass. This involved comparing whether patterns differed in island-mainland pairs extracted from studies testing the island rule (38.6% of cases) versus pairs extracted from studies not testing the island rule (61.4% of cases).

Testing ecological hypotheses explaining insular size shifts

To evaluate the relative role of key mechanisms proposed to influence body size evolution in island fauna, we compiled a further range of variables (Supplementary Table 1, Extended Data Fig. 1). These included island area (linked to both resource limitation and to ecological release from both predation and competition) and spatial isolation (linked to reduced colonization from mainland populations for smaller taxa, i.e. immigration selection⁵⁰). In addition, we included climatic and resource seasonality, which are linked to the starvation resistance hypothesis, and productivity and species diet, each of which are linked to resource limitation. Because body size evolution may be influenced by climate (e.g. Bergmann's rule)^{9,626}, we also included mean temperature, which is linked to body size adaptations for enhancing heat conservation or dissipation (thermoregulation hypothesis). For amphibians, we included precipitation as a proxy for water supply linked to aquatic habitats, moisture and humidity (water availability hypothesis).

We modelled interactions between body size and each of the explanatory variables because we expected these factors to differentially affect species of different sizes, thus producing different effects in small, medium-sized and large species. In line with the ecological release and resource

limitation hypotheses, we expected the slope of the $\ln RR$ -mainland mass relationship to be steeper in smaller islands, isolated from the mainland and with fewer or no predators (Fig. 1). Further, if resource availability is a key factor, we also expected large species to undergo dwarfism on islands with low productivity^{48,49}, and for dwarfism to be accentuated in dietary niches with high energy requirements, including carnivory⁹. In addition, high seasonality in resources and in temperature was expected to result in increased gigantism in small-sized species, because energy reserves increase faster than energy depletion as body size increases (starvation resistance hypothesis)^{9,627}. We hypothesized that smaller species would benefit comparatively more by increasing in size than larger species. Because amphibians are generally small-sized, we also fitted a model for this group with only additive terms (mainland mass + sdNDVI) where seasonality in resources would result in larger body sizes for all species. Finally, mechanisms driven by thermoregulation and water availability predict that body size shifts are associated with temperature and rainfall, respectively. Mean temperature was expected to predominantly affect endotherms and small ectotherms with good thermoregulating abilities (reptiles and anurans) living on cold islands which, compared to similar-sized species on islands with a mild climate, would exhibit more pronounced gigantism to enhance heat conservation. We fitted the effect of temperature as an interactive (mainland mass $\times$ Tmean) or additive term (mainland mass + Tmean) to assess whether only small species or all species would increase in size in low temperature islands (see details in Supplementary Table 1, Extended Data Fig 1, Supplementary Table 7).

Prior to modeling, all the moderators (explanatory variables) were inspected and $\log_{10}$ -transformed if necessary to meet normality assumptions in model errors. We considered a result to be significant when the 95% confidence interval (CI) did not cross zero. We assessed the explained heterogeneity using Omnibus test for moderators (Q_m) and the percentage of variance explained by the moderators using R^2 marginal⁶²⁸. All figures show the relationship between size response ratio and body mass, and how this might be altered by the mechanisms explained above.

All analyses were performed in R 3.5.3⁶²⁹ using the packages *metafor* v2.0⁶³⁰ and *metagear* v0.4⁶³¹ for the meta-regression models and data imputation, *metaDigitise* v1.0⁶³² for data extraction from plots, *ape* v5.2⁶³³ for estimating branch lengths and resolving polytomies, *rotl*

v3.0.4⁶³⁴ for building the phylogenies for our species by searching the Open Tree Taxonomy⁶³⁵ and retrieving the phylogenetic relationships from the Open Tree of Life⁶³⁶, *sf* v0.7-3⁶³⁷ and *raster* v2.7-15⁶³⁸ for spatial analyses, *dplyr* v0.8.0.1⁶³⁹ and *reshape2* v1.4.3⁶⁴⁰ for data manipulation and *ggplot2* v 3.3.0.9000⁶⁴¹ and *ggpubr* v0.1.8⁶⁴² for data visualization. ArcMap 10.5 was used for Figure 2. Silhouettes in figures were extracted from ‘phylopic’ (<https://phylopic.org>). The PRISMA Checklist for systematic reviews is available in Appendix 3.

Data availability

All data are available at https://github.com/anabenlop/Island_Rule and https://figshare.com/projects/Body_size_evolution_in_insular_vertbrates/89102.

Code availability

The code to conduct the analyses is available at https://github.com/anabenlop/Island_Rule.

References

- 1 Foster, J. B. Evolution of mammals on islands. *Nature* **202**, 234-235 (1964).
- 2 Baeckens, S. & Van Damme, R. The island syndrome. *Curr. Biol.* **30**, R338-R339 (2020).
- 3 Whittaker, R. J. & Fernández-Palacios, J. M. *Island biogeography: ecology, evolution, and conservation*. (Oxford University Press, 2007).
- 4 Lomolino, M. V. Body size of mammals on islands: the island rule reexamined. *Am. Nat.* **125**, 310-316 (1985).
- 5 Lomolino, M. V. Body size evolution in insular vertebrates: generality of the island rule. *J. Biogeogr.* **32**, 1683-1699 (2005).
- 6 Meiri, S., Cooper, N. & Purvis, A. The island rule: made to be broken? *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **275**, 141-148 (2008).
- 7 Meiri, S., Dayan, T. & Simberloff, D. The generality of the island rule reexamined. *J. Biogeogr.* **33**, 1571-1577 (2006).
- 8 Meiri, S., Dayan, T. & Simberloff, D. Body size of insular carnivores: little support for the island rule. *Am. Nat.* **163**, 469-479 (2004).
- 9 Lomolino, M. V., Sax, D. F., Palombo, M. R. & van der Geer, A. A. Of mice and mammoths: evaluations of causal explanations for body size evolution in insular mammals. *J. Biogeogr.* **39**, 842-854 (2012).
- 10 Millien, V. Relative effects of climate change, isolation and competition on body-size evolution in the Japanese field mouse, *Apodemus argenteus*. *J. Biogeogr.* **31**, 1267-1276 (2004).
- 11 Angerbjorn, A. Gigantism in island populations of wood mice (*Apodemus*) in Europe. *Oikos* **47**, 47-56 (1986).
- 12 Schillaci, M. A., Meijaard, E. & Clark, T. The effect of island area on body size in a primate species from the Sunda Shelf Islands. *J. Biogeogr.* **36**, 362-371 (2009).

571 13 Radtkey, R. R., Fallon, S. M. & Case, T. J. Character displacement in some
572 Cnemidophorus lizards revisited: a phylogenetic analysis. *Proceedings of the National*
573 *Academy of Sciences* **94**, 9740-9745 (1997).

574 14 McClain, C. R., Durst, P. A., Boyer, A. G. & Francis, C. D. Unravelling the determinants
575 of insular body size shifts. *Biol. Lett.* **9**, 20120989 (2013).

576 15 Clegg, S. M. & Owens, P. The 'island rule' in birds: medium body size and its ecological
577 explanation. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **269**, 1359-1365 (2002).

578 16 Raia, P. & Meiri, S. The island rule in large mammals: Paleontology meets ecology.
579 *Evolution* **60**, 1731-1742 (2006).

580 17 Bromham, L. & Cardillo, M. Primates follow the 'island rule': implications for
581 interpreting *Homo floresiensis*. *Biol. Lett.* **3**, 398-400 (2007).

582 18 Montesinos, R., da Silva, H. R. & de Carvalho, A. L. G. The 'island rule' acting on anuran
583 populations (Bufonidae: *Rhinella ornata*) of the Southern Hemisphere. *Biotropica* **44**,
584 506-511 (2012).

585 19 Boback, S. M. Body size evolution in snakes: evidence from island populations. *Copeia*
586 **2003**, 81-94 (2003).

587 20 Meiri, S. Size evolution in island lizards. *Global Ecol. Biogeogr.* **16**, 702-708 (2007).

588 21 Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C. & Meiri, S. Is the island rule
589 general? Turtles disagree. *Global Ecol. Biogeogr.* **23**, 689-700 (2014).

590 22 Boyer, A. G. & Jetz, W. Biogeography of body size in Pacific island birds. *Ecography*
591 **33**, 369-379 (2010).

592 23 Herczeg, G., Gonda, A. & Merilä, J. Evolution of gigantism in nine-spined sticklebacks.
593 *Evolution* **63**, 3190-3200 (2009).

594 24 Palmer, M. Testing the 'island rule' for a tenebrionid beetle (Coleoptera, Tenebrionidae).
595 *Acta Oecol.* **23**, 103-107 (2002).

596 25 McClain, C. R., Boyer, A. G. & Rosenberg, G. The island rule and the evolution of body
597 size in the deep sea. *J. Biogeogr.* **33**, 1578-1584 (2006).

598 26 Biddick, M., Hendriks, A. & Burns, K. Plants obey (and disobey) the island rule. *Proc.*
599 *Natl. Acad. Sci. USA* **116**, 17632-17634 (2019).

600 27 Lokatis, S. & Jeschke, J. M. The island rule: An assessment of biases and research trends.
601 *Journal of Biogeography* **45**, 289-303 (2018).

602 28 Gurevitch, J., Koricheva, J., Nakagawa, S. & Stewart, G. Meta-analysis and the science
603 of research synthesis. *Nature* **555**, 175 (2018).

604 29 Nakagawa, S. & Santos, E. S. Methodological issues and advances in biological meta-
605 analysis. *Evol. Ecol.* **26**, 1253-1274 (2012).

606 30 Hedges, L. V., Gurevitch, J. & Curtis, P. S. The meta-analysis of response ratios in
607 experimental ecology. *Ecology* **80**, 1150-1156 (1999).

608 31 Atchley, W. R., Gaskins, C. T. & Anderson, D. Statistical properties of ratios. I.
609 Empirical results. *Syst. Zool.* **25**, 137-148 (1976).

610 32 Prairie, Y. T. & Bird, D. F. Some misconceptions about the spurious correlation problem
611 in the ecological literature. *Oecologia* **81**, 285-288 (1989).

612 33 Phillips, L. B., Hansen, A. J. & Flather, C. H. Evaluating the species energy relationship
613 with the newest measures of ecosystem energy: NDVI versus MODIS primary
614 production. *Remote Sens. Environ.* **112**, 3538-3549 (2008).

615 34 Olesen, J. M. & Valido, A. Lizards as pollinators and seed dispersers: an island
616 phenomenon. *Trends Ecol. Evol.* **18**, 177-181 (2003).

- 617 35 Zamora-Camacho, F., Reguera, S. & Moreno-Rueda, G. Bergmann's Rule rules body
618 size in an ectotherm: heat conservation in a lizard along a 2200-metre elevational
619 gradient. *J. Evol. Biol.* **27**, 2820-2828 (2014).
- 620 36 Valenzuela-Sánchez, A., Cunningham, A. A. & Soto-Azat, C. Geographic body size
621 variation in ectotherms: effects of seasonality on an anuran from the southern temperate
622 forest. *Front. Zool.* **12**, 37 (2015).
- 623 37 Ashton, K. G. Do amphibians follow Bergmann's rule? *Can. J. Zool./Rev. Can. Zool.* **80**,
624 708-716 (2002).
- 625 38 Lajeunesse, M. J. Bias and correction for the log response ratio in ecological
626 meta-analysis. *Ecology* **96**, 2056-2063 (2015).
- 627 39 Lomolino, M. V. *et al.* Of mice and mammoths: generality and antiquity of the island
628 rule. *J. Biogeogr.* **40**, 1427-1439 (2013).
- 629 40 Boback, S. M. & Guyer, C. Empirical evidence for an optimal body size in snakes.
630 *Evolution* **57**, 345-451 (2003).
- 631 41 Meiri, S., Raia, P. & Phillimore, A. B. Slaying dragons: limited evidence for unusual
632 body size evolution on islands. *J. Biogeogr.* **38**, 89-100 (2011).
- 633 42 Brown, J. H., Marquet, P. A. & Taper, M. L. Evolution of body size: consequences of an
634 energetic definition of fitness. *Am. Nat.* **142**, 573-584 (1993).
- 635 43 Damuth, J. Cope's rule, the island rule and the scaling of mammalian population density.
636 *Nature* **365**, 748-750 (1993).
- 637 44 Raia, P., Carotenuto, F. & Meiri, S. One size does not fit all: no evidence for an optimal
638 body size on islands. *Global Ecol. Biogeogr.* **19**, 475-484 (2010).
- 639 45 Maurer, B. The evolution of body size in birds. II. The role of reproductive power. *Evol.*
640 *Ecol.* **12**, 935-944 (1998).
- 641 46 Blackburn, T. M. & Gaston, K. J. The distribution of body sizes of the world's bird
642 species. *Oikos*, 127-130 (1994).
- 643 47 Feldman, A., Sabath, N., Pyron, R. A., Mayrose, I. & Meiri, S. Body sizes and
644 diversification rates of lizards, snakes, amphisbaenians and the tuatara. *Global Ecol.*
645 *Biogeogr.* **25**, 187-197 (2016).
- 646 48 McNab, B. K. Minimizing energy expenditure facilitates vertebrate persistence on
647 oceanic islands. *Ecol. Lett.* **5**, 693-704 (2002).
- 648 49 McNab, B. K. Geographic and temporal correlations of mammalian size reconsidered: a
649 resource rule. *Oecologia* **164**, 13-23 (2010).
- 650 50 Durst, P. A. & Roth, V. L. Mainland size variation informs predictive models of
651 exceptional insular body size change in rodents. *Proceedings of the Royal Society B:*
652 *Biological Sciences* **282**, 20150239 (2015).
- 653 51 Tobias, J. A., Ottenburgs, J. & Pigot, A. L. Avian Diversity: Speciation,
654 Macroevolution, and Ecological Function. *Annu. Rev. Ecol., Evol. Syst.* **51**, 533-560
655 (2020).
- 656 52 Li, J.-W. *et al.* Rejecting strictly allopatric speciation on a continental island: prolonged
657 postdivergence gene flow between Taiwan (*Leucodioptron taewanus*, Passeriformes
658 Timaliidae) and Chinese (*L. canorum canorum*) hwameis. *Mol. Ecol.* **19**, 494-507 (2010).
- 659 53 Somveille, M., Rodrigues, A. S. & Manica, A. Why do birds migrate? A macroecological
660 perspective. *Global Ecol. Biogeogr.* **24**, 664-674 (2015).
- 661 54 Blondel, J. Evolution and ecology of birds on islands: trends and prospects. *Vie et milieu*
662 (1980) **50**, 205-220 (2000).

663 55 Grant, P. R. Patterns on islands and microevolution. *Evolution on islands* (1998).
664 56 Novosolov, M. *et al.* Power in numbers. Drivers of high population density in insular
665 lizards. *Global Ecol. Biogeogr.* **25**, 87-95 (2016).
666 57 Santini, L. *et al.* Global drivers of population density in terrestrial vertebrates. *Global*
667 *Ecol. Biogeogr.* **27**, 968-979 (2018).
668 58 Castellano, S. & Giacoma, C. Morphological variation of the green toad, *Bufo viridis*, in
669 Italy: a test of causation. *J. Herpetol.*, 540-550 (1998).
670 59 Ashton, K. G. Body size variation among mainland populations of the western rattlesnake
671 (*Crotalus viridis*). *Evolution* **55**, 2523-2533 (2001).
672 60 Athanassiou, A., van der Geer, A. A. & Lyras, G. A. Pleistocene insular Proboscidea of
673 the Eastern Mediterranean: a review and update. *Quat. Sci. Rev.* **218**, 306-321 (2019).
674 61 Herridge, V. L. & Lister, A. M. Extreme insular dwarfism evolved in a mammoth. *Proc.*
675 *R. Soc. Lond., Ser. B: Biol. Sci.* **279**, 3193-3200 (2012).
676 62 Faurby, S. & Svenning, J.-C. Resurrection of the island rule: human-driven extinctions
677 have obscured a basic evolutionary pattern. *Am. Nat.* **187**, 812-820 (2016).
678 63 Steadman, D. W. *Extinction and biogeography of tropical Pacific birds*. (University of
679 Chicago Press, 2006).
680 64 MacPhee, R. D. & Horowitz, I. New craniodental remains of the Quaternary Jamaican
681 monkey *Xenothrix mcgregori* (Xenotrichini, Callicebinae, Pitheciidae), with a
682 reconsideration of the Aotus hypothesis. *Am. Mus. Novit.* **2004**, 1-51 (2004).
683 65 MacPhee, R. & Fleagle, J. Postcranial remains of *Xenothrix mcgregori* (Primates,
684 Xenotrichidae) and other late Quaternary mammals from long Mile Cave, Jamaica. *Bull.*
685 *Am. Mus. Nat. Hist. N. Y.* **206**, 287-321 (1991).
686 66 Pregill, G. Body size of insular lizards: a pattern of Holocene dwarfism. *Evolution* **40**,
687 997-1008 (1986).
688 67 Hansen, D. M., Donlan, C. J., Griffiths, C. J. & Campbell, K. J. Ecological history and
689 latent conservation potential: large and giant tortoises as a model for taxon substitutions.
690 *Ecography* **33**, 272-284 (2010).
691 68 Boyer, A. G. Extinction patterns in the avifauna of the Hawaiian islands. *Divers. Distrib.*
692 **14**, 509-517 (2008).
693 69 White, A. W., Worthy, T. H., Hawkins, S., Bedford, S. & Spriggs, M. Megafaunal
694 meiolaniid horned turtles survived until early human settlement in Vanuatu, Southwest
695 Pacific. *Proc. Natl. Acad. Sci. USA* **107**, 15512-15516 (2010).
696 70 Alcover, J. A., Sans, A. & Palmer, M. The extent of extinctions of mammals on islands.
697 *J. Biogeogr.* **25**, 913-918 (1998).
698 71 Steadman, D. W. Prehistoric extinctions of Pacific island birds: biodiversity meets
699 zooarchaeology. *Science* **267**, 1123-1131 (1995).
700 72 Pérez-Méndez, N., Jordano, P., García, C. & Valido, A. The signatures of Anthropocene
701 defaunation: cascading effects of the seed dispersal collapse. *Sci. Rep.* **6**, 24820 (2016).
702 73 Heinen, J. H., van Loon, E. E., Hansen, D. M. & Kissling, W. D. Extinction-driven
703 changes in frugivore communities on oceanic islands. *Ecography* **41**, 1245-1255 (2018).
704 74 van der Geer, A. A. E. Changing invaders: trends of gigantism in insular introduced rats.
705 *Environ. Conserv.* **45**, 203-211 (2018).
706 75 van der Geer, A. A. E., Lomolino, M. V. & Lyras, G. 'On being the right size' - Do aliens
707 follow the rules? *J. Biogeogr.* **45**, 515-529 (2018).

- Mathys, B. A. & Lockwood, J. L. Rapid evolution of great kiskadees on Bermuda: an assessment of the ability of the island rule to predict the direction of contemporary evolution in exotic vertebrates. *J. Biogeogr.* **36**, 2204-2211 (2009).
- Quinn, J. F. & Dunham, A. E. On hypothesis testing in ecology and evolution. *Am. Nat.* **122**, 602-617 (1983).
- McGill, B. J. & Nekola, J. C. Mechanisms in macroecology: AWOL or purloined letter? Towards a pragmatic view of mechanism. *Oikos* **119**, 591-603 (2010).
- Pigot, A. L. *et al.* Macroevolutionary convergence connects morphological form to ecological function in birds. *Nat. Ecol. Evol.* **4**, 230-239 (2020).
- Krause, M. A., Burghardt, G. M. & Gillingham, J. C. Body size plasticity and local variation of relative head and body size sexual dimorphism in garter snakes (*Thamnophis sirtalis*). *J. Zool.* **261**, 399-407 (2003).
- Krystufek, B., Tvrtkovic, N., Paunovic, M. & Ozkan, B. Size variation in the Northern white-breasted hedgehog *Erinaceus roumanicus*: latitudinal cline and the island rule. *Mammalia* **73**, 299-306 (2009).
- Kubo, M. O. & Takatsuki, S. Geographical Body Size Clines in Sika Deer: Path Analysis to Discern Amongst Environmental Influences. *Evol. Biol.* **42**, 115-127 (2015).
- Kuchling, G., Rhodin, A. G., Ibarrondo, B. R. & Trainor, C. R. A new subspecies of the snake-neck turtle *Chelodina mccordi* from Timor-Leste (East Timor)(Testudines: Chelidae). *Chelonian Conserv. Biol.* **6**, 213-222 (2007).
- Kuo, C.-Y., Lin, Y.-T. & Lin, Y.-S. Sexual size and shape dimorphism in an agamid lizard, *Japalura swinhonis* (Squamata: Lacertilia: Agamidae). *Zool. Stud.* **48**, 351-361 (2009).
- Kurta, A. & Ferkin, M. The correlation between demography and metabolic rate: a test using the beach vole (*Microtus breweri*) and the meadow vole (*Microtus pennsylvanicus*). *Oecologia* **87**, 102-105 (1991).
- Kutrup, B., Cakir, E., Colak, Z., Bulbul, U. & Karaoglu, H. Age and growth of the green toad, *Bufo viridis* (Laurenti, 1768) from an island and a mainland population in Giresun, Turkey. *J. Anim. Vet. Adv.* **10**, 1469-1472 (2011).
- Kwet, A., Steiner, J. & Zillikens, A. A new species of Adenomera (Amphibia: Anura: Leptodactylidae) from the Atlantic rain forest in Santa Catarina, southern Brazil. *Stud. Neotrop. Fauna Environ.* **44**, 93-107 (2009).
- Lampert, K. P., Bernal, X. E., Rand, A. S., Mueller, U. G. & Ryan, M. J. Island populations of *Physalaemus pustulosus*: History influences genetic diversity and morphology. *Herpetologica* **63**, 311-319 (2007).
- Lawlor, T. E. The evolution of body size in mammals: evidence from insular populations in Mexico. *Am. Nat.* **119**, 54-72 (1982).
- Lee, D. E. *et al.* Growth, Age at Maturity, and Age-Specific Survival of the Arboreal Salamander (*Aneides lugubris*) on Southeast Farallon Island, California. *J. Herpetol.* **46**, 64-71 (2012).
- Li, Y. *et al.* Reduced predator species richness drives the body gigantism of a frog species on the Zhoushan Archipelago in China. *J. Anim. Ecol.* **80**, 171-182 (2011).
- Libois, R. M. & Fons, R. Le mulot des Iles d'Hyeres: un cas de gigantisme insulaire. *Vie Milieu* **40**, 217-222 (1990).
- Lichtenbelt, W. D. V. & Albers, K. B. Reproductive adaptations of the green iguana on a semiarid island. *Copeia*, 790-798 (1993).

- 754 94 Lim, B. The distribution, food habits and parasite patterns of the leopard cat
755 (*Prionailurus bengalensis*) in Peninsular Malaysia. *The Journal Wildlife and Parks* **17**
756 (1999).
- 757 95 Lin, L.-H., Mao, F., Chen, C. & Ji, X. Reproductive traits of the gray ratsnake *Ptyas*
758 *korros* from three geographically distinct populations. *Current Zoology* **58**, 820-827
759 (2012).
- 760 96 Lindell, L. E., Forsman, A. & Merila, J. Variation in number of ventral scales in snakes:
761 effects on body size, growth rate and survival in the adder, *Vipera berus*. *J. Zool.* **230**,
762 101-115 (1993).
- 763 97 Ljubisavljević, K., Džukić, G., Vukov, T. & Kalezić, M. Morphological variability of the
764 Hermann's tortoise (*Testudo hermanni*) in the Central Balkans. *Acta Herpetologica* **7**,
765 253-262 (2012).
- 766 98 Lo Cascio, P. & Corti, C. Indagini sull'ecologia dei rettili sauri della RNO e del SIC
767 "Isola di Lampedusa". *Naturalista sicil* **32**, 319-354 (2008).
- 768 99 Lo Cascio, P. & Pasta, S. Preliminary data on the biometry and the diet of a micro-insular
769 population of *Podarcis wagleriana* (Reptilia: Lacertidae). *Acta Herpetologica* **1**, 147-152
770 (2006).
- 771 100 Lo Valvo, M. & Giacalone, G. Biometrical analyses of a Sicilian green toad, *Bufo siculus*
772 (Stöck et al. 2008), population living in sicily *Int. J. Morphol.* **31**, 681-686 (2013).
- 773 101 Long, E. S., Courtney, K. L., Lippert, J. C. & Wall-Scheffler, C. M. Reduced body size of
774 insular black-tailed deer is caused by slowed development. *Oecologia* **189**, 675-685
775 (2019).
- 776 102 López-Martín, J., Ruiz-Olmo, J. & Padró, I. Comparison of skull measurements and
777 sexual dimorphism between the Minorcan pine marten (*Martes martes minoricensis*) and
778 the Iberian pine marten (*M. m. martes*): A case of insularity. *Mamm. Biol.* **71**, 13-24
779 (2006).
- 780 103 Lötters, S. *et al.* Bio-sketches and partitioning of sympatric reed frogs, genus *Hyperolius*
781 (Amphibia; Hyperoliidae), in two humid tropical African forest regions. *J. Nat. Hist.* **38**,
782 1969-1997 (2004).
- 783 104 Luiselli, L., Filippi, E. & Capula, M. Geographic variation in diet composition of the
784 grass snake (*Natrix natrix*) along the mainland and an island of Italy: the effects of
785 habitat type and interference with potential competitors. *Herpetological Journal* **15**, 221-
786 230 (2005).
- 787 105 Luiselli, L., Petrozzi, F., Mebert, K., Zuffi, M. A. L. & Amori, G. Resource partitioning
788 and dwarfism patterns between sympatric snakes in a micro-insular Mediterranean
789 environment. *Ecol. Res.* **30**, 527-535 (2015).
- 790 106 Luz, A. C., Vicente, L. & Monasterio, C. in *Mainland and insular lizards. A*
791 *Mediterranean perspective* (eds C. Corti, P. Lo Cascio, & Biaggini M) 111-123
792 (Firenze University Press, 2006).
- 793 107 Lymberakis, P., Poulakakis, N., Kaliontzopoulou, A., Valakos, E. & Mylonas, M. Two
794 new species of *Podarcis* (Squamata; Lacertidae) from Greece. *Syst. Biodivers.* **6**, 307-318
795 (2008).
- 796 108 Lynch, J., Conroy, J., Kitchener, A., Jefferies, D. & Hayden, T. Variation in cranial form
797 and sexual dimorphism among five European populations of the otter *Lutra lutra*. *J. Zool.*
798 **238**, 81-96 (1996).

- 799 109 Lyon Jr, M. W. Mammals collected in eastern Sumatra by Dr. WL Abbott during 1903,
800 1906, and 1907, with descriptions of new species and subspecies. *Proc. U. S. Natl. Mus.*
801 (1908).
- 802 110 Lyon, M. W. Mammals of Banka, Mendanau, and Billiton Islands, between Sumatra and
803 Borneo. *Proc. U. S. Natl. Mus.* **31**, 575–612 (1906).
- 804 111 Lyon, M. W. Mammals collected by Dr. WL Abbott on Borneo and some of the small
805 adjacent islands. *Proc. U. S. Natl. Mus.* **40**, 53-146 (1911).
- 806 112 Lyon, M. W. Tree shrews: an account of the mammalian family Tupaiidae. *Proceedings*
807 *of the U. S. National Museum* **45**, 1-188 (1913).
- 808 113 Lyon, M. W. Mammals collected by Dr. WL Abbott on the chain of islands lying off the
809 western coast of Sumatra, with descriptions of twenty-eight new species and subspecies.
810 *Proc. U. S. Natl. Mus.* **52**, 437–462 (1916).
- 811 114 Maddock, S. *Systematics and phylogeography of Seychelles amphibians*. PhD thesis,
812 UCL (University College London), (2016).
- 813 115 Madeira, B. d. C. M. A. *Sexual dimorphism and reproductive phenology of common birds*
814 *in São Tomé island: conservation implications* PhD thesis, Universidade de Lisboa,
815 (2018).
- 816 116 Madsen, T. & Shine, R. Phenotypic plasticity in body sizes and sexual size dimorphism
817 in European grass snakes. *Evolution* **47**, 321-325 (1993).
- 818 117 Mageski, M. *et al.* The island rule in the Brazilian frog *Phyllodytes luteolus* (Anura:
819 Hylidae): incipient gigantism? *Zoologia* **32**, 329-333 (2015).
- 820 118 Magnanou, E., Fons, R., Blondel, J. & Morand, S. Energy expenditure in Crocidurinae
821 shrews (Insectivora): Is metabolism a key component of the insular syndrome? *Comp.*
822 *Biochem. Physiol., A: Mol. Integr. Physiol.* **142**, 276-285 (2005).
- 823 119 Maharadatunkamsi, H. S., Kitchener, D. & Schmitt, L. Relationships between
824 morphology, genetics and geography in the cave fruit bat *Eonycteris spelaea* (Dobson,
825 1871) from Indonesia. *Biol. J. Linn. Soc.* **79**, 511-522 (2003).
- 826 120 Major, R. E. Latitudinal and insular variation in morphology of a small Australian
827 passerine: consequences for dispersal distance and conservation. *Aust. J. Zool.* **60**, 210-
828 218 (2012).
- 829 121 Mallick, S., Driessen, M. & Hocking, G. Biology of the southern brown bandicoot
830 (*Isodon obesulus*) in south-eastern Tasmania. II. Demography. *Aust. Mammal.* **20**, 339-
831 347 (1998).
- 832 122 Malmquist, M. G. Character displacement and biogeography of the pygmy shrew in
833 northern Europe. *Ecology* **66**, 372-377 (1985).
- 834 123 Manier, M. K. Geographic variation in the long-nosed snake, *Rhinocheilus lecontei*
835 (Colubridae): beyond the subspecies debate. *Biol. J. Linn. Soc.* **83**, 65-85 (2004).
- 836 124 Manríquez-Morán, N. L., Cruz, M. V.-S. & Mendez-De La Cruz, F. R. Reproductive
837 biology of the parthenogenetic lizard, *Aspidoscelis cozumela*. *Herpetologica* **61**, 435-439
838 (2005).
- 839 125 Marinelli, L. & Millar, J. S. The ecology of beach-dwelling *Peromyscus maniculatus* on
840 the Pacific Coast. *Canadian Journal of Zoology* **67**, 412-417 (1989).
- 841 126 Martins, M., Araujo, M. S., Sawaya, R. J. & Nunes, R. Diversity and evolution of
842 macrohabitat use, body size and morphology in a monophyletic group of Neotropical
843 pitvipers (Bothrops). *J. Zool.* **254**, 529-538 (2001).

- 127 Martins, M., Arnaud, G. & Avila-Villegas, H. Juvenile recruitment, early growth, and
morphological variation in the endangered Santa Catalina island rattlesnake, *Crotalus*
catalinensis. *Herpetol. Conserv. Biol.* **7**, 376-382 (2012).
- 128 Marunouchi, J., Ueda, H. & Ochi, O. Variation in age and size among breeding
populations at different altitudes in the Japanese newts, *Cynops pyrrhogaster*. *Amphibia-*
Reptilia **21**, 381-396 (2000).
- 129 Matsui, M. & Ota, H. Parameters of fecundity in *Microhyla ornata* from the Yaeyama
Group of the Ryukyu Archipelago. *Japanese Journal of Herpetology* **10**, 73-79 (1984).
- 130 Matsui, M., Shimada, T. & Sudin, A. First record of the tree frog genus *Chiromantis* from
Borneo with the description of a new species (Amphibia: Rhacophoridae). *Zool. Sci.* **31**,
45-51 (2014).
- 131 Matsui, M., Toda, M. & Ota, H. A new species of frog allied to *Fejervarya limnocharis*
from the Southern Ryukyus, Japan (Amphibia: Ranidae). *Curr. Herp.* **26**, 65-79 (2007).
- 132 Mazák, J. H. & Groves, C. P. A taxonomic revision of the tigers (*Panthera tigris*) of
Southeast Asia. *Mamm. Biol.* **71**, 268-287 (2006).
- 133 McCord, W. P. & Iverson, J. B. A new species of *Ocadia* (Testudines: Batagurinae) from
southwestern China. *Proc. Biol. Soc. Wash.* **107**, 52-59 (1994).
- 134 McCord, W. P. & Thomson, S. A. A new species of *Chelodina* (Testudines: Pleurodira:
Chelidae) from northern Australia. *J. Herpetol.* **36**, 255-267 (2002).
- 135 McFadden, K. W. & Meiri, S. Dwarfism in insular carnivores: a case study of the pygmy
raccoon. *J. Zool.* **289**, 213-221 (2013).
- 136 McLaughlin, J. F. & Roughgarden, J. Avian predation on *Anolis* lizards in the
northeastern caribbean: Inter-island contrast. *Ecology* **70**, 617-628 (1989).
- 137 Medway, L. Observations on the fauna of Pulau Tioman and Pulau Tulai. 2. The
mammals *Natl. Mus. Singap. Bull.* **34**, 9-32 (1966).
- 138 Mees, G. F. *A systematic review of the Indo-Australian Zosteropidae (Part III)*. (EJ Brill,
1969).
- 139 Meijaard, E. & Groves, C. A taxonomic revision of the *Tragulus* mouse-deer
(Artiodactyla). *Zool. J. Linn. Soc.* **140**, 63-102 (2004).
- 140 Meijaard, E. & Groves, C. P. Morphometrical relationships between South-east Asian
deer (Cervidae, tribe Cervini): evolutionary and biogeographic implications. *J. Zool.* **263**,
179-196 (2004).
- 141 Meik, J. M., Lawing, A. M. & Pires-daSilva, A. Body size evolution in insular speckled
rattlesnakes (Viperidae: *Crotalus mitchellii*). *Plos One* **5** (2010).
- 142 Melo, M. *et al.* Multiple lines of evidence support the recognition of a very rare bird
species: the Príncipe thrush. *J. Zool.* **282**, 120-129 (2010).
- 143 Melo, M. & O’Ryan, C. Genetic differentiation between Príncipe Island and mainland
populations of the grey parrot (*Psittacus erithacus*), and implications for conservation.
Mol. Ecol. **16**, 1673-1685 (2007).
- 144 Melton, R. Body size and island *Peromyscus*: a pattern and a hypothesis. *Evolutionary*
Theory **6**, 113-126 (1982).
- 145 Menzies, J. & Tyler, M. The systematics and adaptations of some Papuan microhylid
frogs which live underground. *J. Zool.* **183**, 431-464 (1977).
- 146 Merriam, C. H. in *North American Fauna N° 10* 5-34 (Government Printing Office,
1895).

- 889 147 Merriam, C. H. Synopsis of the American shrews of the genus *Sorex*. *North American*
890 *Fauna* **10**, 57-125 (1895).
- 891 148 Merriam, C. H. Six new mammals from Cozumel Island, Yucatan. *Proc. Biol. Soc. Wash.*
892 **14**, 99-104 (1901).
- 893 149 Merriam, C. H. Descriptions of ten new kangaroo rats. *Proc. Biol. Soc. Wash.* **20**, 75-80
894 (1907).
- 895 150 Miller, C. & Miller, T. Population dynamics and diet of rodents on Rangitoto Island, New
896 Zealand, including the effect of a 1080 poison operation. *N. Z. J. Ecol.*, 19-27 (1995).
- 897 151 Miller, G. S. Mammals collected by Dr. WL Abbott in the region of the Indragiri River,
898 Sumatra. *Proc. Acad. Nat. Sci. Phila.* **54**, 143-159 (1902).
- 899 152 Miller, G. S. *Catalogue of the Mammals of Western Europe (Europe Exclusive of Russia)*
900 *in the Collection of the British Museum*. (Order of the Trustees of the British Museum,
901 1912).
- 902 153 Millien, V. & Damuth, J. Climate change and size evolution in an island rodent species:
903 New perspectives on the island rule. *Evolution* **58**, 1353-1360 (2004).
- 904 154 Molina-Borja, M. & Rodriguez-Dominguez, M. A. Evolution of biometric and life-
905 history traits in lizards (*Gallotia*) from the Canary Islands. *J. Zool. Syst. Evol. Res.* **42**, 44-
906 53 (2004).
- 907 155 Monadjem, A., McCleery, R. A. & Collier, B. A. Activity and movement patterns of the
908 tortoise *Stigmochelys pardalis* in a subtropical savanna. *J. Herpetol.* **47**, 237-242 (2013).
- 909 156 Montesinos, R., da Silva, H. R. & Gomes de Carvalho, A. L. The 'island rule' acting on
910 anuran populations (*Bufo*: *Rhinella ornata*) of the Southern Hemisphere. *Biotropica*
911 **44**, 506-511 (2012).
- 912 157 Montgomery, C. E., Boback, S. M., Green, S. E., Paulissen, M. A. & Walker, J. M.
913 *Cnemidophorus lemniscatus* (Squamata: Teiidae) on Cayo Cochino Pequeño, Honduras:
914 extent of island occupancy, natural history, and conservation status. *Herpetol. Conserv.*
915 *Biol.* **6**, 10-24 (2011).
- 916 158 Moratelli, R., Wilson, D. E., Novaes, R. L., Helgen, K. M. & Gutiérrez, E. E. Caribbean
917 Myotis (Chiroptera, Vespertilionidae), with description of a new species from Trinidad
918 and Tobago. *J. Mammal.* **98**, 994-1008 (2017).
- 919 159 Mori, A. & Hasegawa, M. Early growth of *Elaphe quadrivirgata* from an insular gigantic
920 population. *Curr. Herp.* **21**, 43-50 (2002).
- 921 160 Mori, A., Ikeuchi, I. & Hasegawa, M. Calling activity of an anuran assemblage in a
922 temporary pond in a dry forest of Madagascar. *Curr. Herp.* **34**, 140-148 (2015).
- 923 161 Mori, E. *et al.* Skull shape and Bergmann's rule in mammals: hints from Old World
924 porcupines. *J. Zool.* **308**, 47-55 (2019).
- 925 162 Motokawa, M. Geographic variation in the Japanese white-toothed shrew *Crocidura*
926 *dsinezumi*. *Acta Theriologica* **48**, 145-156 (2003).
- 927 163 Motokawa, M. & Lin, L.-K. Geographic variation in the mole-shrew *Anourosorex*
928 *squamipes*. *Mammal Study* **27**, 113-120 (2002).
- 929 164 Motokawa, M., Lin, L.-K., Harada, M. & Hattori, S. Morphometric geographic variation
930 in the Asian lesser white-toothed shrew *Crocidura shantungensis* (Mammalia,
931 Insectivora) in East Asia. *Zool. Sci.* **20**, 789-795 (2003).
- 932 165 Munks, S. The breeding biology of *Pseudocheirus peregrinus viverrinus* on Flinders
933 Island, Bass Strait. *Wildl. Res.* **22**, 521-533 (1995).
- 934 166 MVZ. (University of California, Berkeley).

- 935 167 Myers, S., Brown, G. & Kleindorfer, S. Divergence in New Holland honeyeaters
936 (*Phylidonyris novaehollandiae*): evidence from morphology and feeding behavior. *J.*
937 *Ornithol.* **151**, 287-296 (2010).
- 938 168 Nagorsen, D. Body weight variation among insular and mainland American martens.
939 *Martens, sables, and fishers biology and conservation. Comstock, Ithaca, NY*, 85-97
940 (1994).
- 941 169 Nagy, Z. T., Glaw, F. & Vences, M. Systematics of the snake genera *Stenophis* and
942 *Lycodryas* from Madagascar and the Comoros. *Zool. Scr.* **39**, 426-435 (2010).
- 943 170 Nanova, O. & Prôa, M. Cranial features of mainland and Commander Islands (Russia)
944 Arctic foxes (*Vulpes lagopus*) reflect their diverging foraging strategies. *Polar Res.* **36**, 7
945 (2017).
- 946 171 Napier, P. H. *Catalogue of Primates in the British Museum (Natural History) and*
947 *Elsewhere in the British Isles. Part III: Family Cercopithecidae, subfamily Colobinae.,*
948 Vol. 3 (British Museum (Natural History), 1985).
- 949 172 Nelson, E. Descriptions of new birds from the Tres Marias Islands, western Mexico.
950 *Proc. Biol. Soc. Wash.* **12**, 5-11 (1898).
- 951 173 Nelson, E. W. in *North American Fauna* 1-314 (US Government Printing Office, 1909).
- 952 174 Newman, D. G. Activity, dispersion, and population densities of Hamilton's frog
953 (*Leiopelma hamiltoni*) on Maud and Stephens Islands, Nez Zealands. *Herpetologica* **46**,
954 319-330 (1990).
- 955 175 Nguyen, S. T., Oshida, T., Dang, P. H., Bui, H. T. & Motokawa, M. A new species of
956 squirrel (Sciuridae: Callosciurus) from an isolated island off the Indochina Peninsula in
957 southern Vietnam. *J. Mammal.* **99**, 813-825 (2018).
- 958 176 Nijman, V. Group composition and monandry in grizzled langurs, *Presbytis comata*, on
959 Java. *Folia Primatol.* **88**, 237-254 (2017).
- 960 177 Nor, S. M. *The mammalian fauna on the islands at the northern tip of Sabah, Borneo.*
961 (Field Museum of Natural History., 1996).
- 962 178 Norman, F. & Hurley, V. Gonad measurements and other parameters from Chestnut Teal
963 *Anas castanea* collected in the Gippsland Lakes region, Victoria. *Emu-Austral*
964 *Ornithology* **84**, 52-55 (1984).
- 965 179 Nowak, R. M. & Walker, E. P. *Walker's Mammals of the World*. Vol. 1 (JHU Press,
966 1999).
- 967 180 O'Connell, D. P. *et al.* A sympatric pair of undescribed white-eye species (Aves:
968 Zosteropidae: Zosterops) with different origins. *Zool. J. Linn. Soc.* **186**, 701-724 (2019).
- 969 181 Oates, J. F., Davies, A. G. & Delson, E. in *Colobine Monkeys: Their Ecology, Behaviour*
970 *and Evolution* (eds A. Glyn Davies & John F. Oates) (Cambridge University Press,
971 1994).
- 972 182 Oh, H.-S., Yoshinaga, Y., Kaneko, T., Iida, H. & Mori, T. Taxomic re-examination of the
973 *Apodemus agrarius chejuensis*, comparing external and cranial morphological characters
974 among four Asian *Apodemus* species. *J. Fac. Agr. Kyushu Univ* **47**, 373-386 (2003).
- 975 183 Ohdachi, S., Abe, H., Oh, H. & Han, S. Morphological relationships among populations
976 in the *Sorex caecutiens/shinto* group (Eulipotyphla, Soricidae) in East Asia, with a
977 description of a new subspecies from Cheju Island, Korea. *Mamm. Biol.* **70**, 345-358
978 (2005).

979 184 Okada, S., Izawa, M. & Ota, H. Growth and reproduction of *Gekko hokouensis* (Reptilia :
980 Squamata) on Okinawajima Island of the Ryukyu Archipelago, Japan. *J. Herpetol.* **36**,
981 473-479 (2002).

982 185 O'Keeffe, J., O'Boyle, I. & Fogarty, U. Observations on the pathology of tuberculosis in
983 badgers. Report No. 1898473838, (University College Dublin. Centre for Veterinary
984 Epidemiology and Risk Analysis, Dublin, 1997).

985 186 Olsen, P., Debus, S., Czechura, G. & Mooney, N. Comparative feeding ecology of the
986 grey goshawk *Accipiter novaehollandiae* and brown goshawk *Accipiter fasciatus*. *Aust.*
987 *Bird Watch.* **13**, 178-192 (1990).

988 187 Oneto, F., Ottonello, D. & Salvidio, S. Primi dati sulla biometria di *Euleptes euopaea*
989 (Genè, 1839) dell'isola del Tino (La Spezia, Liguria). *Doriana, Ann. Mus. Civ. Stor. nat*
990 *G. Doria, Genova* **8**, 1-8 (2008).

991 188 Oromi, N. *et al.* Geographical variations in adult body size and reproductive life history
992 traits in an invasive anuran, *Discoglossus pictus*. *Zoology* **119**, 216-223 (2016).

993 189 Osgood, W. H. Natural history of the Queen Charlotte islands. *North American Fauna*, 1-
994 87 (1901).

995 190 Osgood, W. H. Revision of the mice of the American genus *Peromyscus*. *North American*
996 *Fauna* **28**, 1-285 (1909).

997 191 Ota, H. Taxonomic redefinition of *Japalura swinhonis* Günther (Agamidae: Squamata),
998 with a description of a new subspecies of *J. polygonata* from Taiwan. *Herpetologica*,
999 280-294 (1991).

1000 192 Ota, H., Lau, M. W., Weidenhofer, T., Yasukawa, Y. & Bogadek, A. Taxonomic review
1001 of the geckos allied to *Gekko chinensis* Gray 1842 (Gekkonidae Reptilia) from China and
1002 Vietnam. *Trop. Zool.* **8**, 181-196 (1995).

1003 193 Owens, J. R. *Ecology and Behavior of the Bioko Island Drill (Mandrillus leucophaeus*
1004 *poensis)*. PhD thesis, Drexel University, (2013).

1005 194 Pabijan, M., Gehring, P.-S., Koehler, J., Glaw, F. & Vences, M. A new microendemic
1006 frog species of the genus *Blommersia* (Anura: Mantellidae) from the east coast of
1007 Madagascar. *Zootaxa* **2978**, 34-50 (2011).

1008 195 Pafilis, P. *et al.* Reproductive biology of insular reptiles: marine subsidies modulate
1009 expression of the "island syndrome". *Copeia*, 545-552 (2011).

1010 196 Pafilis, P., Kapsalas, G., Lymberakis, P., Protopappas, D. & Sotiropoulos, K. Diet
1011 composition of the Karpathos marsh frog (*Pelophylax cerigensis*): what does the most
1012 endangered frog in Europe eat? *Anim. Biodivers. Conserv.* **42**, 1-8 (2019).

1013 197 Pafilis, P., Meiri, S., Foufopoulos, J. & Valakos, E. Intraspecific competition and high
1014 food availability are associated with insular gigantism in a lizard. *Naturwissenschaften*
1015 **96**, 1107-1113 (2009).

1016 198 Pafilis, P. *et al.* Body size affects digestive performance in a Mediterranean lizard.
1017 *Herpetological Journal* **26**, 199-205 (2016).

1018 199 Pagh, S. *et al.* Methods for the identification of farm escapees in feral mink (*Neovison*
1019 *vison*) populations. *PloS One* **14** (2019).

1020 200 Pahl, L. & Lee, A. Reproductive traits of 2 populations of the common ringtail possum,
1021 *Pseudocheirus peregrinus*, in Victoria. *Aust. J. Zool.* **36**, 83-97 (1988).

1022 201 Palacios, F. & Fernández, J. A new subspecies of hare from Majorca (Balearic Islands).
1023 *Mammalia* **56**, 71-86 (1992).

- 1024 202 Papakosta, M. A. Biometric variation in *Martes foina* from mainland Greece and the
1025 Aegean Islands. *Turkish Journal of Zoology* **41**, 654-663 (2017).
- 1026 203 Parker, W. S. & Pianka, E. R. Comparative ecology of populations of the lizard *Uta*
1027 *stansburiana*. *Copeia*, 615-632 (1975).
- 1028 204 Parrish, G. & Gill, B. Natural history of the lizards of the Three Kings Islands, New
1029 Zealand. *N. Z. J. Zool.* **30**, 205-220 (2003).
- 1030 205 Pasachnik, S. A. *et al.* Body size, demography, and body condition in Uta spiny-tailed
1031 iguanas, *Ctenosaura bakeri*. *Herpetol. Conserv. Biol.* **7**, 391-398 (2012).
- 1032 206 Pearson, D. *et al.* Ecological notes on crowned snakes *Elapognathus coronatus* from the
1033 Archipelago of the Recherche in southwestern Australia. *Aust. Zool.* **31**, 610-617 (2001).
- 1034 207 Pedrono, M. & Markwell, T. Maximum size and mass of the ploughshare tortoise,
1035 *Geochelone yniphora*. *Chelonian Conserv. Biol.* **4**, 190-190 (2001).
- 1036 208 Perez-Ramos, E. *et al.* *Aspidoscelis sexlineata* (Sauria: Teiidae) in Mexico: distribution,
1037 habitat, morphology, and taxonomy. *Southwest. Nat.* **55**, 419-425 (2010).
- 1038 209 Persson, S. *et al.* Influence of age, season, body condition and geographical area on
1039 concentrations of chlorinated and brominated contaminants in wild mink (*Neovison*
1040 *vison*) in Sweden. *Chemosphere* **90**, 1664-1671 (2013).
- 1041 210 Peters, J. A. The snakes of the subfamily Dipsadinae. *Miscellaneous Publications of the*
1042 *Museum of Zoology, University of Michigan* **114** (1960).
- 1043 211 Pocock, R. The civet cats of Asia. Part II. . *J. Bombay Nat. Hist. Soc.* **36**, 629-656 (1933).
- 1044 212 Pocock, R. in *Proceedings of the Zoological Society of London.* 613-684 (Wiley Online
1045 Library).
- 1046 213 Pocock, R. I. *The Fauna Of British India Including Ceylon And Burma.* (Taylor and
1047 Francis Ltd, 1939).
- 1048 214 Pons, J. M., Kirwan, G. M., Porter, R. F. & Fuchs, J. A reappraisal of the systematic
1049 affinities of Socotran, Arabian and East African scops owls (Otus, Strigidae) using a
1050 combination of molecular, biometric and acoustic data. *Ibis* **155**, 518-533 (2013).
- 1051 215 Abe, H. Habitat factors affecting the geographic size variation of Japanese moles.
1052 *Mammal Study* **21**, 71-87 (1996).
- 1053 216 Abramov, A. V., Jenkins, P. D., Rozhnov, V. V. & Kalinin, A. A. Description of a new
1054 species of Crocidura (Soricomorpha: Soricidae) from the island of Phu Quoc, Vietnam.
1055 *Mammalia* **72**, 269-272 (2008).
- 1056 217 Abramov, A. V. & Puzachenko, A. Y. Sexual dimorphism of craniological characters in
1057 Eurasian badgers, *Meles spp.*(Carnivora, Mustelidae). *Zoologischer Anzeiger-A Journal*
1058 *of Comparative Zoology* **244**, 11-29 (2005).
- 1059 218 Adams, N. E., Dean, M. D. & Pauly, G. B. Morphological divergence among populations
1060 of *Xantusia riversiana*, a night lizard endemic to the Channel Islands of California.
1061 *Copeia* **106**, 550-562 (2018).
- 1062 219 Aguilar-Moreno, M. *et al.* Dimorfismo sexual de *Aspidoscelis costata costata* (Squamata:
1063 Teiidae) en el sur del Estado de México, México. *Rev. Chil. Hist. Nat.* **83**, 585-592
1064 (2010).
- 1065 220 Ajtić, R. Morphological, biogeographical and ecological characteristics of Kotschy's
1066 gecko (*Cyrtodactylus kotschy* Steindachner, 1870 Gekkonidae) from the mainland
1067 portion of its distribution range. *Fauna Balkana* **3**, 1-70 (2014).
- 1068 221 Albrecht, G. H., Jenkins, P. D. & Godfrey, L. R. Ecogeographic size variation among the
1069 living and subfossil prosimians of Madagascar. *Am. J. Primatol.* **22**, 1-50 (1990).

- 1070 222 Allen, G. M. *The Mammals of China and Mongolia*. (American Museum of Natural
1071 History, 1938).
- 1072 223 Allen, J. A. A preliminary study of the South American opossums of the genus *Didelphis*.
1073 *Bulletin of the American Museum of Natural History* **16**, 249-279 (1902).
- 1074 224 Allen, J. A. Mammals collected in Alaska and northern British Columbia by the Andrew
1075 J. Stone expedition of 1902. *Bulletin of the American Museum of Natural History* **19**,
1076 521-567 (1903).
- 1077 225 Allen, J. A. Notes on American deer of the genus *Mazama*. *Bulletin of the American*
1078 *Museum of Natural History* **34**, 521-553 (1915).
- 1079 226 Allen, J. A. Review of the South American Sciuridae. *Bulletin of the American Museum*
1080 *of Natural History* **34**, 147-309 (1915).
- 1081 227 Allen, J. A. & Chapman, F. M. On a collection of mammals from the island of Trinidad,
1082 with descriptions of new species. *Bulletin of the American Museum of Natural History* **5**,
1083 203-234 (1898).
- 1084 228 Allen, J. A., Kermode, F. & Andrews, R. C. The white bear of southwestern British
1085 Columbia. *Bulletin of the American Museum of Natural History* **26**, 233-238 (1909).
- 1086 229 Altunisik, A., Kalayci, T. E., Gul, C., Ozdemir, N. & Tosunoglu, M. A
1087 skeletochronological study of the smooth newt *Lissotriton vulgaris* (Amphibia: Urodela)
1088 from an island and a mainland population in Turkey. *Ital. J. Zool.* **81**, 381-388 (2014).
- 1089 230 Amor, N. & Farjallah, S. Morphological variation of the African green toad, *Bufo*
1090 *boulengeri* (Amphibia: Anura) in Tunisia. *Pak. J. Zool.* **43** (2011).
- 1091 231 Anderson, R. P. & Handley, C. O. Dwarfism in insular sloths: Biogeography, selection,
1092 and evolutionary rate. *Evolution* **56**, 1045-1058 (2002).
- 1093 232 Andrade, P. *et al.* Ecomorphological patterns in the Blackcap *Sylvia atricapilla*: insular
1094 versus mainland populations. *Bird Study* **62**, 498-507 (2015).
- 1095 233 Andrews, R. M. Evolution of life histories: a comparison of *Anolis* lizards from matched
1096 island and mainland habitats. *Breviora* **454**, 1-51 (1979).
- 1097 234 Angarita-Sierra, T. & Lynch, J. D. A new species of *Ninia* (Serpentes: Dipsadidae) from
1098 Chocó-Magdalena biogeographical province, western Colombia. *Zootaxa* **4244**, 478-492
1099 (2017).
- 1100 235 Angelici, F., Capizzi, D., Amori, G. & Luiselli, L. Morphometric variation in the skulls
1101 of the crested porcupine *Hystrix cristata* from mainland Italy, Sicily, and northern Africa.
1102 *Mamm. Biol.* **68**, 165-173 (2003).
- 1103 236 Aowphol, A., Rujirawan, A., Taksintum, W., Arsirapot, S. & Mcleod, D. S. Re-
1104 evaluating the taxonomic status of *Chiromantis* in Thailand using multiple lines of
1105 evidence (Amphibia: Anura: Rhacophoridae). *Zootaxa* **3702**, 101-123 (2013).
- 1106 237 Araújo, M. & Martins, M. The defensive strike of five species of lanceheads of the genus
1107 *Bothrops* (Viperidae). *Braz. J. Biol.* **67**, 327-332 (2007).
- 1108 238 Arendt, W. J. & Arendt, A. L. Bill deformity in a Pearly-eyed Thrasher from Montserrat,
1109 West Indies. *Auk* **54**, 324-332 (1986).
- 1110 239 Arntzen, J. & García-París, M. Morphological and allozyme studies of midwife toads
1111 (genus *Alytes*), including the description of two new taxa from Spain. *Contributions to*
1112 *Zoology* **65**, 5-34 (1995).
- 1113 240 Aubret, F. Island colonisation and the evolutionary rates of body size in insular neonate
1114 snakes. *Heredity* **115**, 349-356 (2015).

1115 241 Auer, M. & Taskavak, E. Population structure of syntopic *Emys orbicularis* and
1116 *Mauremys rivulata* in western Turkey. *Biologia* **59**, 81-84 (2004).
1117 242 Aumann, T. *Aspects of the biology of the Brown Goshawk Accipiter fasciatus in south-*
1118 *eastern Australia*. MSc thesis, Monash University, (1986).
1119 243 Baier, F. & Hoekstra, H. E. The genetics of morphological and behavioural island traits
1120 in deer mice. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **286** (2019).
1121 244 Baker-Gabb, D. J. *Comparative ecology and behaviour of swamp harriers Circus*
1122 *approximans, spotted harriers C. assimilis and other raptors in Australia and New*
1123 *Zealand*, Royal Australasian Ornithological Union, (1982).
1124 245 Bakhuis, W. L. Size and sexual differentiation in the lizard *Iguana iguana* on a semi-arid
1125 island. *J. Herpetol.* **16**, 322-325 (1982).
1126 246 Balham, R. W. Grey and mallard ducks in the Manawatu district, New Zealand. *Emu-*
1127 *Austral Ornithology* **52**, 163-191 (1952).
1128 247 Bangma, J. T. *et al.* Perfluorinated alkyl acids in plasma of American alligators (*Alligator*
1129 *mississippiensis*) from Florida and South Carolina. *Environ. Toxicol. Chem.* **36**, 917-925
1130 (2017).
1131 248 Bangma, J. T. *et al.* Variation in perfluoroalkyl acids in the American alligator (*Alligator*
1132 *mississippiensis*) at Merritt Island National Wildlife Refuge. *Chemosphere* **166**, 72-79
1133 (2017).
1134 249 Banks, R. A new insular subspecies of spiny pocket mouse (Mammalia: Rodentia). *Proc.*
1135 *Biol. Soc. Wash.* **80**, 101-104 (1967).
1136 250 Barbo, F. E. *et al.* Another new and threatened species of lancehead genus Bothrops
1137 (Serpentes, Viperidae) from Ilha dos Franceses, Southeastern Brazil. *Zootaxa* **4097**, 511-
1138 529 (2016).
1139 251 Barbour, T. & Allen, G. M. The white-tailed deer of eastern United States. *J. Mammal.* **3**,
1140 65-78 (1922).
1141 252 Bartle, J. & Sagar, P. Intraspecific variation in the New Zealand bellbird *Anthornis*
1142 *melanura*. *Notornis* **34**, 253-306 (1987).
1143 253 Barwick, R. E. *Studies on the scincid lizard Egernia cunninghami (Gray, 1832)*. PhD
1144 thesis, The Australian National University, (1965).
1145 254 Başkale, E., Ulubeli, S. A. & Kaska, Y. Age structures and growth parameters of the
1146 Levantine frog, *Pelophylax bedriagae*, at different localities in Denizli, Turkey. *Acta*
1147 *Herpetologica* **13**, 147-154 (2018).
1148 255 Bee, J. W. & Hall, E. R. *Mammals of northern Alaska on the Arctic Slope*. (Museum of
1149 Natural History, University of Kansas, 1956).
1150 256 Bejakovic, D., Kalezic, M. L., Aleksic, I., Dzukic, G. & CrnobrnjaIsailovic, J. Female
1151 reproductive cycle and clutch traits in the Dalmatian wall lizard (*Podarcis melisellensis*).
1152 *Folia Zoologica* **44**, 371-380 (1995).
1153 257 Bell, B. D. Observations on the ecology and reproduction of the New Zealand
1154 Leiopelmid frogs. *Herpetologica* **34**, 340-354 (1978).
1155 258 Bell, R. C. & Irian, C. G. Phenotypic and genetic divergence in reed frogs across a
1156 mosaic hybrid zone on São Tomé Island. *Biol. J. Linn. Soc.* **128**, 672-680 (2019).
1157 259 Bennett, D. & Hampson, K. Further Observations of Varanus olivaceus on the Polillo
1158 Islands. *Hampson et al, Wildlife and Conservation in the Polillo Islands. Polillo project*
1159 *Final Report, Multimedia CD. Viper Press, Glossop* (2003).

1160 260 Benson, S. B. Two new Pocket Mice: genus *Perognathus*, from the Californias.
1161 *University of California Publications in Zoology* **32**, 449-454 (1930).

1162 261 Bentz, E. J., Rodríguez, M. R., John, R. R., Henderson, R. W. & Powell, R. Population
1163 densities, activity, microhabitats, and thermal biology of a unique crevice-and litter-
1164 dwelling assemblage of reptiles on Union Island, St. Vincent and the Grenadines.
1165 *Herpetol. Conserv. Biol.* **6**, 40-50 (2011).

1166 262 Beovides-Casas, K. & Mancina, C. A. Natural history and morphometry of the Cuban
1167 iguana (*Cyclura nubila* Gray, 1831) in Cayo Siju, Cuba. *Anim. Biodivers. Conserv.* **29**, 1-
1168 8 (2006).

1169 263 Berardo, F. *Habitat preferenziali e dinamica di popolazione di Testudo hermanni nelle*
1170 *aree costiere del Molise*. PhD thesis, Università degli studi del Molise, (2015).

1171 264 Berry, R. The evolution of an island population of the house mouse. *Evolution* **18**, 468-
1172 483 (1964).

1173 265 Berry, R., Jakobson, M. & Peters, J. The house mice of the Faroe Islands: a study in
1174 microdifferentiation. *J. Zool.* **185**, 73-92 (1978).

1175 266 Bertolero, A. in *Enciclopedia Virtual de los Vertebrados Españoles*. (ed A. Salvador,
1176 Marco, A.) (Museo Nacional de Ciencias Naturales, 2015).

1177 267 Bertolero, A., Cheylan, M. & Nougarede, J.-P. Accroissement de la fécondité chez la
1178 tortue d'Hermann *Testudo hermanni hermanni* en condition insulaire: Un contre-exemple
1179 du syndrome insulaire? *Revue d'écologie* **62**, 93-98 (2007).

1180 268 Bischoff, W. Bemerkungen zur innerartlichen Variabilität von *Gallotia atlantica* (Peters
1181 & Doria, 1882) (Lacertidae). *Bonn. Zool. Beitr.* **36**, 489-506 (1985).

1182 269 Bishop, C. A. & Rouse, J. D. Polychlorinated biphenyls and organochlorine pesticides in
1183 plasma and the embryonic development in Lake Erie water snakes (*Nerodia sipedon*
1184 *insularum*) from Pelee Island, Ontario, Canada (1999). *Arch. Environ. Contam. Toxicol.*
1185 **51**, 452-457 (2006).

1186 270 Blackburn, D. C. A new squeaker frog (Arthroleptidae: Arthroleptis) from Bioko island,
1187 Equatorial Guinea. *Herpetologica* **66**, 320-334 (2010).

1188 271 Blondel, J., Perret, P., Anstett, M. C. & Thebaud, C. Evolution of sexual size dimorphism
1189 in birds: test of hypotheses using blue tits in contrasted Mediterranean habitats. *J. Evol.*
1190 *Biol.* **15**, 440-450 (2002).

1191 272 Boback, S. M. A morphometric comparison of island and mainland boas (*Boa*
1192 *constrictor*) in Belize. *Copeia* **2006**, 261-267 (2006).

1193 273 Bosc, V. Inventaire du Phyllodactyle d'Europe *Euleptes europaea* sur l'îlot de Roscana
1194 (Corse du Sud). 1-18 (Conservatoire des Espaces Naturels de Corse, 2009).

1195 274 Boye, P. Notes on the morphology, ecology and geographic origin of the Cyprus long-
1196 eared hedgehog (*Hemiechinus auritus dorotheae*). *Bonn. Zool. Beitr.* **42**, 115-123 (1991).

1197 275 Braithwaite, L. & Miller, B. The mallard, *Anas platyrhynchos*, and mallard-black duck,
1198 *Anas superciliosa rogersi*, hybridization. *Wildl. Res.* **2**, 47-61 (1975).

1199 276 Brandon-Jones, D. The taxonomic affinities of the Mentawai Islands Sureli, *Presbytis*
1200 *potenziani* (Bonaparte, 1856)(Mammalia: Primata: Cercopithecidae). *Raffles Bull. Zool.*
1201 **41**, 331-357 (1993).

1202 277 Brasileiro, C. A., Haddad, C. F., Sawaya, R. J. & Sazima, I. A new and threatened island-
1203 dwelling species of *Cycloramphus* (Anura: Cycloramphidae) from southeastern Brazil.
1204 *Herpetologica* **63**, 501-510 (2007).

- 1205 278 Brecko, J. *et al.* Functional and ecological relevance of intraspecific variation in body
1206 size and shape in the lizard *Podarcis melisellensis* (Lacertidae). *Biol. J. Linn. Soc.* **94**,
1207 251-264 (2008).
- 1208 279 Brisbin Jr, I. L. & Lenarz, M. S. Morphological comparisons of insular and mainland
1209 populations of southeastern white-tailed deer. *J. Mammal.* **65**, 44-50 (1984).
- 1210 280 Bromham, L. & Cardillo, M. Primates follow the 'island rule': implications for
1211 interpreting *Homo floresiensis*. *Biol. Lett.* **3**, 398-400 (2007).
- 1212 281 Brown, J. L., Maan, M. E., Cummings, M. E. & Summers, K. Evidence for selection on
1213 coloration in a Panamanian poison frog: a coalescent-based approach. *J. Biogeogr.* **37**,
1214 891-901 (2010).
- 1215 282 Bruschi, S. *et al.* Comments on the status of the Sardinian-Corsican lacertid lizard
1216 *Podarcis tiliguerta*. *Proc. Calif. Acad. Sci.* **57**, 225-245 (2006).
- 1217 283 Buden, D. W. Morphological variation and distributional ecology of the giant
1218 Micronesian gecko (*Perochirus scutellatus*) of Kapingamarangi Atoll. *Pac. Sci.* **52**, 250-
1219 258 (1998).
- 1220 284 Burt, W. H. Descriptions of heretofore unknown mammals from islands in the Gulf of
1221 California, Mexico. *Transactions of the San Diego Society of Natural History* **16**, 161-
1222 185 (1932).
- 1223 285 Butynski, T. M., de Jong, Y. A. & Hearn, G. W. Body measurements for the monkeys of
1224 Bioko Island, Equatorial Guinea. *Primate Conservation* **24**, 99-105 (2009).
- 1225 286 Cabot, J. & Urdiales, C. The subspecific status of Sardinian warblers *Sylvia*
1226 *melanocephala* in the Canary Islands with the description of a new subspecies from
1227 Western Sahara. *Bull. Br. Ornithol. Club* **125**, 230-240 (2005).
- 1228 287 Camerano, L. Ricerche intorno alla variazione del *Phyllodactylus europaeus* Gené. *Bull.*
1229 *Mus Zool. Anat. comp. R. Univ Torino* **19** 1-28 (1904).
- 1230 288 Camps, D. in *Enciclopedia Virtual de los Vertebrados Españoles* (ed A. Salvador,
1231 Barja, I.) (Museo Nacional de Ciencias Naturales, 2017).
- 1232 289 Capula, M. *et al.* Thermal ecology of the aeolian wall lizard, *Podarcis raffonei*. what
1233 about body temperatures in microinsular lizards. *Scripta Herpetologica. Studies on*
1234 *Amphibians and Reptiles in honour of Benedetto Lanza. Monografie della Societas*
1235 *Herpetologica Italica* **3**, 39-47 (2014).
- 1236 290 Carbone, M. *Caratteristiche della popolazione di Testudo hermanni Gmelin del Parco*
1237 *Naturale della Maremma*. MSc thesis, Università degli studi di Genova, (1988).
- 1238 291 Carrascal, L. M., Moreno, E. & Valido, A. Morphological evolution and changes in
1239 foraging behaviour of island and mainland populations of Blue Tit (*Parus caeruleus*)—a
1240 test of convergence and ecomorphological hypotheses. *Evol. Ecol.* **8**, 25-35 (1994).
- 1241 292 Carretero, M. & Llorente, G. Morphometry in a community of Mediterranean lacertid
1242 lizards, and its ecological relationships. *Hist. Anim.* **2**, 77-99 (1993).
- 1243 293 Case, T. J., Cody, M. L. & Ezcurra, E. *A new island biogeography of the Sea of Cortés*.
1244 (Oxford University Press on Demand, 2002).
- 1245 294 Case, T. J. & Schwaner, T. D. Island/mainland body size differences in Australian
1246 varanid lizards. *Oecologia* **94**, 102-109 (1993).
- 1247 295 Castellano, S., Rosso, A., Doglio, S. & Giacoma, C. Body size and calling variation in
1248 the green toad (*Bufo viridis*). *J. Zool.* **248**, 83-90 (1999).
- 1249 296 Castilla, A. M. & Bauwens, D. Reproductive characteristics of the lacertid lizard
1250 *Podarcis atrata*. *Copeia* **2000**, 748-756 (2000).

- 1251 297 Catzefflis, F., Maddalena, T., Hellwing, S. & Vogel, P. Unexpected findings on the
1252 taxonomic status of East Mediterranean *Crocidura russula* auct. (Mammalia,
1253 Insectivora). *Zeitschrift für Säugetierkunde* **50**, 185-201 (1985).
- 1254 298 Celis-Diez, J. L. *et al.* Population abundance, natural history, and habitat use by the
1255 arboreal marsupial *Dromiciops gliroides* in rural Chiloé Island, Chile. *J. Mammal.* **93**,
1256 134-148 (2012).
- 1257 299 Ceríaco, L. M. *et al.* Description of a new endemic species of shrew (Mammalia,
1258 Soricomorpha) from Príncipe Island (Gulf of Guinea). *Mammalia* **79**, 325-341 (2015).
- 1259 300 Chamberlain, J. The Block Island meadow mouse, *Microtus proVectus*. *J. Mammal.* **35**,
1260 587-589 (1954).
- 1261 301 Charles-Dominique, P. Urine marking and territoriality in *Galago alleni* (Waterhouse,
1262 1837—Lorisoidea, Primates)—a field study by radio-telemetry. *Z. Tierpsychol.* **43**, 113-
1263 138 (1977).
- 1264 302 Chasen, F. & Kloss, C. B. On a collection of mammals from the lowlands and islands of
1265 North Borneo. *Bull. Raffles Mus.* **6**, 1-82 (1931).
- 1266 303 Chen, S.-L. *et al.* Taxonomic status of the Korean populations of the genus *Scincella*
1267 (Squamata: Scincidae). *J. Herpetol.* **35**, 122-129 (2001).
- 1268 304 Cherem, J. J., Olimpio, J. & Ximenez, A. Descrição de uma nova espécie do gênero
1269 *Cavia* Pallas, 1766 (Mammalia-Caviidae) das Ilhas dos Moleques do Sul, Santa Catarina,
1270 Sul do Brasil. *Biotemas* **12**, 95-117 (1999).
- 1271 305 Chondropoulos, B. P. & Lykakis, J. J. Ecology of the Balkan wall lizard, *Podarcis*
1272 *taurica ionica* (Sauria: Lacertidae) from Greece. *Copeia*, 991-1001 (1983).
- 1273 306 Clark, R. J. Herpetofauna of the islands of the Argo-Saronic Gulf, Greece. *Proc. Calif.*
1274 *Acad. Sci.* **35**, 23-36 (1967).
- 1275 307 Clegg, S. M. *et al.* Microevolution in island forms: the roles of drift and directional
1276 selection in morphological divergence of a passerine bird. *Evolution* **56**, 2090-2099
1277 (2002).
- 1278 308 Clover, R. C. Phenetic relationships among populations of *Podarcis sicula* and *P.*
1279 *melisellensis* (Sauria: Lacertidae) from islands in the Adriatic Sea. *Syst. Biol.* **28**, 284-298
1280 (1979).
- 1281 309 Cole, C. J., Dessauer, H. C., Townsend, C. R. & Arnold, M. G. Unisexual lizards of the
1282 genus *Gymnophthalmus* (Reptilia, Teiidae) in the Neotropics: genetics, origin, and
1283 systematics. American Museum novitates; no. 2994. (1990).
- 1284 310 Colyn, M. Données pondérales sur les primates Cercopithecidae d’Afrique Centrale
1285 (Bassin du Zaïre/Congo). *Mammalia* **58**, 483-488 (1994).
- 1286 311 Copley, P., Read, V., Robinson, A. & Watts, C. Preliminary studies of the Nuyts
1287 Archipelago bandicoot *Isoodon obesulus nauticus* on the Franklin Islands, South
1288 Australia. *Bandicoots and Bilbies’*. (Eds JH Seebeck, PR Brown, RL Wallis and CM
1289 Kemper.) pp, 345-356 (1990).
- 1290 312 Cordero Rivera, A., Velo-Antón, G. & Galán, P. Ecology of amphibians in small coastal
1291 Holocene islands: local adaptations and the effect of exotic tree plantations. *Munibe* **25**,
1292 94-103 (2007).
- 1293 313 Corti, C. & Zuffi, M. Aspects of population ecology of *Testudo hermanni hermanni* from
1294 Asinara Island, NW Sardinia (Italy, Western Mediterranean Sea): preliminary data.
1295 *Amphibia-Reptilia* **24**, 441-447 (2003).

1296 314 Cowan, I. M. Insularity in the genus *Sorex* on the north coast of British Columbia. *Proc.*
1297 *Biol. Soc. Wash.* **54**, 95-107 (1941).

1298 315 Cox, D. T. *et al.* Patterns of seasonal and yearly mass variation in West African tropical
1299 savannah birds. *Ibis* **153**, 672-683 (2011).

1300 316 Creer, S., Chou, W. H., Malhotra, A. & Thorpe, R. S. Offshore insular variation in the
1301 diet of the Taiwanese bamboo viper *Trimeresurus stejnegeri* (Schmidt). *Zool. Sci.* **19**,
1302 907-913 (2002).

1303 317 Cruz-Elizalde, R. *et al.* Sexual dimorphism and natural history of the Western Mexico
1304 Whiptail, *Aspidoscelis costata* (Squamata: Teiidae), from Isla Isabel, Nayarit, Mexico.
1305 *North-Western Journal of Zoology* **10**, 374-381 (2014).

1306 318 Cumbo, V., Licata, F., Mercurio, E., Anz, S. & Lo Valvo, M. in *VIII Congresso*
1307 *Nazionale Societas Herpetologica Italica*. 401-404 (Ianieri Edizioni).

1308 319 D'Angelo, S., Galia, F. & Lo Valvo, M. Biometric characterization of two Sicilian pond
1309 turtle (*Emys trinacris*) populations of south-western Sicily. *Revista Española de*
1310 *Herpetología* **22**, 15-22 (2008).

1311 320 Dagosto, M., Gebo, D. L. & Dolino, C. Positional behavior and social organization of the
1312 Philippine tarsier (*Tarsius syrichta*). *Primates* **42**, 233-243 (2001).

1313 321 Darevsky, I. Two new species of the worm-like lizard *Dibamus* (Sauria, Dibamidae),
1314 with remarks on the distribution and ecology of *Dibamus* in Vietnam. *Asiatic*
1315 *Herpetological Research* **4**, 1-12 (1992).

1316 322 David, P., Vidal, N. & Pauwels, O. S. A morphological study of Stejneger's pitviper
1317 *Trimeresurus stejnegeri* (Serpentes, Viperidae, Crotalinae), with the description of a new
1318 species from Thailand. *Russian Journal of Herpetology* **8**, 205-222 (2001).

1319 323 De la Cruz, J. O. & Casas Jr, E. V. Captive Observations and Comparative Morphology
1320 of Philippine Tarsier (*Carlito syrichta*) in Brgy. Hugpa, Biliran, Biliran: A Preliminary
1321 Study. *Philippine Journal of Natural Sciences* **20**, 46-54 (2015).

1322 324 Decker, D. Systematics of the coatis, genus *Nasua* (Mammalia: Procyonidae). *Proc. Biol.*
1323 *Soc. Wash.* **104**, 370-386 (1991).

1324 325 Delany, M. Variation in the long-tailed field-mouse (*Apodemus sylvaticus* (L.)) in north-
1325 west Scotland I. Comparisons of individual characters. *Proceedings of the Royal Society*
1326 *of London. Series B. Biological Sciences* **161**, 191-199 (1964).

1327 326 Delany, M. & Healy, M. Variation in the white-toothed shrews (*Crocidura spp.*) in the
1328 British Isles. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **164**, 63-74 (1966).

1329 327 Delany, M. & Healy, M. Variation in the long-tailed field-mouse (*Apodemus sylvaticus*
1330 (L.)) in the Channel Islands. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **166**, 408-421 (1967).

1331 328 Delaugerre, M. La variation géographique chez *Phyllodactylus europaeus* Gené (Reptilia,
1332 Sauria, Gekkonidae). Etude de la population de l'îlot Sperduto Grande (Sud de la Corse,
1333 réserve naturelle des Iles Lavezzi). *Publications de la Société Linnéenne de Lyon* **54**,
1334 262-269 (1985).

1335 329 Delaugerre, M. Le Phyllodactyle d'Europe sur l'île de Port-Cros: mise en place d'un suivi
1336 géographique; la population de la Gabinière. 38 (Parc national de Port-Cros, Hyères,
1337 2003).

1338 330 Delaugerre, M. Archipel de la Galite «Notes herpétologiques 2008» Petites îles de
1339 Méditerranée 08. (2009).

1340 331 Delaugerre, M. & Dubois, A. La variation géographique et la variabilité
1341 intrapopulationnelle chez *Phyllodactylus europaeus* (Reptilia, Sauria, Gekkonidae).

- 1342 *Bulletin du Museum National d'Histoire Naturelle Section A Zoologie Biologie et*
1343 *Ecologie Animales* **7**, 709-736 (1985).
- 1344 332 Delibes de Castro, M. Sobre las Ginetas de la Isla de Ibiza (*Genetta genetta isabellae* n.
1345 ssp.). *Doñana. Acta Vertebrata* **4**, 139-160 (1977).
- 1346 333 Delibes, M. & Amores, F. The Stone Marten *Martes foina* (Erxleben, 1777)(Mammalia,
1347 Carnivora) from Ibiza (Pitiusic, Balearic Islands). *Miscellanea Zoologica (Spain)* (1986).
- 1348 334 Delson, E., Terranova, C. J., Jungers, W. L., Sargis, E. J. & Jablonski, N. G. Body mass
1349 in Cercopithecidae (Primates, Mammalia): estimation and scaling in extinct and extant
1350 taxa. *Anthropological papers of the AMNH; no. 83.* (2000).
- 1351 335 Dietzen, C., Garcia-del-Rey, E., Castro, G. D. & Wink, M. Phylogeography of the blue tit
1352 (*Parus teneriffae*-group) on the Canary Islands based on mitochondrial DNA sequence
1353 data and morphometrics. *J. Ornithol.* **149**, 1-12 (2008).
- 1354 336 Djong, H. T., Matsui, M., Kuramoto, M., Nishioka, M. & Sumida, M. A new species of
1355 the *Fejervarya limnocharis* complex from Japan (Anura, Dicroglossidae). *Zool. Sci.* **28**,
1356 922-929 (2011).
- 1357 337 Dodd Jr, C. K. Population structure and the evolution of sexual size dimorphism and sex
1358 ratios in an insular population of Florida box turtles (*Terrapene carolina bauri*).
1359 *Canadian Journal of Zoology* **75**, 1495-1507 (1997).
- 1360 338 Donihue, C. M., Brock, K. M., Foufopoulos, J. & Herrel, A. Feed or fight: testing the
1361 impact of food availability and intraspecific aggression on the functional ecology of an
1362 island lizard. *Funct. Ecol.* **30**, 566-575 (2016).
- 1363 339 Driessen, M. M. & Rose, R. K. *Isoodon obesulus* (Peramelemorphia: Peramelidae).
1364 *Mammalian Species* **47**, 112-123 (2015).
- 1365 340 Du, W.-G., Ji, X. & Zhang, Y.-P. Inter-population variation in life-history traits of a
1366 Chinese lizard (*Takydromus septentrionalis*, Lacertidae). *The Herpetological Journal* **16**,
1367 233-237 (2006).
- 1368 341 Du, W.-G., Ji, X., Zhang, Y.-P., Xu, X.-F. & Shine, R. Identifying sources of variation in
1369 reproductive and life-history traits among five populations of a Chinese lizard
1370 (*Takydromus septentrionalis*, Lacertidae). *Biol. J. Linn. Soc.* **85**, 443-453 (2005).
- 1371 342 Dumbell, G. S. *The ecology, behaviour and management of New Zealand Brown Teal, or*
1372 *Pateke (Anas aucklandica chlorotis)* PhD thesis, University of Auckland, (1987).
- 1373 343 Dunham, A. E., Tinkle, D. W. & Gibbons, J. W. Body size in island lizards: a cautionary
1374 tale. *Ecology* **59**, 1230-1238 (1978).
- 1375 344 Dunn, E. R. & Saxe Jr, L. Results of the Catherwood-Chaplin West Indies Expedition,
1376 1948. Part V. Amphibians and reptiles of San Andrés and Providencia. *Proc. Acad. Nat.*
1377 *Sci. Phila.*, 141-165 (1950).
- 1378 345 Dunstone, N. *et al.* Uso del hábitat, actividad y dieta de la güiña (*Oncifelis guigna*) en el
1379 Parque Nacional Laguna San Rafael, XI Región, Chile. *Boletín del Museo Nacional de*
1380 *Historia Natural (Chile)* **51**, 147-158 (2002).
- 1381 346 Elton, S. & Morgan, B. J. Muzzle size, paranasal swelling size and body mass in
1382 *Mandrillus leucophaeus*. *Primates* **47**, 151-157 (2006).
- 1383 347 Esteves, R. G. *Comportamento alimentar e aspectos reprodutivos de Bothrops insularis*
1384 *(Amaral, 1921)(Serpentes: Viperidae), na criação ex-situ no Instituto Vital Brazil* MSc
1385 thesis, Universidade Federal do Estado do Rio de Janeiro, (2016).
- 1386 348 Faaborg, J. & Winters, J. E. Winter resident returns and longevity and weights of Puerto
1387 Rican birds. *Bird-Banding*, 216-223 (1979).

- 1388 349 Fang, Y.-P. & Lee, L.-L. Re-evaluation of the Taiwanese white-toothed shrew, *Crocidura*
1389 *tadae* Tokuda and Kano, 1936 (Insectivora: Soricidae) from Taiwan and two offshore
1390 islands. *J. Zool.* **257**, 145-154 (2002).
- 1391 350 Faraone, F. P. *Indagini sulla variazione fenotipica in ambienti insulari e microinsulari*
1392 *delle specie di Podarcis Wagler, 1820 (Reptilia, Lacertidae) presenti in Sicilia e in*
1393 *alcune isole circumsiciliane*. PhD thesis, Università degli studi di Palermo, (2011).
- 1394 351 Faraone, F. P., Giacalone, G. & Lo Valvo, M. in *Atti VIII Congresso Nazionale Societas*
1395 *Herpetologica Italica*. 247-252.
- 1396 352 Fitch, H. S. Variation in clutch and litter size in New World reptiles. *Univ Kans Mus Nat*
1397 *Hist Misc Publ* **76**, 1-76 (1985).
- 1398 353 Fooden, J. Systematic review of the rhesus macaques, *Macaca mulatta* (Zimmermann,
1399 1780). *Fieldiana Zool* **96**, 180 (2000).
- 1400 354 Fooden, J. & Albrecht, G. H. Latitudinal and insular variation of skull size in
1401 crab-eating macaques (Primates, Cercopithecidae: *Macaca fascicularis*). *Am. J. Phys.*
1402 *Anthropol.* **92**, 521-538 (1993).
- 1403 355 Ford, H. The honeyeaters of Kangaroo Island. *South Australian Ornithologist* **27**, 134–
1404 138 (1976).
- 1405 356 Fornasiero, S., Corti, C., Luiselli, L. & Zuffi, M. A. Sexual size dimorphism,
1406 morphometry and phenotypic variation in the Whip snake *Hierophis viridiflavus* from a
1407 central Mediterranean area. *Revue d'écologie* (2007).
- 1408 357 Forsman, A. Variation in sexual size dimorphism and maximum body size among adder
1409 populations: effects of prey size. *J. Anim. Ecol.*, 253-267 (1991).
- 1410 358 Forsman, A., Merila, J. & Ebenhard, T. Phenotypic evolution of dispersal-enhancing
1411 traits in insular voles. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* **278**, 225-232 (2011).
- 1412 359 Foster, J. *The evolution of the native land mammals of the Queen Charlotte Islands and*
1413 *the problem of insularity*. PhD thesis, University of British Columbia, (1963).
- 1414 360 Fox, N. C. Some morphological data on the Australasian Harrier (*Circus approximans*
1415 *gouldi*) in New Zealand. *Notornis* **24**, 9-19 (1977).
- 1416 361 Franco, M., Quijano, A. & Soto-Gamboa, M. Communal nesting, activity patterns, and
1417 population characteristics in the near-threatened monito del monte, *Dromiciops gliroides*.
1418 *J. Mammal.* **92**, 994-1004 (2011).
- 1419 362 Fukada, H. Growth and maturity of the Japanese rat snake, *Elaphe climacophora*
1420 (Reptilia, Serpentes, Colubridae). *J. Herpetol.*, 269-274 (1978).
- 1421 363 Galán, P. Reproductive characteristics of an insular population of the lizard *Podarcis*
1422 *hispanica* from Northwest Spain (Cies Islands, Galicia). *Copeia* **2003**, 657-665 (2003).
- 1423 364 Ganem, G., Granjon, L., Ba, K. & Duplantier, J. M. Body size variability and water
1424 balance - A comparison between mainland and island populations of *Mastomys huberti*
1425 (Rodentia, Muridae) in Senegal. *Experientia* **51**, 402-410 (1995).
- 1426 365 Garcés-Restrepo, M. F., Giraldo, A. & Carr, J. L. Population ecology and morphometric
1427 variation of the chocoan river turtle (*Rhinoclemmys nasuta*) from two localities on the
1428 Colombian Pacific coast. *Boletín Científico. Centro de Museos. Museo de Historia*
1429 *Natural* **17**, 160-171 (2013).
- 1430 366 Garcia-Porta, J., Smid, J., Sol, D., Fasola, M. & Carranza, S. Testing the island effect on
1431 phenotypic diversification: insights from the Hemidactylus geckos of the Socotra
1432 Archipelago. *Sci. Rep.* **6** (2016).

- 1433 367 Gaulke, M. Overview on the present Knowledge on *Varanus mabitang* Gaulke and Curio,
1434 2001, Including new Morphological and Meristic Data. *Biawak* **4**, 50-58 (2010).
- 1435 368 Geissmann, T. *Evolution of communication in gibbons (Hylobatidae)*. PhD thesis,
1436 Universität Zürich, (1993).
- 1437 369 Giacalone, G., Abbate, M., Fritz, U. & Lo Valvo, M. Preliminary data on distribution,
1438 morphometric and genetic characterization of Hermann's tortoise in Sicily. *Herpetologia*
1439 *Sardiniae*, 282-286 (2008).
- 1440 370 Giagia, E. V. *Cytotaxonomical study of eastern European hedgehog Erinaceus concolor*
1441 *M. in Greece* PhD thesis, University of Patras, (1977).
- 1442 371 Gil Escobedo, L. J. *Ámbitos de Hogar de la Iguana de Órgano Ctenosaura palearis*
1443 *(Sauria: Iguanidae) en el Bosque Tropical Estacionalmente Seco de Cabañas, Zacapa,*
1444 *Guatemala* BSc thesis, Universidade de San Carlos de Guatemala, (2016).
- 1445 372 Gill, B. & McLean, I. G. Morphometrics of the whitehead *Mohoua albicilla* on Little
1446 Barrier Island, New Zealand. *N. Z. J. Zool.* **13**, 267-271 (1986).
- 1447 373 Glaw, F., Hawlitschek, O., Glaw, K. & Vences, M. Integrative evidence confirms new
1448 endemic island frogs and transmarine dispersal of amphibians between Madagascar and
1449 Mayotte (Comoros archipelago). *Science of Nature* **106** (2019).
- 1450 374 Glaw, F., Koehler, J., Townsend, T. M. & Vences, M. Rivaling the World's Smallest
1451 Reptiles: Discovery of Miniaturized and Microendemic New Species of Leaf Chameleons
1452 (Brookesia) from Northern Madagascar. *Plos One* **7** (2012).
- 1453 375 Glaw, F., Nagy, Z. T., Franzen, M. & Vences, M. Molecular phylogeny and systematics
1454 of the pseudoxyrhophiine snake genus Liopholidophis (Reptilia, Colubridae): evolution
1455 of its exceptional sexual dimorphism and descriptions of new taxa. *Zool. Scr.* **36**, 291-300
1456 (2007).
- 1457 376 Glaw, F., Vences, M., Andreone, F. & Vallan, D. Revision of the *Boophis majori* group
1458 (Amphibia : Mantellidae) from Madagascar, with descriptions of five new species. *Zool.*
1459 *J. Linn. Soc.* **133**, 495-529 (2001).
- 1460 377 Goldberg, S. R., Bursey, C. R. & Arreola, J. Gastrointestinal helminths from eight species
1461 of *Aspidoscelis* (Squamata: Teiidae) from Mexico. *West. N. Am. Nat.*, 223-227 (2014).
- 1462 378 Goldman, E. A. & Jackson, H. H. Raccoons of North and Middle America. *North*
1463 *American Fauna*, 1-153 (1950).
- 1464 379 Goltsman, M., Kruchenkova, E. P., Sergeev, S., Volodin, I. & Macdonald, D. W. 'Island
1465 syndrome' in a population of Arctic foxes (*Alopex lagopus*) from Mednyi Island. *J. Zool.*
1466 **267**, 405-418 (2005).
- 1467 380 González Quintero, E. P. *Análisis taxonómico del coyote (Canis latrans) de la Península*
1468 *de Baja California, México*. MSc thesis, Centro de Investigaciones Biológicas del
1469 Noroeste, S.C, (2004).
- 1470 381 González Rossell, A. *Ecología y conservación de la iguana (Cyclura nubila nubila) en*
1471 *Cuba*. PhD thesis, Universidad de Alicante, (2018).
- 1472 382 Granjon, L. & Cheylan, G. Biometric differentiation of black rat (*Rattus rattus*)
1473 populations in the West Mediterranean islands. *Mammalia* **54**, 213-231 (1990).
- 1474 383 Grant, P. *A systematic study of the terrestrial birds of the Tres Marias Islands, Mexico.*
1475 (Peabody Museum of Natural History, Yale University, 1965).
- 1476 384 Grant, P. The coexistence of two wren species of the genus *Thryothorus*. *The Wilson*
1477 *Bulletin*, 266-278 (1966).

- 1478 385 Grant, P. *Further information on the relative length of the tarsus in land birds*. (Yale
1479 Peabody Museum of Natural History, 1966).
- 1480 386 Grant, P. Ecological and morphological variation of Canary Island blue tits, *Parus*
1481 *caeruleus* (Aves: Paridae). *Biol. J. Linn. Soc.* **11**, 103-129 (1979).
- 1482 387 Grinnell, J. The species of the mammalian genus *Sorex* of West-Central California.
1483 *University of California Publications in Zoology* **10**, 179-195 (1913).
- 1484 388 Grinnell, J., Dixon, J. S. & Linsdale, J. M. *The fur bearing mammals of California.*,
1485 (University of California Press, 1937).
- 1486 389 Guarino, F. M., Garcia, G. & Andreone, F. Huge but moderately long-lived: age structure
1487 in the mountain chicken, *Leptodactylus fallax*, from Montserrat, West Indies.
1488 *Herpetological Journal* **24**, 167-173 (2014).
- 1489 390 Gursky, S. Effects of radio transmitter weight on a small nocturnal primate. *Am. J.*
1490 *Primatol.* **46**, 145-155 (1998).
- 1491 391 Gutsche, A. & Streich, W. J. Demography and Endangerment of the Utila Island Spiny-
1492 Tailed Iguana, *Ctenosaura bakeri*. *J. Herpetol.* **43**, 105-113 (2009).
- 1493 392 Hadi, S., Ziegler, T. & Hodges, J. K. Group structure and physical characteristics of
1494 Simakobu monkeys (*Simias concolor*) on the Mentawai Island of Siberut, Indonesia.
1495 *Folia Primatol.* **80**, 74-82 (2009).
- 1496 393 Hagen, B. Zur Kleinsäugerfauna Siziliens. *Bonn. Zool. Beitr.* **5**, 1-15 (1954).
- 1497 394 Hai, B. T., Tu, L. N., Duong, V. T. & Son, N. T. in *HỘI NGHỊ KHOA HỌC TOÀN*
1498 *QUỐC VỀ SINH THÁI VÀ TÀI NGUYÊN SINH VẬT LẦN THỨ 7 - National Scientific*
1499 *Conference on Ecological and Biological Resources.* 670-676.
- 1500 395 Harding, L. E. *Trachypithecus cristatus* (Primates: Cercopithecidae). *Mammalian Species*
1501 **42**, 149-165 (2010).
- 1502 396 Hasegawa, M. Insular radiation in life-history of the lizard *Eumeces okadae* in the Izu
1503 islands, Japan. *Copeia*, 732-747 (1994).
- 1504 397 Hasegawa, M. & Moriguchi, H. Geographic variation in food habits, body size and life
1505 history traits of the snakes on the Izu Islands, p. 414-432. *Current herpetology in East*
1506 *Asia. M. Matui, T. Hikida, and RC Goris (eds.). Herpetological Society of Japan,*
1507 *Tokyo.[Google Scholar]* (1989).
- 1508 398 Hawlitschek, O., Nagy, Z. T. & Glaw, F. Island evolution and systematic revision of
1509 Comoran snakes: why and when subspecies still make sense. *PLoS One* **7** (2012).
- 1510 399 Heaney, L. R. Island area and body size of insular mammals: evidence from the
1511 tri-colored squirrel (*Callosciurus prevosti*) of Southeast Asia. *Evolution* **32**, 29-44
1512 (1978).
- 1513 400 Heaney, L. R. & Timm, R. M. Systematics and distribution of shrews of the genus
1514 *Crociodura* (Mammalia: Insectivora) in Vietnam. (1983).
- 1515 401 Hedges, S. B. & Thomas, R. At the lower size limit in amniote vertebrates: a new
1516 diminutive lizard from the West Indies. *Caribb. J. Sci.* **37**, 168-173 (2001).
- 1517 402 Heinsohn, G. E. Ecology and reproduction of the Tasmanian bandicoots (*Perameles*
1518 *gunni* and *Isoodon obesulus*). (1966).
- 1519 403 Hemelaar, A. Age, growth and other population characteristics of *Bufo bufo* from
1520 different latitudes and altitudes. *J. Herpetol.*, 369-388 (1988).
- 1521 404 Heo, J.-H. *et al.* Can an invasive prey species induce morphological and behavioral
1522 changes in an endemic predator? Evidence from a South Korean snake (*Oocatochus*
1523 *rufodorsatus*). *Asian Herpetological Research* **5**, 245-254 (2014).

- 1524 405 Hernández Gallegos, O., López Moreno, A. E., Méndez de la Cruz, F. R. & Walker, J.
1525 Home range of the parthenogenetic lizard *Aspidoscelis maslini* (Fritts, 1969), on a beach
1526 strand. (2018).
- 1527 406 Hernandez-Gallegos, O., Esthela Lopez-Moreno, A., Fernando Mendez-Sanchez, J.,
1528 Lloyd Rheubert, J. & Roberto Mendez-de la Cruz, F. Home range of *Aspidoscelis*
1529 *cozumela* (Squamata: Teiidae): a parthenogenetic lizard microendemic to Cozumel
1530 Island, Mexico. *Rev. Biol. Trop.* **63**, 771-781 (2015).
- 1531 407 Hernandez-Salinas, U., Ramirez-Bautista, A., Pavon, N. P. & Rosas Pacheco, L. F.
1532 Morphometric variation in island and mainland populations of two lizard species from the
1533 Pacific Coast of Mexico. *Rev. Chil. Hist. Nat.* **87**, 1-9 (2014).
- 1534 408 Herrel, A., Cottam, M. D., Godbeer, K., Sanger, T. & Losos, J. B. An ecomorphological
1535 analysis of native and introduced populations of the endemic lizard *Anolis maynardi* of
1536 the Cayman Islands. *Breviora* **522**, 1-10 (2011).
- 1537 409 Herrel, A., Spithoven, L., Van Damme, R. & De Vree, F. Sexual dimorphism of head size
1538 in *Gallotia galloti*: testing the niche divergence hypothesis by functional analyses. *Funct.*
1539 *Ecol.* **13**, 289-297 (1999).
- 1540 410 Hershkovitz, P. Mammals of Northern Colombia, Preliminary Report No. 6: Rabbits
1541 (Leporidae), with Notes on the Classification and Distribution of the South American
1542 Forms. *Proc. U. S. Natl. Mus.* (1950).
- 1543 411 Hervias-Parejo, S. *et al.* Small size does not restrain frugivory and seed dispersal across
1544 the evolutionary radiation of Galapagos lava lizards. *Current Zoology* **65**, 353-361
1545 (2019).
- 1546 412 Heyer, M. M., Heyer, W. R. & de Sá, R. O. *Leptodactylus pentadactylus*. *Catalogue of*
1547 *American Amphibians and Reptiles (CAAR)* (2012).
- 1548 413 Heyer, W., Rand, A., Cruz, C., Peixoto, O. L. & Nelson, C. E. Frogs of Boracéia.
1549 *Arquivos de Zoologia (Museu de Zoologia da Universidade Sao Paulo)* **31**, 231-410
1550 (1990).
- 1551 414 Higgins, P. J. *Handbook of Australian, New Zealand and Antarctic birds. Volume 4:*
1552 *Parrots to Dollarbird.*, Vol. 4 (Oxford University Press, 1999).
- 1553 415 Higgins, P. J. & Davies, S. J. J. F. *Handbook of Australian, New Zealand and Antarctic*
1554 *Birds. Volume 3: Snipe to Pigeons.* Vol. 3 (Oxford University Press, 1996).
- 1555 416 Higgins, P. J. & Peter, J. M. *Handbook of Australian, New Zealand and Antarctic Birds.*
1556 *Volume 6: Pardalotes to Shrike-thrushes.* Vol. 6 (Oxford University Press, 2002).
- 1557 417 Higgins, P. J., Peter, J. M. & Cowling, S. J. *Handbook of Australian, New Zealand and*
1558 *Antarctic Birds. Volume 7: Boatbill to Starlings.* Vol. 7 (Oxford University Press, 2006).
- 1559 418 Higgins, P. J., Peter, J. M. & Steele, W. K. *Handbook of Australian, New Zealand and*
1560 *Antarctic Birds. Volume 5: Tyrant-flycatchers to Chats.* Vol. 5 (Oxford University Press,
1561 2001).
- 1562 419 Hill, J. The Robinson collection of Malaysian mammals. *Bull. Raffles Mus.* **29**, 6-22
1563 (1960).
- 1564 420 Hitchmough, R. A. & McCallum, J. The mammals, birds, reptiles and freshwater fish of
1565 the eastern island group of the Bay of Islands. *Tane* **26**, 127-134 (1980).
- 1566 421 Hofmeyr, M. D. Egg production in *Chersina angulata*: an unusual pattern in a
1567 Mediterranean climate. *J. Herpetol.* **38**, 172-179 (2004).
- 1568 422 Hollingsworth, B. D. The systematics of chuckwalla (*Sauromalus*) with a phylogenetic
1569 analysis of other iguanid lizards. *Herpetol. Monogr.*, 38-191 (1998).

- 1570 423 How, R. A., Cowan, M. A., Teale, R. J. & Schmitt, L. H. Environmental correlates of
1571 reptile variation on the Houtman Abrolhos archipelago, eastern Indian Ocean. *J.*
1572 *Biogeogr.* (2020).
- 1573 424 Huang, W.-S. Ecology and reproductive patterns of the agamid lizard *Japalura swinhonis*
1574 on an east Asian island, with comments on the small clutch sizes of island lizards. *Zool.*
1575 *Sci.* **24**, 181-188 (2007).
- 1576 425 Hummelinck, P. W. *Studies on the fauna of Curaçao, Aruba, Bonaire, and the*
1577 *Venezuelan Islands: No. 6.* (1940).
- 1578 426 Hutterer, R. Variation and evolution of the Sicilian shrew: taxonomic conclusions and
1579 description of a possibly related species from the Pleistocene of Morocco (Mammalia:
1580 Soricidae). *Bonn. Zool. Beitr.* **42**, 241-251 (1991).
- 1581 427 Huyghe, K., Vanhooydonck, B., Herrel, A., Tadić, Z. & Van Damme, R. Morphology,
1582 performance, behavior and ecology of three color morphs in males of the lizard *Podarcis*
1583 *melisellensis*. *Integr. Comp. Biol.* **47**, 211-220 (2007).
- 1584 428 Iliopoulou-Georgoudaki, J. The relationship between climatic factors and forearm length
1585 of bats: evidence from the chiropterofauna of Lesvos island (Greece–East Aegean).
1586 *Mammalia* **50**, 475-482 (1986).
- 1587 429 Inger, R. F., Stuart, B. L. & Iskandar, D. T. Systematics of a widespread southeast Asian
1588 frog, *Rana chalconota* (Amphibia: Anura: Ranidae). *Zool. J. Linn. Soc.* **155**, 123-147
1589 (2009).
- 1590 430 Iverson, J. B., Hines, K. N. & Valiulis, J. M. The nesting ecology of the Allen Cays rock
1591 iguana, *Cyclura cychlura inornata* in the Bahamas. *Herpetol. Monogr.* **18**, 1-36 (2004).
- 1592 431 Jacobs, D. Morphological divergence in an insular bat, *Lasiurus cinereus semotus*. *Funct.*
1593 *Ecol.*, 622-630 (1996).
- 1594 432 Jenkins, P. & Veitch, C. Sexual dimorphism and age determination in the North Island
1595 saddleback (*Philesturnus carunculatus rufaster*). *N. Z. J. Zool.* **18**, 445-450 (1991).
- 1596 433 Jenkins, P. D., Abramov, A. V., Bannikova, A. A. & Rozhnov, V. V. Bones and genes:
1597 resolution problems in three Vietnamese species of Crocidura (Mammalia,
1598 Soricomorpha, Soricidae) and the description of an additional new species. *ZooKeys*, 61
1599 (2013).
- 1600 434 Jenssen, T. A. *et al.* Morphological characteristics of the lizard *Anolis carolinensis* from
1601 South Carolina. *Herpetologica*, 401-411 (1995).
- 1602 435 Jessop, R. Biometrics and moult of red-capped plovers in Victoria, Tasmania, South
1603 Australia and North West Australia. *Stilt* **17** 29-35 (1990).
- 1604 436 Jessop, T. S. *et al.* Maximum body size among insular Komodo dragon populations
1605 covaries with large prey density. *Oikos* **112**, 422-429 (2006).
- 1606 437 Ji, X. & Wang, Z. W. Geographic variation in reproductive traits and trade-offs between
1607 size and number of eggs of the Chinese cobra (*Naja atra*). *Biol. J. Linn. Soc.* **85**, 27-40
1608 (2005).
- 1609 438 Ji, X., Xie, Y. Y., Sun, P. Y. & Zheng, X. Z. Sexual dimorphism and female reproduction
1610 in a viviparous snake, *Elaphe rufodorsata*. *J. Herpetol.* **31**, 420-422 (1997).
- 1611 439 Jimenez, J. E. & McMahon, E. in *Canids: foxes, wolves, jackals, and dogs: status survey*
1612 *and conservation action plan* (eds C. Sillero-Zubiri, M. Hoffmann, & D.W Macdonald)
1613 (IUCN, 2004).

- 1614 440 Joubert, L. & Cheylan, M. La tortue d'Hermann de Corse: résultat des recherches menées
1615 en 1985 et 1986. *Travaux scientifiques—Parc naturel régional et réserves naturelles de*
1616 *Corse*, 1-54 (1989).
- 1617 441 Judd, F. W. & Ross, R. K. Year-to-year variation in clutch size of island and mainland
1618 populations of *Holbrookia propinqua* (Reptilia, Lacertilia, Iguanidae). . *J. Herpetol.* **12**,
1619 203-207 (1978).
- 1620 442 Kang, C. *et al.* Differential predation drives the geographical divergence in multiple traits
1621 in aposematic frogs. *Behav. Ecol.* **28**, 1122-1130 (2017).
- 1622 443 Keast, A. Competitive interactions and the evolution of ecological niches as illustrated by
1623 the Australian honeyeater genus *Meliphaga* (Meliphagidae). *Evolution*, 762-784
1624 (1968).
- 1625 444 Keehn, J. E., Nieto, N. C., Tracy, C. R., Gienger, C. M. & Feldman, C. R. Evolution on a
1626 desert island: body size divergence between the reptiles of Nevada's Anaho Island and the
1627 mainland around Pyramid Lake. *J. Zool.* **291**, 269-278 (2013).
- 1628 445 Kim, D.-I. *et al.* Patterns of morphological variation in the Schlegel's Japanese gecko
1629 (*Gekko japonicus*) across populations in China, Japan, and Korea. *Journal of Ecology*
1630 *and Environment* **43**, 34 (2019).
- 1631 446 Kim, S.-I., Oshida, T., Lee, H., Min, M.-S. & Kimura, J. Evolutionary and
1632 biogeographical implications of variation in skull morphology of raccoon dogs
1633 (*Nyctereutes procyonoides*, Mammalia: Carnivora). *Biol. J. Linn. Soc.* **116**, 856-872
1634 (2015).
- 1635 447 King, R. B. Body size variation among island and mainland snake populations.
1636 *Herpetologica* **45**, 84-88 (1989).
- 1637 448 King, R. B. Variation in brown snake (*Storeria dekayi*) morphology and scalation: Sex,
1638 family, and microgeographic differences. *J. Herpetol.* **31**, 335-346 (1997).
- 1639 449 King, T. The birds of the Lesio-Louna and Lefini Reserves, Batéké Plateau, Republic of
1640 Congo. *Malimbus* **33**, 1-41 (2011).
- 1641 450 Kitchener, D., Hisheh, S., Schmitt, L. & Maryanto, I. Morphological and genetic
1642 variation in *Aethalops alecto* (Chiroptera, Pteropodidae) from Java, Bali and Lombok Is,
1643 Indonesia. *Mammalia* **57**, 255-272 (1993).
- 1644 451 Kitchener, D. & Schmitt, L. Morphological and genetic variation in *Suncus murinus*
1645 (Soricidae: Crocidurinae) from Java, Lesser Sunda islands, Maluku and Sulawesi,
1646 Indonesia. *Mammalia* **58**, 433-452 (1994).
- 1647 452 Kitchener, D., Schmitt, L., Hisheh, S., How, R. & Cooper, N. Morphological and genetic
1648 variation in the bearded tomb bats (Taphozous: Emballonuridae) of Nusa Tenggara,
1649 Indonesia. *Mammalia* **57**, 63-84 (1993).
- 1650 453 Klauber, L. M. The gopher snakes of Baja California: with descriptions of a new
1651 subspecies of *Pituophis catenifer*. *Transactions of the San Diego Society of Natural*
1652 *History* **11**, 1-40 (1946).
- 1653 454 Klauber, L. M. A new gopher snake (*Pituophis*) from Santa Cruz Island, California
1654 *Transactions of the San Diego Society of Natural History* **11**, 41-48 (1946).
- 1655 455 Klauber, L. M. Some new and revived subspecies of rattlesnakes. *Transactions of the San*
1656 *Diego Society of Natural History* **11**, 61-116 (1949).
- 1657 456 Klauber, L. M. *Rattlesnakes: their habits, life histories, and influence on mankind.*
1658 (University of California Press, 1997).
- 1659 457 Kloss, C. B. On mammals collected in Siam. *J Nat Hist Soc Siam* **4**, 333-407 (1919).

- 1660 458 Klütsch, C., Misof, B., Grosse, W. R. & Moritz, R. Genetic and morphometric
1661 differentiation among island populations of two *Norops* lizards (Reptilia: Sauria:
1662 Polychrotidae) on independently colonized islands of the Islas de Bahía (Honduras). *J.*
1663 *Biogeogr.* **34**, 1124-1135 (2007).
- 1664 459 Knapp, C. R., Iverson, J. B. & Owens, A. K. Geographic variation in nesting behavior
1665 and reproductive biology of an insular iguana (*Cyclura cychlura*). *Can. J. Zool./Rev. Can.*
1666 *Zool.* **84**, 1566-1575 (2006).
- 1667 460 Kohno, H. Reptiles in a seabird colony: herpetofauna of Nakanokamishima Island of the
1668 Yaeyama group, Ryukyu Archipelago. *Island Studies in Okinawa* **9**, 73-89 (1991).
- 1669 461 Koopman, K. F. & Diamond, J. Zoogeography of mammals from islands off the
1670 northeastern coast of New Guinea. *Am. Mus. Novit.* **2690** (1979).
- 1671 462 Kraus, C., Trillmich, F. & Künkele, J. Reproduction and growth in a precocial small
1672 mammal, *Cavia magna*. *J. Mammal.* **86**, 763-772 (2005).
- 1673 463 Proetz, D., Ruthensteiner, B., Scherz, M. D. & Glaw, F. Systematic revision of the
1674 Malagasy chameleons *Calumma boettgeri* and *C. linotum* (Squamata: Chamaeleonidae).
1675 *Zootaxa* **4048**, 211-231 (2015).
- 1676 464 Qualls, C. P., Shine, R., Donnellan, S. & Hutchinson, M. The evolution of viviparity
1677 within the Australian scincid lizard *Lerista bougainvillii*. *J. Zool.* **237**, 13-26 (1995).
- 1678 465 Quin, D. G., Smith, A. P. & Norton, T. W. Eco-geographic variation in size and sexual
1679 dimorphism in sugar gliders and squirrel gliders (Marsupialia: Petauridae). *Aust. J. Zool.*
1680 **44**, 19-45 (1996).
- 1681 466 Rader, J. A., Dillon, M. E., Chesson, R. T., Sabat, P. & del Rio, C. M. Morphological
1682 divergence in a continental adaptive radiation: South American ovenbirds of the genus
1683 *Cinclodes*. *Auk* **132**, 180-190 (2015).
- 1684 467 Radočaj, M., Jelić, D., Karaica, D. & Kapelj, S. Morphological and reproductive traits of
1685 the insular population of *Podarcis siculus* (Reptilia: Lacertidae) from Krk Island
1686 (Croatia) *Hyla* **2**, 5-22 (2011).
- 1687 468 Raia, P. *et al.* The blue lizard spandrel and the island syndrome. *BMC Evol. Biol.* **10**, 289
1688 (2010).
- 1689 469 Rajaratnam, R. *Ecology of leopard cat (Prionailurus bengalensis) in Tabin Wildlife*
1690 *Reserve, Sabah, Malaysia*. PhD thesis, Universiti Kebangsaan Malaysia (2000).
- 1691 470 Rausch, R. L. Geographic variation in size in North American brown bears, *Ursus arctos*
1692 L., as indicated by condylobasal length. *Can. J. Zool./Rev. Can. Zool.* **41**, 33-45 (1963).
- 1693 471 Reboucas, R., da Silva, H. R., Sanuy, D. & Sole, M. Sexual maturity and growth of male
1694 toads (*Rhinella ornata*): A comparison between insular and mainland populations. *Zool.*
1695 *Anz.* **283**, 12-19 (2019).
- 1696 472 Reboucas, R., da Silva, H. R. & Sole, M. Frog size on continental islands of the coast of
1697 Rio de Janeiro and the generality of the island rule. *PloS One* **13** (2018).
- 1698 473 Redfield, J. Distribution, abundance, size, and genetic variation of *Peromyscus*
1699 *maniculatus* on the Gulf Islands of British Columbia. *Can. J. Zool./Rev. Can. Zool.* **54**,
1700 463-474 (1976).
- 1701 474 Renaud, S. *et al.* Morphometrics and genetics highlight the complex history of Eastern
1702 Mediterranean spiny mice. *Biol. J. Linn. Soc.* (2020).
- 1703 475 Reynolds, R. G., Niemiller, M. L. & Fitzpatrick, B. M. Genetic analysis of an endemic
1704 archipelagic lizard reveals sympatric cryptic lineages and taxonomic discordance.
1705 *Conserv. Genet.* **13**, 953-963 (2012).

- 1706 476 Rhodin, A. G. Chelid turtles of the Australasian archipelago: a new species of *Chelodina*
1707 from Roti island, Indonesia. *Breviora* **498**, 1-31 (1994).
- 1708 477 Rickart, E. A., Heaney, L. R., Goodman, S. M. & Jansa, S. Review of the Philippine
1709 genera *Chrotomys* and *Celaenomys* (Murinae) and description of a new species. *J.*
1710 *Mammal.* **86**, 415-428 (2005).
- 1711 478 Ridgway, R. *The birds of North and Middle America: A descriptive catalogue of the*
1712 *higher groups, genera, species, and subspecies of birds known to occur in North*
1713 *America, from the Arctic lands to the Isthmus of Panama, the West Indies and other*
1714 *islands of the Caribbean sea, and the Galapagos Archipelago.* (US Government Printing
1715 Office, 1904).
- 1716 479 Rivas, G. A. *et al.* A distinctive new species of *Gonatodes* (Squamata:
1717 *Sphaerodactylidae*) from Isla La Blanquilla, Venezuela, with remarks on the distribution
1718 of some other Caribbean sphaerodactylid lizards. *South Am. J. Herpetol.* **8**, 5-18 (2013).
- 1719 480 Robertson, H., Whitaker, A. & Fitzgerald, B. Morphometrics of forest birds in the
1720 Orongorongo Valley, Wellington, New Zealand. *N. Z. J. Zool.* **10**, 87-97 (1983).
- 1721 481 Rocha, C. F. & Vrcibradic, D. Reproductive traits of two sympatric viviparous skinks
1722 (*Mabuya macrorhyncha* and *Mabuya agilis*) in a Brazilian restinga habitat.
1723 *Herpetological Journal* **9**, 43-53 (1999).
- 1724 482 Rodríguez-Cabal, M. A., Amico, G. C., Novaro, A. J. & Aizen, M. A. Population
1725 characteristics of *Dromiciops gliroides* (Philippi, 1893), an endemic marsupial of the
1726 temperate forest of Patagonia. *Mamm. Biol.* **73**, 74-76 (2008).
- 1727 483 Rog, S., Ryan, M. J., Mueller, U. & Lampert, K. P. Evidence for morphological and
1728 genetic diversification of túngara frog populations on islands. *Herpetol. Conserv. Biol.* **8**,
1729 228-239 (2013).
- 1730 484 Rogers, K., Rogers, A. & Rogers, D. Bander's Aid. *Supplement Number One. Royal*
1731 *Australasian Ornithologists' Union Report* (1990).
- 1732 485 Ross, R. K. & Judd, F. W. Comparison of lipid cycles of *Holbrookia propinqua* from
1733 Padre island and mainland Texas. *J. Herpetol.* **16**, 53-60 (1982).
- 1734 486 Rowe-Rowe, D. & Crafford, J. Density, body size, and reproduction of feral house mice
1735 on Gough Island. *S. Afr. J. Zool.* **27**, 1-5 (1992).
- 1736 487 Runemark, A., Sagonas, K. & Svensson, E. I. Ecological explanations to island
1737 gigantism: dietary niche divergence, predation, and size in an endemic lizard. *Ecology* **96**,
1738 2077-2092 (2015).
- 1739 488 Sabater Pi, J. Contribution to the ecology of *Colobus polykomos satanas* (Waterhouse,
1740 1838) of Rio Muni, Republic of Equatorial Guinea. *Folia Primatol.* **19**, 193-207 (1973).
- 1741 489 Sagonas, K., Pafilis, P. & Valakos, E. D. Effects of insularity on digestion: living on
1742 islands induces shifts in physiological and morphological traits in island reptiles. *The*
1743 *Science of Nature* **102**, 55 (2015).
- 1744 490 Sahimi, H. N. M., Chubo, J. K., Mohd, M., Saripuddin, N. B. & Ab Rahim, S. S. The
1745 distribution and population density of Bornean tarsier, *Tarsius bancanus borneanus*
1746 (Elliot) in secondary and rehabilitated forests of Universiti Putra Malaysia, Bintulu
1747 Sarawak Campus, Sarawak, Malaysia. *Tropical Life Sciences Research* **29**, 139 (2018).
- 1748 491 Salvidio, S., Cresta, P. & Dolmen, D. The common toad *Bufo bufo* population of Hitra
1749 island, central Norway. *Fauna Norvegica Series A* **14**, 51-55 (1993).

- 1750 492 Sanches, V. Q. A. & Grings, D. R. Daily movement and habitat use of *Iguana iguana*
1751 (Linnaeus, 1758) in an urban second growth Amazonian forest fragment in Brazil.
1752 *Herpetology Notes* **11**, 93-96 (2018).
- 1753 493 Sanderson, J., Sunquist, M. E. & Iriarte, A. W. Natural history and landscape-use of
1754 guignas (*Oncifelis guigna*) on Isla Grande de Chiloé, Chile. *J. Mammal.* **83**, 608-613
1755 (2002).
- 1756 494 Sargis, E. J., Millien, V., Woodman, N. & Olson, L. E. Rule reversal: Ecogeographical
1757 patterns of body size variation in the common treeshrew (Mammalia, Scandentia). *Ecol.*
1758 *Evol.* **8**, 1634-1645 (2018).
- 1759 495 Sá-Sousa, P., Almeida, A., Rosa, H., Vicente, L. & Crespo, E. Genetic and
1760 morphological relationships of the Berlenga wall lizard (*Podarcis bocagei berlengensis*:
1761 Lacertidae). *J. Zool. Syst. Evol. Res.* **38**, 95-102 (2000).
- 1762 496 Sá-Sousa, P. & Harris, D. J. *Podarcis carbonelli* Pérez-Mellado, 1981 is a distinct
1763 species. *Amphibia-Reptilia* **23**, 459-468 (2002).
- 1764 497 Sá-Sousa, P., Vicente, L. & Crespo, E. Morphological variability of *Podarcis hispanica*
1765 (Sauria: lacertidae) in Portugal. *Amphibia-Reptilia* **23**, 55-69 (2002).
- 1766 498 Scali, S. *et al.* in *11th Ordinary General Meeting Societas Europaea Herpetologica*
1767 (*SEH*). 161-166.
- 1768 499 Scheffer, V. B. & Dalquest, W. W. A new shrew from Destruction Island, Washington. *J.*
1769 *Mammal.* **23**, 333-335 (1942).
- 1770 500 Schifter, H., Cunningham-van Someren, G. & van Someren, G. C. The Avifauna of the
1771 North Nandi Forest, Kenya. *Annalen des Naturhistorischen Museums in Wien. Serie B für*
1772 *Botanik und Zoologie* **100**, 425-479 (1998).
- 1773 501 Schlotfeldt, B. E. & Kleindorfer, S. Adaptive divergence in the Superb Fairy-wren
1774 (*Malurus cyaneus*): a mainland versus island comparison of morphology and foraging
1775 behaviour. *Emu* **106**, 309-319 (2006).
- 1776 502 Schultz, A. H. Observations on the growth, classification and evolutionary specialization
1777 of gibbons and siamangs. *Human Biology* **5**, 212 (1933).
- 1778 503 Schultz, A. H. The relative weight of the testes in primates. *The Anatomical Record* **72**,
1779 387-394 (1938).
- 1780 504 Schwaner, T. D. & Sarre, S. D. Body size of tiger snakes in Southern Australia, with
1781 particular reference to *Notechis ater serventyi* (Elapidae) on Chappell Island. *J. Herpetol.*
1782 **22**, 24-33 (1988).
- 1783 505 Schwaner, T. D. & Sarre, S. D. Body size and sexual dimorphism in mainland and island
1784 tiger snakes. *J. Herpetol.* **24**, 320-322 (1990).
- 1785 506 Schwartz, A. Three new mammals from southern Florida. *J. Mammal.* **33**, 381-385
1786 (1952).
- 1787 507 Schwartz, A. Snakes of the genus *Alsophis* in Puerto Rico and the Virgin Islands. *Studies*
1788 *on the Fauna of Curaçao and other Caribbean Islands* **23**, 177-227 (1966).
- 1789 508 Scott, D. Notes on the Eastern Hognose Snake, *Heterodon platyrhinos* Latreille
1790 (Squamata, Colubridae), on a Virginia Barrier Island. *Brimleyana*, 51-55 (1986).
- 1791 509 Selcer, K. W. & Judd, F. W. Variation in the reproductive ecology of *Holbrookia*
1792 *propinqua* (Sauria, Iguanidae). *Texas Journal of Science* **34**, 125-135 (1982).
- 1793 510 Senczuk, G., García, A., Colangelo, P., Annesi, F. & Castiglia, R. Morphometric and
1794 genetic divergence in island and mainland populations of *Anolis nebulosus* (Squamata:

- 1795 Polychrotidae) from Jalisco (Mexico): an instance of insular gigantism. *Ital. J. Zool.* **81**,
1796 204-214 (2014).
- 1797 511 Seock, M., Nam, K.-B. & Yoo, J.-C. Distribution and movement tendencies of short-
1798 tailed viper snakes (*Gloydius saxatilis*) by altitude. *Asian Herpetological Research* **8**, 39-
1799 47 (2017).
- 1800 512 Serrano-Cardozo, V. H., Ramírez-Pinilla, M. P., Ortega, J. E. & Cortes, L. A. Annual
1801 reproductive activity of *Gonatodes albogularis* (Squamata: Gekkonidae) living in an
1802 anthropic area in Santander, Colombia. *South Am. J. Herpetol.* **2**, 31-38 (2007).
- 1803 513 Shaidani, N.-I. *The biogeographic origins and trophic ecology of Maine's island red-
1804 backed salamanders (Plethodon cinereus)* MSc thesis, University of Maine, (2017).
- 1805 514 Sharples, C. M., Fa, J. E. & BELL, D. J. Geographical variation in size in the European
1806 rabbit *Oryctolagus cuniculus* (Lagomorpha: Leporidae) in western Europe and North
1807 Africa. *Zool. J. Linn. Soc.* **117**, 141-158 (1996).
- 1808 515 Shekelle, M. Observations of wild Sangihe island tarsiers *Tarsius sangirensis*. *Asian
1809 Primates Journal* **3**, 18-23 (2013).
- 1810 516 Shimada, T., Maeda, S. & Sakakibara, M. A morphological study of *Cynops
1811 pyrrhogaster* from the Chita Peninsula: rediscovery of the "extinct" Atsumi race endemic
1812 to peninsular regions of Aichi prefecture, Central Japan. *Curr. Herp.* **35**, 38-52 (2016).
- 1813 517 Shine, R. Venomous snakes in cold climates: ecology of the Australian genus *Drysdalia*
1814 (Serpentes: Elapidae). *Copeia*, 14-25 (1981).
- 1815 518 Shine, R., Sun, L.-x., Zhao, E. & Bonnet, X. A review of 30 years of ecological research
1816 on the Shedao pitviper, *Gloydius shedaoensis*. *Herpetological Natural History* **9**, 1-14
1817 (2003).
- 1818 519 Sicuro, F. L. & Oliveira, L. F. B. Variations in leopard cat (*Prionailurus bengalensis*)
1819 skull morphology and body size: sexual and geographic influences. *PeerJ* **3**, e1309
1820 (2015).
- 1821 520 Siler, C. D., McVay, J. D., Diesmos, A. C. & Brown, R. M. A new species of fanged
1822 frog, genus *Limnonectes* (Amphibia: Anura: Dicroglossidae) from southeast Mindanao
1823 Island, Philippines. *Herpetologica* **65**, 105-114 (2009).
- 1824 521 Siliceo-Cantero, H. H., García, A., Reynolds, R. G., Pacheco, G. & Lister, B. C.
1825 Dimorphism and divergence in island and mainland Anoles. *Biol. J. Linn. Soc.* **118**, 852-
1826 872 (2016).
- 1827 522 Simberloff, D., Dayan, T., Jones, C. & Ogura, G. Character displacement and release in
1828 the small Indian mongoose, *Herpestes javanicus*. *Ecology* **81**, 2086-2099 (2000).
- 1829 523 Sinclair, E. Morphological variation among populations of the quokka, *Setonix
1830 brachyurus* (Macropodidae: Marsupialia), in Western Australia. *Aust. J. Zool.* **46**, 439-
1831 449 (1998).
- 1832 524 Siqueira, C. C., Van Sluys, M., Ariani, C. V. & Rocha, C. F. Feeding ecology of *Thoropa
1833 miliaris* (Anura, Cycloramphidae) in four areas of Atlantic rain forest, southeastern
1834 Brazil. *J. Herpetol.* **40**, 520-525 (2006).
- 1835 525 Slábová, M. & Frynta, D. Morphometric variation in nearly unstudied populations of the
1836 most studied mammal: The non-commensal house mouse (*Mus musculus domesticus*) in
1837 the Near East and Northern Africa. *Zoologischer Anzeiger-A Journal of Comparative
1838 Zoology* **246**, 91-101 (2007).
- 1839 526 Sleeman, D., Cussen, R., Southey, A. & O'Leary, D. The Badgers *Meles meles* (L.) of
1840 Coney Island, Co Sligo. *The Irish Naturalists' Journal*, 10-18 (2002).

- 1841 527 Sleeman, D. P., Davenport, J., Cussen, R. E. & Hammond, R. F. The small-bodied
1842 badgers (*Meles meles* (L.)) of Rutland Island, Co. Donegal. *The Irish Naturalists'*
1843 *Journal*, 1-6 (2009).
- 1844 528 Sleeman, P., Cussen, R., O'Donoghue, T. & Costello, E. Badgers (*Meles meles*) on
1845 Fenit island, and their presence or absence on other Islands in co. Kerry, Ireland. *Small*
1846 *carnivore conservation* **10** (2001).
- 1847 529 Smith, F. A. Evolution of body size among woodrats from Baja California, Mexico.
1848 *Funct. Ecol.*, 265-273 (1992).
- 1849 530 Smith, R. J. & Jungers, W. L. Body mass in comparative primatology. *J. Hum. Evol.* **32**,
1850 523-559 (1997).
- 1851 531 Smith, T. B. & Temple, S. A. Feeding habits and bill polymorphism in hook-billed kites.
1852 *The Auk* **99**, 197-207 (1982).
- 1853 532 Society, F. o. M.-L. I. (2008).
- 1854 533 Sody, H. Notes on some primates, carnivora, and the babirusa from the Indo-Malayan
1855 and Indo-Australian regions. *Treubia* **20**, 121-190 (1949).
- 1856 534 Song, L., Fa-Hong, Y. & Xue-Fei, L. Cranial morphometric study of four giant flying
1857 squirrels (Petaurista)(Rodentia: Sciuridae) from China. *Zool. Res.* **33**, 119-126 (2012).
- 1858 535 Sparkman, A. M. *et al.* Convergence in reduced body size, head size, and blood glucose
1859 in three island reptiles. *Ecol. Evol.* **8**, 6169-6182 (2018).
- 1860 536 Stager, K. E. The avifauna of the Tres Marias islands, Mexico. *The Auk* **74**, 413-432
1861 (1957).
- 1862 537 Stamenković, S. & Matić, R. Morphological correlates of prey consumed by *Podarcis*
1863 *melisellensis* (Braun, 1877) and *P. siculus* (Rafinesque, 1810)(Sauria, Lacertidae) from
1864 two mainland regions in the eastern Adriatic area. *Archives of Biological Sciences* **65**,
1865 1015-1025 (2013).
- 1866 538 Steven, D. M. Recent evolution in the genus *Clethrionomys*. *Symp. Soc. Exp. Biol.* **7**,
1867 310-319 (1953).
- 1868 539 Stoddart, D. M. & Braithwaite, R. W. A strategy for utilization of regenerating heathland
1869 habitat by the brown bandicoot (*Isodon obesulus*; Marsupialia, Peramelidae). *J. Anim.*
1870 *Ecol.*, 165-179 (1979).
- 1871 540 Storer, R. W. Variation in the red-tailed hawks of southern Mexico and Central America.
1872 *The Condor* **64**, 77-78 (1962).
- 1873 541 Strickland, D. & Norris, D. R. An example of phenotypic adherence to the island rule? -
1874 Anticosti gray jays are heavier but not structurally larger than mainland conspecifics.
1875 *Ecol. Evol.* **5**, 3687-3694 (2015).
- 1876 542 Struhsaker, T. T. *The red colobus monkeys: variation in demography, behavior, and*
1877 *ecology of endangered species*. (Oxford University Press, 2010).
- 1878 543 Stuart-Smith, J. F., Stuart-Smith, R. D., Swain, R. & Wapstra, E. Size dimorphism in
1879 *Rankinia Tympanocryptis diemensis* (Family Agamidae): sex-specific patterns and
1880 geographic variation. *Biol. J. Linn. Soc.* **94**, 699-709 (2008).
- 1881 544 Stubbs, D. & Swingland, I. R. The ecology of a Mediterranean tortoise (*Testudo*
1882 *hermanni*): a declining population. *Can. J. Zool./Rev. Can. Zool.* **63**, 169-180 (1985).
- 1883 545 Swarth, H. The lemming of Nunivak Island, Alaska. *Proc. Biol. Soc. Wash* **44**, 101-104
1884 (1931).
- 1885 546 Swarth, H. S. *Birds and mammals of the 1909 Alexander Alaska expedition*. Vol. 7 (The
1886 University Press, 1911).

- 1887 547 Takada, Y., Sakai, E., Uematsu, Y. & Tateishi, T. Morphometric variation of house mice
1888 (Mus musculus) on the Izu Islands. *Mammal Study* **24**, 51-65 (1999).
- 1889 548 Takada, Y., Sakai, E., Uematsu, Y. & Tateishi, T. Morphological variation of large
1890 Japanese field mice, *Apodemus speciosus* on the Izu and Oki Islands. *Mammal Study* **31**,
1891 29-40 (2006).
- 1892 549 Takada, Y., Yamada, H. & Tateishi, T. Morphometric variation of Japanese wild mice on
1893 islands. *Journal of the Mammalogical Society of Japan* **19**, 113-128 (1994).
- 1894 550 Takenaka, T. & Hasegawa, M. Female-biased mortality and its consequence on adult sex
1895 ratio in the freshwater turtle *Chinemys reevesii* on an island. *Curr. Herp.* **20**, 11-17
1896 (2001).
- 1897 551 Tamarin, R. H. Dispersal, population regulation, and K-selection in field mice. *The*
1898 *American Naturalist* **112**, 545-555 (1978).
- 1899 552 Tanaka, K. Phenotypic plasticity of body size in an insular population of a snake.
1900 *Herpetologica* **67**, 46-57 (2011).
- 1901 553 Tanaka, K. & Ota, H. Natural history of two colubrid snakes, *Elaphe quadrivirgata* and
1902 *Rhabdophis tigrinus*, on Yakushima Island, southwestern Japan. *Amphibia-Reptilia* **23**,
1903 323-331 (2002).
- 1904 554 Taverne, M. *et al.* Diet variability among insular populations of Podarcis lizards reveals
1905 diverse strategies to face resource-limited environments. *Ecol. Evol.* **9**, 12408-12420
1906 (2019).
- 1907 555 Taylor, E. H. The lizards of Thailand. *University of Kansas Science Bulletin* **44**, 687-
1908 1077 (1963).
- 1909 556 Taylor, H. L. & Cooley, C. R. A multivariate analysis of morphological variation among
1910 parthenogenetic teiid lizards of the *Cnemidophorus cozumela* complex. *Herpetologica*,
1911 67-76 (1995).
- 1912 557 Terada, C., Tatsuzawa, S. & Saitoh, T. Ecological correlates and determinants in the
1913 geographical variation of deer morphology. *Oecologia* **169**, 981-994 (2012).
- 1914 558 Terborgh, J., Faaborg, J. & Brockmann, H. J. Island colonization by Lesser Antillean
1915 birds. *The Auk* **95**, 59-72 (1978).
- 1916 559 Tessa, G., Crottini, A., Giacoma, C., Guarino, F. M. & Randrianirina, J. E. Comparative
1917 longevity and age at sexual maturity in twelve rainforest frogs of the genera Boophis,
1918 Gephyromantis, and Mantidactylus (Anura: Mantellidae) from Madagascar.
1919 *Phyllomedusa: Journal of Herpetology* **16**, 13-21 (2017).
- 1920 560 Thomas, D. & Broughton, E. *Status of three Canadian caribou populations north of 70 in*
1921 *winter 1977*. (Canadian Wildlife Service, 1978).
- 1922 561 Thouless, C. & Bassri, K. A. Taxonomic status of the Farasan Island gazelle. *J. Zool.*
1923 **223**, 151-159 (1991).
- 1924 562 Tsai, T. Sexual dimorphism of Chinese green tree viper, *Trimeresurus stejnegeri*
1925 *stejnegeri*. *Biol Bull Nat Taiwan Normal Univ* **33**, 13-22 (1998).
- 1926 563 Uller, T. *et al.* Genetic differentiation predicts body size divergence between island and
1927 mainland populations of common wall lizards (*Podarcis muralis*). *Biol. J. Linn. Soc.* **127**,
1928 771-786 (2019).
- 1929 564 Ursin, E. *Geographical variation in Apodemus sylvaticus and A. flavicollis (Rodentia,*
1930 *Muridae) in Europe, with special reference to Danish and Latvian populations.*
1931 (Munksgaard, 1956).

- 1932 565 Valakos, E. D. & Polymeni, R. M. The food of *Cyrtodactylus kotschy* (Steindachner,
1933 1870) (Sauria - Gekkonidae) during the wet season in the Mediterranean insular
1934 ecosystems of the Aegean. *Herpetological Journal* **1**, 474-477 (1990).
- 1935 566 Van Heezik, Y. M., Cooper, J. & Seddon, P. J. Population characteristics and
1936 morphometrics of angulate tortoises on Dassen Island, South Africa. *J. Herpetol.*, 447-
1937 453 (1994).
- 1938 567 Van Weenen, J. *Aspects of the ecology of Cunningham's skink, Egernia cunninghami, on*
1939 *West Island, South Australia*. BSc (Honours) thesis, University of Adelaide, (1995).
- 1940 568 Vanek, J. P. & Burke, R. L. Insular dwarfism in female eastern hog-nosed snakes
1941 (*Heterodon platirhinos*; Dipsadidae) on a barrier island. *Can. J. Zool./Rev. Can. Zool.* **98**,
1942 157-164 (2020).
- 1943 569 Vanzolini, P. E. & Reboucas-Spieker, R. Distribution and differentiation of animals along
1944 the coast and in continental islands of the state of São Paulo, Brazil. 3. Reproductive
1945 differences between *Mabuya caissara* and *Mabuya macrorhyncha* (Sauria, Scincidae).
1946 *Papéis Avulsos de Zoologia, São Paulo* **29**, 95-109 (1976).
- 1947 570 Vasconcelos, R. & Carranza, S. Systematics and biogeography of *Hemidactylus*
1948 *homoeolepis* Blanford, 1881 (Squamata: Gekkonidae), with the description of a new
1949 species from Arabia. *Zootaxa* **3835**, 501-527 (2014).
- 1950 571 Velo-Anton, G. & Cordero-Rivera, A. Ethological and phenotypic divergence in insular
1951 fire salamanders: diurnal activity mediated by predation? *Acta Ethologica* **20**, 243-253
1952 (2017).
- 1953 572 Vences, M. Erste untersuchungen über die fortpflanzungsbiologie des Iberischen
1954 Scheibenzünglers *Discoglossus galganoi* (Amphibia: Anura: Discoglossidae). *Acta*
1955 *Biologica Benrodis* **6**, 89-98 (1994).
- 1956 573 Vences, M., Glaw, F., Mercurio, V. & Andreone, F. Review of the Malagasy tree snakes
1957 of the genus *Stenophis* (Colubridae). *Salamandra* **40**, 161-179 (2004).
- 1958 574 Vences, M., Köhler, J., Pabijan, M. & Glaw, F. Two syntopic and microendemic new
1959 frogs of the genus *Blommersia* from the east coast of Madagascar. *African Journal of*
1960 *Herpetology* **59**, 133-156 (2010).
- 1961 575 Ventura, J. & Fuster, M. J. L. Morphometric analysis of the black rat, *Rattus rattus*, from
1962 Congreso Island (Chafarinas Archipelago, Spain). *Orsis: organismos i sistemes* **15**, 91-
1963 102 (2000).
- 1964 576 Verdon, E. & Donnelly, M. A. Population structure of Florida box turtles (*Terrapene*
1965 *carolina bauri*) at the southernmost limit of their range. *J. Herpetol.* **39**, 572-577 (2005).
- 1966 577 Vershinin, A. The biology and trapping of the ermine in Kamchatka. *Byull. Mosk. o-va*
1967 *Ispyt. Prir. Otd. Biol* **77**, 16-26 (1972).
- 1968 578 Vervust, B., Grbac, I. & Van Damme, R. Differences in morphology, performance and
1969 behaviour between recently diverged populations of *Podarcis sicula* mirror differences in
1970 predation pressure. *Oikos* **116**, 1343-1352 (2007).
- 1971 579 Villeneuve, A. R. Habitat selection and population density of the world's smallest
1972 chameleon, *Brookesia micra*, on Nosy Hara, Madagascar. *Herpetol. Conserv. Biol.* **12**,
1973 334-341 (2017).
- 1974 580 Vogel, P. & Sofianidou, T. The shrews of the genus *Crocidura* on Lesbos, an eastern
1975 Mediterranean island. *Bonn. Zool. Beitr.* **46**, 339-347 (1996).
- 1976 581 Vogrin, M. Sexual Dimorphism in *Podarcis sicula campestris*. *Turkish Journal of*
1977 *Zoology* **29**, 189-191 (2005).

1978 582 Vrcibradic, D. & Rocha, C. F. Observations on the natural history of the lizard *Mabuya*
1979 *macrorhyncha* Hoge (Scincidae) in Queimada Grande island, Sao Paulo, Brazil. *Rev.*
1980 *Bras. Zool.* **22**, 1185-1190 (2005).

1981 583 Walker, R., Woods-Ballard, A. J. & Rix, C. E. Population density and seasonal activity of
1982 the threatened Madagascar spider tortoise (*Pyxis arachnoides arachnoides*) of the
1983 southern dry forests; South West Madagascar. *Afr. J. Ecol.* **46**, 67 (2008).

1984 584 Wang, Y., Li, Y., Wu, Z. & Murray, B. Insular shifts and trade-offs in life-history traits
1985 in pond frogs in the Zhoushan Archipelago, China. *J. Zool.* **278**, 65-73 (2009).

1986 585 Watkins, G. G. Proximate causes of sexual size dimorphism in the iguanian lizard
1987 *Microlophus occipitalis*. *Ecology* **77**, 1473-1482 (1996).

1988 586 Wayne, R. K. *et al.* A morphologic and genetic study of the island fox, *Urocyon*
1989 *littoralis*. *Evolution* **45**, 1849-1868 (1991).

1990 587 West, A. G. *Variation in mandible shape and body size of house mice Mus musculus*
1991 *across the New Zealand archipelago: A Trans-Tasman comparison using geometric*
1992 *morphometrics* MSc thesis, University of Waikato, (2017).

1993 588 Whitehead, V. *Population dynamics of the eastern box turtle (Terrapene carolina*
1994 *carolina) in the Maryville College Woods, Maryville, Tennessee: a report of a senior*
1995 *study*, Maryville College, (2017).

1996 589 Woodman, N. *et al.* A new southern distributional limit for the Central American rodent
1997 *Peromyscus stirtoni*. *Caribb. J. Sci.* **38**, 281-284 (2002).

1998 590 Wu, Z., Li, Y. & Murray, B. R. Insular shifts in body size of rice frogs in the Zhoushan
1999 Archipelago, China. *J. Anim. Ecol.* **75**, 1071-1080 (2006).

2000 591 Xu, F., Adler, G. H. & Li, Y. Covariation in insular life-history traits of the rice frog
2001 (*Fejervarya limnocharis*) in Eastern China. *Asian Herpetological Research* **4**, 28-35
2002 (2013).

2003 592 Yabe, T. Population structure and male melanism in the Reeves' turtle, *Chinemys*
2004 *reevesii*. *Japanese Journal of Herpetology* **15**, 131-137 (1994).

2005 593 Yasukawa, Y., Ota, H. & Iverson, J. B. Geographic variation and sexual size dimorphism
2006 in *Mauremys mutica* (Cantor, 1842)(Reptilia: Bataguridae), with description of a new
2007 subspecies from the southern Ryukyus, Japan. *Zool. Sci.* **13**, 303-317 (1996).

2008 594 Yurgenson, P. B. Ermines of the Far East territory. *Byull. Mosk. o-va Ispyt. Prir. Otd.*
2009 *Biol* **45**, 239-243 (1936).

2010 595 Yusefi, G. H., Kiabi, B. H., Khalatbari, L., Faizolahi, K. & Monteiro, N. M.
2011 Morphological analysis of Brandt's hedgehog (*Paraechinus hypomelas*) reflects the
2012 isolation history of Persian Gulf islands and has implications for taxonomy. *Biol. J. Linn.*
2013 *Soc.* **119**, 497-510 (2016).

2014 596 Yustian, I. *Ecology and conservation status of Tarsius bancanus saltator on Belitung*
2015 *Island, Indonesia.* (Cuvillier Verlag, 2007).

2016 597 Ziegler, T., Vences, M., Glaw, F. & Böhme, W. Remarks on the genital morphology of
2017 the Malagasy snake genus *Liophidium* (Reptilia, Serpentes, Colubridae). *Acta biologica*
2018 *Benrodis* **8** (1996).

2019 598 Zihlman, A. L., Mootnick, A. R. & Underwood, C. E. Anatomical contributions to
2020 hylobatid taxonomy and adaptation. *Int. J. Primatol.* **32**, 865-877 (2011).

2021 599 Zuffi, M. & Gariboldi, A. Sexual dimorphism of the European pond terrapin, *Emys*
2022 *orbicularis* (L., 1758) from Italy. *Scientia herpetologica* **1995**, 124-129 (1995).

2023 600 Zuffi, M. A., Sacchi, R., Pupin, F. & Cencetti, T. Sexual size and shape dimorphism in
2024 the Moorish gecko (*Tarentola mauritanica*, Gekkota, Phyllodactylidae). *North-Western*
2025 *Journal of Zoology* **7**, 189-197 (2011).

2026 601 Zuffi, M. A. L., Odetti, F., Batistoni, R. & Mancino, G. Geographic variation of sexual
2027 size dimorphism and genetics in the European pond turtle, *Emys orbicularis* and *Emys*
2028 *trinacris*, of Italy. *Ital. J. Zool.* **73**, 363-372 (2006).

2029 602 Zug, G. R., Hedges, S. B. & Sunkel, S. Variation in reproductive parameters of three
2030 neotropical snakes, *Coniophanes fissidens*, *Dipsas catesbyi*, and *Imantodes cenchoa*.
2031 *Smithson. Contrib. Zool.* **300** (1979).

2032 603 Higgins, J. P. *et al.* *Cochrane handbook for systematic reviews of interventions*. (John
2033 Wiley & Sons, 2019).

2034 604 Santini, L., Benítez-López, A., Ficetola, G. F. & Huijbregts, M. A. Length–mass
2035 allometries in amphibians. *Integr. Zool.* **13**, 36-45 (2018).

2036 605 Lislevand, T., Figuerola, J. & Székely, T. Avian body sizes in relation to fecundity,
2037 mating system, display behavior, and resource sharing. *Ecology* **88**, 1605-1605 (2007).

2038 606 Van Valkenburgh, B. in *Body size in mammalian paleobiology: estimation and biological*
2039 *implications* (eds J. Damuth & Bruce J. MacFadden) Ch. 10, 181-206 (Cambridge
2040 University Press, 1990).

2041 607 Jones, K. E. *et al.* PanTHERIA: a species-level database of life history, ecology, and
2042 geography of extant and recently extinct mammals. *Ecology* **90**, 2648-2648 (2009).

2043 608 Scharf, I. *et al.* Late bloomers and baby boomers: ecological drivers of longevity in
2044 squamates and the tuatara. *Global Ecol. Biogeogr.* **24**, 396-405 (2015).

2045 609 Regis, K. W. & Meik, J. M. Allometry of sexual size dimorphism in turtles: a comparison
2046 of mass and length data. *PeerJ* **5**, e2914 (2017).

2047 610 Gosler, A., Greenwood, J., Baker, J. & Davidson, N. The field determination of body size
2048 and condition in passerines: a report to the British Ringing Committee. *Bird Study* **45**, 92-
2049 103 (1998).

2050 611 Rising, J. D. & Somers, K. M. The measurement of overall body size in birds. *The Auk*
2051 **106**, 666-674 (1989).

2052 612 Jetz, W., Thomas, G., Joy, J., Hartmann, K. & Mooers, A. The global diversity of birds in
2053 space and time. *Nature* **491**, 444-448 (2012).

2054 613 Wan, X., Wang, W., Liu, J. & Tong, T. Estimating the sample mean and standard
2055 deviation from the sample size, median, range and/or interquartile range. *BMC Med. Res.*
2056 *Methodol.* **14**, 135 (2014).

2057 614 Bracken, M. in *Effective care of the newborn infant* (ed J.C. Sinclair and M.B. Bracken)
2058 13-20 (Oxford University Press, 1992).

2059 615 Pettorelli, N. *et al.* The Normalized Difference Vegetation Index (NDVI): unforeseen
2060 successes in animal ecology. *Clim. Res.* **46**, 15-27 (2011).

2061 616 Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1°km spatial resolution climate surfaces
2062 for global land areas. *Int. J. Climatol.* **37**, 4302-4315 (2017).

2063 617 Blackburn, T. M., Gaston, K. J. & Loder, N. Geographic gradients in body size: a
2064 clarification of Bergmann's rule. *Divers. Distrib.* **5**, 165-174 (1999).

2065 618 Wilman, H. *et al.* EltonTraits 1.0: Species-level foraging attributes of the world's birds
2066 and mammals: Ecological Archives E095-178. *Ecology* **95**, 2027-2027 (2014).

2067 619 Meiri, S. Traits of lizards of the world: Variation around a successful evolutionary
2068 design. *Global Ecol. Biogeogr.* **27**, 1168-1172 (2018).

2069 620 Benítez-López, A. *et al.* The impact of hunting on tropical mammal and bird populations.
2070 *Science* **356**, 180-183 (2017).
2071 621 Solé, M. & Rödder, D. in *Amphibian Ecology and Conservation: a handbook of*
2072 *techniques*. (ed C. Kenneth Dodd Jr) 167-184 (Oxford University Press, 2010).
2073 622 McClain, C. R., Durst, P. A. P., Boyer, A. G. & Francis, C. D. Unravelling the
2074 determinants of insular body size shifts. *Biol. Lett.* **9**, 5pp.-5pp. (2013).
2075 623 Rozzi, R. Space-time patterns of body size variation in island bovids: The key role of
2076 predatory release. *J. Biogeogr.* **45**, 1196-1207 (2018).
2077 624 Hadfield, J. & Nakagawa, S. General quantitative genetic methods for comparative
2078 biology: phylogenies, taxonomies and multi-trait models for continuous and categorical
2079 characters. *J. Evol. Biol.* **23**, 494-508 (2010).
2080 625 Lajeunesse, M. J. On the meta-analysis of response ratios for studies with correlated and
2081 multi-group designs. *Ecology* **92**, 2049-2055 (2011).
2082 626 McClain, C. R., Durst, P. A. P., Boyer, A. G. & Francis, C. D. Unravelling the
2083 determinants of insular body size shifts. *Biol. Lett.* **9** (2013).
2084 627 Lindstedt, S. L. & Boyce, M. S. Seasonality, fasting endurance, and body size in
2085 mammals. *The American Naturalist* **125**, 873-878 (1985).
2086 628 Nakagawa, S. & Schielzeth, H. A general and simple method for obtaining R² from
2087 generalized linear mixed-effects models. *Methods Ecol. Evol.* **4**, 133-142 (2013).
2088 629 R-Core-Team. *R: A language and environment for statistical computing.*, (R Foundation
2089 for Statistical Computing, 2018).
2090 630 Viechtbauer, W. Conducting meta-analyses in R with the metafor package. *J. Stat. Softw.*
2091 **36**, 1-48 (2010).
2092 631 Lajeunesse, M. J. Facilitating systematic reviews, data extraction and meta-analysis with
2093 the metagear package for R. *Methods Ecol. Evol.* **7**, 323-330 (2016).
2094 632 Pick, J. L., Nakagawa, S. & Noble, D. W. Reproducible, flexible and high-throughput
2095 data extraction from primary literature: The metaDigitise r package. *Methods Ecol. Evol.*
2096 **10**, 426-431 (2019).
2097 633 Paradis, E., Claude, J. & Strimmer, K. APE: analyses of phylogenetics and evolution in R
2098 language. *Bioinformatics* **20**, 289-290 (2004).
2099 634 Michonneau, F., Brown, J. W. & Winter, D. J. rotl: an R package to interact with the
2100 Open Tree of Life data. *Methods Ecol. Evol.* **7**, 1476-1481 (2016).
2101 635 Rees, J. A. & Cranston, K. Automated assembly of a reference taxonomy for
2102 phylogenetic data synthesis. *Biodivers. Data J.* (2017).
2103 636 Hinchliff, C. E. *et al.* Synthesis of phylogeny and taxonomy into a comprehensive tree of
2104 life. *Proc. Natl. Acad. Sci. USA* **112**, 12764-12769 (2015).
2105 637 Pebesma, E. Simple features for R: standardized support for spatial vector data. *The R*
2106 *Journal* **10**, 439-446 (2018).
2107 638 Hijmans, R. J. *et al.* Package 'raster'. *R package* (2015).
2108 639 Wickham, H., Francois, R., Henry, L. & Müller, K. dplyr: A Grammar of Data
2109 Manipulation. R package version 0.4. 3. *R Found. Stat. Comput., Vienna.* [https://CRAN.](https://CRAN.R-project.org/package=dplyr)
2110 *R-project.org/package=dplyr* (2015).
2111 640 Wickham, H. reshape2: Flexibly reshape data: a reboot of the reshape package. *R*
2112 *package version 1* (2012).
2113 641 Wickham, H. *ggplot2: elegant graphics for data analysis.* (Springer, 2016).

2114 642 Kassambara, A. ggpubr:“ggplot2” based publication ready plots. R package version 0.1.
2115 6. (2017).

2116

2117 **Acknowledgements**

2118 We are grateful to K. B. Aubry, J. E. Keehn, S. Michaelides and D. Strickland for sharing their
2119 data with us, S. Meiri for pointing out valuable sources of measurement data, and to A. Sánchez-
2120 Tojar, P. Peres-Neto and S. Nakagawa for useful discussion on the analytical framework. ABL
2121 was supported by a Juan de la Cierva-Incorporación grant (IJCI-2017-31419) from the Spanish
2122 Ministry of Science, Innovation and Universities. LS and MAJH were supported by the ERC
2123 project (62002139 ERC – CoG SIZE 647224). We thank numerous biological collections, in
2124 particular the Natural History Museum, Tring, for providing access to specimens. Bird trait data
2125 collection was supported by Natural Environment Research Council grant nos. NE/I028068/1
2126 and NE/P004512/1 (to JAT).

2127 **Author Contributions**

2128 AB-L conceived and coordinated the research, led the analyses and wrote the first draft; AB-L,
2129 LS, JG-Z, MAJH and JAT helped to develop the conceptual framework; LS compiled the
2130 environmental rasters; JAT, PW and BM provided morphometric data. All authors contributed to
2131 the data collection from the literature and to the writing of the final manuscript.

2132 **Competing interests**

2133 The authors declare no competing interests.

2134

Figures and Tables

Figure 1. Conceptual figure showing body size evolution in island populations. According to the island rule, changes in body size of island populations are dependent on the body mass of mainland relatives, with small species tending to increase in size on islands (gigantism) and large species tending to decrease in size (dwarfism). By plotting the log response ratio ($\ln RR$) between insular mass and mainland mass, against mainland mass, we can test if insular populations adhere to the rule (intercept > 0 and slope < 0) (blue line). Mechanisms proposed to drive ‘island rule’ effects are mainly based on reduced predation, inter- and intra-specific competition, and food availability, suggesting that the relationship will steepen in small, remote islands (red line).

Figure 2. Location of island populations included in our analyses for mammals (N = 1058, blue), birds (N = 695, orange), reptiles (N = 547, yellow), and amphibians (N = 179, green). The size of each point indicates the number of species sampled on each island; some points overlap. See Fig. S1 for a 4-panel figure with the location of insular populations separated for each taxonomic group.

Figure 3. Relationship between $\ln RR$ (log response ratio between island mass and mainland body mass) and body mass in the mainland for (a) mammals (N = 1058), (b) birds (N = 695), (c) reptiles (N = 547) and (d) amphibians (N = 179). Models were fitted using phylogenetic multi-level meta-regression models with mainland body mass as moderator, and observation-level ID, source ID, species ID and phylogeny as random effects. $\ln RR > 0$ indicates gigantism; $\ln RR < 0$ indicates dwarfism; and $\ln RR = 0$ indicates stasis (no shift in body size from mainland to island populations). The size of points represents the inverse of the sampling variance for each paired island-mainland response ratio in the model. Shaded areas represent 95% confidence intervals. Note that y-axes have different scales.

Figure 4. The effect of island area and spatial isolation on insular size shifts in terrestrial vertebrates for (a) mammals (N = 1058), (b) birds (N = 695), (c) reptiles (N = 547) and (d) amphibians (N = 179). Continuous variables are represented at the 10% and 90% quantile for each extreme (close vs remote islands; small vs large islands). $\ln RR > 0$ indicates gigantism; $\ln RR < 0$ indicates dwarfism; and $\ln RR = 0$ indicates stasis (no shift in body size from mainland to island populations). Shaded areas represent 95% confidence intervals.

Table 1. Parameter estimates for the phylogenetic meta-regression models testing the generality of the island rule in terrestrial vertebrates. k : number of island-mainland comparisons ($\ln RR$), Q_m : test of moderators ($\log_{10}(\text{mainland mass})$). R^2_m : marginal R^2 , estimated percentage of heterogeneity explained by the moderator (fixed effects). R^2_c : conditional R^2 , percentage of heterogeneity attributable to fixed and random effects.

Class	k	Intercept (CI)	Slope (CI)	Q_m (p-value)	R^2_m	R^2_c
Mammals	1058	0.208 (0.052 – 0.365)	-0.088 (-0.122 – -0.055)	27.30 ($p < 0.001$)	11.4	56.4
Birds	695	0.216 (0.117 – 0.315)	-0.104 (-0.145 – -0.064)	25.40 ($p < 0.001$)	7.0	43.4
Reptiles	547	0.410 (0.006 – 0.814)	-0.305 (-0.419 – -0.190)	27.21 ($p < 0.001$)	17.6	66.6
Amphibians	179	0.195 (0.012 – 0.377)	-0.107 (-0.320 – 0.107)	0.96 ($p = 0.328$)	1.4	67.1

a

b

c

d

a

b

c

d