Influence of delayed density and ultraviolet radiation on caterpillar granulovirus infection and mortality

^{1,2,*} Adam Pepi, ²Vincent Pan, and ²Richard Karban

¹Graduate Group in Ecology, University of California, Davis, CA 95616

²Department of Entomology & Nematology, University of California, Davis, CA 95616

* Corresponding author. Email: adampepi@gmail.com

Keywords:

Arctiinae; baculovirus; Bodega Marine Lab; disease dynamics; latitudinal gradient; population cycles

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Abstract

1. Infectious disease is an important potential driver of population cycles, but this must occur through delayed density-dependent infection and resulting fitness effects. Delayed density-dependent infection by baculoviruses can be caused by environmental persistence of viral occlusion bodies, which can be influenced by environmental factors. In particular, ultraviolet radiation is potentially important in reducing the environmental persistence of viruses by inactivating viral occlusion bodies.
2. Delayed density-dependent viral infection has rarely been observed empirically at the population level although theory predicts that it is necessary for these pathogens to drive population cycles. Similarly, field studies have not examined the potential effects of ultraviolet radiation on viral infection rates in natural animal populations. We tested if viral infection is delayed density-dependent with the potential to drive cyclic dynamics and if ultraviolet radiation influences viral infection.
3. We censused 18 moth populations across nearly 9° of latitude over two years and quantified the effects of direct and delayed density and ultraviolet radiation on granulovirus infection rate, infection severity, and survival to adulthood. Caterpillars were collected from each population in the field and reared in the laboratory.
4. We found that infection rate, infection severity, and survival to adulthood exhibited delayed density-dependence. Ultraviolet radiation in the previous summer decreased infection severity, and increased survival probability of the virus. Structural equation modelling found that the effect of lagged density on moth survival was mediated through infection rate and infection severity, and was 2.5 fold stronger than the effect of ultraviolet radiation on survival through infection severity.

5. Our findings provide clear evidence that delayed density dependence can arise through viral infection rate and severity in insects, which supports the role of viral disease as a potential mechanism, among others, that may drive insect population cycles. Furthermore, our findings support predictions that ultraviolet radiation can modify viral disease dynamics in insect populations, most likely through attenuating viral persistence in the environment.

Introduction

Infectious disease plays an important role in population dynamics, potentially regulating populations and driving cycles. Many insect populations undergo dramatic cyclic fluctuations, and cyclic, delayed density-dependent disease outbreaks have been proposed as an explanation (Anderson & May, 1980, 1981). However, many other mechanisms have also been proposed to explain insect population cycles (Myers & Cory, 2013), particularly specialist parasitoids (Berryman, 1996). For some species of Lepidoptera, empirical evidence has suggested that baculoviruses are most likely the cause of cyclicity (Myers, 2000; Myers & Cory, 2013, 2016), though this has not been formally tested.

Delayed density-dependent feedbacks are required to drive cyclic population dynamics (Turchin, 2003). Detection of delayed density-dependence involving many ecological factors has been common (Turchin, 1990), and there have been several observations that viral epizootics follow high densities of some insect species (Cory & Myers, 2003; Fuxa, 2004; Myers, 2000; Myers & Cory, 2016). However, direct demonstrations of delayed density-dependent viral infection rate and infection-induced mortality are less common (Burthe et al., 2006; Fleming, Kalkmakoff, Archibald, & Stewart, 1986; Rothman, 1997). Observational studies over relatively small areas (>15km) using space-for-time substitutions of local densities have shown delayed density-dependent incidence of viral infection in voles (Burthe et al., 2006) and soil-dwelling hepialid caterpillars (Fleming et al., 1986). Experimental manipulation of western tent caterpillar (*Malacosoma californicum pluviale*) densities at the tree level showed delayed density-dependent infection rates by a nucleopolyhedrovirus (Rothman, 1997).

Besides host density, other aspects of the local environment may affect viral transmission and dynamics in the field (Cory & Myers, 2003). In particular, ultraviolet radiation has been shown to inactivate viruses of all kinds (Sagripanti & Lytle, 2007), including baculovirus occlusion bodies (Griego, Martignoni, & Claycomb, 1985; Witt & Stairs, 1975). In field studies, examination of the presence of baculovirus occlusion bodies on shaded vs. unshaded foliage suggested that sunlight on leaves may inactivate viruses (Olofsson, 1988). A study of baculovirus transmission in forest tent caterpillars (*Malacosoma disstria*) found depressed transmission rates on forest edges as opposed to the forest interior, which was attributed to sunlight inactivating virus on leaves at forest edges (Roland & Kaupp, 1995). A field study of two strains of gypsy moth NPV showed variable resistance of virus to ultraviolet, in which inactivation rate from ambient ultraviolet was greater in a more potent strain than a less potent one (Akhanaev et al., 2017). In another study of forest tent caterpillars using tree ring analyses, it was suggested periods of weaker ultraviolet radiation increased the effect of forest tent caterpillar outbreaks on tree growth (Haynes, Tardif, & Parry, 2018). In this example, less ultraviolet radiation may have allowed better viral persistence and transmission and produced more severe caterpillar outbreaks.

In the present study, we examined the effects of host density, delayed-density dependence, and ultraviolet radiation on the survival, infection rate, and infection severity of an undescribed granulovirus (see supplement) in Ranchman's tiger moth (*Arctia virginialis*). Ranchman's tiger moth (Erebidae:Arctiinae) is a univoltine species with a long larval stage – usually lasting from late summer until the following spring. Pupation occurs in late spring, and adults emerge in early summer, mate and lay eggs during the summer. Larvae hatch mid-summer, and feed cryptically in litter and undergrowth until they become larger in winter months. Caterpillars are generalists and feed on a variety of herbaceous plants, with a preference for alkaloid-containing hosts (Karban, Karban, Huntzinger, Pearse, & Crutsinger, 2010), and occur mostly in open grassland, shrubland, or savannah. Caterpillars are parasitized by tachinid parasitoids, though caterpillars sometimes survive parasitism (English-Loeb,

Karban, & Brody, 1990). Parasitism has been found to have little role in population dynamics, leaving the observed oscillatory dynamics largely unexplained (Karban & de Valpine, 2010). Caterpillars also sometimes show symptoms of granulovirus infection, which include poor growth, inactivity, watery, opaque frass, regurgitation of milky fluid, and death. Granulovirus is a dsDNA virus that persists outside of the host in the environment within a protective protein shell as an occlusion body (OB). Horizontal transmission can occur when caterpillars consume OBs on food sources, which dissolve in caterpillars' alkaline gut and release infective virus particles (Vega & Kaya, 2012). Virus particles infect the caterpillar starting from the epithelial tissue and then move to other parts of the body, including the trachea, fat bodies, and hemolymph (Barrett, Brownwright, Primavera, & Palli, 1998), potentially causing death of the caterpillar. In Ranchman's tiger moth (*Arctia virginialis*), granulovirus, like parasitoids, does not uniformly kill its hosts.

To test the potential role of granulovirus in delayed-density dependent population dynamics in Ranchman's tiger moth and the influence of ultraviolet radiation on the persistence and transmission of virus in moth populations, we censused 18 populations over two years along a gradient of latitude and ultraviolet radiation and reared caterpillars from these populations in the laboratory. To identify the mechanisms through which granulovirus affects moth population dynamics, we measured granulovirus infection rate, severity, and survival of caterpillars to adulthood. We analyzed the effect of caterpillar density and ultraviolet radiation at the study sites on viral infection and survival. We also tested if viral infection mediated the effects of caterpillar density and ultraviolet radiation on caterpillar survival using a structural equation model.

Methods

Censuses

Caterpillars were counted visually along 4 m wide linear transects of varying length (78-300m) at 18 sites along a ~1000 km latitudinal gradient of the Pacific coast in California, Oregon, and Washington (Figure 1). Transects were surveyed in 2019 and 2020, starting from 1 March through 30

May. We visited sites in each year from south to north, in accordance with the phenology of caterpillar development so that caterpillars were surveyed when most were 4th or 5th instars (sampling dates in Table S1). Transects were surveyed at a constant walking speed and all by the same observer (A. Pepi). Density was estimated from caterpillar counts over the transect area (4m x length). During the second year, sites were revisited within ten days from the calendar date of the first visit; however, this species is a slow-growing caterpillar (roughly ten month development period), so density estimates were likely not overly sensitive to small deviations in sampling date. In 2020, up to ca-. 30 caterpillars per site (or less if fewer were found) were collected haphazardly and brought back to the laboratory for rearing.

Rearing

Caterpillars were reared individually in 6 oz plastic souffle cups with fabric lids and kept in an incubator at 18 °C and 75% RH. Caterpillars were fed every 3-4 days with washed organic romaine lettuce and yellow bush lupine (*Lupinus arboreus*) leaves collected from a part of Bodega Marine Reserve without *Arctia virginalis*. Parasitoids that emerged from caterpillars were counted and identified to family. Caterpillars that died during rearing were frozen at -20 °C for subsequent dissection, except for those that clearly died due to emerged parasitoids. Caterpillars were reared until pupation (~50 days), and pupae were placed together by site of origin (no more than ~10 per site) in 30 cm x 30 cm flight cages and allowed to emerge as moths, at room temperature. Pupae and moths were reared until one month after the last moth emerged and sprayed once or twice weekly with water to prevent desiccation. Adults do not feed. Dead pupae and moths were stored dry at room temperature for subsequent dissection. Individuals that reached adulthood fully formed were counted as having survived; individuals that died as caterpillars or failed to emerge from pupae or expand wings were counted as not surviving.

Dissection and viral assays

After death, each individual was dissected to assess infection status. Two to six tissue samples per insect were taken from the abdomen of adults and pupae or fat bodies of caterpillars, broken up

with forceps, smeared on a glass slide with water, and examined under a light microscope at 200x magnification with phase contrast. Visible occlusion bodies (OB) ($<0.15\mu\text{m}$) were confirmed by adding a drop of 1 M NaOH to the slide, which dissolves the viral protein coat, turning the OB transparent (Lacey & Solter, 2012)(Figure S1). Tissue samples were taken until a positive NaOH test was obtained; if no virus was found after six tissue samples, individuals were classified as uninfected (see supplemental methods for more detail).

Infection severity was rated as uninfected, low, medium, high, or very high, based on the number of samples required to discover virus, the density of OBs in tissue smear samples, and the coloration of hemolymph and fat body (Figure S2). Low severity infections took many samples to identify and had low densities of OBs. Medium severity infections took fewer samples to identify and had moderate densities of OBs. High severity samples had infections in all samples with high densities of OBs, with organ systems in the body still intact and identifiable. Very high severity infections had infections in all samples with very high densities of OBs, with internal organ systems unidentifiable.

Ultraviolet irradiation

Daily values for ultraviolet radiation in 2019, specifically ‘Erythemal Daily Dose’ in Watts/m² from OMI/Aura satellite data were accessed from NASA GES DISC (Hovila, Arola, & Tamminen, 2014). Values were extracted for each sampling location and the average value calculated from 360 to 180 days before the collection date of caterpillars (~March-September at most sites; see Table S1 for collection dates). This timing coincided with the season during which viral OBs would have persisted in the environment outside of hosts.

Statistical analysis

First, we analyzed the density-dependence of infection rate (proportion of infected individuals) and the influence of ultraviolet radiation on infection rate. The proportion of all individuals (N=208) that were infected was analyzed in response to log caterpillar density per m² from transects in 2019 (density in the previous year) and 2020 (density in the same year) using beta-binomial generalized

linear mixed models with site as a random effect. To test for the effect of ultraviolet radiation, we implemented a model with both ultraviolet radiation and log density, to control for the effect of density on infection rate.

Second, we analyzed the density-dependence of infection severity, and the influence of ultraviolet radiation on infection severity. We analyzed infection severity (N=198 with severity rated) in response to log density (same year and previous year) using cumulative link mixed models with site as a random effect. As before, to test for the effect of ultraviolet radiation, we implemented a model with both ultraviolet radiation and log density, to control for the effect of density on infection severity. To test for effects of density and ultraviolet on infection severity independent of infection status, we implemented models using only infected individuals (N=184). This allowed us to test whether different processes influenced infection rate vs. infection severity. We implemented cumulative link mixed models with only log density, and both log density and ultraviolet radiation as predictors as before.

Third, we tested for density-dependent survival and effects of ultraviolet radiation on survival. We analyzed survival to emergence as fully formed moths (N=208) in response to caterpillar density, tachinid parasitoid load, and ultraviolet radiation in a beta-binomial generalized linear mixed model with site as a random effect.

Lastly, we tested our structural hypothesis that viral infection and infection severity were the mechanisms through which lagged density and ultraviolet radiation affected survival in moth populations. There are multiple possible pathways through which density, ultraviolet radiation, infection rate, and infection severity might influence survival and thus population dynamics. To test our specific structural hypothesis, we constructed a piecewise structural equation model which tested whether infection rate and infection severity mediated the effects of ultraviolet radiation and caterpillar density on survival. To do this, we transformed infection severity into a continuous variable between 0 and 1, by converting to numerical categories from 1-5, dividing by 5, and subtracting 0.1. We constructed 3 component sub-models of our structural equation model: a beta-binomial model of

infection rate predicted by log density and ultraviolet radiation, a beta model of infection severity rate predicted by infection rate, log density and ultraviolet radiation, and a beta-binomial model of survival predicted by infection rate and infection severity (Figure 5). All sub-models included site as a random effect, and only individuals that were not parasitized and were scored for infection severity were included in the analysis (N=198). Standardized coefficients were calculated using the latent-theoretic method (Grace, Johnson, Lefcheck, & Byrnes, 2018).

Beta-binomial and beta generalized linear mixed models were fitted using glmmTMB (Brooks et al., 2017). Cumulative link mixed models were fitted using the ordinal package (Christensen, 2019). Piecewise structural equation models were fit using piecewiseSEM v. 1.2.1 (Lefcheck, 2016), and component models were fit using glmmTMB. Ultraviolet radiation was logged and scaled to improve model convergence. We checked for multicollinearity in models using VIFs; multicollinearity was low to moderate in all models (<6) except for same year density in models of infection severity (VIF from 10.5-11.5). Since we included same year density as a control variable, we kept these models as is despite high VIFs. Plots were generated using ggplot2 (Wickham, 2009), ggeffects (Lüdtke, 2018), and viridis (Garnier, 2018). All analysis and plotting were conducted in R v. 3.6.3 (R Development Core Team 2020).

Results

Infection rate

The probability of infection increased with log density in the previous year in models of infection rate (Table 1, Figure 2a). Infection rate had little relation to log density in the current year or to ultraviolet radiation over the season when environmental OBs could have been exposed (Table 1, Figure 2b).

Infection severity

Log density in the previous year increased infection severity in models of infection severity that included both infected and uninfected caterpillars but this effect was weakened when ultraviolet

radiation was included in the model (Table 1, Figure 3a,b). This effect of ultraviolet radiation may have been due to low power or to a negative correlation between log density in the previous year and ultraviolet radiation (Pearson's r : -0.53). Ultraviolet radiation decreased infection severity when included (Table 1, Figure 3c,d). In models that included only infected caterpillars, density in the current and previous years had little effect on infection severity, and ultraviolet radiation had a negative effect on infection severity (Table 1).

Survival to adult emergence

Survival to adult emergence decreased with log density in the previous year and with the number of emerged parasites. Survival increased in response to density in the current year, and in response to ultraviolet radiation (Table 1, Figure 4).

Structural equation model

We hypothesized a causal model linking ultraviolet radiation and disease to caterpillar survival (Fig. 5). Shipley's d-separation test (Shipley, 2000) indicated that our structural equation model was correctly specified (Fig. 5, Fisher's $C = 7.38$, $df = 6$, $P = 0.288$), meaning that there were no missing paths in our causal model. The model results showed that the negative effect of log density in the previous year on survival was mediated by infection rate and infection severity and that the effect of ultraviolet radiation on caterpillar survival was mediated by infection severity (Fig 5). The relative indirect positive effect size of log density in the previous year was ~2.5x greater than the positive effect of ultraviolet radiation on survival (-0.085 vs. 0.035, calculated by multiplying path coefficients).

Discussion

Overall, our results show that log density during the previous year increased infection rate and infection severity and decreased survival to adult emergence, suggesting a strong delayed density-dependent effect of disease on caterpillar population dynamics. The effect of delayed density on both infection rate and infection severity in univariate models and the result of the structural equation model suggest that the effect of density on survival acts through infection severity as well as infection rate.

This is likely due to dose-dependent infection severity, in which caterpillars that are exposed to more viral occlusion bodies develop more severe infections that result in death (Cabodevilla et al., 2011; Eberle, Asser-Kaiser, Sayed, Nguyen, & Jehle, 2008; Matthews, Smith, & Edwards, 2002).

In contrast, log density in the current year, representing direct density-dependence, had no effect on infection rate, a marginal negative effect on infection severity, and a positive effect on survival to adult emergence. Though inconsistent, this positive effect may have been due to ascertainment bias: sites that had larger viral outbreaks may have already declined in density over the course of the larval season (summer 2019-spring 2020). Therefore, density at the time of collection had potentially a reversed relationship to environmental OB density: sites that had declined from earlier high densities would have higher levels of OB, resulting in more severe infections and a higher probability of death. Overall, our results indicated negative delayed density dependence and weak or positive direct density-dependence consistent with the second-order, oscillatory population dynamics observed in the system over the long term (Pepi, Holyoak & Karban, unpublished analyses).

Ultraviolet radiation by contrast had no detectable effect on infection rate but reduced infection severity and increased survival. The reduction in infection severity was even greater when only infected individuals were considered. The effect of ultraviolet radiation on infection severity and survival but not infection results may represent dose-dependent effects of exposure to OBs. Specifically, at sites with higher ultraviolet radiation, more viral OBs were likely inactivated before caterpillars could be exposed to them, so caterpillars received smaller doses of virus and thus became less severely infected and were more likely to survive.

The observational nature of our study gives it some strengths and weaknesses. Our study is one of few to show delayed-density dependent viral infection in the field, and the only study to show effects of radiation on viral infection in a natural population of insects. Our results provide validation in a field system to laboratory studies of virus dynamics and inactivation effects of ultraviolet radiation on viruses (e.g., Bjørnstad et al., 1998; Witt & Stairs, 1975). Despite highly variable conditions between

sites, which were spread across a gradient of over 1000 km, we were able to detect clear effects of delayed density on infection rate, severity, and survival to adulthood, and of ultraviolet radiation on infection severity and survival to adulthood. We were also able to use structural equation modelling to show that infection rate and severity mediate the effect of delayed density and ultraviolet radiation on survival and strengthen the inferences made from our observational study.

Density in the previous year was a strong predictor of infection rates in populations (Fig 2). This finding makes sense because there must have been a large enough population of hosts in the previous year for the disease to spread and produce sufficient inoculum to persist into the current year. Density in the previous year also affected infection severity (Fig 3), but this effect became much weaker when ultraviolet radiation was included in models and even weaker when only infected caterpillars were considered. In the structural equation model, infection severity was well explained by infection rate because individuals must be infected to have high infection severity. This result may be due to the conflation of infection rate and infection severity in the model since the same dose-dependent mechanism may have caused them. However, in all models of infection severity that included both ultraviolet radiation and delayed density, the effect of delayed density, beyond whether individuals were infected or not, was weaker than the effects of ultraviolet radiation. Thus when effects on infection severity were detected they may have in fact been generated by the spurious negative relationship between delayed density and ultraviolet radiation. In contrast to infection severity, ultraviolet radiation had no significant effect on infection rate. Ultimately, both delayed density and ultraviolet radiation had strong and opposing effects on survival to adulthood, though the effect of density in the previous year was much greater in magnitude. The effects of delayed density on survival were mediated through infection with limited evidence of direct effect on infection severity. In contrast the effects of ultraviolet radiation on survival were mediated entirely through effects on infection severity.

298 A possible explanation for the finding that infection rates are primarily determined by host
 299 density in the previous year but not by ultraviolet radiation is that viral infection may be determined in
 300 part through vertical transmission (Burden, Griffiths, Cory, Smith, & Sait, 2002; Cabodevilla et al.,
 301 2011; Cory, 2015; Cory & Myers, 2003). In particular, vertical transmission is likely to occur in *A.*
 302 *virginalis* through persistent sub-lethal infections. We regularly observed OB in egg samples in adult
 303 dissections, and thus eggs of adults with sublethal infections are likely contaminated with virus.
 304 Vertical transmission represents a mechanism through which density in the previous year might
 305 influence infection rates in the following year but is not subject to influence from ultraviolet radiation.
 306 In this way, vertical transmission may maintain higher infection rates after high-density years with
 307 epizootics, even if ultraviolet radiation attenuates inoculum in the environment (Cabodevilla et al.,
 308 2011). Sublethal infections may represent another means by which granulovirus affects population
 309 dynamics by imposing fitness costs on adults (Cabodevilla et al., 2011; Matthews et al., 2002;
 310 Rothman, 1997), although we did not measure this in the present study.

311 In summary, we demonstrated population-level delayed density-dependent effects on viral
 312 infection rate, infection severity, and survival to the reproductive stage, showing how viral infection
 313 may drive cyclic dynamics in insects. We also showed for the first time that ultraviolet radiation may
 314 influence disease dynamics and ultimately population dynamics of insects, through decreased infection
 315 severity and increased survival rates. This provides support for the proposal that viral epizootics may
 316 be an important mechanism driving cyclic dynamics of insects and Lepidoptera in particular (Myers &
 317 Cory, 2013). Our findings also suggest that ultraviolet radiation may be an important factor to consider
 318 as a driver of insect viral disease in the context of global change. In particular, long-term changes in
 319 atmospheric transmittance of solar radiation, or “global dimming and brightening,” have been observed
 320 and are potentially anthropogenic in origin (Wild, 2016). Such long-term changes may have the
 321 potential to influence insect viral disease and population dynamics through changing levels of
 322 ultraviolet attenuation of viral inoculum (Haynes et al., 2018).

Acknowledgements

We would like to thank Marcel Holyoak, Jenny Cory, and Jay Rosenheim for helpful comments on the manuscript. We would also like to thank California State Parks, Ben Becker at Point Reyes National Seashore, Kristen Ward at Golden Gate National Recreation Area, Jackie Sones at Bodega Marine Reserve, Brendan Leigh at Humboldt Bay NWR, Karlee Jewell at North Coast Landtrust, Diane Steeck at the City of Eugene, Grey Wolf at the City of Salem, Oregon Metro Parks, and Christian Haaning at the City of Portland for assistance and access to field sites. We would also like to thank Harry Kaya for assistance with identifying the granulovirus, Jay Rosenheim for the use of incubators, Neal Williams for the use of compound microscopes, and Claire Beck and Jasmine Daragahi for help with rearing caterpillars. The research was funded by NSF-LTREB-1456225 and an NSF REU supplement (DEB-2018169).

Author contributions

AP, VP, and RK conceived the study. AP and VP collected the data and conducted the analyses. AP wrote the manuscript. All authors contributed critically to the drafts and approved final publication.

Data Availability

Data will be archived on KNB upon acceptance.

References

- Akhanaev, Y. B., Belousova, I. A., Ershov, N. I., Nakai, M., Martemyanov, V. V., & Glupov, V. V. (2017). Comparison of tolerance to sunlight between spatially distant and genetically different strains of *Lymantria dispar* nucleopolyhedrovirus. *Plos One*, 12(12), e0189992.
- Anderson, R. M., & May, R. M. (1980). Infectious Diseases and Population Cycles of Forest Insects. *Science*, 210(4470), 658–661.
- Anderson, R. M., & May, R. M. (1981). The population dynamics of microparasites and their invertebrate hosts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 291(1054), 451–519.

348 Barrett, J. W., Brownwright, A. J., Primavera, M. J., & Palli, S. R. (1998). Studies of the
349 nucleopolyhedrovirus infection process in insects by using the green fluorescence protein as a
350 reporter. *Journal of Virology*, 72(4), 3377–3382.

351 Berryman, A. A. (1996). What causes population cycles of forest Lepidoptera? *Trends in Ecology and*
352 *Evolution*, 11(1), 28–32. doi: 10.1016/0169-5347(96)81066-4

353 Bjørnstad, O. N., Begon, M., Stenseth, N. C., Falck, W., Sait, S. M., & Thompson, D. J. (1998).
354 Population dynamics of the indian meal moth: Demographic stochasticity and delayed regulatory
355 mechanisms. *Journal of Animal Ecology*, 67(1), 110–126. doi: 10.1046/j.1365-2656.1998.00168.x

356 Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., ... Bolker,
357 B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated
358 generalized linear mixed modeling. *R Journal*, 9(2), 378–400. doi: 10.32614/rj-2017-066

359 Burden, J. P., Griffiths, C. M., Cory, J. S., Smith, P., & Sait, S. M. (2002). Vertical transmission of
360 sublethal granulovirus infection in the Indian meal moth, *Plodia interpunctella*. *Molecular*
361 *Ecology*, 11(3), 547–555.

362 Burthe, S., Telfer, S., Lambin, X., Bennett, M., Carslake, D., Smith, A., & Begon, M. (2006). Cowpox
363 virus infection in natural field vole *Microtus agrestis* populations: Delayed density dependence
364 and individual risk. *Journal of Animal Ecology*, 75(6), 1416–1425. doi: 10.1111/j.1365-
365 2656.2006.01166.x

366 Cabodevilla, O., Villar, E., Virto, C., Murillo, R., Williams, T., & Caballero, P. (2011). Intra- and
367 intergenerational persistence of an insect nucleopolyhedrovirus: Adverse effects of sublethal
368 disease on host development, reproduction, and susceptibility to superinfection. *Applied and*
369 *Environmental Microbiology*, 77(9), 2954–2960. doi: 10.1128/AEM.02762-10

370 Christensen, R. H. B. (2019). *ordinal—Regression Models for Ordinal Data*.

371 Cory, J. S. (2015). Insect virus transmission: different routes to persistence. *Current Opinion in Insect*
372 *Science*, 8, 130–135.

- 373 Cory, J. S., & Myers, J. H. (2003). The Ecology and Evolution of Insect Baculoviruses. *Annual Review*
374 *of Ecology, Evolution, and Systematics*, 34, 239–272. doi:
375 10.1146/annurev.ecolsys.34.011802.132402
- 376 Eberle, K. E., Asser-Kaiser, S., Sayed, S. M., Nguyen, H. T., & Jehle, J. A. (2008). Overcoming the
377 resistance of codling moth against conventional *Cydia pomonella* granulovirus (CpGV-M) by a
378 new isolate CpGV-II2. *Journal of Invertebrate Pathology*, 98(3), 293–298. doi:
379 10.1016/j.jip.2008.03.003
- 380 English-Loeb, G. M., Karban, R., & Brody, A. K. (1990). Arctiid larvae survive attack by a tachinid
381 parasitoid and produce viable offspring. *Ecological Entomology*, 15(3), 361–362.
- 382 Fleming, S. B., Kalmakoff, J., Archibald, R. D., & Stewart, K. M. (1986). Density-dependent virus
383 mortality in populations of *Wiseana* (Lepidoptera: Hepialidae). *Journal of Invertebrate Pathology*,
384 48(2), 193–198. doi: 10.1016/0022-2011(86)90123-0
- 385 Fuxa, J. R. (2004). Ecology of insect nucleopolyhedroviruses. *Agriculture, Ecosystems & Environment*,
386 103(1), 27–43.
- 387 Garnier, S. (2018). *viridis: Default Color Maps from “matplotlib.”* Retrieved from [https://cran.r-](https://cran.r-project.org/package=viridis)
388 [project.org/package=viridis](https://cran.r-project.org/package=viridis)
- 389 Grace, J. B., Johnson, D. J., Lefcheck, J. S., & Byrnes, J. E. K. (2018). Quantifying relative
390 importance: computing standardized effects in models with binary outcomes. *Ecosphere*, 9(6),
391 e02283.
- 392 Griego, V. M., Martignoni, M. E., & Claycomb, A. E. (1985). Inactivation of nuclear polyhedrosis virus
393 (Baculovirus subgroup A) by monochromatic UV radiation. *Applied and Environmental*
394 *Microbiology*, 49(3), 709–710. doi: 10.1128/aem.49.3.709-710.1985
- 395 Haynes, K. J., Tardif, J. C., & Parry, D. (2018). Drought and surface-level solar radiation predict the
396 severity of outbreaks of a widespread defoliating insect. *Ecosphere*, 9(8). doi: 10.1002/ecs2.2387
- 397 Hovila, J., Arola, A., & Tamminen, J. (2014). *OMI/Aura Surface UVB Irradiance and Erythema Dose*

398 *Daily L2 Global Gridded 0.25 degree x 0.25 degree V3*,. NASA Goddard Space Flight Center,
399 Goddard Earth Sciences Data and Information Services Center. doi:
400 10.5067/Aura/OMI/DATA2028

401 Karban, R., & de Valpine, P. (2010). Population dynamics of an Arctiid caterpillar-tachinid parasitoid
402 system using state-space models. *Journal of Animal Ecology*, 79(3), 650–661. doi: 10.1111/j.1365-
403 2656.2010.01664.x

404 Karban, R., Karban, C., Huntzinger, M., Pearse, I. S., & Crutsinger, G. M. (2010). Diet mixing
405 enhances the performance of a generalist caterpillar , *Platypreria virginalis*. *Ecological*
406 *Entomology*, 35, 92–99. doi: 10.1111/j.1365-2311.2009.01162.x

407 Lacey, L. A., & Solter, L. F. (2012). Initial handling and diagnosis of diseased invertebrates. In L. A.
408 Lacey (Ed.), *Manual of techniques in invertebrate pathology* (2nd ed., pp. 1–14). London, United
409 Kingdom: Elsevier Ltd.

410 Lefcheck, J. S. (2016). piecewiseSEM: Piecewise structural equation modelling in r for ecology,
411 evolution, and systematics. *Methods in Ecology and Evolution*, 7(5), 573–579. doi: 10.1111/2041-
412 210X.12512

413 Lüdtke, D. (2018). ggeffects: Tidy data frames of marginal effects from regression models. *Journal of*
414 *Open Source Software*, 3(26), 772.

415 Matthews, H. J., Smith, I., & Edwards, J. P. (2002). Lethal and sublethal effects of a granulovirus on
416 the tomato moth *Lacanobia oleracea*. *Journal of Invertebrate Pathology*, 80(2), 73–80. doi:
417 10.1016/S0022-2011(02)00100-3

418 Myers, J. H. (2000). Population fluctuations of the western tent caterpillar in southwestern British
419 Columbia. *Population Ecology*, 42(3), 231–241. doi: 10.1007/PL00012002

420 Myers, J. H., & Cory, J. S. (2013). Population Cycles in Forest Lepidoptera Revisited. *Annual Review*
421 *of Ecology, Evolution, and Systematics*, 44(1), 565–592. doi: 10.1146/annurev-ecolsys-110512-
422 135858

423 Myers, J. H., & Cory, J. S. (2016). Ecology and evolution of pathogens in natural populations of
424 Lepidoptera. *Evolutionary Applications*, 9(1), 231–247. doi: 10.1111/eva.12328

425 Olofsson, E. (1988). Environmental persistence of the nuclear polyhedrosis virus of the European pine
426 sawfly in relation to epizootics in Swedish Scots pine forests. *Journal of Invertebrate Pathology*,
427 52(1), 119–129. doi: 10.1016/0022-2011(88)90110-3

428 Roland, J., & Kaupp, W. J. (1995). Reduced transmission of forest tent caterpillar (Lepidoptera:
429 Lasiocampidae) nuclear polyhedrosis virus at the forest edge. *Environmental Entomology*, 24(5),
430 1175–1178. doi: 10.1093/ee/24.5.1175

431 Rothman, L. D. (1997). Immediate and delayed effects of a viral pathogen and density on tent
432 caterpillar performance. *Ecology*, 78(5), 1481–1493. doi: 10.1890/0012-
433 9658(1997)078[1481:IADEOA]2.0.CO;2

434 Sagripanti, J. L., & Lytle, C. D. (2007). Inactivation of influenza virus by solar radiation.
435 *Photochemistry and Photobiology*, 83(5), 1278–1282. doi: 10.1111/j.1751-1097.2007.00177.x

436 Shipley, B. (2000). A New Inferential Test for Path Models Based on Directed Acyclic Graphs A New
437 Inferential Test for Path Models Based on Directed Acyclic Graphs. *Structural Equation*
438 *Modelling*, 7(2), 206–218. doi: 10.1207/S15328007SEM0702

439 Turchin, P. (1990). Rarity of density dependence or population regulation with lags? *Nature*,
440 344(6267), 660–663. doi: 10.1038/344660a0

441 Turchin, P. (2003). *Complex Population Dynamics: A Theoretical/Empirical Synthesis*. Princeton:
442 Princeton University Press.

443 Vega, F. E., & Kaya, H. K. (2012). *Insect pathology*. Academic press.

444 Wickham, H. (2009). *ggplot2: elegant graphics for data analysis*. New York: Springer.

445 Wild, M. (2016). Decadal changes in radiative fluxes at land and ocean surfaces and their relevance for
446 global warming. *Wiley Interdisciplinary Reviews: Climate Change*, 7(1), 91–107. doi:
447 10.1002/wcc.372

Witt, D. J., & Stairs, G. R. (1975). The effects of ultraviolet irradiation on a Baculovirus infecting
Galleria mellonella. *Journal of Invertebrate Pathology*, 26(3), 321–327. doi: 10.1016/0022-
2011(75)90229-3

Figures & Tables

Table 1. Results of univariate models.

Model structure	Parameter	Logit $\beta \pm 1$ SE	z	P
Infection ~ log density	Log density current year	-0.20±0.38	-0.52	0.603
	Log density previous year	0.55±0.26	2.10	0.036
Infection ~ log density + ultraviolet	Log density current year	-0.10±0.39	-0.25	0.802
	Log density previous year	0.92±0.46	1.998	0.046
	Ultraviolet radiation	0.66±0.58	1.13	0.259
Severity ~ log density	Log density current year	-0.28±0.19	-1.49	0.135
	Log density previous year	0.26±0.11	2.35	0.019
Severity ~ log density + ultraviolet	Log density current year	-0.28±0.18	-1.61	0.101
	Log density previous year	0.16±0.11	1.37	0.170
	Ultraviolet radiation	-0.35±0.17	-2.04	0.041
Severity infected ~ log density	Log density current year	-0.27±0.19	-1.39	0.163
	Log density previous year	0.018±0.11	1.66	0.096
Severity infected ~ log density + ultraviolet	Log density current year	-0.28±0.18	-1.52	0.129
	Log density previous year	0.06±0.12	0.48	0.631
	Ultraviolet radiation	-0.44±0.18	-2.40	0.017
Survival ~ log density + ultraviolet	Log density current year	1.15±0.51	2.25	0.024
	Log density previous year	-0.75±0.31	-2.39	0.017
	Ultraviolet radiation	0.64±0.32	1.97	0.049
	Emerged parasites	-5.39±1.58	-3.42	0.0006

Figure 1. Map of study site locations in Washington, Oregon, and California, overlaid on a map of average annual insolation (GHI), in kWh/m².

Figure 2. Infection rate in response to (a) density in the previous year and (b) ultraviolet radiation as scaled log erythemal daily dose in the summer before collection (2019) from beta-binomial generalized linear mixed models.

Figure 3. Infection severity in response to (a-b) density in the previous year, and (c-d) ultraviolet radiation as scaled log erythemal daily dose in the summer before collection cumulative link mixed models. The left column (a,c) show model predicted rate of infection at each severity class with 95% confidence intervals, and the right column (b,d) shows the proportion of the total population predicted to be infected at each severity class by color.

Figure 4. Survival to adult emergence in response to (a) density in the previous year, and (b) ultraviolet radiation as scaled log erythemal daily dose in betabinomial generalized linear mixed models. Size of points is scaled to the number of individuals reared from each site.

Figure 5. Path diagram of piecewise structural equation model. Size of arrows is proportional to the standardized path coefficient (shown next to lines), and line type indicates P-value of the path parameter.

542 **Supplement**

543 **Table S1.**

Site	Latitude	Longitude	Number reared	Date Collected
Franklin Point, CA	37.155	-122.35	29	3/1/2020
Pescadero Marsh, CA	37.265	-122.41	16	3/1/2020
Golden Gate NRA, CA	37.844	-122.55	28	3/13/2020
Point Reyes South, CA	38.038	-122.99	15	3/13/2020
Point Reyes North, CA	38.088	-122.97	29	3/13/2020
Point Reyes Kehoe, CA	38.153	-122.94	2	3/13/2020
Bodega Marine Reserve, CA	38.319	-123.06	16	3/27/2020
Manchester State Park, CA	38.959	-123.72	6	3/27/2020
Loyalton, CA	39.700	-120.30	6	5/30/2020
Humboldt NWR, CA	40.670	-124.21	20	4/10/2020
Freshwater Farms, CA	40.786	-124.09	11	4/10/2020
Klamath River, Yreka, CA	41.829	-122.61	11	4/10/2020
Checkermallow Access, OR	44.072	-123.20	12	4/11/2020
Fairview Wetland, OR	44.895	-123.00	20	4/11/2020
Graham Oaks, OR	45.299	-122.80	30	4/11/2020
Powell Butte, OR	45.488	-122.50	33	4/11/2020
Salish Ponds, OR	45.531	-122.45	1	4/11/2020
West Rocky Prairie, WA	46.887	-122.87	2	4/12/2020

544

545

Figure S1. Fat body tissue smear from a caterpillar infected with GV under a light microscope at 200x magnification with phase contrast. A drop of NaOH was added which moved through the sample. The red arrows point to the boundary between NaOH solution and water. The yellow arrows point to the OBs (Figure A) which turned transparent (Figure B) after they were dissolved by the strong base.

Figure S2. Ventral dissection of late instar *A. virginalis* larva in the order of infection severity, ranging from (A) uninfected, (B) low severity, (C) medium severity, (D) high severity, to (E) very high severity. The caterpillar's gut in figure A is missing. Caterpillars with higher infection severity have a cloudier hemolymph and a darker and more yellowish fat body. They also often have large clusters of infected tissues next to the midgut (e.g. Figure D).