1 Original Article

2

3 **Weak effect of *Gypsy* retrotransposon bursts on *Sonneratia alba***
 4 **salt stress gene expression**

5

6 Yushuai Wang, Aimei Dai, Tian Tang*

7

8 State Key Laboratory of Biocontrol and Guangdong Key Laboratory of Plant Resources,
 9 School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, Guangdong, People's
 10 Republic of China

11

12 * Correspondence: lsstt@mail.sysu.edu.cn

13 Running Title: Weak effect of TE bursts on gene expression

14

15 **Abstract**

16 **Background and Aims**

17 Transposable elements (TEs) are an important source of genetic diversity and can be co-opted
18 for the regulation of host genes. However, to what extent the pervasive TE colonization of
19 plant genomes has contributed to stress adaptation remains controversial. Plants inhabiting
20 harsh environments in nature provide a unique opportunity to answer this question.

21 **Methods**

22 We compared TE compositions and their evolutionary dynamics in the genomes of two
23 mangrove species: the pioneer *Sonneratia alba* and its less salt-tolerant relative *S. caseolaris*.
24 Age distribution, strength of purifying selection and the removal rate of LTR (long terminal
25 repeat) retrotransposons were estimated. Phylogenetic analysis of LTR retrotransposons and
26 their distribution in the genome of *S. alba* were surveyed. Small RNA sequencing and
27 whole-genome bisulfite sequencing was conducted using leaves of *S. alba*. Expression
28 pattern of LTR retrotransposons and their nearby genes were examined using RNA-seq data
29 of *S. alba* under different salt treatments.

30 **Key Results**

31 *S. alba* possesses more TEs than *S. caseolaris*. Particularly, many more young *Gypsy* LTR
32 retrotransposons have accumulated in *S. alba* than in *S. caseolaris* despite an increase in
33 purifying selection against TE insertions. The top two most abundant *Gypsy* families in *S.*
34 *alba* preferentially insert in gene-poor regions. They are under relaxed epigenetic repression,
35 probably due to the presence of CHROMO domains in their 3'-ends. Although a considerable

36 number of TEs in *S. alba* showed differential expression under salt stress, only four copies
 37 were significantly correlated with their nearby genes in expression levels. One such TE-gene
 38 pair involves *Abscisic acid 8'-hydroxylase 3* functioning in abscisic acid catabolism.

39 **Conclusions**

40 This study sheds light on the evolutionary dynamics and potential function of TEs in an
 41 extremophile. Our results suggest that the conclusion on co-option of TEs should be cautious
 42 even though activation of TEs by stress might be prevalent.

43

44 **Keywords:** Transposable elements (TEs), Genome evolution, TE co-option, Stress adaptation,
 45 Mangroves

46

47 INTRODUCTION

48 Transposable elements (TEs) are genetically diverse mobile sequences that can move to
 49 new sites in genomes by a “copy-and-paste” mechanism via an RNA intermediate (Class I,
 50 retrotransposons) or a “cut-and-paste” mechanism (Class II, transposons) (Wicker et al., 2007,
 51 Bourque et al., 2018). TE insertions often disrupt gene function or regulation and induce
 52 ectopic/nonallelic recombination events (Chuong et al., 2017). As a result, their activities are
 53 repressed by host epigenetic silencing and small interfering RNAs (siRNAs) (Fultz et al.,
 54 2015). Bursts of TE abundance are further subject to mutational decay and are eliminated by
 55 genome purging mechanisms in the long run. In many higher eukaryotes TEs constitute more
 56 than half of the DNA (Fedoroff, 2012) and play a critical role in the evolution of genome size
 57 and organization (Lisch, 2013). It is thus important to dissect the evolutionary dynamics of
 58 TEs in the genome.

59 While TEs are often deleterious or effectively neutral, there is growing evidence that TE
 60 activity can help the host adapt to environmental challenges (Casacuberta and Gonzalez,
 61 2013, Horvath et al., 2017). High copy number and mutagenicity make TEs a major source of
 62 genetic innovation that can be activated by environmental cues (Capy et al., 2000,
 63 Grandbastien, 2015, Rey et al., 2016). This may accelerate mutation rates and generate
 64 variability upon which natural selection could act. Indeed, recent findings show that TEs can
 65 facilitate rapid adaptation to novel environments in species with limited genetic variation,
 66 such as *Capsella rubella* the annual inbreeding forb (Niu et al., 2019) and *Cardiocondyla*
 67 *obscurior* the invasive inbreeding ant (Schrader et al., 2014). Meanwhile, some TEs are
 68 known to carry stress-responsive regulatory elements. These elements, once activated by

69 stress, may have a large role in remodeling regulatory networks and reshaping transcriptomes
70 (Naito et al., 2009, Cavrak et al., 2014, Grandbastien, 2015, Chuong et al., 2017). Despite
71 many reports that individual TE integration events result in changes in gene expression or
72 even phenotypes under stress conditions (Butelli et al., 2012, Lisch, 2013), there is empirical
73 evidence that regulatory function of TEs is not always strong (de Souza et al., 2013, Simonti
74 et al., 2017).

75 Mangroves are a group of phylogenetically diverse woody species that invaded and
76 adapted to the extreme and extremely fluctuating environments in the tropical intertidal zone
77 (Tomlinson, 1986). These species have undergone multiple population bottlenecks during
78 their evolutionary history and typically harbor extremely low standing genetic variation
79 (Zhou et al., 2011, Guo et al., 2018). A recent study reported massive and convergent TE load
80 reduction in mangroves, when compared to non-mangrove species (Lyu et al., 2018).
81 However, we found a continuous host-TE battle masked by the TE load reduction in
82 *Rhizophora apiculata*, the tall-stilted mangrove, suggesting TE activities may enable
83 mangroves to maintain genetic diversity (Wang et al., 2018). It still remains unclear how
84 often TE activities are significant to the evolution of mangrove genomes and to what extent
85 TEs may have influenced gene expression during mangrove stress adaptation.

86 In this study, we surveyed the evolutionary dynamics and potential regulatory functions
87 of TEs in *Sonneratia alba* Sm., the pioneer mangrove (Tomlinson, 1986), in comparison with
88 its relative *S. caseolaris* (L.) Engl.. Although both are mangroves, *S. alba* is more salt-tolerant
89 than *S. caseolaris* and often grows on the seaward side of mangrove forests with maximal
90 growth in 5 to 50% sea water (Ball and Pidsley, 1995). In contrast, *S. caseolaris* is common in

the inner parts of mangrove forests (Bunt et al., 1982) and in some instances has also been found growing in fully fresh water (Rahim and Bakar, 2018). Here we report recent amplification of *Gypsy* LTR (long terminal repeat) retrotransposons in *S. alba*. Nevertheless, this burst of *Gypsy* retrotransposons has only a weak effect on salt stress gene expression. Our results suggest that the conclusion on co-option of TEs should be cautious even though activation of TEs by stress might be prevalent.

97

98 MATERIALS AND METHODS

99 Annotation and classification of transposable elements

Whole genome sequences of *Sonneratia alba*, *Sonneratia caseolaris*, and *Eucalyptus grandis* were retrieved from previous studies (Myburg et al., 2014, Lyu et al., 2018). LTR retrotransposons were *de novo* mined and classified using work flows described previously (Wang et al., 2018). The annotation of long interspersed nuclear elements (LINE), short interspersed nuclear elements (SINE), and DNA transposons (MULE, hAT, PIF, CACTA and TcMar) was carried out as follows. First, sequences of the most conserved coding domain of each TE family were retrieved from the NCBI database (Coordinators, 2018) and used as queries against each genome using the TARGeT pipeline (Han et al., 2009) with the *e*-value cutoff of 0.001. Then, the identified homologs together with their 5 kb up- and downstream sequences were aligned within individual TE families using MAFFT (Katoh et al., 2002) and resolved into lineages using phylogenetic analyses as implemented in SiLiX (Miele et al., 2011). The terminals of each putative element were determined manually according to

sequence similarity with its closely related elements. Miniature inverted-repeat transposable elements (MITEs) and Helitron copies were identified using MITE_HUNTER (Han and Wessler, 2010) and HelitronScanner (Xiong et al., 2014) with default parameters, respectively. Finally, all *de novo* identified TEs were added to RepBase (v.23.06) (Bao et al., 2015) for each genome separately and used to annotate TE contents in each species using RepeatMasker (v.4.0.0) (Smit et al., 2013-2015).

Dating LTR retrotransposon elements

Age of each LTR retrotransposon element was estimated based on sequence divergence between their LTR pair (SanMiguel et al., 1998). The pairs from individual LTR retrotransposons were aligned using MUSCLE (v.3.8.31) with default parameters (Edgar, 2004). Sequence divergence k between each pair of LTRs was estimated using Kimura's two-parameter model (Kimura, 1980). The age (T) of each LTR retrotransposons was estimated using the formula $T = k / 2r$, where r is the substitution rate of 1.3×10^{-8} substitutions per site per year (Ma and Bennetzen, 2004).

Distribution of TEs across genomic regions

The insertion frequency of TEs categories in intergenic regions, introns, and coding regions in *S. alba* and *S. caseolaris* were estimated according to the procedure described before (Wright et al., 2003). Briefly, the genome sequences and annotated gene positions of *S. alba* and *S. caseolaris* were obtained from Lyu et al. (2018). For each species, each scaffold (>100 kb) was submitted to a BLAST search against our TE database to determine the positions of each TE superfamily along the entire set of scaffolds. TE positions were then

verified manually to eliminate all redundancies and record the presence of nested insertions. To estimate gene density, sequences were scanned using 100 kb sliding windows with 50 kb steps across each scaffold. Gene density of each window was calculated as the fraction of the total length occupied by coding sequences.

Phylogenetic analyses of LTR retrotransposons

Multiple alignments of full-length sequences of intact *Copia* or *Gypsy* retrotransposons were conducted using MAFFT (Katoh et al., 2002). Maximum likelihood phylogenetic trees were inferred with FastTree (Price et al., 2009) using a generalized time-reversible (GTR) model of nucleotide substitution. The generated trees were visualized with Figtree (v.1.4.2) (Rambaut, 2012). Resampling of estimated log-likelihoods (RELL)-like local support values was performed using FastTree (Price et al., 2009) with default parameters.

Identification and sequence analysis of the chromodomain in LTR retrotransposons

Chromodomains residing in LTR retrotransposons were identified by tBLASTx (NCBI) searches against the *nr* database (NCBI) with default parameters (Boratyn et al., 2012). Multiple alignments of the chromodomain amino acid sequences were performed using MAFFT (Katoh et al., 2002). Tertiary structures of the chromodomains peptides were predicted using the Phyre2 server with default parameters (Kelley et al., 2015). Known chromodomain amino acid sequences were retrieved from the NCBI database (Coordinators, 2018), including DmHP1 (GenBank accession number, M57574) from *Drosophila melanogaster*, MAGGY (L35053) and Pyret (AB062507) from *Magnaporthe grisea*, Cft1 (AF051915) from *Cladosporium fulvum*, PpatensLTR1 (XM_001752430) from

Physcomitrella patens, Tf2-12 (RVW30894) from *Vitis vinifera*, sr35 (AC068924), rn8 (AK068625) and RIRE3 (AC119148) from *Oryza sativa*, Tekay (AF050455) from *Zea mays*, RetroSor2 (AF061282) from *Sorghum bicolor*, and Tma (AF147263) from *Arabidopsis thaliana*.

Epigenome sequencing and analyses

Young leaves of *S. alba* were collected from Qinlan Harbour, Hainan, China. Genomic DNA and total RNA were extracted separately using the modified CTAB protocol (Yang et al., 2008). Bisulfite (BS-seq) and small RNA sequencing were conducted with two biological replicates each according to previously published procedures (Wen et al., 2016, Wang et al., 2018).

Bisulfite sequencing reads were filtered for quality using Trimmomatic (v.0.32) (Bolger et al., 2014) and mapped to the *S. alba* genome using Bismark (v.0.16.3) (Krueger and Andrews, 2011) with default parameters. Only uniquely mapping reads were retained. Methylation levels of each LTR retrotransposon copy were calculated as the proportion of Cs among Cs and Ts ($\#C/(\#C+\#T)$) by methylation context (CG, CHG and CHH) using custom Perl scripts. Only sites covered by more than three reads were used for the calculation. Results were visualized using R (v.3.1.3) (<https://www.R-project.org>).

Small RNA sequencing reads were quality controlled following procedures published previously (Wen et al., 2016) while structural non-coding RNAs, such as ribosomal RNA (rRNA), transfer RNA (tRNA), small nuclear RNA (snRNA), small nucleolar RNA (snoRNA), and known microRNAs (miRNAs) were removed. The remaining reads were considered as putative siRNAs and aligned to *S. alba* genomes using Bowtie (Langmead et al.,

2009) with no mismatches allowed. SiRNA density of a particular LTR retrotransposon element was calculated as read counts per kilo base pair per element. Multiply-mapping reads were weighted by the number of their mapping locations. All statistical calculations were performed in R (v.3.1.3).

Expression analyses of TEs and TE-flanking genes under salt treatments

Transcriptome data from *S. alba* under the 0, 250mM, and 500mM NaCl treatments were obtained from Feng et al. (2020) (NCBI Sequence Read Archive database accession No. PRJNA615770). Sequencing adapters and low quality reads were filtered using Trimmomatic (v.0.32) (Bolger et al., 2014). Clean reads were mapped to coding genes or intact LTR retrotransposons separately using Bowtie (Langmead et al., 2009) allowing no more than two mismatches. We calculated the expression level of each LTR retrotransposon element as reads per kilobase of transcript. Expression levels of individual genes were analyzed using HTSeq (v.0.6.1) (Anders et al., 2015) with the parameter -s set to “no” and normalized to Reads Per Kilobase per Million mapped reads (RPKM). Only genes located within 3 kb up- or downstream of the insertion of an intact or truncated LTR retrotransposon were considered in our analysis. The expression differences of TEs or genes across salt treatments were identified using DESeq2 (Love et al., 2014) requiring Benjamini-Hochberg multiple testing corrected (Benjamini and Hochberg, 1995) FDR < 0.05 and ≥ 2 -fold change.

Identification of *cis*-acting regulatory elements (CREs) of LTR retrotransposons

The CREs within LTRs of the consensus sequence of each LTR retrotransposon family or within the whole sequence of each truncated LTR retrotransposon were identified using

PlantCARE (http://bioinformatics.psb.ugent.be/webtools/plantcare/html/search_CARE.html) (Lescot et al., 2002). *Gypsy* and *Copia* families that have more than ten copies were considered in this analysis.

MiRNA target prediction

Plant mature miRNAs and their precursors were retrieved from miRBase (Release 22) (Kozomara et al., 2019). The identification of the known and novel miRNAs was performed by miRDeep2 software package (Friedlander et al., 2012) with the default parameters. The expression level of each identified miRNAs was normalized to Reads Per Million mapped reads (RPM). MiRNAs that with a RPM > 1 in both biological replicates were considered expressed and retained for further analysis. The prediction of the putative target genes for the known and novel miRNAs was done with the psRNATarget (Dai and Zhao, 2011) using an expectation score of 2.5 as the cutoff in the prediction. All LTR retrotransposons including the intact and truncated elements and solo-LTRs were used in these analyses.

Accession numbers

Sequencing data from this article can be found in the GenBank data libraries under accession number PRJNA636918.

RESULTS

TEs are more abundant in *S. alba* than in *S. caseolaris*.

S. alba and *S. caseolaris* diverged about ~11.6 million years (Myr) ago (Yang et al.,

2016). The whole genome sequences of *S. caseolaris* (~208Mb, 1C = 259 Mb) and *S. alba* (~207Mb, 1C = 284 Mb) have recently been *de novo* assembled (Lyu et al., 2018). We used the whole genome of *Eucalyptus grandis* (Myburg et al., 2014) as the outgroup following a previous study (Xu et al., 2017) and studied the evolutionary dynamics of TEs in *S. alba* in comparison with *S. caseolaris*. We annotated TEs in each species considering both TE structure and sequence homology and clustered them into families using the same procedure (see *MATERIALS AND METHODS*).

Consistent with the report of TE load reduction in mangroves (Lyu et al., 2018), we found TEs only accounted for 5.3% (11,091,467 bp) and 11.0% (22,805,675 bp) of the *S. caseolaris* and *S. alba* genomes, respectively, far less than the proportion of TEs in the *E. grandis* genome (27.7%, 191,425,687 bp, Figure 1A). In all three species, LTR retrotransposons are the dominant TE components (Figure 1A). Nevertheless, *Gypsy* retrotransposons are more abundant than *Copia* retrotransposons in the two *Sonneratia* species while the opposite pattern is observed in the outgroup *E. grandis* (Figure 1A). The expansion of *Gypsy* retrotransposons is more prominent in *S. alba* (12.7 Mb or 6.1%) than in *S. caseolaris* (3.3 Mb or 1.6%, Figure 1B). We identified 947 intact LTR retrotransposon copies comprising 180 *Copia* and 767 *Gypsy* elements in *S. alba* and 125 intact LTR retrotransposon copies consisting of 51 *Copia* and 74 *Gypsy* elements in *S. caseolaris* (Supplementary Data: Table S1). Although mangroves have experienced an overall reduction in TE load (Lyu et al., 2018), more *Gypsy* retrotransposons have accumulated in *S. alba* than in *S. caseolaris*.

***Gypsy* retrotransposons experienced recent bursts in *S. alba*.**

To explore the dynamics of TE accumulation, we first estimated ages of the intact *Gypsy* and *Copia* retrotransposons by comparing LTR sequences flanking each element and assuming a molecular clock of 1.3×10^8 (Ma and Bennetzen, 2004). We found these retrotransposons were much younger in *S. alba* (*Copia*: 1.47 Myr old; *Gypsy*: 0.87 Myr old) than in *S. caseolaris* (*Copia*: 3.85 Myr old and *Gypsy*: 3.99 Myr old). These differences are highly significant (Mann-Whitney *U*-test, both $P < 0.001$, Figure 1C). The recent burst of TE activity is most striking for *Gypsy* elements in *S. alba*. Young (< 1 Myr old) *Gypsy* elements account for 54.4% of the intact elements in *S. alba* but only 8.1% in *S. caseolaris* (Figure 1D). By contrast, the proportions of young *Copia* elements are more similar between *S. alba* (37.9%) and *S. caseolaris* (25.5%, Figure 1D). Therefore, there are more recent originations of the *Gypsy* element in *S. alba* than in *S. caseolaris*.

We then inferred the intensity of purifying selection against deleterious TE insertions by comparing TE distribution in relation to coding regions. We found TE insertions are strongly under-represented in coding regions compared to introns and intergenic regions (Table 1), consistent with the prediction of the deleterious insertion model (Finnegan, 1992). The abundance of all TE categories regardless of their locations are significantly lower in *S. alba* than in *S. caseolaris* (Table 1), suggesting TEs are under stronger purifying selection in the former species. However, the abundance of *Gypsy* elements in introns and intergenic regions was comparable between *S. alba* and *S. caseolaris* (Mann-Whitney *U*-test, all $P > 0.05$), suggesting that the insertions of *Gypsy* into introns and intergenic regions have experienced relaxation of purifying selection in *S. alba*. It is also possible that purifying selection against *Gypsy* insertions into coding regions is even stronger than that for other TE families in *S.*

264 *alba*; otherwise, we should see more *Gypsy* insertions in the coding regions of that species.

265 Finally, we analyzed the removal rate of LTR elements by estimating the ratio of
266 solo-LTRs to intact elements (S:I). Unequal intra-element homologous recombination
267 resulting in solo-LTRs is known to be one of the major mechanisms eliminating LTR
268 retrotransposons (Devos et al., 2002). As shown in Table 2, the S:I ratio is the lowest for
269 *Gypsy* elements in *S. alba* (0.34), followed by *Copia* in *S. alba* (1.79) and then the *Copia* and
270 *Gypsy* elements in *S. caseolaris* (2.45 and 3.72). Taken together, these results suggest that
271 despite strong purifying selection, high proliferation activity and low elimination rate jointly
272 contributed to the rapid accumulation of *Gypsy* elements in the *S. alba* genome.

273 **Recent bursts of two *Gypsy* families in *S. alba* are associated with integration site**
274 **preference.**

275 To compare the evolutionary histories of *Copia* and *Gypsy* retrotransposons in the two
276 *Sonneratia* species, we conducted phylogenetic analyses using full length sequences of intact
277 elements for the *Copia* and *Gypsy* superfamilies separately. The maximum-likelihood
278 phylogenetic trees grouped LTR retrotransposon copies of *S. alba* and *S. caseolaris* into 26
279 *Copia* and 23 *Gypsy* families, half of which (13 out of 26 for *Copia* and 11 out of 23 for
280 *Gypsy*) were shared by both genomes (Supplementary Data: Figure S1). Remarkably, we
281 found that two *Gypsy* families (RLG_1 and RLG_8) are ~10 times more abundant in *S. alba*
282 (RLG_1: 226 copies and RLG_8: 273 copies) than in *S. caseolaris* (RLG_1: 20 and RLG_8:
283 22, Supplementary Data: Figure S1). These two *Gypsy* families account for 64.6% of the
284 young copies (< 1 Myr) and 65.1% of the total intact *Gypsy* elements in *S. alba* (Figure 2A).
285 Furthermore, three RLG_1 elements and four RLG_8 elements have identical 5' and 3' LTR

sequences and perfect 5-bp Target Site Duplications (TSDs), suggesting they are newly originated copies. Therefore, recent bursts of RLG_1 and RLG_8 in *S. alba* are the major contributors to the observed expansion of *Gypsy* elements.

Plant LTR retrotransposons are preferentially associated with large heterochromatic regions that flank functional centromeres (Kumar and Bennetzen, 1999). We wondered whether such integration site preference, leading to loose epigenetic repression, might have contributed to the expansion of RLG_1 and RLG_8 families. We scanned scaffolds of the *S. alba* genome using 100 kb sliding windows with a step-size of 50 kb and estimated gene density in each window (see *MATERIALS AND METHODS*). As expected, about half of the RLG_1 copies (50.9%) and 2/5 of the RLG_8 copies (40.7%) are located in gene-poor regions (gene density < 0.05) in *S. alba*. In contrast, the proportion is only 8.1% and 12.5% for the third and fourth most abundant LTR retrotransposon families (Figure 2B).

Further analysis of the structure of RLG_1 and RLG_8 detected a chromodomain (CHD) in the 3' end of both families. CHD was first identified in the heterochromatin protein-1 (HP1) of *Drosophila melanogaster* (Paro and Hogness, 1991) and considered to be involved in chromatin remodeling and the formation of heterochromatin (Paro and Hogness, 1991). Sequence analyses of CHD proteins from diverse organisms classified CHD of the RLG_1 and RLG_8 families into Group I chromodomains (Figure 3A). Both of them possess a highly conservative protein secondary structure (Figure 3B). CHD in *Gypsy* elements, also named chromoviruses, has been experimentally demonstrated to promote the integration of newly emerging retroelements into heterochromatic regions (Gao et al., 2008).

Expanded *Gypsy* families are under relaxed epigenetic repression

If CHD indeed promotes the insertions of RLG_1 and RLG_8 into heterochromatin regions, we would expect that epigenetic repression of these elements is relaxed in the host genome. In plants, CHH methylation mediated by 24-nt siRNAs is responsible for the establishment of *de novo* methylation against the emergence of new TE copies (Law and Jacobsen, 2010). We thus estimated correlations between gene density, CHH methylation level, and 24-nt siRNA abundance for the RLG_1 and RLG_8 copies in *S. alba*. Gene density of the window in which each RLG_1 and RLG_8 copy resides was used in this analysis.

For both RLG_1 and RLG_8, copies residing in the gene-poor regions (gene density < 0.05) exhibited a significantly lower level of CHH methylation (RLG_1: median, hereafter, 10.6% and RLG_8: 15.9%) than the rest (RLG_1: 44.0% and RLG_8: 37.7%, Mann-Whitney *U*-test, all $P < 0.001$, Figure 4A). The 24-nt siRNA targeting pattern is in accordance with the difference in methylation levels, i.e., a lower level of 24-nt siRNA abundance on TEs residing in gene-poor regions than the other TEs (4.0 vs. 7.5 reads per element per kilobase for RLG_1 and 2.9 vs. 5.8 for RLG_8, Mann-Whitney *U*-test, all $P < 0.001$; Figure 4A). These results support the idea that members of RLG_1 and RLG_8 have experienced relaxed epigenetic repression associated with their insertion preference. Gene-poor regions might represent a “safe haven” for these elements to coexist with host cells.

Weak effect of TE expansion on salt stress gene expression in *S. alba*

Despite the overall preference for integration into gene-poor regions, there is still a considerable fraction of young copies (< 1 Myr) of RLG_1 (21.8%) and RLG_8 (45.2%) inserted into genomic regions with gene density > 0.15 (Figure 4B). It is possible that recently amplified TEs can affect nearby gene expression by carrying repressive epigenetic

markers or creating new regulatory elements (Naito et al., 2009, Chuong et al., 2017). Recently, Feng et al. (2020) reported a comparative transcriptome study of *S. alba* treated with 0, 250mM, and 500mM NaCl. They used the 250mM NaCl condition as a control and 0 and 500mM NaCl as stresses with low or high salt (Feng et al., 2020). Using these data, we asked whether intact LTR retrotransposons in *S. alba* can be activated by salt stress and how TE expansion might affect gene expression under stress conditions.

Many more of the 767 and 180 intact *Gypsy* and *Copia* retrotransposons were expressed in roots (*Gypsy*: 420 and *Copia*: 15) than in leaves (*Gypsy*: 169 and *Copia*: 11) (Figure 5A). This is coincident with the fact that roots are directly exposed to the saline environment. An intact retrotransposon was considered to be expressed if it had more than 5 reads in both biological replicates under at least one of the low, medium, or high salinity conditions. There was more upregulation ($\log_2$ fold change > 1) than downregulation ($\log_2$ fold change > -1) for these retroelements in all comparisons, most prominently for the comparison between 500mM and 250mM NaCl in roots (Figure 5A). However, only four retrotransposons were significantly mis-regulated due to large variation among replicates.

We further checked changes of expression in genes with intact TEs in introns, within 3kb upstream of the transcription start site, or downstream of the transcription termination site for the top four most abundant LTR retrotransposon families in *S. alba*. Gene expression level was calculated as reads per kb per million reads (RPKM). For all the four families, genes with neighboring TEs were expressed at a significantly lower level than the whole transcriptome under all conditions (Mann-Whitney *U*-test, all $P < 0.001$ in both leaves and roots; Figure 5B), supporting the view that TEs carry repressive epigenetic marks that may

reduce flanking gene expression (Naito et al., 2009, Chuong et al., 2017). While a few TE-adjacent genes were significantly mis-regulated ($FDR < 0.05$ and fold change ≥ 2) by salt (Figure 5C and Supplementary Data: Table S2), this set of loci was not enriched in mis-expressed genes relative to the whole transcriptome in any of our comparisons (Fisher exact test, all $P > 0.05$, Supplementary Data: Table S3). We also estimated correlations between TE expression levels and their nearby genes in leaves or roots across the three salt conditions (Supplementary Data: Table S4). Only two significantly correlated pairs were identified in leaves: a RLC_4 copy and its 1307-bp upstream gene of unknown function (Pearson's correlation, $r = 0.99$, adjusted $P < 0.05$), and a RLG_8 copy and its 505-bp downstream gene *Abscisic acid 8'-hydroxylase 3* (Pearson's correlation, $r = 0.99$, adjusted $P < 0.05$). *Abscisic acid 8'-hydroxylase* catalyzes the first step in the oxidative degradation of (+)-ABA (Krochko et al., 1998). Among the four ABA 8'-hydroxylase genes in *Arabidopsis*, *CYP707A3* supports the drought tolerance of plants via maintenance of ABA levels (Umezawa et al., 2006).

Besides intact LTR retrotransposons, truncated elements and solo-LTRs may also affect expression of nearby genes. We then examined expression changes of all the truncated retrotransposons (3,279) and solo-LTRs (586) as well as expression changes of their nearby genes in *S. alba* as described above. Similar with the intact elements, the truncated LTR retrotransposons were expressed more in roots (615) than in leaves (355), and exhibited more upregulation ($\log_2$ fold change > 1) than downregulation ($\log_2$ fold change > -1) upon high salinity treatment (Supplementary Data: Figure S2). In contrast, solo-LTRs were rarely expressed (Supplementary Data: Figure S2). A total of 24 truncated retrotransposons and one

solo-LTR were significantly mis-expressed in roots or leaves between treatments. Two TE-gene pairs were found to have significant expression correlation across conditions of different salt treatments. One pair was identified in leaves, involving a truncated RLG_6 copy (RLG_6_T1) and its 1306-bp downstream gene *thylakoid soluble phosphoprotein of 9 kDa* (*TSP9*) (Pearson's correlation, $r = 0.99$, adjusted $P < 0.01$). *TSP9*, is a plant-specific protein in the photosynthetic thylakoid membrane that regulates light harvest in *Arabidopsis* (Carlberg et al., 2003; Fristedt et al., 2009). The other pair was identified in roots, involving another truncated RLG_6 copy (RLG_6_T2) and its 705-bp upstream gene *glutamate-tRNA ligase*, also known as *glutamyl-tRNA synthetase* (Pearson's correlation, $r = -0.99$, adjusted $P < 0.05$).

Taken together, we identified four TE-gene pairs with significant expression correlation in *S. alba* that involve both intact and truncated LTR retrotransposons. Among the *S. caseolaris* orthologs of these four genes, only *TSP9* had a truncate RLG_6 copy inserted in its downstream region. However, none of the RLG_6 copies was expressed in leaves of *S. caseolaris* under normal condition (Wang et al., unpublished data). Orthologs of the other three genes in *S. caseolaris* had no insertion of RLC_4, RLG_8 or RLG_6 (BLASTn, all $e\text{-value} > 0.05$) in their flanking 10-kb up- or downstream region.

Potential *cis*- and *trans*-acting roles of TEs on gene expression in *S. alba*

It is very well known that LTRs contain CREs that can confer stress-inducible expression pattern on TEs and their flanking genes (Cavrak et al., 2014). To infer the potential *cis*-acting role of LTR retrotransposons, we predicted CREs in LTRs for each family using PlantCARE

(Lescot et al., 2002). Consensus sequences of *Gypsy* and *Copia* families that have more than ten copies were used in this analysis. We identified abundant CREs responsive to various biotic and abiotic stresses in LTRs, whereas none of these CREs is responsive to salt stress (Figure 6). RLG_1 and RLG_8 contained CREs responsive to anoxic, drought, low temperature, pathogen and fungal elicitor, which are not more abundant than CREs residing in LTRs of other families (Figure 6). Interestingly, the LTRs of RLG_8 harboring three ABA-responsive elements (ABREs) involved in ABA and drought responsiveness (Figure 6) is coincident with a RLG_8 copy positively co-expressed with *Abscisic acid 8'-hydroxylase 3* the gene nearby (see above). ABREs and MYB binding sites (MBSs) involved in drought inducibility were also found in the other three retrotransposons that showed significant expression correlation with their flanking genes (Figure 6). These results suggest that *cis*-regulatory role of LTR retrotransposons in *S. alba*, although rare, might occur via cross talk between drought and salt stress signaling pathways.

Besides *cis*-acting regulation, TEs can also function in *trans*, either resembling microRNA (miRNA) genes (Li et al., 2011) or acting as miRNA sponges competing with target transcripts of the same miRNA (Cho and Paszkowski, 2017; Cho, 2018; Palazzo and Koonin, 2020). To test these possibilities in *S. alba*, we annotated known and novel miRNAs using small RNA sequencing data described above with miRDeep2 (Friedländer et al., 2012) and predicted miRNA targets using psRNATarget (Dai and Zhao, 2011). An expectation score of 2.5 was used as the cutoff in the target prediction. All LTR retrotransposons including the intact and truncated elements and solo-LTRs were used in these analyses. We identified 105 known and 14 novel miRNAs expressed (Reads per million mapped reads, RPM > 1) in both

biological replicates; none was derived from TEs (Supplementary Data: Table S5). We then used the criteria described by Cho and Paszkowski (2017) to identify TE-gene pairs with potential miRNA competition. Among the 65 LTR retrotransposons predicted to contain potential miRNA binding sites, only 21 were expressed in roots or leaves of *S. alba* under either condition of salt treatments (Supplementary Data: Table S6). Requiring co-expression of both TEs and genes, we identified six TE-gene pairs with sequence matching in sense orientation and miRNA-binding sites within the matching regions. Nevertheless, none of them showed significantly expression correlation across different salt treatment conditions (Pearson correlation, all $P > 0.05$, Supplementary Data: Table S7). Therefore, *trans*-acting role of LTR retrotransposons is negligible in *S. alba*.

DISCUSSION

In her seminal studies of maize, McClintock proposed that TEs are “controlling elements” that may reprogram genome responses to environmental challenges (McClintock, 1984). Although still controversial, this idea has been supported by increasing empirical evidence (Naito et al., 2009, Cavrak et al., 2014, Horvath et al., 2017). TEs can be activated by stress, generate genetic variation, and modulate gene expression (Feschotte, 2008, Henaff et al., 2014). While induced TE activity has long been thought to benefit the host, it is puzzling that mangroves have experienced convergent reduction of TE loads despite being adapted to extreme environments with limited standing genetic variation (Guo et al., 2018, Lyu et al., 2018). How often TE activities are regulated by environmental cues and how these activities may contribute to a significant adaptive genome response in natural environments

440 remain intriguing questions.

441 In asking whether divergent habitats promote differential TE accumulation, we found
 442 TEs that were accumulating more in *S. alba* than in *S. caseolaris* despite the overall tendency
 443 of unloading the TE burden in mangroves (Figure 1A,B). This result, together with the fact
 444 that *S. alba* faces higher salinity than *S. caseolaris*, is consistent with the view that TEs tend
 445 to be active in plants growing in challenging habitats. The large fraction of young copies in *S.*
 446 *alba* (Figure 1C,D) suggests that TE originations are more frequent in *S. alba* than *S.*
 447 *caseolaris*. As young elements do not have enough time to degenerate, genome purging
 448 mechanisms should have little effect, if any, on their copy number. Although we do not have
 449 direct evidence that increased mobility in *S. alba* is triggered by environmental stress,
 450 transcriptome analyses suggest that more TEs are constitutively expressed in roots than
 451 leaves (Figure 5A), supporting this idea. Interestingly, purifying selection against TE
 452 insertions was even stronger in *S. alba* than in *S. caseolaris* (Table 1). The high origination
 453 rate and strong purifying selection might have led to fast turnover of TEs in *S. alba*, echoing
 454 a previous report of constant TE-host conflict masked by unloading of the TE burden in *R.*
 455 *apiculata* (Wang et al., 2018). Difference in unequal homologous recombination, as inferred
 456 by the S:I ratio, (Table 2) may further contribute to the differential TE accumulation between
 457 the two *Sonneratia* species.

458 The recent burst of TE abundance is particularly prominent for the *Gypsy*
 459 retrotransposons in *S. alba*. We found that the two most abundant *Gypsy* families in *S. alba*
 460 (RLG_1 and RLG_8) mainly accumulate in gene-poor regions. This is likely due to a balance
 461 between TE integration site preference and post-integration selection processes (Sultana et al.,

2017). The presence of the CHD domain in RLG_1 and RLG_8 likely drive heterochromatic integration site preference. However, the same integration site preference should be maintained in *S. caseolaris* if the CHD domain is sufficient to locate TE insertions to the same genomic region. Therefore, integration site preference alone cannot explain the difference in TE originations between species. Alternatively, if the two families differed in integration site preference between species, e.g. because of divergence of host proteins involved in TE tethering, the amplified copies should be much older than our observation (< 1 Myr), given that *S. alba* and *S. caseolaris* have diverged for ~11.6 Myr. We propose that the differential accumulation of RLG_1 and RLG_8 between species was probably triggered by environmental stress. It is possible that severe environments reactivated TEs more in *S. alba* than in *S. caseolaris* and then heterochromatin provided a “safe haven” for the accidentally inserted copies, allowing them to proliferate rapidly with relaxed epigenetic repression. The synergy between these mechanisms may constitute a self-reinforcing and self-perpetuating cycle of TE origination leading to long-term maintenance of genetic diversity. This hypothesis needs to be rigorously tested using data on stress-activated retrotransposition. We do see more up- than down-regulation of RLG_1 and RLG_8 during salt stress (Supplementary Data: Figure S3), suggesting stress can activate TE transcription. The presence of CREs responsive to anoxia and drought but not to salt in the LTRs of RLG_1 and RLG_8 suggests that the cross-talk between different stress signaling pathways might play a significant role in the potential TE activation in *S. alba*.

Although stress-induced TE activity in *S. alba* is plausible, our transcriptome analyses suggest that TE expansion into regulatory sequences is rare. This is evident in the observation

that TEs are mainly expressed in roots whereas the vast majority of differentially expressed TE-adjacent genes were found in leaves (Figure 5). Comparative transcriptome analyses between *S. alba* and *S. caceolaris* also suggest that TE expansion in *S. alba* influences between-species expression divergence mainly via carrying repressive epigenetic markers rather than creating new regulatory elements (Supplementary Note). There are only four potential cases of TE co-option in *S. alba* (Supplementary Data: Table S4). In one case, unlike constitutive expression of *CYP707A3* in *Arabidopsis*, *Abscisic acid 8'-hydroxylase 3* in *S. alba* and the RLG_8 copy upstream of it exhibited salt-induced expression in leaves only under 500mM NaCl, leading to a significant positive correlation in their expression levels (Supplementary Data: Table S4). This result together with the presence of ABREs in the LTRs of RLG_8 suggests that the RLG_8 copy might have been integrated into a stress-responsive gene regulatory network involving ABA catabolism. In *Arabidopsis*, *cyp707a3* mutant plants exhibit enhanced drought tolerance (Umezawa et al., 2006). Our result might also suggest that the strong salt tolerance of *S. alba* is achieved via reduced ABA degradation. We were unable to make any biological inferences in the other case of the RLC_4 copy, since the gene next to it has no known function. The rest two cases both involved truncated RLG_6 copies that possess tens of MBSs involved in drought-inducibility (Figure 6). The positive expression correlation of one RLG_6 copy with *TSP9* in leaves and the negative expression correlation of another RLG_6 copy with *glutamate-tRNA ligase* in roots suggest that TEs might mediate the cross-talk between photosynthesis and nitrogen metabolism during the adaptation of *S. alba* to mangrove swamps.

It should be noted that a retrotransposon is potentially responsive to diverse stress

conditions. As we only examined one stress condition (salinity), it is possible that co-option of TEs under other stress conditions was missed in this study. Nevertheless, it has been shown that enhancers often evolve from ancient TE sequences (Lynch et al., 2015, Villar et al., 2015). The evolutionary age of a TE was found to correlate positively with its likelihood of contributing to a regulatory element in human and mouse (Simonti et al., 2017). As the vast majority of retrotransposons in *S. alba* are relatively young, they may not have had enough time to get integrated into gene regulatory networks unless strong selection favors maintenance of the TE insertions, such as the *Hopscotch* insertion increasing *tb1* expression in maize (Studer et al., 2011). Considering the small effective population size of *S. alba* (Zhou et al., 2007, Zhou et al., 2011), natural selection may not be efficient enough to drive rapid fixation of TE insertions even though some TEs are capable of modulating expression of nearby genes.

In summary, our results together with a previous study (Wang et al., 2018) suggest that expansion of *Gypsy* retrotransposons, probably triggered by stress, is prevalent in mangroves despite the overall tendency of unloading the TE burden. The extent of TE proliferation across families is influenced by integration site preference and is subject to epigenetic repression and genome purging mechanisms. Although genetic variation generated by TE activities may be beneficial to host adaption, TE exaptation in *S. alba* is rare at least in terms of salt stress. Further experiments are needed to test the hypothesis of rapid transcriptional regulatory rewiring mediated by TE mobilization even though this hypothesis is appealing for extremophiles in which stress-induced reactivation of TEs is thought to be frequent.

FUNDING

This study was funded by the National Key Research and Development Program of China (2017YFC0506101), the National Science Foundation of China (31770246 and 31970245), the Science and Technology Program of Guangzhou (201707020035), the Fundamental Research Funds for the Central Universities (20lgpy117), the program of Guangdong Key Laboratory of Plant Resources (PlantKF05), the China Postdoctoral Science Foundation (2019TQ0391) and the Chang Hungta Science Foundation of Sun Yat-sen University.

LITERATURE CITED

- Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31: 166-169.
- Ball MC, Pidsley SM. 1995. Growth Responses to Salinity in Relation to Distribution of Two Mangrove Species, *Sonneratia alba* and *S. lanceolata*, in Northern Australia. *Functional Ecology* 9: 77.
- Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mobile DNA* 6: 11.
- Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* 57: 289-300.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30: 2114-2120.
- Boratyn GM, Schaffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. 2012. Domain

enhanced lookup time accelerated BLAST. *Biology Direct* 7: 12.

Bourque G, Burns KH, Gehring M, *et al.* 2018. Ten things you should know about transposable elements. *Genome Biology* 19: 199.

Bunt JS, Williams WT, Clay HJ. 1982. River water salinity and the distribution of mangroves species along several rivers in north Queensland. *Australian Journal of Botany* 30: 401-412.

Butelli E, Licciardello C, Zhang Y, *et al.* 2012. Retrotransposons control fruit-specific, cold-dependent accumulation of anthocyanins in blood oranges. *Plant Cell* 24: 1242-1255.

Capy P, Gasperi G, Biemont C, Bazin C. 2000. Stress and transposable elements: co-evolution or useful parasites? *Heredity (Edinb)* 85 (Pt 2): 101-106.

Carlberg I, Hansson M, Kieselbach T, Schröder WP, Andersson B, Vener AV. 2003. A novel plant protein undergoing light-induced phosphorylation and release from the photosynthetic thylakoid membranes. *Proceedings of the National Academy of Sciences of the United States of America*. 100: 757-762.

Casacuberta E, Gonzalez J. 2013. The impact of transposable elements in environmental adaptation. *Molecular Ecology* 22: 1503-1517.

Cavrak VV, Lettner N, Jamge S, Kosarewicz A, Bayer LM, Mittelsten Scheid O. 2014. How a retrotransposon exploits the plant's heat stress response for its activation. *PLoS Genetics* 10: e1004115.

Cho J, Paszkowski J. 2017. Regulation of rice root development by a retrotransposon acting as a microRNA sponge. *Elife*. 6: e30038.

572 Cho J. 2018. Transposon-Derived Non-coding RNAs and Their Function in Plants. *Frontiers*
573 *in plant science* 9: 600.

574 Chuong EB, Elde NC, Feschotte C. 2017. Regulatory activities of transposable elements:
575 from conflicts to benefits. *Nature Reviews Genetics* 18: 71-86.

576 Coordinators NR. 2018. Database resources of the National Center for Biotechnology
577 Information. *Nucleic Acids Research* 46: D8-D13.

578 Dai X, Zhao PX. 2011. psRNATarget: a plant small RNA target analysis server. *Nucleic Acids*
579 *Research*. 39(Web Server issue): W155-W159.

580 de Souza FS, Franchini LF, Rubinstein M. 2013. Exaptation of transposable elements into
581 novel cis-regulatory elements: is the evidence always strong? *Molecular Biology and*
582 *Evolution* 30: 1239-1251.

583 Devos KM, Brown JK, Bennetzen JL. 2002. Genome size reduction through illegitimate
584 recombination counteracts genome expansion in Arabidopsis. *Genome Research* 12:
585 1075-1079.

586 Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
587 throughput. *Nucleic Acids Research* 32: 1792-1797.

588 Fedoroff NV. 2012. Presidential address. Transposable elements, epigenetics, and genome
589 evolution. *Science* 338: 758-767.

590 Feng X, Xu S, Li J, *et al.* 2020. Molecular adaptation to salinity fluctuation in tropical
591 intertidal environments of a mangrove tree *Sonneratia alba*. *BMC Plant Biology* 20:
592 178.

593 Feschotte C. 2008. Transposable elements and the evolution of regulatory networks. *Nature*

594 *Reviews Genetics* 9: 397-405.

595 Finnegan DJ. 1992. *Transposable elements. The Genome of Drosophila melanogaster*, 1st edn.

596 New York: Academic Press.

597 Friedländer MR, Mackowiak SD, Li N, Chen W, Rajewsky N. 2012. miRDeep2 accurately

598 identifies known and hundreds of novel microRNA genes in seven animal clades.

599 *Nucleic Acids Research* 40: 37-52.

600 Fristedt R, Carlberg I, Zygadlo A, *et al.* 2009. Intrinsically unstructured phosphoprotein TSP9

601 regulates light harvesting in *Arabidopsis thaliana*. *Biochemistry*. 48: 499-509.

602 Fultz D, Choudury SG, Slotkin RK. 2015. Silencing of active transposable elements in plants.

603 *Current Opinion in Plant Biology* 27: 67-76.

604 Gao X, Hou Y, Ebina H, Levin HL, Voytas DF. 2008. Chromodomains direct integration of

605 retrotransposons to heterochromatin. *Genome Research* 18: 359-369.

606 Grandbastien MA. 2015. LTR retrotransposons, handy hitchhikers of plant regulation and

607 stress response. *Biochimica et Biophysica Acta* 1849: 403-416.

608 Guo Z, Li X, He Z, *et al.* 2018. Extremely low genetic diversity across mangrove taxa

609 reflects past sea level changes and hints at poor future responses. *Global Change*

610 *Biology* 24: 1741-1748.

611 Han Y, Burnette JM, 3rd, Wessler SR. 2009. TARGeT: a web-based pipeline for retrieving

612 and characterizing gene and transposable element families from genomic sequences.

613 *Nucleic Acids Research* 37: e78.

614 Han Y, Wessler SR. 2010. MITE-Hunter: a program for discovering miniature inverted-repeat

615 transposable elements from genomic sequences. *Nucleic Acids Research* 38: e199.

616 Henaff E, Vives C, Desvoyes B, *et al.* 2014. Extensive amplification of the E2F transcription
617 factor binding sites by transposons during evolution of Brassica species. *The Plant*
618 *Journal* 77: 852-862.

619 Horvath V, Merenciano M, Gonzalez J. 2017. Revisiting the Relationship between
620 Transposable Elements and the Eukaryotic Stress Response. *Trends in Genetics* 33:
621 832-841.

622 Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple
623 sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30:
624 3059-3066.

625 Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre2 web portal for
626 protein modeling, prediction and analysis. *Nature Protocols* 10: 845-858.

627 Kimura M. 1980. A simple method for estimating evolutionary rates of base substitutions
628 through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*
629 16: 111-120.

630 Kozomara A, Birgaoanu M, Griffiths-Jones S. 2019. miRBase: from microRNA sequences to
631 function. *Nucleic Acids Research*. 47: D155-D162.

632 Krochko JE, Abrams GD, Loewen MK, Abrams SR, Cutler AJ. 1998. (+)-Absciscic acid
633 8'-hydroxylase is a cytochrome P450 monooxygenase. *Plant Physiology* 118:
634 849-860.

635 Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for
636 Bisulfite-Seq applications. *Bioinformatics* 27: 1571-1572.

637 Kumar A, Bennetzen JL. 1999. Plant retrotransposons. *Annual Review of Genetics* 33:

638 479-532.

639 Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient
640 alignment of short DNA sequences to the human genome. *Genome Biology* 10: R25.

641 Law JA, Jacobsen SE. 2010. Establishing, maintaining and modifying DNA methylation
642 patterns in plants and animals. *Nature Reviews Genetics* 11: 204-220.

643 Lescot M, Déhais P, Thijs G, *et al.* 2002. PlantCARE, a database of plant cis-acting regulatory
644 elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic
645 Acids Research* 30: 325-327.

646 Li Y, Li C, Xia J, Jin Y. 2011. Domestication of Transposable Elements into MicroRNA
647 Genes in Plants. *PLoS ONE* 6: e19212.

648 Lisch D. 2013. How important are transposons for plant evolution? *Nature Reviews Genetics*
649 14: 49-61.

650 Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
651 RNA-seq data with DESeq2. *Genome Biology* 15: 550.

652 Lynch VJ, Nnamani MC, Kapusta A, *et al.* 2015. Ancient transposable elements transformed
653 the uterine regulatory landscape and transcriptome during the evolution of
654 mammalian pregnancy. *Cell Reports* 10: 551-561.

655 Lyu H, He Z, Wu CI, Shi S. 2018. Convergent adaptive evolution in marginal environments:
656 unloading transposable elements as a common strategy among mangrove genomes.
657 *New Phytologist* 217: 428-438.

658 Ma J, Bennetzen JL. 2004. Rapid recent growth and divergence of rice nuclear genomes.
659 *Proceedings of the National Academy of Sciences of the United States of America* 101:

12404-12410.

McClintock B. 1984. The significance of responses of the genome to challenge. *Science* 226: 792-801.

Miele V, Penel S, Duret L. 2011. Ultra-fast sequence clustering from similarity networks with SiLiX. *BMC Bioinformatics* 12: 116.

Myburg AA, Grattapaglia D, Tuskan GA, *et al.* 2014. The genome of *Eucalyptus grandis*. *Nature* 510: 356-362.

Naito K, Zhang F, Tsukiyama T, *et al.* 2009. Unexpected consequences of a sudden and massive transposon amplification on rice gene expression. *Nature* 461: 1130-1134.

Niu XM, Xu YC, Li ZW, *et al.* 2019. Transposable elements drive rapid phenotypic variation in *Capsella rubella*. *Proceedings of the National Academy of Sciences of the United States of America* 116: 6908-6913.

Palazzo AF, Koonin EV. 2020. Functional Long Non-coding RNAs Evolve from Junk Transcripts. *Cell* 183: 1151-1161.

Paro R, Hogness DS. 1991. The Polycomb protein shares a homologous domain with a heterochromatin-associated protein of *Drosophila*. *Proceedings of the National Academy of Sciences of the United States of America* 88: 263-267.

Price MN, Dehal PS, Arkin AP. 2009. FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Molecular Biology and Evolution* 26: 1641-1650.

Rahim AC, Bakar MFA. 2018. *Pidada—Sonneratia caseolaris*. *Exotic Fruits*, 1st edn. New York: Academic Press.

682 Rambaut A. 2012. FigTree v.1.4.2: Tree Figure Drawing Tool.
683 <http://tree.bio.ed.ac.uk/software/figtree>. Accessed 23 Jun. 2020.

684 Rey O, Danchin E, Mirouze M, Loot C, Blanchet S. 2016. Adaptation to Global Change: A
685 Transposable Element-Epigenetics Perspective. *Trends in Ecology and Evolution* 31:
686 514-526.

687 SanMiguel P, Gaut BS, Tikhonov A, Nakajima Y, Bennetzen JL. 1998. The paleontology of
688 intergene retrotransposons of maize. *Nature Genetics* 20: 43-45.

689 Schrader L, Kim JW, Ence D, *et al.* 2014. Transposable element islands facilitate adaptation
690 to novel environments in an invasive species. *Nature Communications* 5: 5495.

691 Simonti CN, Pavlicev M, Capra JA. 2017. Transposable Element Exaptation into Regulatory
692 Regions Is Rare, Influenced by Evolutionary Age, and Subject to Pleiotropic
693 Constraints. *Molecular Biology and Evolution* 34: 2856-2869.

694 Smit A, Hubley R, Green P. 2013-2015. RepeatMasker Open-4.0.
695 <http://www.repeatmasker.org/>. Accessed 23 Jun. 2020.

696 Studer A, Zhao Q, Ross-Ibarra J, Doebley J. 2011. Identification of a functional transposon
697 insertion in the maize domestication gene *tb1*. *Nature Genetics* 43: 1160-1163.

698 Sultana T, Zamborlini A, Cristofari G, Lesage P. 2017. Integration site selection by
699 retroviruses and transposable elements in eukaryotes. *Nature Reviews Genetics* 18:
700 292-308.

701 Tomlinson P. 1986. The botany of mangroves, 2nd edn. Cambridge: Cambridge University
702 Press.

703 Umezawa T, Okamoto M, Kushiro T, *et al.* 2006. CYP707A3, a major ABA 8'-hydroxylase

involved in dehydration and rehydration response in *Arabidopsis thaliana*. *The Plant Journal* 46: 171-182.

Villar D, Berthelot C, Aldridge S, *et al.* 2015. Enhancer evolution across 20 mammalian species. *Cell* 160: 554-566.

Wang Y, Liang W, Tang T. 2018. Constant conflict between Gypsy LTR retrotransposons and CHH methylation within a stress-adapted mangrove genome. *New Phytologist* 220: 922-935.

Wen M, Lin X, Xie M, *et al.* 2016. Small RNA transcriptomes of mangroves evolve adaptively in extreme environments. *Scientific Reports* 6: 27551.

Wicker T, Sabot F, Hua-Van A, *et al.* 2007. A unified classification system for eukaryotic transposable elements. *Nature Reviews Genetics* 8: 973-982.

Wright SI, Agrawal N, Bureau TE. 2003. Effects of recombination rate and gene density on transposable element distributions in *Arabidopsis thaliana*. *Genome Research* 13: 1897-1903.

Xiong W, He L, Lai J, Dooner HK, Du C. 2014. HelitronScanner uncovers a large overlooked cache of Helitron transposons in many plant genomes. *Proceedings of the National Academy of Sciences of the United States of America* 111: 10263-10268.

Xu S, He Z, Guo Z, *et al.* 2017. Genome-Wide Convergence during Evolution of Mangroves from Woody Plants. *Molecular Biology and Evolution* 34: 1008-1015.

Yang G, Zhou R, Tang T, Shi S. 2008. Simple and efficient isolation of high-quality total RNA from *Hibiscus tiliaceus*, a mangrove associate and its relatives. *Preparative Biochemistry and Biotechnology* 38: 257-264.

726 Yang Y, Duke NC, Peng F, *et al.* 2016. Ancient Geographical Barriers Drive Differentiation
727 among *Sonneratia caseolaris* Populations and Recent Divergence from *S. lanceolata*.
728 *Frontiers in Plant Science* 7: 1618.

729 Zhou R, Zeng K, Wu W, *et al.* 2007. Population genetics of speciation in nonmodel
730 organisms: I. Ancestral polymorphism in mangroves. *Molecular Biology and*
731 *Evolution* 24: 2746-2754.

732 Zhou R, Ling S, Zhao W, *et al.* 2011. Population genetics in nonmodel organisms: II. natural
733 selection in marginal habitats revealed by deep sequencing on dual platforms.
734 *Molecular Biology and Evolution* 28: 2833-2842.

735

Tables

Table 1. Location of TE insertions in relation to coding regions in *S. alba* and *S. caseolaris*.

	<i>S. alba</i>			<i>S. caseolaris</i>		
	Intergenic	Intron	Coding	Intergenic	Intron	Coding
<i>Copia</i>	13.5***(1.43) [†]	5.8**(0.31)	4.1***(0.06)	20.0(0.95)	8.4(0.09)	8.0(0.07)
<i>Gypsy</i>	38.0(4.12)	12.4(0.74)	7.4***(0.18)	36.7(1.07)	13.8(0.15)	15.3(0.14)
DNA transposon	55.8***(0.91)	23.2**(0.31)	17.2***(0.23)	83.1(0.85)	29.7(0.24)	30.1(0.26)
Others	112.4***(5.10)	41.0(1.33)	14.7***(0.25)	146.2(4.43)	41.0(0.81)	31.6(0.41)

[†]Upper values in each cell are the number of elements per megabase of DNA in each category and values in parentheses are the proportion of DNA in each location occupied by TEs. Significance levels are shown for comparisons between *S. alba* and *S. caseolaris*, calculated using the Mann-Whitney *U*-test. **, $P < 0.01$; ***, $P < 0.001$.

742

743 **Table 2.** Comparison of the ratios of solo-LTRs to intact elements (S:I) for *Copia* and *Gypsy*

744 LTR retrotransposons between *S. alba* and *S. caseolaris*.

	Superfamily	Solo-LTR (S)	Intact Element (I)	S:I
<i>S. alba</i>	<i>Copia</i>	322	180	1.79
	<i>Gypsy</i>	264	767	0.34
<i>S. caseolaris</i>	<i>Copia</i>	120	49	2.45
	<i>Gypsy</i>	283	76	3.72

745

746

747 **Figure legends**

748 **Figure 1.** TE composition in *Sonneratia alba* and *S. caseolaris*. (A) Bar plot shows the
749 proportions of transposon groups in *S. alba*, *S. caseolaris*, and *Eucalyptus grandis* (the
750 outgroup). Tree topology is adopted from Lyu et al. (2018). (B) Histogram shows the
751 distribution of transposon contents across families between *S. alba* and *S. caseolaris*. (C)
752 Histogram shows the copy number of *Copia* and *Gypsy* elements in *S. alba* and *S. caseolaris*
753 grouped by age. (D) Line graph shows the relative proportion of total *Copia* and *Gypsy*
754 elements in different age bins.

755 **Figure 2.** Relative proportion and chromosome distribution of LTR retrotransposon families
756 in *S. alba*. (A) Proportions of intact copy numbers from different LTR retrotransposon
757 families ranked in a descending order. Note that RLG_1 and RLG_8 are the top two abundant
758 LTR retrotransposons in *S. alba*. (B) Chromosome distribution of coding genes (the upper
759 row), the top four abundant LTR retrotransposon families (RLG_1, RLG_8 and RLG_4 of
760 *Gypsy* and RLC_6 of *Copia*), and all TEs along the seven largest scaffolds of the *S. alba*
761 genome. The putative heterochromatin regions (gene density < 0.05) are indicated with the
762 grey shadow. Note that the distributions of RLG_1 and RLG_8 elements track the genomic
763 locations of putative heterochromatic regions but no such pattern is observed for RLG_4 and
764 RLC_6.

765 **Figure 3.** Amino acid sequence alignment and structure prediction of the chromodomain. (A)
766 Multiple alignments of amino acid sequences of known chromodomains and chromo-like
767 motifs from the RLG_1 and RLG_8 families. A schematic representation of β -strands

(arrows) and α -helix (cylinder) is on the bottom. The GenBank accession numbers for the sequences used are: DmHP1(KNA_A)-M57574 from *Drosophila melanogaster*; MAGGY-L35053 and Pyret-AB062507 from *Magnaporthe grisea*; Cft1-AF051915 from *Cladosporium fulvum*; PpatensLTR1-XM_001752430 from *Physcomitrella patens*; Tf2-12-RVW30894 from *Vitis vinifera*; sr35-AC068924, rn8-AK068625 and RIRE3-AC119148 from *Oryza sativa*; Tekay-AF050455 from *Zea mays*; RetroSor2-AF061282 from *Sorghum bicolor*; Tma-AF147263 from *Arabidopsis thaliana*. (B) Predicted three-dimensional structure of chromodomains from the DmHP1 protein (M57574) and the RLG_1 and RLG_8 families.

Figure 4. Epigenetic repression and insertional locations of RLG_1 and RLG_8 in *S. alba*.

(A) Correlations between CHH methylation level, 24-nt siRNA abundance of the RLG_1 and RLG_8 elements, and gene density of the 100 Kb region in which each retroelement is located in the *S. alba* genome. Each dot represents an intact RLG_1 or RLG_8 element and is colored in red to black gradient according to gene density of the genomic region in which the element is located. (B) Age distribution of the RLG_1 or RLG_8 elements along genomic regions with different gene density. Contour lines represent estimated two dimensional kernel density.

Figure 5. Expression patterns of TEs and their flanking genes under salt treatments in *S. alba*.

(A) Fold changes of the expression levels of intact *Gypsy* and *Copia* retrotransposons across salt treatments (the upper: 0 mM vs 250 mM NaCl, the lower: 250 mM vs 500 mM NaCl). (B) Expression levels of genes neighboring copies of the four most abundant LTR retrotransposon families in comparison with the whole transcriptome. (C) Fold expression

change of genes next to copies of the four most abundant LTR retrotransposon families across salt treatments (the upper: 0 mM vs 250 mM NaCl, the lower: 500 mM vs 250 mM NaCl). Genes with significant mis-expression ($FDR < 0.05$ and ≥ 2 -fold change) between treatments are indicated in red.

Figure 6. Summary of *cis*-acting regulatory elements (CREs) in the LTRs of *Gypsy* and *Copia* families with more than ten copies in *S. alba*. Also shown are CREs in the full length of the two truncated elements (RLG_6_T1 and RLG_6_T2) which significantly correlated with their nearby genes in expression levels. Color depths and numbers indicate the abundance of specific CREs.

[Supplementary Information]

Figure S1. Maximum likelihood (ML) phylogenetic trees of *Copia* and *Gypsy* LTR retrotransposon families.

Figure S2. Expression fold changes of the truncated LTR retrotransposons and solo-LTRs as well as their nearest genes across salt treatments in leaves and roots. TEs or genes with significant mis-expression (Benjamini-Hochberg FDR < 0.05) between treatments are indicated in red. Green lines indicate two-fold expression changes.

Figure S3. Expression level fold change of intact RLG_1 and RLG_8 retrotransposons across salt treatments.

Table S1. List of all intact LTR retrotransposons in *Sonneratia alba* and *S. caseolaris*. (Excel)

Table S2. TE-adjacent genes in *Sonneratia alba* that were significantly mis-regulated by salt treatment. (Excel)

Table S3. The number of TE-adjacent genes in *Sonneratia alba* that were significantly mis-regulated by salt treatment.

Table S4. Correlations between expression levels of TEs and their nearby genes in *Sonneratia alba*. (Excel)

Table S5. Annotation of known and new miRNAs in leaves of *S. alba*. (Excel)

Table S6. LTR retrotransposons predicted to be miRNA targets in *S. alba*. (Excel)

Table S7. Pairs of LTR retrotransposons and genes with potential miRNA competition in *S.*

820 *alba*. Pearson correlation was calculated for expression levels of each pair across 0mM, 250
821 mM and 500mM salt treatments in roots and leaves, respectively. *cor*, Pearson correlation
822 coefficient; P, P value.

A **B**

A

chromodomains

Dml_HP1 (KNA_A)

EYAVEKIIDRRV-----RKGMVEYYL-KWKGYPETENTWEPENNL-DCQDLIQQYEA-----

Group I chromodomains of LTR retrotransposons

MAGGY

EYVEEILDSFWETRGRGGRRLKY-IVRWAGY--SEPTTEPADYLENAAQLVKNFHRR-----

Pyret

TYNVKQIFDHRNRHG-----KIKY-FIKWENYGHEKNIWEPLNHLQDCQEPRLQYYQE-----

Cft1

EFEVEKILDKKGQR-----Y-LVKWKGYDESENTWEPRINLANCYQLLRQFQKW-----

PpatensLTR1

EFEVEEILDSRRCRN-----KLEY-LIHWRGYDISECTWEPSKNLANASAKVKFFAFV-----

Tf2-12 (RVW30894)

DREIEKILDHRTMGHSRKNRRTDf-LVQWKGISETEASWERDVT LWQFEKEVQAYWQSKSTR

RLG_1

DKEVEDILDHRVLGYSKMNRTEY-LVKWKDLPESEATWEKDVT LWQYEDLIRRYQESHSTR

RLG_8

EREVDRVLDRRVLGYSKKNRRTEY-LVLWKGLPMSEASWEKEATLWQFEDKIQEYLHSIPTR

Group II chromodomains of LTR retrotransposons

sr35

---PFQFLDKRLVKKGN-RSVLQL-LTHWYHSSPSESTWEDMEDLFARFPRALAW-----

rn8

--WPELILDRRLVKKGN-AAHVQV-LIKWSSLSADDA TWEDYDVLKTRFPSAPAWGQA-----

RIRE3

VEKPIRILETSERRTRN-RVIRFC-KVQWSNHSEESTWEREDELKSAHPHLF-----

Tekay

SEYPVRILETSRRITRS-KVINMC-KVQWSHHSEDEATWEREDELRAEFP-----

Retrosor2

REYPVRILDRATKETRS-STFPMC-KVLWSNHTEREATWEKESELQLRYPYPYLFERYV-----

Tma

EARPVRVLERRIKELRR-KKIPLI-KVLWDCDGVTEETWEPDARMKARFKKCF-----

B

Dml_PH1

RLG_1

RLG_8

A **B**

Promoter element		Intact														Truncated		Function
		RLG_8	RLG_1	RLG_6	RLC_4	RLG_9	RLG_15	RLC_22	RLG_3	RLG_13	RLG_5	RLC_11	RLG_19	RLG_2	RLC_7	RLG_6_T1	RLG_6_T2	
Promoter element	TATA-box	14	8	32	10	0	7	9	13	16	2	2	10	8	3	24	22	core promoter element around -30 of transcription start
	CAAT-box	6	19	21	14	5	5	4	21	30	2	9	4	6	12	31	28	common cis-acting element in promoter and enhancer regions
Stress responsiveness	W box	2	1	1	1	0	0	0	1	0	0	0	0	1	0	1	1	wounding and pathogen responsiveness
	Box-W1	2	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	fungal elicitor responsive element
	WUN-motif	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	wounding and pathogen responsive elements
	box E	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	cis-element for induction upon fungal elicitation
	CGTCA-motif	0	0	2	0	0	0	0	1	2	0	0	0	0	0	1	1	cis-acting regulatory element involved in the MeJA-responsiveness
	TGACG-motif	0	0	2	0	0	0	0	1	2	0	0	0	0	0	1	1	
	GCC box	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	wounding and pathogen responsiveness
	GC-motif	3	1	3	0	1	0	0	0	0	0	0	0	0	0	1	1	enhancer-like element involved in anoxic specific inducibility
	ARE	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0	2	cis-acting regulatory element essential for the anaerobic induction
	ABRE	3	0	1	1	0	0	0	0	8	0	0	0	2	0	3	2	abscisic acid (ABA) and drought responsiveness
	MBS	5	0	2	0	0	5	0	3	3	4	3	1	0	0	14	11	MYB binding site involved in drought-inducibility
	CCAAT-box	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	MYBHv1 binding site, stress responses and development
	LTR	1	1	1	0	1	0	0	1	0	0	0	0	0	0	2	1	cis-acting element involved in low-temperature responsiveness