Integrated population genomic analysis and numerical simulation to estimate larval dispersal of *Acanthaster cf. solaris* between Ogasawara and other Japanese regions

Mizuki Horoiwa¹, Takashi Nakamura², Hideaki Yuasa³, Rei Kajitani³, Yosuke Ameda⁴,
Tetsuro Sasaki⁴, Hiroki Taninaka⁵, Taisei Kikuchi⁶, Takehisa Yamakita⁷, Atsushi Toyoda⁸,
Takehiko Itoh³, Nina Yasuda^{9,*}

¹ Graduate School of Agriculture, University of Miyazaki, Miyazaki, Japan

² School of Environment and Society, Tokyo Institute of Technology, Tokyo, Japan

³ School of Life Science and Technology, Department of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan

⁴ Institute of Boninology, Chichijima, Ogasawara, Tokyo, Japan

⁵ Interdisciplinary Graduate School of Agriculture and Engineering, University of Miyazaki, Miyazaki,

⁶ Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

⁷ Japan Agency for Marine-Earth Science and Technology, Kanagawa, Japan

⁸ Department of Genomics and Evolutionary Biology, National Institute of Genetics, Shizuoka, Japan

⁹ Faculty of Agriculture, University of Miyazaki, Miyazaki, Japan

* Correspondence:

Corresponding Author

nina27@cc.miyazaki-u.ac.jp

Keywords: crown-of-thorn starfish, larval dispersal, secondary outbreak, population genetics, Ogasawara, Kuroshio current, oceanographic dispersal simulation

Word count: 4928

Number of figures and tables: 4

Article type: Full paper

25 Abstract

26 The estimation of larval dispersal of marine species occurring on an ecological timescale is
 27 significant for conservation. In 2018, a semi-population outbreak of crown of thorns starfish,
 28 *Acanthaster cf. solaris* was observed on a relatively isolated oceanic island, Ogasawara. The aim of
 29 this study was to assess whether this population outbreak was caused by large-scale larval
 30 recruitment (termed secondary outbreak) from the Kuroshio region. We estimated larval dispersal of
 31 the coral predator *A. cf. solaris* between the Kuroshio and Ogasawara regions using both population
 32 genomic analysis and oceanographic dispersal simulation. Population genomic analysis revealed
 33 overall genetically homogenized patterns among Ogasawara and other Japanese populations,
 34 suggesting that the origin of the populations in the two regions is the same. In contrast, a simulation
 35 of 26-year oceanographic dispersal indicated that larvae are mostly self-seeded in Ogasawara
 36 populations and have difficulty reaching Ogasawara from the Kuroshio region within one generation.
 37 However, a connectivity matrix produced by the larval dispersal simulation assuming a Markov chain
 38 indicated gradual larval dispersal migration from the Kuroshio region to Ogasawara in a stepping-
 39 stone manner over multiple years. These results suggest that, while large-scale larval dispersal from
 40 an outbreak of the Kuroshio population spreading to the Ogasawara population within one generation
 41 is unlikely. This study also highlighted the importance of using both genomic and oceanographic
 42 methods to estimate larval dispersal, which provides significant insight into larval dispersal that
 43 occurs on ecological and evolutionary timescales.

44

1 Introduction

Many benthic marine invertebrates in coral reef ecosystems exhibit planktonic larval dispersal during their early life history. Such larval dispersal connects different populations and forms a meta-population structure (Shanks, 2009). Thus, for effective conservation and management of benthic marine invertebrates, assessment of larval dispersal is important (Botsford et al., 2009; Almany et al 2009). Because it is challenging to directly track the movement of tiny, numerous larvae in the field, several different indirect methods, including population genomic analysis (e.g. Yasuda et al., 2015; Arndt and Smith, 2002), oceanographic simulation (e.g. Miyake et al., 2009; Storlazzi et al., 2017), plankton-netting in the field (e.g. Yasuda et al., 2015; Suzuki et al., 2016), and drifting buoys (e.g. Fukuda and Hanamura, 1996), have been employed to estimate larval dispersal. Although each method has advantages and disadvantages, the first two methods can be applied to relatively large spatial scales and thus have been used more frequently than the other methods. On the one hand, the data obtained by population genomic analysis reflect the biological processes of larval dispersal and recruitment. However, they not only include larval dispersal occurring on an ecological timescale but also reflect historical gene flow and genetic breaks caused by past climate change and plate tectonics (Benzie, 1999; Crandall et al., 2014). Oceanographic simulation, on the other hand, can capture snapshots of larval dispersal, while even a sophisticated biophysical model cannot accurately simulate larval behavior, recruitment, and survival in the ocean (White et al., 2019). It is therefore important to integrate different methods to estimate larval dispersal (Marko and Hart, 2017), although few studies have attempted to do so (but see Schunter et al., 2011; Alberto et al., 2011; Nakabayashi et al., 2019; Taninaka et al., 2019).

In this study, we targeted the crown-of-thorns starfish, *Acanthaster cf. solaris*, to estimate larval dispersal. Chronic outbreaks of the coral predator *A. cf. solaris* and its siblings have been the major management issue in coral reef ecosystems in the Indian and Pacific oceans (Baird et al., 1990; Baird et al., 2013). With high fecundity and a larval stage that lasts a few to several weeks (Yamaguchi, 1977; Lucas, 1973), a population outbreak can cause successive population outbreaks, termed secondary outbreaks (reviewed in Birkeland and Lucas, 1990; Pratchett et al., 2014; Yasuda, 2018), via larval dispersal in neighboring regions, especially in the western Pacific region. Assessment of the spatial range of secondary outbreaks of *A. cf. solaris* is necessary for coral reef conservation.

Many population genomic and phylogeographic studies of *A. cf. solaris* have been conducted using various molecular markers (e.g., Benzie, 1992, 1999; Vogler et al, 2013; Timmers et al., 2011; Yasuda et al., 2009; 2015). These studies have indicated strong gene flow over a long distance along western boundary currents such as the Kuroshio Current region (genetic homogeneity from across the Ryukyu Islands to the temperate Pacific coast of mainland Japan, Yasuda et al., 2009) and the Eastern Australian Current region (the Great Barrier Reef, Nash et al., 1988; Benzie et al., 1999; Vogler et al., 2013; Harrison et al., 2017), while relatively limited gene flow and genetic differentiation were observed among distant Pacific islands (Yasuda et al., 2009; Timmers et al., 2012; Vogler et al., 2013). Overall, the studies found strong gene flow and genetic homogeneity in regions where strong currents are associated, although genetic data strongly reflect historical gene flow that was formed during past climate change, and the spatial extent of secondary outbreaks could be overestimated (Benzie 1999, Yasuda et al. 2009, Yasuda et al. 2015). Larval dispersal simulation studies in the Great Barrier Reef (GBR) have indicated southward larval dispersal along the Eastern Australian Current, which is consistent with historically observed patterns of secondary outbreaks of *A. cf. solaris*. (Dight et al., 1990a, b, James and Scandol, 1992; Scandol and James, 1992). Plankton netting survey has also showed evidence of large-scale larval dispersal of *A. cf. solaris* in the GBR (Uthicke et al., 2015). While intensive connectivity studies of *A. cf. solaris* have been conducted at

the Indo-Pacific scale and in some localized regions where secondary outbreaks were suspected (e.g., the Kuroshio region and the GBR), connectivity between such regions and relatively isolated oceanic island regions is yet to be examined.

In 2018, a population outbreak of *A. cf. solaris* on Ogasawara Island was reported (Biodiversity Center of Japan, 2019), which was the second population outbreak on Ogasawara since 1979 (Kurata, 1984). Ogasawara Island is an oceanic island situated ca. 1000 km south of the Tokyo urban center and is out of the mainstream of the Kuroshio Current. It is therefore still unknown whether this population outbreak is attributed to a secondary outbreak in the Kuroshio region.

The aim of this study was to estimate larval dispersal between the Kuroshio region and Ogasawara to determine whether the population outbreak in Ogasawara was caused by a secondary outbreak in the Kuroshio region. To meet this aim, we used both a method of population genomic analysis called multiplexed ISSR genotyping by sequencing (MIG-seq) and oceanographic larval dispersal simulation based on the Global HYbrid Coordinate Ocean Model (HYCOM).

2 Material and Methods

2.1 Sampling and DNA extraction

A total of 187 starfish samples were collected from six Kuroshio regions (Tatsukushi, Miyazaki, Sakurajima, Onna Village, Miyako, and Sekisei Lagoon) and Ogasawara via scuba diving (Fig. 1). The tube feet were preserved in 99.5% ethanol prior to DNA extraction. Genomic DNA extraction was carried out using a modified heat alkaline method (Meeker et al., 2007; Nakabayashi et al., 2019).

2.2 Whole-genome sequencing and nuclear genome assembly as the reference

We obtained a new high-quality whole-genome sequence for mapping analysis. Whole-genome sequencing of the sample collected from Miyazaki (BioSample ID, SAMD00056692) was performed using Sequel, which is a single-molecule real-time sequencer from Pacific Biosciences (PacBio; Menlo Park, CA, USA), according to the manufacturer's protocol. Previously published whole-genome short-read sequencing data (Wada et al., 2020) were downloaded from the BioProject PRJDB4009 in the ENA/Genbank/DDBJ database. Briefly, we obtained two paired-end (PE) libraries (insert sizes: 300 and 500 bp) and six mate-pair (MP) libraries (insert sizes: 3k, 5k, 8k, 10k, 12k, and 15k).

For the *in silico* procedures in this section, default parameters were used except where otherwise noted. Illumina PE and MP reads were filtered and trimmed using Platanus_trim v1.0.7, (http://platanus.bio.titech.ac.jp/pltanus_trim). We assembled trimmed PE and MP reads using Platanus v1.2.5 (Kajitani et al., 2014) with the commands "assemble," "scaffold," and "gap_close," resulting in the scaffolds.

In addition to the short-read assembly, PacBio long-read assembly was conducted using Canu v1.7 (Koren et al., 2017). We polished the Canu contigs with the PacBio long reads using Pbalig v0.3.1 (<https://www.pacb.com/support/software-downloads/>) and Arrow in the GenomicConsensus package v2.2.2 (<https://github.com/PacificBiosciences/GenomicConsensus>). We additionally polished the contigs with Illumina (Illumina, San Diego, CA, USA) short reads using Bowtie2 v2.3.5.1 (Langmead and Salzberg, 2012) and Pilon v1.22 (Walker et al., 2014). The short read-based polishing was iterated until the number of corrected sites did not increase (number of iterations: 18).

Gaps of the short read-based scaffolds (Platanus result) were filled with polished long read-based contigs using an in-house program. The procedure is as follows (Suppl. Fig. 1):

(1) Fixed-length regions flanking the gaps in scaffolds were extracted.

(2) The flanking sequences were aligned (queried) to the long read-based contigs using Minimap2 v2.17 (Li, 2018) with the options of "-c -k 19."

(3) The alignment results were filtered out if sequence identity <90%, query coverage <25%, or multiple best-scoring hits were identified.

(4) Each gap was filled with a region between the alignments in the contig if the following conditions were satisfied:

(i) The flanking region pair was aligned to the same contig.

(ii) The distance between alignments <50 kbp.

(iii) The strands of the alignments were consistent.

(iv) There was no 'N' in the corresponding regions in the contig.

(5) Steps (1) to (4) were iterated eight times. Here, the specific length of the flanking regions was applied for each iteration (500, 1k, 5k, 10k, 20k, 40k, 80k, and 160k).

(6) Step (5) was iterated twice.

Finally, to eliminate contamination, we performed homology searches of the assembly against the NCBI genome database (human, bacteria, and virus), the mitochondrial genome of *A. planci* (accession no. NC007788), and the chromosome sequence of the COTS symbiont (COTS27, accession no. AP019861) using BLASTN v2.7.1 (Altschul et al., 1990). The assembled sequences (scaffolds or contigs) were eliminated as contamination if sequence identity $\geq 90\%$ and query coverage $\geq 50\%$ (Suppl. Table 1). We assessed the completeness of the draft genome using BUSCO v.4.0.6 with the Metazoa Odb10 dataset (Seppey et al. 2019). K-mer-based ($k = 17$) estimations of the sizes of the genome and duplicated regions were also conducted using Jellyfish v.2.2.10 (Marçais and Kingsford, 2011) and GenomeScope v.2.0 (Ranallo-Benavidez et al. 2020).

2.3 MIG-seq Analysis

MIG-seq is a population genomic method that can easily detect a moderate number of neutral single nucleotide polymorphisms (SNPs) (Suyama and Matsuki, 2015). Previous studies using MIG-seq on marine species successfully discovered species boundaries and genetic structures that were undetectable using traditional genetic markers such as mitochondrial DNA and nuclear ribosomal internal transcribed spacer 2 (Nakabayashi et al. 2019; Takata et al. 2019).

We used eight pairs of multiplex ISSR primers (MIG-seq primer set 1; Suyama and Matsuki, 2015) with the Multiplex PCR Assay Kit Ver. 2 (TaKaRa Bio Inc., Shiga, Japan) for the first PCR; we slightly modified the protocol by Suyama and Matsuki (2015) and used a total reaction volume of 7 μ L in a T100™ Thermal Cycler (Bio-Rad, Hercules, CA, USA) under the following conditions:

initial denaturation at 94 °C for 1 min; followed by 25 cycles of 94 °C for 30 s, 38 °C for 1 min, and 72 °C for 1 min; and final extension at 72°C for 10 min. The samples were indexed in the second PCR, which was performed with PrimeSTAR GXL DNA polymerase (TaKaRa Bio Inc.) using a total reaction volume of 12 µL in a thermal cycler under the following conditions: 21 cycles of 98 °C for 10 s, 54 °C for 15 s, and 68 °C for 1 min. PCR products were run on an agarose gel, and 350–800 bp PCR products were extracted from the gel. The gel-extracted DNA was pooled and sequenced using MiSeq Control Software v2.0.12 (Illumina) with the MiSeq Reagent v3 150 cycle kit (Illumina). Image analysis and base calling were performed using Real-Time Analysis Software v1.17.21 (Illumina).

To eliminate low-quality reads and primer sequences from the raw data, we used the FASTX-Toolkit v0.0.14 (Gordon and Hannon, 2012; http://hannonlab.cshl.edu/fastx_toolkit/index.html) with a fastq-quality-filter setting of $-q\ 30 -p\ 40$. We removed adapter sequences for the Illumina MiSeq run from both the 5' end (GTCAGATCGGAAGAGCACACGTCTGAACTCCAGTCAC) and the 3' end (CAGAGATCGGAAGAGCGTCGTGTAGGGAAAGAC) using Cutadapt v1.13 (Martin, 2011), and then we excluded short reads less than 80 bp. We used Stacks v2.2 (Catchen et al., 2013; Rochette and Catchen, 2017) to stack the reads and extract the SNPs. We used the newly sequenced *A. cf. solaris* genome as a reference, as described below. We used the Gstacks option (-rm, -pcr, -duplicates) in Stacks v2.2 to remove PCR duplicates by randomly discarding all but one pair of each set of reads. We used population software in Stacks to prepare the dataset. We used the minimum percentage of individuals required to process a locus across all data (r) equal to 0.95. After excluding individuals missing more than 10% of SNP data, we used a single SNP option to avoid strong linkage between loci for Structure analysis. We used all SNPs for the remaining genetic analyses.

2.4 Population Genetics

BayeScan v2.1 (Narum and Hess, 2011) was used to detect possible SNPs under natural selection with a default setting. Genetic diversity (expected and observed heterozygosities, H_E and H_O , respectively), Hardy-Weinberg Equilibrium, global F_{ST} including and excluding Ogasawara samples, and pairwise F_{ST} values across different populations were calculated using GenAlEx v6.502 (Peakall and Smouse, 2012) with 9999 permutations.

To test whether the Ogasawara population is genetically different from other populations, a hierarchical analysis of molecular variance (AMOVA) was conducted with prior grouping (Ogasawara vs other populations) using Arlequin v3.5 (Excoffier and Lischer, 2010).

Bayesian clustering analysis was conducted using Structure software v2.3.4 (Prichards et al., 2000) with 500,000 burn-in followed by 500,000 steps of Markov chain Monte Carlo simulations and 10 iterations. We used admixture- and allele frequency-correlated models that were expected to best explain the data set. The LOCPRIOR model (Hubisz et al. 2009) was used with Ogasawara vs other populations and without prior regional grouping, and the number of putative clusters (K) was set from 1 to 7. The online software Structure Harvester (Earl and von Holdt, 2012) was used to summarize the likelihood value. Structure plots were summarized using the online software CLUMPAK (Kopelman et al., 2015). To further visualize the genetic relationships among different populations, principal coordinate analysis (PCoA) was conducted using GenAlEx v6.502 based on pairwise genetic distances among populations.

PGDspider v2.0.8.3 (Lischer & Excoffier, 2012) was used to convert the data formats.

2.5 Oceanographic larval dispersal simulation

We conducted a Lagrangian larval dispersal simulation to elucidate the oceanographic connectivity between the regions. For this simulation, we used Connectivity Modeling System v2.0 (Paris et al., 2013) with the ocean analysis/reanalysis products produced by the Global Ocean Forecasting System 3.0 of Global HYCOM (Chassignet et al., 2007), which are 1/12° resolution (~9 km) daily products archived from 1993 to 2018 (<https://www.hycom.org/>). The surface water current and sea surface temperature (SST) in the Global HYCOM products ranged from 115°E–155°E longitude and 15°N–40°N latitude. Thirty source and sink sites fitting the purpose of this study were chosen to create source sink polygons (see Figure 1A). The particles regarded as COTS larvae were numerically released from ocean areas inside the polygons and tracked. Once the particle entered a polygon, the particle was removed and judged as larval settlement success, and the connection between the source polygon and sink polygon was counted. The particle release timing (start time of COTS mass spawning) and duration of the spawning at each site were determined by the following rules using a history of SST at the site provided by Global HYCOM analysis/reanalysis products based on Yasuda et al. (2010): (1) If SST at the site rose above 28 °C, the particles were released until SST dropped below 26 °C or 30 days passed; (2) If SST at the site did not reach 28 °C, the particles were released while SST was higher than 26 °C. A total of 100 particles per day were released from each source polygon under the above conditions. Finally, we obtained a connectivity matrix that was created based on larval dispersal simulations of 26 years (1993 to 2018).

Because the connectivity based on the larval dispersal simulation is a one-generation direct linkage, indicating that stepping-stone migration had not been considered, we conducted back-tracing of the origin site through a discrete-time Markov chain using the connectivity matrix. To this end, we first set the following form of the connectivity matrix $\mathbf{N}(m \times m)$:

$$\mathbf{N} = \begin{matrix} & \begin{matrix} \text{source} \\ n_{1,1} & \cdots & n_{1,j} & \cdots & n_{1,m} \\ \vdots & \ddots & & & \vdots \\ n_{i,1} & & n_{i,j} & & n_{i,m} \\ \vdots & & & \ddots & \vdots \\ n_{m,1} & \cdots & n_{m,j} & \cdots & n_{m,m} \end{matrix} \\ \begin{matrix} \vdots \\ n_{i,1} \\ \vdots \\ n_{m,1} \end{matrix} & \begin{matrix} \vdots \\ n_{i,j} \\ \vdots \\ n_{m,j} \end{matrix} & \begin{matrix} \vdots \\ n_{i,m} \\ \vdots \\ n_{m,m} \end{matrix} \end{matrix} \begin{matrix} \\ \\ \text{sink} \\ \end{matrix} \quad (1)$$

The matrix elements were normalized by the total number of sink particles as follows:

$$\mathbf{P} = \begin{matrix} & \begin{matrix} p_{1,1} & \cdots & p_{1,j} & \cdots & p_{1,m} \\ \vdots & \ddots & & & \vdots \\ p_{i,1} & & p_{i,j} & & p_{i,m} \\ \vdots & & & \ddots & \vdots \\ p_{m,1} & \cdots & p_{m,j} & \cdots & p_{m,m} \end{matrix} \\ \begin{matrix} p_{1,1} \\ \vdots \\ p_{i,1} \\ \vdots \\ p_{m,1} \end{matrix} & \begin{matrix} p_{1,j} \\ \vdots \\ p_{i,j} \\ \vdots \\ p_{m,j} \end{matrix} & \begin{matrix} p_{1,m} \\ \vdots \\ p_{i,m} \\ \vdots \\ p_{m,m} \end{matrix} \end{matrix}, \quad (2)$$

where

$$p_{i,j} = \frac{n_{i,j}}{\sum_{j=1}^m n_{i,j}}. \quad (3)$$

The normalized connectivity matrix can be regarded as the transition matrix of the Markov chain. The n -th power of the transition matrix $\mathbf{P}^n$ represents the n time-step back-trace to the origin source.

3 Results

3.1 Sequencing and SNP metrics

In total, 11,461,866 raw reads ranging from 9,190 to 281,474 reads per sample were obtained for 187 *A. cf. solaris* samples. After filtering out the low-quality reads, 6,125,996 reads ranging from 5,996 to 107,468 reads per sample remained. On average, 189,536 reads per individual remained after two-step filtering. We excluded seven samples missing more than 10% data and used a total of 180 samples for analysis. Using all SNP options, we obtained a total of 464 SNPs, which were used for all subsequent analyses except for Structure analysis. For Structure analysis, 115 SNPs using a single SNP option were used. BayeScan indicated that all loci were neutral (q -values >0.05).

3.2 Genetic diversity and genetic differentiation

Genetic diversity (H_O and H_E) of the studied populations, including that from Ogasawara, were comparable and ranged from 0.039–0.044 and 0.039–0.045 for H_O and H_E , respectively (Suppl. Table 2). Global F_{ST} values were significant ($P < 0.001$) but small both for groups including ($F_{ST} = 0.005$) and excluding the Ogasawara population ($F_{ST} = 0.006$). Likewise, AMOVA with prior grouping (Ogasawara vs other populations) indicated that F_{CT} was low and not significant (0.00167, $P = 0.219$). Pairwise F_{ST} values across the studied populations were also small, and only two pairs (Sekisei Lagoon [SKS] and Tatsukushi [TTK], Miyazaki [MYZ] and Ogasawara [OGS]) became significant after sequential Bonferroni correction ($\alpha < 0.05$) (Table 1).

Mean log likelihood values (mean $\ln P(D)$) were highest at $K = 1$ without prior grouping and comparable at K from 1 to 7 with prior grouping, with the lowest standard deviation value of mean $\ln P(D)$ found at $K = 1$ (Suppl. Table 3). Structure bar plots with and without prior geographic grouping suggested genetic homogeneity across the populations (Fig. 2A, Suppl. Fig. 2). PCoA indicated that, along the x-axis, SKS, MYK and Sakurajima [SKR] were divided from the others, although it was not related to geographic location or the year of sampling (Fig. 2B).

3.3 The larval dispersal simulation

The results of the larval dispersal simulations are shown as a connectivity matrix (Fig. 1B). The diagonal elements of the matrix indicate self-seeding. Basically, site numbers from 1 to 20 were assigned from south to north along the Kuroshio Current; site numbers from 21 to 25 were assigned from Nii-Jima and Shikine islands, which are closer to Japan's main island, to the Ogasawara and Iwo-Jima Islands, which are further; and site numbers from 26 to 30 were assigned along the Tsushima Warm Current. The linkage of the site numbers for numerical simulation to site names for

genomic analysis was as follows: 5 = Sekisei Lagoon (SKS), 6 = Miyako (MYK), 7 = Onna Village (ONN), 11 = Miyazaki (MYZ), 13 = Tatsukushi (TTK), 25 = Ogasawara (OGS), and 27 = Sakurajima (SKR). The results of this analysis confirmed that the larvae were transported from south to north by following the Kuroshio Current. However, they also confirmed that self-seeding dominated in the Ogasawara area, and it is very rare for larvae from the other sites to reach the Ogasawara area directly or for larvae from the Ogasawara area to reach the other sites directly.

Figure 3 shows the back-tracing results obtained using Markov chains. The bar of each site indicates the ratio of the number of particles released from the source site (laws of transition matrix $\mathbf{P}^n$). The results indicate that the origin sites gradually converged with some of the upstream sites of the Kuroshio Current. The population in the Ogasawara area is not well mixed with that of the Kuroshio region, but the 200–500 time-steps, which may be regarded as representing over a few hundred generations, confirmed that the populations in the Kuroshio region and the Ogasawara region are almost mixed.

4 Discussion

This study examined larval dispersal of the coral predator *A. cf. solaris* between the Kuroshio region and Ogasawara using population genomic analysis and oceanographic simulation to test the secondary outbreak hypothesis. Contrasting results were obtained using different methods. Population genomic analysis indicated genetic homogeneity between the populations of the two regions, and oceanographic numerical simulation indicated that multiple-generation stepping-stone migration was required to disperse larvae from the Kuroshio region to Ogasawara. These results indicate that the Ogasawara population of *A. cf. solaris* has the same genetic origin as that of the Kuroshio region, but a large amount of one-generation larval dispersal, such as a secondary outbreak from the Kuroshio region to Ogasawara, is unlikely. Rather, the data suggest that the population outbreak in Ogasawara might have been caused by successful self-seeding. This study highlights the importance of an integrated approach for estimating larval dispersal, including its timescale, for conservation purposes.

4.1 Potential of secondary outbreak between Ogasawara and Kuroshio region

In this study, all population genomics results indicated genetic similarity and no genetic structure between *A. cf. solaris* populations in the Ogasawara and Kuroshio regions. The AMOVA and Structure results indicate that the origins of *A. cf. solaris* populations in the Ogasawara and Kuroshio regions are genetically similar. Low F_{ST}/F_{CT} values between the Ogasawara and Kuroshio regions indicate (1) a contemporary, large amount of larval dispersal, (2) accumulation of stepping-stone migrants over multiple years, or (3) recent separation of a population with no or limited ongoing gene flow. Based on the oceanographic numerical simulation analysis results and historical records of *A. cf. solaris* in Ogasawara, the latter two are more likely than contemporary, large amounts of larval dispersal.

The result of the 26-year oceanographic simulation suggests that the chance of direct one-generation larval dispersal between the Kuroshio region and Ogasawara is very low, although stepping-stone migration (e.g. via Hachijyo-jima in Izu islands) between the two regions over multiple generations is physically possible. The Markov chain simulation, which simulated stepping-stone migration over multiple generations, indicated that the Ogasawara population is dominated by migrants from the Kuroshio region after 200–500 generations. This implies that the accumulation of larvae from low

levels of migration between the Kuroshio and Ogasawara regions over multiple years would result in genetic homogeneity. Given that there are no observed reports of population outbreaks of *A. cf. solaris* in stepping-stone regions (e.g., Izu islands) in the last 40 years (Yasuda 2018), genetic homogeneity between the Ogasawara and Kuroshio regions observed by *F*-statistics and STRUCTURE analysis could be attributed to stepping-stone larval migration over multiple generations. Crandall et al. (2018) demonstrated observed results of the marine species along Hawaii archipelagos where *F_{ST}* is not significantly different from zero were not actually panmictic but having weak hierarchical structure of isolation-by-distance caused by stepping stone migration. Coalescent simulation including *A. cf. solaris* samples from Izu islands would confirm this hypothesis.

Alternatively, it is possible that the observed strong gene flow resulted from recent colonization. According to a literature survey, there was no record of *A. cf. solaris* in the Ogasawara region before 1945, and the first record of this species in Ogasawara was in 1968 (Kurata, 1984; Yasuda, 2018). Since then, the density of *A. cf. solaris* in Ogasawara has been low (Yasuda 2018). However, a lack of a record of *A. cf. solaris* in Ogasawara before 1945 does not necessarily mean a complete absence of this species in the region, given that almost no studies or surveys were conducted before 1945. In addition, recent migration with a small population size would predict low genetic diversity due to the founder effect, although we observed almost the same genetic diversity in the Ogasawara and Kuroshio regions. Therefore, high genetic diversity in Ogasawara and strong gene flow between the Ogasawara and Kuroshio regions is more likely caused by stepping-stone larval migration over multiple generations, as suggested by the oceanographic simulation.

Whichever may be the case, the oceanographic simulation suggests that contemporary, one-generation larval dispersal between the Ogasawara and Kuroshio regions is limited, and thus direct secondary population outbreaks that require a large amount of larval dispersal within a generation from the Kuroshio region to Ogasawara are unlikely.

Kurata (1984) speculated that *A. cf. solaris* in Ogasawara might have colonized either from the Kuroshio region or from Mariana regions such as Guam. According to a population genetics analysis using microsatellite markers, genetic differentiation between the Kuroshio and Guam regions has been identified (Tusso et al., 2016), implying that colonization and/or secondary population outbreaks from Guam to the Ogasawara region are less likely than from the Kuroshio region.

Only one previous study examined the genetic connectivity of coral reef organisms between the Kuroshio and Ogasawara regions. A reef-building coral species, *Acropora* sp., which has a shorter (average 3–4 days) larval stage than that of *A. cf. solaris*, showed genetic differentiation between the Ogasawara and Kuroshio regions (Nakajima et al., 2012). This implies that larval dispersal from the Kuroshio region to the Ogasawara region of coral reef organisms with short larval stages is quite limited.

4.2 Possible causes of population outbreak in Ogasawara

Oceanographic simulation suggests that many larvae are self-recruited in Ogasawara compared with other *A. cf. solaris* populations in the Kuroshio region (Suppl. Table 4). This fact indicates that the observed population outbreak in Ogasawara in 2018 was caused by a sudden increase in regional self-seeding (which is termed “primary outbreak” as opposed to “secondary outbreak”; Birkeland and Lucas, 1990). While several hypotheses have been proposed to explain the primary outbreak of *A. cf. solaris*, the causes of primary outbreaks are still unclear (Pratchett et al., 2014). While the larval survival hypothesis in association with an elevated level of nutrients is widely accepted in GBR,

causes of outbreak would also depend on several regional environmental conditions being met simultaneously (Okaji et al., 2019): elevated levels of larval nutrition and subsequent higher survival rates; lower predation on larvae and juveniles; and physical processes acting on larvae that increase their survival, such as weather conditions and current patterns (Keesing and Halford, 1992; Okaji et al., 2019).

The population outbreak in Ogasawara in 2018 could be attributed to successful local larval recruitment in 2016, given that it takes two years for *A. cf. solaris* larvae to become adults (Birkeland and Lucas 1990). Oceanographic simulation also suggests that the number of self-recruited larvae in Ogasawara was larger (397) in 2016, which exceeded the average (252) over 26 years (Suppl. Table 5). This might be partly due to the smaller number of typhoons during the larval period (July to October, estimated by the spawning period associated with water temperature; Yasuda et al., 2010); in 2016, three typhoons passed through the Ogasawara region, which is below the 70-year average (4.47) (data from Japan Meteorological Agency, Suppl. Table 6). A previous field survey discovered dense clouds of *A. cf. planci* larvae before one typhoon, but they were scattered and disappeared from the coral reef area after the typhoon (Suzuki et al., 2016), implying that typhoons prevent mass settlement of larvae. However, in some years with a higher number of self-recruits and fewer typhoons, no population outbreak was observed in Ogasawara. Thus, other factors such as higher success of fertilization, higher nutrient levels during the larval stage, and less predation during the post-settlement stage might also be associated with the demographic fluctuation of *A. cf. solaris* in the Ogasawara area.

For conservation of corals in the Ogasawara area, caution should be taken two years after the observation of local retention of ocean current systems and preferable conditions for the increase in *A. cf. solaris*, such as a large amount of terrestrial run-off and a fewer number of typhoons during spawning periods. Local monitoring of juvenile starfish would be useful for predicting population outbreaks (Okaji et al. 2019).

4.3 Multiple approach to estimate larval dispersal in marine organisms

This study estimated larval dispersal of *A. cf. solaris* between Kuroshio and Ogasawara using two different approaches: population genomic analysis and larval dispersal simulation. Gene flow estimation by the population genomic approach often reflects both contemporary and evolutionary processes, rendering the interpretation of the amount of contemporary dispersal difficult because of the long evolutionary timescale. Genetic homogeneity can be observed due to a large amount of contemporary larval dispersal (secondary population outbreak), or stepping-stone migrants over multiple generations, or recent divergence with no contemporary dispersal. Our study is unique in that it included an oceanographic model with Markov chain simulation. Such a simulation was useful for testing these alternative hypotheses. A simple oceanographic simulation averaged over 26 years first rejected the secondary population outbreak hypothesis by indicating limited physical larval transport from the Kuroshio to the Ogasawara region within a generation. This result supports the idea that stepping-stone migrants over multiple generations is responsible for the genetic homogeneity observed in the F_{ST} and STRUCTURE analysis between the Kuroshio and Ogasawara populations.

This may give the impression that oceanographic simulation alone is enough for testing the secondary population hypothesis; however, it is generally imperfect for estimating larval dispersal. Our model regarded larvae as neutral passive particles without any of the biological characteristics of larvae that may influence actual larval dispersal, such as vertical movement, natural selection, food

availability, and mortality. In addition, the model only simulates the dispersal process, which does not include post-settlement survival. In some marine species, genetic connectivity is quite limited despite having long pelagic larval duration possibly caused by several biological features of larvae and juveniles (e.g. Barber et al. 2000; Barber et al. 2002). Thus, a comparison with genetic connectivity that reflects all the biological characteristics of the species is also important. While several previous studies have reported good agreement between oceanographic models and genetic analysis and have provided robust and straightforward interpretations of larval dispersal occurring on an ecological timescale (e.g. Galindo et al. 2006; White et al. 2010; Sunday et al. 2014; Taninaka et al. 2019), only a few have discussed the usefulness of oceanographic modeling for identifying alternative hypotheses or provided additional insights into the causes of observed population genetic structure (Galindo et al. 2010; Crandall et al. 2014). By applying Markov chain simulation in the numerical simulation, we could also include the effect of migration over multiple generations, which is easier to compare with genetic results than a single generation simulation result in terms of time-scale. This method underlines the usefulness and importance of using both genetic and oceanographic methods to estimate larval dispersal and provides significant insight into larval dispersal that occurs on ecological and evolutionary timescales.

5 Author Contributions

NY conceived the study. MH, NY, RK, HY, and TY drafted the manuscript. RK, HY, TI, and AT sequenced, assembled, and reconstructed the reference genome. NY, YA, and TS collected the *Acanthaser* samples. MH, HT, HY, TK, and NY conducted the MIG-seq analysis. TN and TY conducted oceanographic simulations. All authors edited the draft and approved it for publication.

6 Funding

Funding for this study was provided by the Environmental Research and Technology Development Fund (4RF-1501 and 4-1304) of the Ministry of the Environment, Japan; a Grant-in-Aid for Young Scientists (A) (17H04996), JSPS KAKENHI 221S0002, 16H06279 (PAGS); and a Grant-in-Aid for Research Fellows (19J21342, 18J23317) from the Japan Society for the Promotion of Science (JSPS).

7 Acknowledgments

We are grateful for Ms. Eriko Nagahiro for her help in the molecular experiment.

8 Figure legends

Figure 1 (A): The site numbers for numerical simulation including sampling sites for genomic analysis (5 = Sekisei Lagoon [SKS], 6 = Miyako [MYK], 7 = Onna Village [ONN], 11 = Miyazaki [MYZ], 13 = Tatsukushi [TTK], 25 = Ogasawara [OGS], and 27 = Sakurajima [SKR]). **(B):** The connectivity matrix of the simulation of 26-year larval dispersal. The sink location is indicated on the x-axis, and the source location is indicated on the y-axis. Colors indicate numbers of particles.

Figure 2 (A): Bar plot from Structure analysis at $K = 2$. **A1:** without prior geographic grouping. **A2:** with prior geographic grouping (Ogasawara vs other regions), indicating the population structure estimated by Structure v2.3.4 (Pritchard et al., 2000). The x-axis indicates individuals, and the y-axis indicates the proportion of hypothetical clusters, shown in different colors. **(B):** Results of PCoA based on a covariance matrix with standardized data. The first and second axes explain 70.03% and 21.20% of the variation, respectively.

Figure 3 The ratio of the number of particles originating from each source site. **(A)** Simulation of 26-year larval dispersal (see sites, Fig. 1A and connectivity matrix, Fig. 1B). The ratio determined by Markov chains with **(B)** time-step $n = 2$, **(C)** $n = 10$, **(D)** $n = 50$, **(E)** $n = 200$, and **(F)** $n = 500$. Site numbers are assigned as follows: 5 = Sekisei Lagoon (SKS), 6 = Miyako (MYK), 7 = Onna Village (ONN), 11 = Miyazaki (MYZ), 13 = Tatsukushi (TTK), 25 = Ogasawara (OGS), and 27 = Sakurajima (SKR).

9 Reference

- Alberto, F., Raimondi, P.T., Reed, D.C., Watson, J.R., Siegel, D.A., Mitarai, S., et al. (2011). Isolation by oceanographic distance explains genetic structure for *Macrocystis pyrifera* in the Santa Barbara Channel. *Mol. Ecol.* 20, 2543-2554. doi: 10.1111/j.1365-294X.2011.05117. x.
- Almany, G.R., Connolly, S.R., Heath, D.D., Hogan, J.D., Jones, G.P., McCook, L.J., et al. (2009). Connectivity, biodiversity conservation and the design of marine reserve networks for coral reefs. *Coral Reefs* 28, 339-351. doi: /10.1007/s00338-009-0484-x.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. *J. Mol. Biol.* 215, 403-410. doi: 10.1016/S0022-2836(05)80360-2.
- Arandt, A., and Smith, M.J. (2002). Genetic diversity and population structure in two species of sea cucumber: differing patterns according to mode of development. *Mol. Ecol.* 7, 1053-1064. doi: /10.1046/j.1365-294x.1998.00429.x.
- Baird, A.H., Pratchett, M.S., Hoey, A.S., Herdiana, Y., and Campbell, S.J. (2013). *Acanthaster planci* is a major cause of coral mortality in Indonesia. *Coral Reefs* 32, 803-812. doi: 10.1007/s00338-013-1025-1.
- Barber, P. H., Palumbi, S. R., Erdmann, M. V., & Moosa, M. K. (2000). A marine Wallace's line?. *Nature*, 406(6797), 692-693.
- Barber, P. H., Palumbi, S. R., Erdmann, M. V., & Moosa, M. K. (2002). Sharp genetic breaks among populations of *Haptosquilla pulchella* (Stomatopoda) indicate limits to larval transport: patterns, causes, and consequences. *Mol. Ecol.* 11, 659-674
- Benzie, J.A.H. (1992). Review of the genetics, dispersal and recruitment of crown-of-thorns starfish (*Acanthaster planci*). *Aust. Mar. Freshw. Res.* 43, 597-610. doi: 10.1071/MF9920597.

- 473 Benzie, J.A.H. (1999). Genetic Structure of Coral Reef Organisms: Ghosts of Dispersal Past. *Am.*
474 *Zool.* 39, 131-145. doi: /10.1093/icb/39.1.131.
- 475 Benzie, J.A.H. (1999). Major Genetic Differences between Crown-of-Thorns Starfish (*Acanthaster*
476 *planci*) Populations in the Indian and Pacific Oceans. *Evolution* 53, 1782-1795. doi:
477 10.1111/j.1558-5646.1999.tb04562.x.
- 478 Biodiversity Center of Japan (2019). Reconnaissance report of coral reefs, monitoring 1000. Ministry
479 of the Environment Japan.62-67. (In Japanese with English Summary)
480 https://www.biodic.go.jp/moni1000/findings/reports/pdf/h30_coral_reef.pdf
- 481 Birkeland, C., and Lucas, J. (1990). *Acanthaster planci*: Major Management Problem of Coral Reefs.
482 (Boca Ranton, Ann Arbor: CRC Press), 272.
- 483 Botsford, L.W., White, J.W., Coffroth, M.A., Paris, C.B., Planes, S., Shearer, T.L., et al. (2009).
484 Connectivity and resilience of coral reef metapopulations in marine protected areas: matching
485 empirical efforts to predictive needs. *Coral Reefs* 28, 327-337. doi: /10.1007/s00338-009-0466-z.
- 486 Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A., and Cresko, W.A. (2013) Stacks: an analysis
487 tool set for population genomics. *Mol. Ecol.* 22, 3124-3140. doi: 10.1111/mec.12354.
- 488 Chassignet, E.P., Hurlburt, H.E., Smedstad, O.M., Halliwell, G.R., Hogan, P.J., Wallcraft, A.J.,
489 Baraille, R. and Bleck, R. (2007). The HYCOM (HYbrid Coordinate Ocean Model) data
490 assimilative system. *J. Mar. Syst.* 65, 60-83. doi: 10.1016/j.jmarsys.2005.09.016.
- 491 Crandall, E.D., Treml, E.A., Liggins, L., Gleeson, L., Yasuda, N., Barber, PH., et al. (2014). Return
492 of the ghosts of dispersal past: historical spread and contemporary gene flow in the blue sea star
493 *Linckia laevigata*. *B. Mar. Sci.* 90, 399-425. doi: /10.5343/bms.2013.1052.
- 494 Crandall, E. D., Toonen, R. J., ToBo Laboratory, & Selkoe, K. A. (2019). A coalescent sampler
495 successfully detects biologically meaningful population structure overlooked by F - statistics.
496 *Evol. Appl.* 12, 255-265.
- 497 Dight, I.J., James, M.K., and Bode, L. (1990a). Modelling the larval dispersal of *Acanthaster planci*
498 I. Large scale hydrodynamics, Cairns Section, Great Barrier Reef Marine Park. *Coral Reefs* 9,
499 115-123. doi: 10.1007/BF00258222.

500 Dight, I.J., James, M.K., and Bode, L. (1990b). Modelling the larval dispersal of *Acanthaster planci*.
501 II. Patterns of reef connectivity. Coral Reefs 9, 125-134. doi: 10.1007/BF00258224.

502 Earl, D.A., and vonHoldt, B.M. (2012). STRUCTURE HARVESTER: a website and program for
503 visualizing STRUCTURE output and implementing the Evanno method. Conserv. Genet. Resour.
504 4, 359-361 doi: 10.1007/s12686-011-9548-7.

505 Excoffier, L., and Lischer, H.E.L. (2010). Arlequin suite ver 3.5: a new series of programs to perform
506 population genetics analyses under Linux and Windows. Mol. Ecol. Resour. 10, 564-567. doi:
507 10.1111/j.1755-0998.2010.02847.x.

508 Fukuda, M., and Hanamura, Y. (1996). Small scale distribution and transport of the Japanese
509 anchovy, *Engraulis japonica*, larvae determined from a free-drifting buoy survey in Iyo-Nada,
510 Seto inland sea [Japan] [1995]. J. Agric. Sci. Technol. 28, 9-20.

511 Galindo, H. M., Olson, D. B., and Palumbi, S. R. (2006). Seascape genetics: a coupled
512 oceanographic-genetic model predicts population structure of Caribbean corals. Curr. Biol. 16,
513 1622-1626.

514 Galindo, H. M., Pfeiffer - Herbert, A. S., McManus, M. A., Chao, Y. I., Chai, F. E. I., and Palumbi,
515 S.R. (2010). Seascape genetics along a steep cline: using genetic patterns to test predictions of
516 marine larval dispersal. Mol. Ecol. 19, 3692-3707.

517 Gordon, A., and Hannon, G.J. (2012). FASTX-Toolkit. FASTQ/A short-reads pre-processing tools.
518 http://hannonlab.cshl.edu/fastx_toolkit/. [Unpublished].

519 Harrison, H.B., Pratchett, M.S., Messmer, V., Saenz-Agudelo, P., and Berumen, M.L. (2017).
520 Microsatellites Reveal Genetic Homogeneity among Outbreak Populations of Crown-of-Thorns
521 Starfish (*Acanthaster cf. solaris*) on Australia's Great Barrier Reef. Diversity 9, 16. doi:
522 10.3390/d9010016.

523 Hubisz, M., Falush, D., Stephens, M., and Pritchard, J. (2009). Inferring weak population structure
524 with the assistance of sample group information. Mol. Ecol. Resour. 9, 1322-1332. doi:
525 10.1111/j.1755-0998.2009.02591.x.

James, M.K., and Scandol, J.P. (1992). Larval dispersal simulations: correlation with the crown-of-thorns starfish outbreaks database. *Mar. Freshw. Res.* 43, 569-581. doi: 10.1071/MF9920569.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9, 357-359. doi: 10.1038/nmeth.1923.

Lucas, J.S. (1973). Reproductive and larval biology of *Acanthaster planci* (L.) in Great Barrier Reef waters. *Micronesica-Series 9*, 197-203.

Kajitani, R., Toshimoto, K., Noguchi, H., Toyoda, A., Ogura, Y., Okuno, M., et al. (2014). Efficient de novo assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome Res.* 24, 1384-1395. doi: 10.1101/gr.170720.113.

Keesing, J.K., and Halford, A.R. (1992). Importance of postsettlement processes for the population dynamics of *Acanthaster planci* (L.). *Aust. Mar. Freshw. Res.* 43, 635-651. doi: 10.1071/MF9920635.

Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A., and Mayrose, I. (2015). Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Mol. Ecol. Resour.* 15, 1179-1191. doi: 10.1111/1755-0998.12387.

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., and Phillippy, A. M. (2017). Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res.* 27, 722-736. doi: 10.1101/gr.215087.116.

Kurata, Y. (1984). Crown-of-thorns starfish of Ogasawara Island. *Marine park information* 61, 7-9. In Japanese.

Lischer, H.E.L., and Excoffier, L. (2012). PGDSpider: an automated data conversion tool for connecting population genetics and genomics programs. *Bioinformatics* 28, 298-299. doi: 10.1093/bioinformatics/btr642.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34, 3094–3100. doi: 10.1093/bioinformatics/bty191.

Marçais, G., and Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics* 27, 764–770. doi: 10.1093/bioinformatics/btr011.

- 553 Marko, P.B., and Hart, M.W. (2017). Genetic Analysis of Larval Dispersal, Gene Flow, and
554 Connectivity. *Evolutionary Ecology of Marine Invertebrate Larvae*.
555 doi:/10.1093/oso/9780198786962.003.0012.
- 556 Martin, M. (2011). Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads.
557 EMBnet 17. doi: 10.14806/ej.17.1.200
- 558 Meeker, N.D., Hutchinson, S.A., Ho, L., and Trede, N.S. (2007). Method for isolation of PCR-ready
559 genomic DNA from zebrafish tissues. *Biotechniques* 43, 610-614. doi: 10.2144/000112619
- 560 Miyake, Y., Kimura, S., Kawamura, T., Horii, T., Kurogi, H., and Kitagawa, T. (2009). Simulating
561 larval dispersal processes for abalone using a coupled particle-tracking and hydrodynamic model:
562 implications for refugium design. *Mar. Ecol. Prog. Ser.* 387, 205-222. doi: 10.3354/meps08086.
- 563 Nakabayashi, A., Yamakita, T., Nakamura, T., Aizawa, H., Kitano, Y.F., Iguchi, A., et al. (2019).
564 The potential role of temperate Japanese regions as refugia for the coral *Acropora hyacinthus* in
565 the face of climate change. *Sci. Rep.* 9, 1892. doi: 10.1038/s41598-018-38333-5.
- 566 Nakajima, Y., Nishikawa, A., Iguchi, A., and Saka, K. (2012). Regional genetic differentiation
567 among northern high-latitude island populations of a broadcast-spawning coral. *Coral Reefs* 31,
568 1125-1133. doi: 10.1007/s00338-012-0932-x.
- 569 Narum, S.R., and Hess, J.E. (2011). Comparison of FST outlier tests for SNP loci under selection.
570 *Mol. Ecol. Resour.* 11, 184-194. doi: 10.1111/j.1755-0998.2011.02987.x.
- 571 Nash, W.J., Goddard, M. and Lucas, J.S. (1988). Population genetic studies of the crown-of-thorns
572 starfish, *Acanthaster planci* (L.), in the Great Barrier Reef region. *Coral Reefs* 7, 11-18. doi:
573 10.1007/BF00301976.
- 574 Okaji, K., Ogasawara, K., Yamakita, E., Kitamura, M., Kumagai, N., Nakatomi, N., et al. (2019).
575 Comprehensive management program of crown-of-thorns starfish outbreaks in Okinawa. *The*
576 *Japanese Coral Reef Society* 21, 91-110. doi: 10.3755/jcrs.21.91. In Japanese.
- 577 Paris, C.B., Helgers, J., van Sebille, E., and Srinivasan, A. (2013) Connectivity Modeling System: A
578 probabilistic modeling tool for the multi-scale tracking of biotic and abiotic variability in the
579 ocean. *Environ. Model. Softw.* 42, 47-54. doi: 10.1016/j.envsoft.2012.12.006.

580 Peakall, R., and Smouse, P.E. (2012). GenAlEx 6.5: genetic analysis in Excel. Population genetic
581 software for teaching and research—an update. *Bioinformatics* 28, 2537-2539. doi:
582 10.1093/bioinformatics/bts460.

583 Pratchett, M.S., Caballes, Ciemon, F., Rivera-Posada, Jairo, A., et al. (2014). Limits to understanding
584 and managing outbreaks of crown-of-thorns starfish (*Acanthaster* spp.). *Oceanogr. Mar. Biol.* 52,
585 133-200. doi: 10.1201/b17143-4.

586 Pritchard, J.K., Stephens, M., and Donnelly, P. (2000). Inference of population structure using
587 multilocus genotype data. *Genetics* 155, 945-959. doi: 10.1093/genetics/155.2.945.

588 Ranallo-Benavidez, T.R., Jaron, K.S., and Schatz, M.C. (2020). GenomeScope 2.0 and Smudgeplot
589 for reference-free profiling of polyploid genomes. *Nat. Commun.* 11, 1432. doi: 10.1038/s41467-
590 020-14998-3.

591 Rochette, N., and Catchen, J. (2017). Deriving genotypes from RAD-seq short-read data using
592 Stacks. *Nat. Protoc.* 12, 2640-2659. doi: 10.1038/nprot.2017.123.

593 Scandol, J.P., and James, M.K. (1992). Hydrodynamics and larval dispersal: a population model of
594 *Acanthaster planci* on the Great Barrier Reef. *Mar. Freshw. Res.* 43, 583-595. doi:
595 10.1071/MF9920583.

596 Schunter, C., Carreras-Carbonell, J., Macpherson, E., Tintoré, J., Vidal-Vijande, E., Pascual, A., et al.
597 (2011). Matching genetics with oceanography: directional gene flow in a Mediterranean fish
598 species. *Mol. Ecol.* 20, 5167-5181. doi: 10.1111/j.1365-294X.2011.05355.x.

599 Seppey M., Manni M., and Zdobnov E.M. (2019). BUSCO: Assessing Genome Assembly and
600 Annotation Completeness. In: *Gene Prediction. Methods Mol. Biol* 1962, eds. M. Kollmar
601 (Humana, New York, NY). doi: 10.1007/978-1-4939-9173-0_14.

602 Shanks, A.L. (2009). Pelagic larval duration and dispersal distance revisited. *Biol. Bull.* 216, 373-
603 385. doi: 10.1086/BBLv216n3p373.

604 Storlazzi, C.D., van Ormondt, M., Chen, Y-L., and Elias, E.P.L. (2017). Modeling Fine-Scale Coral
605 Larval Dispersal and Interisland Connectivity to Help Designate Mutually-Supporting Coral Reef

606 Marine Protected Areas: Insights from Maui Nui, Hawaii. *Front. Mar. Sci.* 4. doi:
607 10.3389/fmars.2017.00381.

608 Sunday, J. M., Popovic, I., Palen, W. J., Foreman, M. G. G., and Hart, M. W. (2014). Ocean
609 circulation model predicts high genetic structure observed in a long - lived pelagic developer.
610 *Mol. Ecol.* 23, 5036-5047.

611 Suyama, Y., and Matsuki, Y. (2015). MIG-seq: an effective PCR-based method for genome-wide
612 single-nucleotide polymorphism genotyping using the next-generation sequencing platform. *Sci.*
613 *Rep.* 5, 16963. doi: 10.1038/srep16963

614 Suzuki, G., Yasuda, N., Ikehara, K., Fukuoka, K., Kameda, T., Kai, S., et al. (2016). Detection of a
615 High-Density Brachiolaria-Stage Larval Population of Crown-of-Thorns Sea Star (*Acanthaster*
616 *planci*) in Sekisei Lagoon (Okinawa, Japan). *Diversity* 8, 9. doi: 10.3390/d8020009.

617 Takata, K., Taninaka, H., Nonaka, M., Iwase, F., Kikuchi, T., Suyama, Y., et al. (2019). Multiplexed
618 ISSR genotyping by sequencing distinguishes two precious coral species (Anthozoa:
619 Octocorallia: Coralliidae) that share a mitochondrial haplotype. *Peer J-Life and Environment*. doi:
620 10.7717/peerj.7769.

621 Taninaka, H., Bernardo, L.P.C., Saito, Y., Nagai, S., Ueno, M., Kitano, Y.F., et al. (2019). Limited
622 fine-scale larval dispersal of the threatened brooding corals *Heliopora* spp. as evidenced by
623 population genetics and numerical simulation. *Conserv. Genet.* 20, 1449-1463. doi:
624 10.1007/s10592-019-01228-7.

625 Timmers, M.A., Andrews, K.R., Bird, C.E., deMaintenton, M.J., Brainard, R.E., and Toonen, R.J.
626 (2011). Widespread Dispersal of the Crown-of-Thorns Sea Star, *Acanthaster planci*, across the
627 Hawaiian Archipelago and Johnston Atoll. *Mar. Sci.* doi: 10.1155/2011/934269.

628 Timmers, M.A., Bird, C.E., Skillings, D.J., Smouse, P.E., and Toonen, R.J. (2012). There's No Place
629 Like Home: Crown-of-Thorns Outbreaks in the Central Pacific Are Regionally Derived and
630 Independent Events. *PLoS ONE* 7, e31159. doi: 10.1371/journal.pone.0031159.

631 Tusso, S., Morcinek, K., Vogler, C., Schupp, P.J., Caballes, C.F., Vargas, S., et al. (2016). Genetic
632 structure of the crown-of-thorns seastar in the Pacific Ocean, with focus on Guam. *Peer J*
633 27168979. doi: 10.7717/peerj.1970.

634 Uthicke, S., Doyle, J., Duggan, S., Yasuda, N., and McKinnon, A.D. (2015). Outbreak of coral-eating
635 Crown-of-Thorns creates continuous cloud of larvae over 320 km of the Great Barrier Reef. *Sci.*
636 *Rep.* 5, 16885. doi: 10.1038/srep16885.

637 Vogler, C., Benzie, J.A.H., Tenggardjaja, K., Barber, P.H., and Wörheide, G. (2013).
638 Phylogeography of the crown-of-thorns starfish: genetic structure within the Pacific species.
639 *Coral Reefs* 32, 515-525. doi: 10.1007/s00338-012-1003-z.

640 Wada, N., Yuasa, H., Kajitani, R., Gotoh, Y., Ogura, Y., Yoshimura, D., et al. (2020). A ubiquitous
641 subcuticular bacterial symbiont of a coral predator, the crown-of-thorns starfish, in the Indo-
642 Pacific. *Microbiome* 8, 123. doi: 10.1186/s40168-020-00880-3.

643 Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., et al. (2014). Pilon: An
644 Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly
645 Improvement. *PLoS ONE* 9, e112963. doi: 10.1371/journal.pone.0112963.

646 White, C., Selkoe, K. A., Watson, J., Siegel, D. A., Zacherl, D. C., and Toonen, R. J. (2010). Ocean
647 currents help explain population genetic structure. *Proc. R. Soc. B.* 277, 1685-1694.

648 White, J.W., Carr, M.H. Caselle, J.E. Palumbi, S.R. Warner, R.R. Menge, B.A., et al. (2019).
649 Empirical approaches to measure connectivity. *Oceanography* 32, 60-61. doi:
650 10.5670/oceanog.2019.31.

651 Yamaguchi, M. (1977). Larval Behavior and Geographic Distribution of Coral Reef Asteroids in the
652 Indo-West Pacific. *Micronesica-Series* 13, 283-296.

653 Yasuda, N., Kajiwar, K., Nagai, S., Ikehara, K., and Nadaoka, K. (2015). First report of field
654 sampling and identification of crown-of-thorns starfish larvae. *Galaxea, Journal of Coral Reef*
655 *Studies* 17, 15-16. doi: 10.3755/galaxea.17.15.

656 Yasuda, N., Nagai, S., Hamaguchi, M., Okaji, K., Gerard, K., and Nadaoka, K. (2009). Gene flow of
657 *Acanthaster planci* (L.) in relation to ocean currents revealed by microsatellite analysis. *Mol.*
658 *Ecol.* 18, 1574-1590. doi: 10.1111/j.1365-294X.2009.04133.x.

659 Yasuda, N., Ogasawara, K., Kajiwar, K., Ueno, M., Oki, K., Taniguchi, H., et al. (2010). Latitudinal
660 differentiation in the reproduction patterns of the crown-of-thorns starfish *Acanthaster planci*

through the Ryukyu Island Archipelago. Plankton Benthos Res. 5, 156-164. doi: 10.3800/pbr.5.156.

Yasuda, N., Taquet, C., Nagai, S., Yoshida, T., and Adjeroud, M. (2015). Genetic connectivity of the coral-eating sea star *Acanthaster planci* during the severe outbreak of 2006–2009 in the Society Islands, French Polynesia. Mar. Ecol. 36, 668-678. doi: 10.1111/maec.12175.

Yasuda, N. (2018). Distribution Expansion and Historical Population Outbreak Patterns of Crown-of-Thorns Starfish, *Acanthaster planci* sensu lato, in Japan from 1912 to 2015. Coral Reef Studies of Japan. Coral Reefs of the World 13, 125-148. doi: 10.1007/978-981-10-6473-9_9.

9. Conflict of Interest

To the best of our knowledge, the named authors have no conflict of interest, financial or otherwise.

Data Availability Statement

The whole genome sequencing data (BioSample ID, SAMD00056692) and MIG-seq data (BioSample: SSUB016768, SAMD00282440-SAMD00282626) were submitted to DDBJ and available.

Table Caption

Table 1. Pairwise F_{ST} values (below) and their probability values (above). Significantly differentiated pairs after sequential Bonferroni correction ($\alpha < 0.05$) are shown in bold face.

	TTK	MYZ	SKR	ONN	MYK	SKS	OGS
TTK		0.295	0.224	0.105	0.003	0.000	0.107
MYZ	0.003		0.400	0.142	0.023	0.000	0.141
SKR	0.004	0.001		0.465	0.484	0.21	0.473
ONN	0.006	0.002	0.000		0.370	0.031	0.447
MYK	0.018	0.007	0.000	0.001		0.355	0.045
SKS	0.028	0.014	0.002	0.004	0.001		0.004
OGS	0.006	0.003	0.000	0.000	0.005	0.008	

12. Supplementary Captions

Suppl. Fig. 1 Schematic model of gap-filling with long read-based contigs. Flanking-region lengths: 500, 1k, 5k, 10k, 20k, 40 kbp, 80 kbp, 160 kbp (multiple values are applied). Gap-closing step is iterated two times.

Suppl. Table 1. Statistics of the draft genome.

Suppl. Table 2 Sample information and genetic diversity inbreeding coefficient. N = number of individuals analyzed. H_o = observed heterozygosity, H_e = expected heterozygosity, F = inbreeding coefficient.

Suppl. Table 3. The results from Structure Harvester to calculate ΔK . K is the number of hypothetical clusters assumed in Structure analysis.

Suppl. Figure 2. Structure analysis at comparable K from 3 to 7 (1 = Tatsukushi [TTK], 2 = Miyazaki [MYZ], 3 = Sakurajima [SKR], 4 = Onna Village [ONN], 5 = Miyako [MYK], 6 = Sekisei Lagoon [SKS], 7 = Ogasawara [OGS]). **(A)**: without prior and **(B)**: with prior grouping for Ogasawara. In $K = 6$, each pattern was supported in five out of ten iterations. The x-axis indicates sampling sites. The y-axis indicates the probability of membership.

Suppl. Table 4. The total number of arrived particles over 26 years. Bold face numbers indicate self-seeding. Self-seeding at Ogasawara is highlighted in yellow. Vertical columns indicate source populations, and horizontal rows indicate sink sites.

Suppl. Table 5. The average number of arrived particles over 26 years. Bold face numbers indicate self-seeding. Self-seeding at Ogasawara is highlighted in yellow. Vertical columns indicate source populations, and horizontal rows indicate sink sites.

Suppl. Table 6. The number of typhoons approaching Ogasawara and Izu Islands over a 70-year span (1951–2020).

A1

A2

TTK MYZ SKR ONN MYK SKS OGS

B

Principal Coordinates (PCoA)