Testing for parallel genomic and epigenomic footprints of adaptation to urban life in a passerine bird

Authors:

Aude E. Caizergues^{1*}, Jeremy Le Luyer², Arnaud Grégoire¹, Marta Szulkin³, Juan-Carlos Senar⁴, Anne Charmantier^{1†}, Charles Perrier^{5†}

¹ CEFE, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

² Ifremer, IRD, Institut Louis Malardé, Univ Polynésie Française, EIO, F-98719 Taravao, Tahiti, Polynésie Française

³ Centre of New Technologies, University of Warsaw, S. Banacha 2c, 02-097 Warsaw, Poland

⁴ Museu de Ciències Naturals de Barcelona, Parc Ciutadella, 08003 Barcelona, Spain

⁵ CBGP, INRAe, CIRAD, IRD, Montpellier SupAgro, Univ. Montpellier, Montpellier, France

† *shared senior authorship*

* **Corresponding author:** Aude E. Caizergues, 1919 route de Mende, 34293 Montpellier cedex 5, FRANCE

Email : audeemiliecaizergues@gmail.com

Author Contributions: AEC, CP and JLL carried out analyses. CP, AC and AG conceived the study. AEC, CP and AC wrote the manuscript. AC funded the study. AC, AG, MS, JCS, AEC, CP conducted the sampling. All the authors contributed to improve the manuscript and gave final approval for publication.

Competing Interest Statement: We declare we have no competing interests.

Keywords: adaptation, DNA methylation, epigenomics, genomics, urbanization

28 **Abstract (250 words):**

29 Identifying the molecular mechanisms involved in rapid adaptation to novel environments and
30 determining their predictability are central questions in Evolutionary Biology and pressing issues due
31 to rapid global changes. Complementary to genetic responses to selection, faster epigenetic variations
32 such as modifications of DNA methylation may play a substantial role in rapid adaptation. In the
33 context of rampant urbanization, joint examinations of genomic and epigenomic mechanisms are still
34 lacking. Here, we investigated genomic (SNP) and epigenomic (CpG methylation) responses to urban
35 life in a passerine bird, the Great tit (*Parus major*). To test whether urban evolution is predictable (*i.e.*
36 parallel) or involves mostly non-parallel molecular processes among cities, we analysed three distinct
37 pairs of city and forest Great tit populations across Europe. Results reveal a polygenic response to
38 urban life, with both many genes putatively under weak divergent selection and multiple differentially
39 methylated regions (DMRs) between forest and city great tits. DMRs mainly overlapped transcription
40 start sites and promotor regions, suggesting their importance in the modulation gene expression. Both
41 genomic and epigenomic outliers were found in genomic regions enriched for genes with biological
42 functions related to nervous system, immunity, behaviour, hormonal and stress responses.
43 Interestingly, comparisons across the three pairs of city-forest populations suggested little parallelism
44 in both genetic and epigenetic responses. Our results confirm, at both the genetic and epigenetic levels,
45 hypotheses of polygenic and largely non-parallel mechanisms of rapid adaptation in new environments
46 such as urbanized areas.

47

48 INTRODUCTION

49 Identifying mechanisms involved in rapid adaptation to new environmental conditions is a
 50 central question in evolutionary biology and a is pressing task in the context of global changes of the
 51 Anthropocene ¹. The vast majority of studies investigating mechanisms involved in rapid adaptation to
 52 new environments have focused on phenotypic plasticity on the one hand and on genetic response to
 53 selection on the other hand. At their crossroad, recent work underlines the potential role of epigenetics
 54 in rapid adaptation to new environments ². In particular, environmental variations can induce
 55 differences in DNA methylation patterns and hence modulate genes' expression and upper-level
 56 phenotypes ^{3,4}. Such methylation-linked phenotypic variations can occur during an individual's
 57 lifetime, especially early on during the organism's development ^{5,6}. Although methylation changes
 58 acquired across an individual's lifetime may often be non-heritable ⁷ but see 8,9, epigenetically induced
 59 phenotypic shifts may nevertheless enhance individuals fitness in new environments. Moreover,
 60 during the course of evolution, divergent genetic variants regulating epigenetic modifications may also
 61 be selected for, hence promoting the evolution of divergent epigenotypes and epigenetically-linked
 62 phenotypic variation ¹⁰. While epigenetic studies focused on human diseases and medical topics are
 63 now abundant, studies in an ecological context are still rare ¹¹. Nevertheless, a few epigenetic studies
 64 in natural populations revealed that DNA methylation shifts might play a determinant role in local
 65 adaptation to environmental variation ¹². There is hence an urgent need for further empirical
 66 investigations of simultaneously rapid genetic and epigenetic evolution in response to environmental
 67 change ¹³.

68 Urbanization rapidly and irreversibly changes natural habitats into human-made environments
 69 and is considered as a major threat to biodiversity ¹⁴. For species who appear to cope with
 70 urbanisation, urban habitats present a myriad of novel environmental conditions compared to the
 71 habitat where they evolved, including high levels of chemical, light and sound pollution, high
 72 proportion of impervious surfaces, high habitat fragmentation, low vegetation cover and high human
 73 densities ^{15,16}. Such extreme environmental changes compared to natural areas are expected to result in
 74 numerous new selection pressures on city-dwelling species ¹⁷. Accordingly, rates of recent phenotypic
 75 change, concerning multiple types of traits related to behaviour, morphology, phenology and

76 physiology, seem to be greater in urban areas than in any other habitat types, including non-urban
77 anthropogenic contexts^{18,19}. The exploration of the molecular mechanisms implicated in urban-driven
78 phenotypic changes has only begun, with both genetic^{20–22}, and epigenetic investigations^{23–25}. For
79 instance, DNA methylation variations have been associated in vertebrates with high levels of traffic-
80 related air pollution²⁶. Yet, epigenetic studies have been performed at relatively small genomic
81 resolution. In addition, very little is known about the level of parallelism and hence of the
82 predictability of genetic and epigenetic evolution in response to urbanisation in distinct cities^{27,28}. So
83 far, there are situations ranging from local adaptation despite strong gene flow (e.g. in the red-tailed
84 bumblebee *Bombus lapidaries*²⁹) to restricted gene flow and independent colonization in different
85 cities by a few founders, followed by adaptation (e.g. in the burrowing owl *Athene cunicularia*,³⁰).
86 Providentially, recent genomic tools and virtually limitless amount of cities offer unique opportunities
87 for comparing at high genomic resolution simultaneously individuals' genomic and epigenomic
88 responses in several cities to study the parallelism and predictability in molecular mechanisms
89 implicated in rapid adaptation to urbanization^{31,32}.

90 In this study, we used genome-wide and epigenome-wide sequencing approaches to compare
91 genetic and epigenetic responses among three pairs of great tit *Parus major* populations in urban and
92 forest habitats. At the European level, population monitoring of Great tits revealed parallel phenotypic
93 shifts in city birds compared to their forest conspecifics, with in particular smaller and lighter urban
94 birds laying earlier and smaller clutches^{33–36}. We investigated both average genomic (SNPs) and
95 epigenomic (CpG methylation) differentiation and we searched for particular genomic footprints of
96 divergent selection as well as differentially methylated regions, between forest and urban populations.
97 Our results show that despite limited genetic differentiation and few genomic footprints of divergent
98 selection between forest and urban populations, urban life was associated with numerous differentially
99 methylated regions notably associated with neural development, behaviour and immunity. Hence, this
100 study shows the potential role of epigenetic response in rapid adaptation to changing environments in
101 urban areas. Importantly, we found little parallelism between cities in both genomic and epigenomic
102 responses to urbanization, possibly confirming the hypothesis that multiple evolutionary ways exist to
103 independently cope with similar novel environmental conditions.

RESULTS

Little genetic and epigenetic average differentiation between urban and rural populations

Genetic. We used a redundancy analysis (RDA) on 74,137 SNPs obtained by RAD-sequencing to document genetic variation among the studied great tits populations, with location (Barcelona, Montpellier or Warsaw), habitat (urban vs. forest), and sex as explanatory variables. The model was highly significant ($P < 0.001$) but explained only a small fraction (*i.e.* less than 2%) of the total variance: $R^2 = 0.018$ (Figure 1C, SI Table S1). All three variables were significant (location: $P = 0.001$; habitat: $P = 0.004$; sex: $P = 0.001$). Partial RDA revealed that the net variation explained by the habitat ($R^2 = 0.004$, $P = 0.001$) was inferior to the net variation explained by location ($R^2 = 0.012$, $P = 0.001$) but higher than sex ($R^2 = 0.002$, $P = 0.004$, Table S1). As expected, when removing the Z chromosome from the data, sex became non-significant ($P = 0.260$), whereas the effects of other variables remained significant and of similar magnitude (Table S2).

Genome-wide differentiation between populations, measured with Weir and Cockerham's F_{ST} ³⁷ was relatively low on average (mean $F_{ST} = 0.019$), in the order of 1 to 2% between habitats for each location ($F_{ST}^{Barcelona(rur-urb)} = 0.018 \pm 0.001$; $F_{ST}^{Montpellier(rur-urb)} = 0.012 \pm 0.001$; $F_{ST}^{Warsaw(rur-urb)} = 0.018 \pm 0.001$) (see SI Table S3), suggesting relatively high gene flow and limited genetic drift among populations. Mean F_{ST} on allosomes was lower (mean $F_{ST} = 0.014$, Table S4) than on the Z chromosome (mean $F_{ST} = 0.022$, Table S5).

Methylation. Similarly to genetic data, we performed an RDA on methylation level of 157,741 CpG sites to describe epigenetic variation among individuals in relation to location, habitat and sex. The model was significant but explained less than 1% of the total variance ($R^2 = 0.007$, $P = 0.001$). All variables contributed significantly (location : $P = 0.001$, habitat: $P = 0.03$, sex: $P = 0.001$, Table S6). Partial RDA revealed that location and sex explained a similar proportion of the total variance which was higher than habitat (location: $R^2 = 0.003$, $P = 0.001$; sex: $R^2 = 0.003$, $P = 0.001$; habitat: $R^2 = 0.001$, $P = 0.3$). When removing the Z chromosome from analyses, results remained similar (Table

S7), showing that difference in methylation was not entirely driven by potential diverging patterns associated with sexual chromosomes.

We then investigated more finely whether individual methylation on CpG cytosines varied across location, habitat (urban vs forest), sex, and location×habitat interaction, using an ANOVA, run on autosomes and Z chromosome separately (Fig 1B, Table S8). For autosomes, we detected a significant effect of location ($F = 3.319$, $P = 0.044$), with Montpellier individuals showing lower methylation levels than Warsaw ones (Fig 1B, Tukey test: $P = 0.04$) and no other difference between pairs of cities. Also, no significant effect of sex ($P = 0.263$) or habitat ($P = 0.478$) was found, suggesting that urbanization did not have an important overall effect on global methylation levels. For the Z chromosome, we found a strong difference between sexes, with homogametic males showing 2.98% more methylated Z than heterogametic females (Fig 1B; $P = 1.45 \times 10^{-15}$), while no significant difference between location ($P = 0.577$) or habitat ($P = 0.915$) was found.

Non-parallel yet strong genomic footprints of divergent selection between urban and forest populations and evidence for polygenic adaptation

We used two methods to investigate outlier SNPs potentially under divergent selection between forest and urban populations: an F_{ST} -outlier based method (Bayescan³⁸, v2.1, default parameters), and a multivariate method (using an RDA,³⁹) aiming respectively at identifying strong outliers indicating footprints of differentiation for each population pair and weaker outliers across the several population pairs studied at once. First, Bayescan identified 15 outliers for Barcelona, 11 for Montpellier and 10 for Warsaw, overall distributed on 15 chromosomes and associated to 13 genes in 5kb upstream or downstream regions (Figure 2, q -value < 0.1 , see Figure S1). None of these outliers was shared between the three population pairs, revealing no convergence between cities. Second, using the multivariate approach based on a RDA (following the procedure for constrained RDA described earlier), we investigated the habitat effect and extracted a list of 1163 loci with outlier loading score following Forester *et al.*³⁹ method (Figure 2, see Figure S2 for threshold), suggesting a higher number of loci undergoing weaker and polygenic selection. These 1163 loci were associated with 561 genes.

To investigate the potentially statistically enriched gene ontologies (GO) for the set of genes associated to RDA outliers, we used topGO R package⁴⁰ with GO referenced for the chicken *Gallus gallus* (ggallus_gene_ensembl). GO analyses revealed the existence of overrepresented ontologies ($P < 0.05$ and at least 3 genes per GO). Among the most promising GO terms we found functions related to the nervous system (GO:0048846, axon extension involved in axon guidance; GO:0035418 protein localization to synapse; GO:0021987 cerebral cortex development; GO:0007274, neuromuscular synaptic transmission), the blood system (GO:0045777 positive regulation of blood pressure; GO:0042311, vasodilatation), hormonal response (GO:0071277, cellular response to estrogen stimuli) and stress (GO:0033555, multicellular response to stress), revealing potential important functions involved in adaptation to urban habitats. Detailed GO results are presented in SI Table S9 and Figure S3.

Evidence for differentially methylated regions in urban environments

We identified a total of 224 distinct DMRs between urban and forest great tits: 80 for Barcelona, 68 for Montpellier and 93 for Warsaw. Only 14 DMRs (6.25%) were found repeatedly in at least two comparisons, and only 3 were common to the three cities. 7 (of these 14 parallel DMRs were in the same direction of methylation in urban compared to forest areas (Figure 3, Figure S4 A). Barcelona urban birds presented significantly more hypomethylated than hypermethylated DMRs ($\chi^2 = 11.25$, $P < 0.001$), but no difference was found for Montpellier ($\chi^2 = 0.941$, $P = 0.332$) nor for Warsaw ($\chi^2 = 0.011$, $P = 0.917$). DMRs were distributed across all the 32 chromosomes as well as on 37 unplaced scaffolds. 203 of the 224 different DMRs (91%) overlapped genes or 5kb flanking regions, among which 52% were directly located in gene bodies, promoter or TSS sequences (35.3% in gene bodies and 47.3% in promoters/TSS) suggesting their putative functional role in gene expression and regulation and/or splicing events.

Following the procedure previously described, GO analyses on the pooled genes list revealed an overrepresentation of modules associated to the nervous system (GO:2000300, regulation of synaptic vesicle exocytosis; GO:0050804 modulation of synaptic transmission), immunity (GO:005728, negative regulation of inflammatory response; GO:0050852, T cell receptor signaling

pathway), metabolic activity (GO:006816 calcium ion transport; GO:0055072, iron ion homeostasis, GO:0043087, regulation of GTPase activity), behaviour (GO:0007626, locomotory behavior) and endocrine processes (GO:0044060: regulation of endocrine process). All enriched GO are presented in SI Figure S5 and Table S10.

We also searched for DMRs between sexes, following the same procedure. We identified 206 DMRs associated with sex, of which 58 for Barcelona, 81 for Montpellier and 99 for Warsaw. On a total of 206 DMRs, 181 (57.3%) were on genes or in a 5kb upstream/downstream region around genes. GO analyses revealed enrichment of genes involved in development, growth and morphogenesis, among others (see detailed enriched GO Table S10 & Figure S5).

Almost twice more sex DMRs were shared between locations (11,7%) than between habitats (6.25%, see Figure S4 A & B ; z-test: χ^2 -squared = 3.885, P = 0.049). When taking into account the direction of methylation difference, 7 sex DMRs were shared between at least two cities (9.7%) which was three times more than for habitat DMRs (3.1%; z-test: X-squared = 7.904, P = 0.005).

DISCUSSION

The urban sprawl is a worldwide phenomenon deeply affecting the environment and thus requiring fast adaptive responses in city dwellers. While a large body of literature already describes a myriad of phenotypic shifts in urban populations of numerous species^{35,41,42}, the molecular bases of these shifts and their evolutionary implications remain yet to be documented and understood. This study uses genomic and epigenomic analyses to decipher the potential molecular bases implicated in phenotypic shifts and adaptation in several urban populations of a passerine bird, the Great tit. Note that this species shows largely parallel phenotypic shifts across its range in terms of morphology and life history (Thompson *et al.*, *sub*). Genomic analyses revealed weak yet significant average differentiation between urban and forest populations, suggesting ongoing gene flow and limited drift in urban populations. These analyses also identified a limited number of loci putatively under strong selection, non-repeated between pairs, and numerous loci supposedly under weaker selection, compatible with a polygenic model of evolution. From the epigenomic side, we found weak average

differentiation of the methylome between urban and forest birds, suggesting an absence of genome-wide epigenetic deregulations, which is notably in line with an absence of strong genetic differentiation. In turn, we identified several strongly differentially methylated regions between urban and forest birds, mostly non-repeated between pairs and hence potentially implicated with local evolution of urban populations. Genes associated with either genomic footprints of divergent selection or differentially methylated regions had relatively similar functions, related to the nervous system, metabolism, immunity and behaviour, that have been repeatedly convicted in other studies⁴⁰. Hence, by identifying non-repeated genetic and epigenetic responses among replicated forest-urban population pairs, our findings support the hypothesis of mostly non-parallel rapid *de novo* adaptation to similar environments via both genetic and epigenetic mechanisms. Our results are in line with accumulating evidence that polygenic adaptation and epigenetic reprogramming may be involved in quick phenotypic shifts in response to rapidly emerging constraints such as urbanization.

Overall genetic differentiation between populations was relatively low (F_{ST} ranging from 0.9% to 3.4%), although higher than what has been found at a much larger scale across the species distribution (e.g. around 1% of F_{ST} between UK and Spanish or French and Spanish populations⁴¹). Low but significant differentiation levels are in line with previously documented divergences between city and forest great tit populations^{20,45}, which altogether suggests important gene flow, large effective population sizes and limited genetic drift at multiple spatial scales⁴⁶. This overall genetic context is particularly suitable to search for genomic footprints of divergent selection between urban and forest populations, which would easily be identifiable above the neutral level of genetic differentiation.

We found a limited number of strong footprints of divergent selection, which is in line with previous results of Perrier et al.²⁰ in Montpellier, and of Salmon et al.²² across nine European cities. Similarly to low levels of parallelism in allele frequency changes between cities observed by previous studies^{22,47}, and despite similarities in phenotypic shifts in the European cities³⁵, none of these outliers were shared between cities. This result suggests limited parallel evolution, supporting a scenario of independent *de novo* evolution between cities and/or different selection pressures between cities. Indeed, there may be multiple evolutionary solutions to the same environmental challenges⁴⁸ and

multiple traits are linked to the same functional outcome⁴⁹. Besides, the identification of numerous outliers by the multivariate framework applied at the scale of all six sampling sites supports a model of polygenic urban adaptation implicating multiple genes, biological pathways and phenotypic traits⁵⁰. Polygenic adaptation may be expected in urban habitats since the multiple new environmental conditions in cities most probably result in many novel selective pressures acting on a multitude of functional traits⁵¹, and because many of these traits may be quantitative, and genetically correlated⁵². Further polygenic analyses on more samples and more markers (i.e. whole genome data) are however required to increase our capacity to estimate more precisely the potential effect of each genetic variant implicated in the adaptation to life in the city^{53,54}.

Several genomic footprints of divergent selection between urban and forest environments were in, or in the vicinity of, genes that have already been described as playing a role in neuronal development, behaviour or cognitive abilities. In particular, the NR4A2 gene plays an important role in recognition of novel objects and memory in mice⁵⁵. Reaction to novel objects and novel food is known as one of the main factors determining the capacity of a species to thrive in an urban environment⁴¹. The DCX gene is related to neuronal plasticity⁵⁶ and experimental approaches revealed that artificial light at night induces an overexpression of this gene linked to a change in behaviour and expression of depressive-like behaviour in crows⁵⁷. Finally, the CHRNA1 gene is associated with aggressive behaviour in chicken⁵⁸, and higher aggressiveness is commonly observed and hypothesized as adaptive in urban bird populations⁵⁹. Besides, the gene ontology enrichment analysis, performed on the entire set of genes identified via the outlier genome scan, reinforced these findings since multiple enriched GO terms were associated with the nervous system and stress response as well as hormonal response. These results are informative on the type of traits involved in avian urban adaptation in cities and corroborate previous results from^{22,60,61} suggesting that natural selection repeatedly acted on neuronal, behavioural and cognitive traits that could contribute to the phenotypic shifts described in urban great tits (i.e. more aggressive and exploratory birds, with higher breath rate :^{62, 63} & Caizergues *et al. in prep*).

Contrary to the common prediction that living in cities is likely to influence epigenomes^{24,25}, no genome wide pattern of differentiation in methylation between urban and forest great tits was detected. However, we observed a difference in mean methylation level between birds from Warsaw and Barcelona on their autosomes, as well as between males and females on the Z chromosome, showing that methylation differences were identifiable. In addition, we found no mean difference in methylation level between habitats, revealing that urbanization did not strongly affect overall methylation levels in a specific direction. This overall low differentiated methylation context is perfectly suitable to investigate more localized zones that could differ in their levels of methylation. Note that the strong methylation contrast between males and females, particularly on sex chromosomes (Figure 1B), is in line with previous reports in vertebrates^{64,65} showing that methylation plays a major role in sex differentiation via regulation of gene expression and genetic imprinting.

Despite the non-significant effect of habitat on overall methylation levels, we found a large number of DMRs between pairs of forest and urban populations, suggesting that urbanization did affect particular regions of the genome. DMR were less likely to occur within a gene body than by chance, but it was not the case for promoter or TSS regions. This latter result contrasts with Watson and collaborators²⁵ who recently found an under-representation of DMR in both gene body and regulatory regions in urban great tits from Malmö (Sweden). Across the three cities, 35.3% of DMR were directly localized in gene bodies, and 47.3% in TSS or promoter regions, suggesting a potential role in gene expression modulation. Direction of methylation patterns did not follow any absolute pattern (no over-representation of hypo- or hypermethylated DMR in urban birds, Figure 3), in line with Watson and collaborators' analyses on blood sample. However, birds in Barcelona presented significantly more hypomethylated DMR than hypermethylated ones, a result that warrants further investigations.

Only a limited number of urbanization-linked DMR were shared between two or more locations. In contrast, three times more sex linked DMRs were found in two locations or more. This comparison suggests that urbanization-linked epigenetic modifications most probably do not occur in a parallel way across cities, but rather that each city might have its particular epigenetic response. Indeed, in the study field of urban evolutionary biology, cities are often regarded as valuable replicates

of human induced habitats^{31,66}, and it is often expected that parallel adaptive responses to similar selective pressures will occur. This expectation is particularly strong when phenotypes show parallel changes, as is the case for the Great tit, which is consistently smaller and lays earlier and smaller broods in the various cities where it has been studied, compared to forest habitats^{67, 68}. However, as discussed above, parallel adaptation to similar environmental conditions should not be expected in the case of independent evolution, especially for multilocus traits. Additionally, cities are different from each other because of a wide array of climatic, cultural, historical and socioeconomic factors¹⁵. In fact, besides the obvious differences in cities' climatic conditions depending on their position on the globe, land use, fragmentation and pollutants levels can also largely vary across cities⁶⁹. In a general way, pollutants are known to affect DNA methylation and result in both hypo and hypermethylation, but the patterns of change observed largely rely on the pollutant involved⁷⁰. In this case, differences in cohorts of pollutants present in cities could be responsible for differences in patterns of methylations. Taken together, these results highlight the importance of questioning the assumption that cities are replicated environments that can be considered similar.

As mentioned earlier, increasing evidence suggests that DNA methylation can be associated with environmental and stress factors (env:¹², stress:⁷¹) especially during early development⁷². Here, we found four genes (POMC, ADAMTS3, PAPD4 et GCC1), associated with DMR that were previously described in great tits as undergoing major changes in methylation levels in case of exposure to higher levels of pollutants⁷³. Notably, the functions of these genes remain to be determined, and they could thus be interesting to target in future studies. In addition, the past literature has repeatedly found SERT and DRD4 as two major genes involved in urban specific avian human avoidance (or wariness) behaviours (see for example Riyahi *et al.*²³ in the Great tit, Garroway and Sheldon⁷⁴ in the blackbird, Van Dongen *et al.*⁷⁵ in the black swan & Mueller *et al.*⁷⁶ in the burrowing owl) (see SI Figure S8 to S12 to see patterns of methylation associated with 6 classical great tit linked candidate genes). In this study, while urban great tits show higher levels of aggressiveness in at least two of the cities (⁴³ & Caizergues *et al. in prep*) we found no DMR associated with these two genes in either of the three forest-city pairs. However, we found a significant urban-related change in methylation linked to the DRD3 gene, belonging to the same gene family as

DRD4 and known to be similarly involved in chicken aggressive behaviour⁷⁷. In line with these results, GO analyses revealed enrichment in genes associated with neuronal functions, behaviour, but also blood, immune and endocrine systems (Table S9, Figure S3), revealing the potential need of physiological adjustments in urban habitats. Surprisingly, a recent study on great tit differences of methylation between city and forest habitats in another European city found no GO enrichment in blood, while some in liver tissue²⁵ (note that they investigated DMSs, Differentially Methylated Sites, which differs from DMRs identified here). These contrasted results highlight the fact that methylation patterns highly depend on the analysed tissues¹¹, and show, once more, that urban linked methylation might not be similar from one city to another. In addition, it has been demonstrated that DNA methylation can undergo seasonal variation⁷⁸. Hence, analyses on multiple tissues and life-stages replicated in multiple pairs of urban and forest populations will allow to draw a broader view of the impact of urbanization on global methylation patterns and to understand replicated parallel occurrence across cities. However, tissue-specific and age-specific analyses in multiple individuals across several pairs of urban and forest environments poses major technical, budget and ethical limitations and should be coordinated very carefully. Additionally, specific drivers of shifts in methylation remain to be disentangled to understand which environmental factors are responsible for which change in methylation. To do so, experimental settings manipulating environmental factors such as performed by Mäkinen and collaborators⁷³ would be particularly useful. More integratively, information on how shifts in methylation patterns affect phenotype, fitness, and adaptation, often remain elusive⁷⁹. To our knowledge, a limited number of studies attempted to link methylation and expression levels in natural population contexts^{11,44}, and even fewer in urban habitats (but see e.g.^{24,25}). Hence, future work might need to tackle the question of the origin and adaptive significance of these variations in a controlled framework.

CONCLUSION

In this study, we found genomic footprints of selection and differentially methylated regions associated with urbanization, suggesting that genomic as well as epigenetic processes could play an

important role in the rapid adaptation to the urban habitat. To our knowledge, our study is the first to use replicated pairs of cities and forest habitats when studying urban linked methylation, offering a great opportunity to investigate convergent responses to anthropogenic environmental conditions. Taken together, our results revealed limited parallelism between cities regarding selection as well as methylation patterns, suggesting that cities might not present exactly similar environmental conditions or that different genetic pathways are involved in adaptation to urban environmental conditions, while still associated with similar biological functions. Furthermore, we highlight the need to unravel both environmental origins and evolutionary implications of methylation shifts, to understand to which extent environmental induced methylation can contribute to adaptation.

MATERIAL AND METHODS

Study sites and sampling

Three pairs of great tit populations in urban and forest environments were sampled in the three European cities of Barcelona (Spain), Montpellier (France) and Warsaw (Poland). For each location, 10 individuals were sampled within the city and 10 individuals were sampled from nearby forest. Blood samples were collected from breeding individuals during spring between 2016 and 2018 (except 2 individuals for Barcelona city collected in 2014 and 2015) and kept in alcohol or queen's buffer. Samples had balanced sex ratio (5 males and 5 females for each population) except for the forest population of Barcelona where 6 females and 4 males were sampled.

DNA extraction, RAD-seq and Reduced-Representation Bisulfite Sequencing

We used QIAGEN DNeasy blood and tissue kit to extract genomic DNA from blood samples following the provided instructions for nucleated blood samples. DNA was quantified using a NanoDrop ND8000 spectrophotometer and a Qubit 2.0 fluorometer with the DNA HS assay kit (Life Technologies). DNA quality was examined on agarose gels. We then performed RAD-sequencing and

RRBS-sequencing using standard protocols. For RAD-sequencing (restriction-site-associated DNA sequencing⁸⁰), the library preparation was done by the Montpellier GenomiX (MGX) platform (CNRS, Montpellier), using the enzyme SbfI. Each individual was identified using a unique six nucleotides tag, individuals were randomly multiplexed in equimolar proportions by libraries of 37 individuals. Each library was sequenced on a lane of an Illumina HiSeq 2500. Paired-end sequencing was used to produce 150 bp reads. This generated an average of 4.9M reads per individual. The DNA of 60 individuals were processed twice to test for reliability of the genotyping process. The RRBS-sequencing started with DNA digestion using MspI restriction enzymes, which cuts CCGG sites and targets regions that are CG rich, permitting to have a high proportion sequences in promoter regions. Each individual was identified using a unique six nucleotides tag. Individuals were randomly multiplexed in equimolar proportions by libraries of 10 individuals. Bisulfite treatment hence converted unmethylated cytosines into uracil, then converted to thymine after PCR amplification. Each library was then sequenced on a lane of an Illumina HiSeq 2500. Paired-end sequencing was used to produce 50 bp reads. This generated an average of 19.3M reads per individual.

Genomic analyses

Fastp v. 0.19.7⁸¹ was used to trim the RAD-seq reads, keeping reads with a minimum quality of 15 before mapping individual sequences against the reference genome of the Great tit (⁴⁴, GenBank assembly accession: GCA_001522545.3) with BWA v0.7.17⁸². Genotyping was conducted with stacks v2.41⁸³ “*gstacks*” and “*population*” functions, using “*snp*” model, filtering for mapping quality > 10, alpha = 0.05, minor allele frequency > 0.05 and observed heterozygosity < 0.65. Loci were retained if present in at least 90% of individuals in each population. Loci with extreme low or high coverage were removed (5% extremes filtered out using vcftools v0.1.15⁸⁴). After filtering, 74,137 SNPs were retained for subsequent population genomic analyses.

To document genomic variation among urban and rural great tits from the three locations we used a redundancy analyses (RDA), with location (Barcelona, Montpellier and Warsaw), environment (urban or rural) and sex as explanatory variables. Partial RDA was also produced to test for each

variable effect (environment, location or sex) alone after controlling for all other variables. The effect of a given factor was considered significant with a p-value < 0.05.

To estimate genome-wide differentiation between populations we used Weir and Cockerham's F_{ST} ³⁷ computed using the StAMPP R package⁸⁵. Average F_{ST} was estimated using all SNPs, and confidence intervals were assessed using 1000 bootstrap replicates.

To identify SNPs potentially under divergent selection between urban and forest habitats we first used Bayescan v2.1³⁸. Analyses were run for each pair of populations (Barcelona, Montpellier & Warsaw) separately, with default parameter option. As recommended by³⁸ we considered SNPs as outliers when they displayed a q-value above the 0.1 threshold.

To detect weaker footprints of divergent selection typically expected in polygenic adaptations in response to complex environmental heterogeneity, a multivariate method was used³⁹. Following a similar procedure as described above for the RDA analysis, we used a constrained RDA to investigate the effect of habitat (forest versus city) and to identify outliers SNPs displaying more than 3 times SD from the mean score on the constrained axis.

Methylation calling and analyses

The RRBS reads were first trimmed using fastp software v0.19.7⁸¹, and quality filtered keeping only reads with a quality > 15. BISMARCK software v0.20.0⁸⁶ was used for mapping reads on the masked reference genome with default parameters and 1 maximum allowed mismatches (-N 1). Methylation information for cytosine in a CpG context with sufficient coverage ($\geq 10x$) was extracted. An anova on mean individual methylation was run to test for location, habitat and sex effect on overall methylation levels.

Similarly, to genetic differentiation, an RDA was performed to describe epigenomic variation in function of location, habitat and sex. A partial RDA was also conducted to test for the habitat effect alone.

To identify differentially methylated regions (DMR) logistic regression including sex as covariate was performed after tiling regions of 1000pb for each location using MethylKit R package “*calculateDiffMeth*” function. Only DMR having > 10% overall difference between forest and urban habitat and q-value < 0.001 were kept. We then investigated if DMRs overlapped genes in 5 kb upstream and downstream regions using bedtools.

Genes associated to genomic outliers and DMRs, and gene ontology analyses

We investigated if genomic outliers and DMR overlapped genes in 5 kb upstream and downstream regions using BEDTools v2.28.0⁸⁷. Gene ontology analyses were performed with the R package topGO⁴⁰ to identify potential statistically enriched to investigate if on the list of all genes associated to RDA genomic outliers and the list of all genes associated to DMR.

ACKNOWLEDGMENTS

We thank all the people that contributed to fieldwork and blood sample collection, in particular the members of the PLT platform of the CEFÉ, as well as the fieldwork teams of Barcelona and Warsaw. We also thank Patricia Sourouille and the CEFÉ GEMEX platform for laboratory support, and Enrique Ortega-Aboud as well as Rémi Allio for their bioinformatics support. This research was funded by the European Research Council (Starting grant ERC-2013-StG-337365-SHE to AC) and the OSU-OREME, the Polish National Science Centre (Grants Sonata BIS 2014/14/E/NZ8/00386 and Opus 2016/21/B/NZ8/03082 to MS), and the Ministry of Economy and Competitiveness, Spanish Research Council (CGL-2016-79568-C3-3-P).

BIBLIOGRAPHY

1. Malhi, Y. (2017). The Concept of the Anthropocene. *Annu. Rev. Environ. Resour.* 42, 77–104.
2. Liu, Q.A. (2013). The impact of climate change on plant epigenomes. *Trends Genet.* 29, 503–505.

- 450 3. Jaenisch, R., and Bird, A. (2003). Signals, Epigenetic regulation of gene expression: how the
451 genome integrates intrinsic and environmental. *Nat. Genet.* 33, 245–254.
- 452 4. Duncan, E.J., Gluckman, P.D., and Dearden, P.K. (2014). Epigenetics, plasticity, and
453 evolution: How do we link epigenetic change to phenotype? *J. Exp. Zool. Part B Mol. Dev.*
454 *Evol.* 322, 208–220.
- 455 5. Waterland, R.A., and Jirtle, R.L. (2003). Transposable Elements: Targets for Early Nutritional
456 Effects on Epigenetic Gene Regulation. *Mol. Cell. Biol.* 23, 5293–5300.
- 457 6. Weaver, I.C.G., Cervoni, N., Champagne, F.A., D'Alession, A.C., Sharma, S., Seckl, J.R.,
458 Dymov, S., Szyf, M., and Meaney, M.J. (2004). Epigenetic programming by maternal
459 behavior. *Nat. Neurosci.* 7, 847–854.
- 460 7. Reik, W., Dean, W., and Walter, J. (2001). Epigenetic reprogramming in mammalian
461 development. *Science* (80-.). 293, 1089–1093.
- 462 8. Crews, D., Gore, A.C., Hsu, T.S., Dangleben, N.L., Spinetta, M., Schallert, T., Anway, M.D.,
463 and Skinner, M.K. (2007). Transgenerational epigenetic imprints on mate preference. *Proc.*
464 *Natl. Acad. Sci. U. S. A.* 104, 5942–5946.
- 465 9. Janowitz Koch, I., Clark, M.M., Thompson, M.J., Deere-Machemer, K.A., Wang, J., Duarte,
466 L., Gnanadesikan, G.E., McCoy, E.L., Rubbi, L., Stahler, D.R., et al. (2016). The concerted
467 impact of domestication and transposon insertions on methylation patterns between dogs and
468 grey wolves. *Mol. Ecol.* 25, 1838–1855.
- 469 10. Richards, C.L., Bossdorf, O., and Pigliucci, M. (2010). What Role Does Heritable Epigenetic
470 Variation Play in Phenotypic Evolution? *Bioscience* 60, 232–237.
- 471 11. Derks, M.F.L., Schachtschneider, K.M., Madsen, O., Schijlen, E., Verhoeven, K.J.F., and van
472 Oers, K. (2016). Gene and transposable element methylation in great tit (*Parus major*) brain
473 and blood. *BMC Genomics* 17, 332.

- 474 12. Foust, C.M., Preite, V., Schrey, A.W., Alvarez, M., Robertson, M.H., Verhoeven, K.J.F., and
475 Richards, C.L. (2016). Genetic and epigenetic differences associated with environmental
476 gradients in replicate populations of two salt marsh perennials. *Mol. Ecol.* 25, 1639–1652.
- 477 13. Danchin, E., Charmantier, A., Champagne, F.A., Mesoudi, A., Pujol, B., and Blanchet, S.
478 (2011). Beyond DNA: integrating inclusive inheritance into an extended theory of evolution.
479 *Nat. Rev. Genet.* 12, 475–486.
- 480 14. Brondizio, E.S., Settele, J., and Díaz, S. (2019). IPBES. 2019 Global assessment report on
481 biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on
482 Biodiversity and Ecosystem Services".
- 483 15. Szulkin, M., Garroway, C.J., Corsini, M., Kotarba, A.Z., and Dominoni, D.M. (2020). How to
484 Quantify Urbanization when Testing for Urban Evolution? In *Urban Evolutionary Biology*, M.
485 Szulkin, J. Munshi-South, and A. Charmantier, eds. (Oxford University Press), pp. 13–35.
- 486 16. Grimm, N.B., Faeth, S.H., Golubiewski, N.E., Redman, C.L., Wu, J., Bai, X., and Briggs, J.M.
487 (2008). Global Change and the Ecology of Cities. *Science* (80-.). 319, 756–760.
- 488 17. Szulkin, M., Munshi-South, J., and Charmantier, A. (2020). *Urban Evolutionary Biology*
489 (Oxford University Press).
- 490 18. Thompson, K.A., Rieseberg, L.H., and Schluter, D. (2018). Speciation and the City. *Trends*
491 *Ecol. Evol.* 33, 815–826.
- 492 19. Alberti, M., Correa, C., Marzluff, J.M., Hendry, A.P., Palkovacs, E.P., Gotanda, K.M., Hunt,
493 V.M., Apgar, T.M., and Zhou, Y. (2017). Global urban signatures of phenotypic change in
494 animal and plant populations. *Proc. Natl. Acad. Sci. U. S. A.* 114, 8951–8956.
- 495 20. Perrier, C., Lozano del Campo, A., Szulkin, M., Demeyrier, V., Gregoire, A., and Charmantier,
496 A. (2018). Great tits and the city: Distribution of genomic diversity and gene–environment
497 associations along an urbanization gradient. *Evol. Appl.* 11, 593–613.

- 498 21. Mueller, J.C., Partecke, J., Hatchwell, B.J., Gaston, K.J., and Evans, K.L. (2013). Candidate
499 gene polymorphisms for behavioural adaptations during urbanization in blackbirds. *Mol. Ecol.*
500 22, 3629–3637.
- 501 22. Salmón, P., Jacobs, A., Ahrén, D., Biard, C., Dingemanse, N., Dominoni, D., Helm, B.,
502 Lundberg, M., Senar, J.C., Sprau, P., et al. (2020). Repeated genomic signatures of adaptation
503 to urbanisation in a songbird across Europe. *bioRxiv*, 2020.05.05.078568.
- 504 23. Riyahi, S., Sánchez-Delgado, M., Calafell, F., Monk, D., and Senar, J.C. (2015). Combined
505 epigenetic and intraspecific variation of the *DRD4* and *SERT* genes influence novelty seeking
506 behavior in great tit *Parus major*. *Epigenetics* 10, 516–525.
- 507 24. McNew, S.M., Beck, D., Sadler-Riggleman, I., Knutie, S.A., Koop, J.A.H., Clayton, D.H., and
508 Skinner, M.K. (2017). Epigenetic variation between urban and rural populations of Darwin’s
509 finches. *BMC Evol. Biol.* 17, 183.
- 510 25. Watson, H., Powell, D., Salmón, P., Jacobs, A., and Isaksson, C. (2020). Urbanization is
511 associated with modifications in DNA methylation in a small passerine bird. *Evol. Appl.*,
512 eea.13160.
- 513 26. Ding, R., Jin, Y., Liu, X., Ye, H., Zhu, Z., Zhang, Y., Wang, T., and Xu, Y. (2017). Dose- and
514 time- effect responses of DNA methylation and histone H3K9 acetylation changes induced by
515 traffic-related air pollution. *Sci. Rep.* 7, 1–9.
- 516 27. Santangelo, J.S., Rivkin, L.R., and Johnson, M.T.J. (2018). The evolution of city life.
517 *Proceedings. Biol. Sci.* 285, 20181529.
- 518 28. Rivkin, L.R., Santangelo, J.S., Alberti, M., Aronson, M.F.J., de Keyser, C.W., Diamond, S.E.,
519 Fortin, M.-J., Frazee, L.J., Gorton, A.J., Hendry, A.P., et al. (2018). A roadmap for urban
520 evolutionary ecology. *Evol. Appl.*
- 521 29. Theodorou, P., Radzevičiūtė, R., Kahnt, B., Soro, A., Grosse, I., and Paxton, R.J. (2018).

- 522 Genome-wide single nucleotide polymorphism scan suggests adaptation to urbanization in an
523 important pollinator, the red-tailed bumblebee (*Bombus lapidarius* L.). *Proc. R. Soc. B Biol.*
524 *Sci.* 285.
- 525 30. Mueller, J.C., Kuhl, H., Boerno, S., Tella, J.L., Carrete, M., and Kempenaers, B. (2018).
526 Evolution of genomic variation in the burrowing owl in response to recent colonization of
527 urban areas. *Proc. R. Soc. B Biol. Sci.* 285, 20180206.
- 528 31. Santangelo, J.S., Miles, L.S., Breitbart, S.T., Murray-Stoker, D., Rivkin, L.R., Johnson, M.T.J.,
529 and Ness, R.W. (2020). Urban Environments as a Framework to Study Parallel Evolution. In
530 Urban Evolutionary Biology, M. Szulkin, J. Munshi-South, and A. Charmantier, eds. (Oxford
531 University Press), pp. 36–53.
- 532 32. Perrier, C., Caizergues, A.E., and Charmantier, A. (2020). Adaptation Genomics in Urban
533 Environments. In Urban Evolutionary Biology, M. Szulkin, J. Munshi-South, and A.
534 Charmantier, eds. (Oxford University Press), pp. 72–88.
- 535 33. Corsini, M., Schöll, E.M., Di Lecce, I., Chatelain, M., Dubiec, A., and Szulkin, M. (2020).
536 Growing in the city: Urban evolutionary ecology of avian growth rates. *Evol. Appl.*
- 537 34. Biard, C., Brischoux, F., Meillère, A., Michaud, B., Nivière, M., Ruault, S., Vaugoyeau, M.,
538 and Angelier, F. (2017). Growing in Cities: An Urban Penalty for Wild Birds? A Study of
539 Phenotypic Differences between Urban and Rural Great Tit Chicks (*Parus major*). *Front. Ecol.*
540 *Evol.* 5.
- 541 35. Chamberlain, D., Hatchwell, B., and Gaston, K.J. (2009). Avian productivity in urban
542 landscapes: A review and meta-analysis Avian productivity in urban landscapes: a review
543 and meta-analysis. *Ibis (Lond. 1859).*, 1–18.
- 544 36. Caizergues, A.E., Charmantier, A., Lambrechts, M.M., Perret, S., Demeyrier, V., Lucas, A.,
545 and Grégoire, A. (2021). An avian urban morphotype: how the city environment shapes great
546 tit morphology at different life stages. *Urban Ecosyst.*, 1–13.

- 547 37. Weir, B.S., and Cockerham, C.C. (1984). Estimating F-Statistics for the Analysis of Population
548 Structure. *Evolution* (N. Y). 38, 1358.
- 549 38. Foll, M., and Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate
550 for both dominant and codominant markers: A Bayesian perspective. *Genetics* 180, 977–993.
- 551 39. Forester, B.R., Jones, M.R., Joost, S., Landguth, E.L., and Lasky, J.R. (2016). Detecting spatial
552 genetic signatures of local adaptation in heterogeneous landscapes. *Mol. Ecol.* 25, 104–120.
- 553 40. Alexa, A., and Rahnenführer, J. (2009). Gene set enrichment analysis with topGO.
- 554 41. Lowry, H., Lill, A., and Wong, B.B.M. (2013). Behavioural responses of wildlife to urban
555 environments. *Biol. Rev.* 88, 537–549.
- 556 42. Sepp, T., McGraw, K.J., Kaasik, A., and Giraudeau, M. (2017). A review of urban impacts on
557 avian life-history evolution: Does city living lead to slower pace of life? *Glob. Chang. Biol.*, 1–
558 18.
- 559 43. Riyahi, S., Björklund, M., Mateos-Gonzalez, F., and Senar, J.C. (2017). Personality and
560 urbanization: behavioural traits and DRD4 SNP830 polymorphisms in great tits in Barcelona
561 city. *J. Ethol.* 35, 101–108.
- 562 44. Laine, V.N., Gossmann, T.I., Schachtschneider, K.M., Garroway, C.J., Madsen, O., Verhoeven,
563 K.J.F., De Jager, V., Megens, H.J., Warren, W.C., Minx, P., et al. (2016). Evolutionary signals
564 of selection on cognition from the great tit genome and methylome. *Nat. Commun.* 7, 1–9.
- 565 45. Senar, J.C., and Björklund, M. (2020). Recent spread of blue tits into the Barcelona urban
566 environment: morphological differences and the role of balanced dispersal. *Evol. Ecol.*, 1–17.
- 567 46. Kvist, L., Martens, J., Higuchi, H., Nazarenko, A.A., Valchuk, O.P., and Orell, M. (2003).
568 Evolution and genetic structure of the great tit (*Parus major*) complex. *Proc. R. Soc. London.*
569 *Ser. B Biol. Sci.* 270, 1447–1454.

- 570 47. Reid, N.M., Proestou, D.A., Clark, B.W., Warren, W.C., Colbourne, J.K., Shaw, J.R.,
571 Karchner, S.I., Hahn, M.E., Nacci, D., Oleksiak, M.F., et al. (2016). The genomic landscape of
572 rapid repeated evolutionary adaptation to toxic pollution in wild fish. *Science* (80-.). 354,
573 1305–1308.
- 574 48. Losos, J.B. (2011). Convergence, adaptation, and constraint. *Evolution* (N. Y). 65, 1827–1840.
- 575 49. Thompson, C.J., Ahmed, N.I., Veen, T., Peichel, C.L., Hendry, A.P., Bolnick, D.I., and Stuart,
576 Y.E. (2017). Many-to-one form-to-function mapping weakens parallel morphological
577 evolution. *Evolution* (N. Y). 71, 2738–2749.
- 578 50. Boyle, E.A., Li, Y.I., and Pritchard, J.K. (2017). An Expanded View of Complex Traits: From
579 Polygenic to Omnigenic. *Cell* 169, 1177–1186.
- 580 51. Shochat, E., Warren, P.S., Faeth, S.H., McIntyre, N.E., and Hope, D. (2006). From patterns to
581 emerging processes in mechanistic urban ecology. *Trends Ecol. Evol.* 21, 186–191.
- 582 52. Lande, R. (1979). Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain:
583 Body Size Allometry. *Evolution* (N. Y). 33, 402.
- 584 53. Zhou, X., and Stephens, M. (2012). Genome-wide efficient mixed-model analysis for
585 association studies. *Nat. Genet.* 44, 821–824.
- 586 54. Robinson, M.R., Wray, N.R., and Visscher, P.M. (2014). Explaining additional genetic
587 variation in complex traits. *Trends Genet.* 30, 124–132.
- 588 55. McNulty, S.E., Barrett, R.M., Vogel-Ciernia, A., Malvaez, M., Hernandez, N., Davatolhagh,
589 M.F., Matheos, D.P., Schiffman, A., and Wood, M.A. (2012). Differential roles for Nr4a1 and
590 Nr4a2 in object location vs. object recognition long-term memory. *Learn. Mem.* 19, 588–592.
- 591 56. Kim, Y.H., Peregrine, J., and Arnold, A.P. (2006). The distribution of expression of
592 doublecortin (DCX) mRNA and protein in the zebra finch brain. *Brain Res.* 1106, 189–196.

- 593 57. Taufique, S.K.T., Prabhat, A., and Kumar, V. (2018). Illuminated night alters hippocampal
594 gene expressions and induces depressive □like responses in diurnal corvids. *Eur. J. Neurosci.*
595 48, 3005–3018.
- 596 58. Buitenhuis, B., Hedegaard, J., Janss, L., and Sørensen, P. (2009). Differentially expressed
597 genes for aggressive pecking behaviour in laying hens. *BMC Genomics* 10, 544.
- 598 59. Sprau, P., and Dingemanse, N.J. (2017). An Approach to Distinguish between Plasticity and
599 Non-random Distributions of Behavioral Types Along Urban Gradients in a Wild Passerine
600 Bird. *Front. Ecol. Evol.* 5, 92.
- 601 60. Sih, A., and Del Giudice, M. (2012). Linking behavioural syndromes and cognition: A
602 behavioural ecology perspective. *Philos. Trans. R. Soc. B Biol. Sci.* 367, 2762–2772.
- 603 61. Sol, D., Lapiedra, O., and González-Lagos, C. (2013). Behavioural adjustments for a life in the
604 city. *Anim. Behav.* 85, 1101–1112.
- 605 62. Torné-Noguera, A., Pagani-Núñez, E., and Senar, J.C. (2014). Great Tit (*Parus major*) breath
606 rate in response to handling stress: urban and forest birds differ. *J. Ornithol.* 155, 315–318.
- 607 63. Senar, J.C., Garamszegi, L.Z., Tilgar, V., Biard, C., Moreno-Rueda, G., Salmón, P., Rivas,
608 J.M., Sprau, P., Dingemanse, N.J., Charmantier, A., et al. (2017). Urban Great Tits (*Parus*
609 *major*) Show Higher Distress Calling and Pecking Rates than Rural Birds across Europe. *Front.*
610 *Ecol. Evol.* 5, 163.
- 611 64. Waters, S.A., Capraro, A., McIntyre, K.L., Graves, J.A.M., and Waters, P.D. (2018). The
612 methylome of vertebrate sex chromosomes. *Genes (Basel).* 9, 230.
- 613 65. Teranishi, M., Shimada, Y., Hori, T., Nakabayashi, O., Kikuchi, T., Macleod, T., Pym, R.,
614 Sheldon, B., Solovei, I., Macgregor, H., et al. (2001). Transcripts of the MHM region on the
615 chicken Z chromosome accumulate as non-coding RNA in the nucleus of female cells adjacent
616 to the DMRT1 locus. *Chromosom. Res.* 9, 147–165.

- 617 66. Donihue, C.M., and Lambert, M.R. (2015). Adaptive evolution in urban ecosystems. *Ambio*
618 44, 194–203.
- 619 67. Caizergues, A.E., Grégoire, A., and Charmantier, A. (2018). Urban versus forest ecotypes are
620 not explained by divergent reproductive selection. *Proc. R. Soc. B Biol. Sci.* 285, 20180261.
- 621 68. Seress, G., Sándor, K., Evans, K.L., and Liker, A. (2020). Food availability limits avian
622 reproduction in the city: An experimental study on great tits *Parus major*. *J. Anim. Ecol.*, 1365-
623 2656.13211.
- 624 69. Cárdenas Rodríguez, M., Dupont-Courtade, L., and Oueslati, W. (2016). Air pollution and
625 urban structure linkages: Evidence from European cities. *Renew. Sustain. Energy Rev.* 53, 1–9.
- 626 70. Head, J.A. (2014). Patterns of DNA methylation in animals: An ecotoxicological perspective.
627 In *Integrative and Comparative Biology* (Oxford University Press), pp. 77–86.
- 628 71. Sun, L., Miao, X., Cui, J., Deng, J., Wang, X., Wang, Y., Zhang, Y., Gao, S., and Yang, K.
629 (2018). Genome-wide high-resolution mapping of DNA methylation identifies epigenetic
630 variation across different salt stress in Maize (*Zea mays* L.). *Euphytica* 214, 1–15.
- 631 72. Meaney, M.J., and Szyf, M. (2005). Environmental programming of stress responses through
632 DNA methylation: Life at the interface between a dynamic environment and a fixed genome.
633 *Dialogues Clin. Neurosci.* 7, 103–123.
- 634 73. Mäkinen, H., van Oers, K., Eeva, T., Laine, V.N., and Ruuskanen, S. (2019). The effect of
635 experimental lead pollution on DNA methylation in a wild bird population. *bioRxiv*, 851998.
- 636 74. Garroway, C.J., and Sheldon, B.C. (2013). Urban behavioural adaptation. *Mol. Ecol.* 22, 3430–
637 3432.
- 638 75. Van Dongen, W.F.D., Robinson, R.W., Weston, M.A., Mulder, R.A., and Guay, P.J. (2015).
639 Variation at the DRD4 locus is associated with wariness and local site selection in urban black
640 swans. *BMC Evol. Biol.* 15, 253.

- 641 76. Mueller, J.C., Carrete, M., Boerno, S., Kuhl, H., Tella, J.L., and Kempnaers, B. (2020). Genes
642 acting in synapses and neuron projections are early targets of selection during urban
643 colonization. *Mol. Ecol.* 29, 3403–3412.
- 644 77. Li, Z., Zheng, M., Abdalla, B.A., Zhang, Z., Xu, Z., Ye, Q., Xu, H., Luo, W., Nie, Q., and
645 Zhang, X. (2016). Genome-wide association study of aggressive behaviour in chicken. *Sci.*
646 *Rep.* 6, 1–11.
- 647 78. Viitaniemi, H.M., Verhagen, I., Visser, M.E., Honkela, A., van Oers, K., and Husby (2019).
648 Seasonal variation in genome-wide DNA methylation patterns and the onset of seasonal timing
649 of reproduction in great tits. *Genome Biol. Evol.* 11, 970–983.
- 650 79. Sepers, B., van den Heuvel, K., Lindner, M., Viitaniemi, H., Husby, A., and van Oers, K.
651 (2019). Avian ecological epigenetics: pitfalls and promises. *J. Ornithol.* 160, 1183–1203.
- 652 80. Baird, N.A., Etter, P.D., Atwood, T.S., Currey, M.C., Shiver, A.L., Lewis, Z.A., Selker, E.U.,
653 Cresko, W.A., and Johnson, E.A. (2008). Rapid SNP Discovery and Genetic Mapping Using
654 Sequenced RAD Markers. *PLoS One* 3, e3376.
- 655 81. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ
656 preprocessor. In *Bioinformatics* (Oxford University Press), pp. i884–i890.
- 657 82. Li, H., and Durbin, R. (2010). Fast and accurate long-read alignment with Burrows–Wheeler
658 transform. *Bioinformatics* 26, 589–595.
- 659 83. Rochette, N.C., Rivera-Colón, A.G., and Catchen, J.M. (2019). Stacks 2: Analytical methods
660 for paired-end sequencing improve RADseq-based population genomics. *Mol. Ecol.* 28, 4737–
661 4754.
- 662 84. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker,
663 R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al. (2011). The variant call format and VCFtools.
664 *Bioinformatics* 27, 2156–2158.

665 85. Pembleton, L., Cogan, N., and Forster, J. (2013). StAMPP: an R package for calculation of
666 genetic differentiation and structure of mixed-ploidy level populations. *Mol. Ecol. Resour.* *13*,
667 946–952.

668 86. Krueger, F., and Andrews, S.R. (2011). Bismark: a flexible aligner and methylation caller for
669 Bisulfite-Seq applications. *Bioinformatics* *27*, 1571–1572.

670 87. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
671 genomic features. *Bioinformatics* *26*, 841–842.

672

FIGURES

Figure 1: (A) Great tit blood sample locations in Europe (in urban and forest sites in and near Barcelona, Montpellier & Warsaw). (B) Distribution of mean percent of methylation on autosomes and on the Z chromosome, compared between habitats (forest versus urban), sexes and locations. (C) Redundancy analysis (RDA) on genomic data (74,137 filtered SNPs). (D) RDA on methylation levels (based on methylation levels observed at 157,741 positions). In (C) and (D), triangles represent rural habitats, circles represent urban habitats, empty and solid symbols represent females and males respectively. *** P -value < 0.001 , ** P -value < 0.01 and * P -value < 0.05 , related to the explanatory factors.

Figure 2: Manhattan plot of mean F_{ST} between urban and forest populations along the Great tit genome for (A) Barcelona, (B) Montpellier and (C) Warsaw. Dark orange (A), purple (B) and light orange points (C) represent significant outlier SNPs identified by the F_{ST} -outlier test Bayescan for each population pair, given with their associated genes in 5 kb. Green points represent outliers found with the multivariate RDA approach. A few SNPs were identified by both the Bayescan and the RDA methods and signalled as a green point circled with the colour used for the considered pair.

SUPPLEMENTARY INFORMATION

Testing for parallel genomic and epigenomic footprints of adaptation to urban life in a passerine bird

Authors:

Aude E. Caizergues^{1*}, Jeremy Le Luyer², Arnaud Grégoire¹, Marta Szulkin³, Juan-Carlos Senar⁴, Anne Charmantier^{1†}, Charles Perrier^{5†}

¹ CEFÉ, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

² Ifremer, IRD, Institut Louis Malardé, Univ Polynésie Française, EIO, F-98719 Taravao, Tahiti, Polynésie Française

³ Centre of New Technologies, University of Warsaw, S. Banacha 2c, 02-097 Warsaw, Poland

⁴ Museu de Ciències Naturals de Barcelona, Parc Ciutadella, 08003 Barcelona, Spain

⁵ CBGP, INRAe, CIRAD, IRD, Montpellier SupAgro, Univ. Montpellier, Montpellier, France

† *shared senior authorship*

*** Corresponding author:** Aude E. Caizergues, 1919 route de Mende, 34293 Montpellier cedex 5, FRANCE

Email : audeemiliecaizergues@gmail.com

SUPPLEMENTARY TABLES

Table S1: Redundancy analysis (RDA) performed on the genetic data including the Z chromosome.

	adjusted R-squared	P-value	RDA1	RDA2	RDA3	RDA4
Full RDA	0.018	0.001	% variance explained by axes			
			0.024	0.021	0.021	0.019
Variables			Biplot scores			
City – Montpellier		0.001	0.57	-0.819	0.059	-0.039
City – Warsaw			0.381	0.883	0.214	0.17
Habitat – Urban		0.001	-0.21	-0.1	0.968	-0.089
Sex – Male		0.004	0.184	0.143	-0.0358	-0.972
Partial RDA for city	0.012	0.001	% variance explained by axe			
			0.025		0.022	
Variable			Biplot scores			
City – Montpellier		0.001	0.63		-0.776	
City – Warsaw			0.356		0.934	
Partial RDA for habitat	0.004	0.001	% variance explained by axe			
					0.022	
Variable			Biplot score			
Habitat – Urban		0.001			0.999	
Partial RDA for sex	0.002	0.004	% variance explained by axe			
					0.02	
Variable			Biplot score			
Sex – Male		0.005			-0.999	

Table S2: Redundancy analysis (RDA) performed on the genetic data without Z chromosome.

	adjusted R-squared	P-value	RDA1	RDA2	RDA3	RDA4
Full RDA	0.016	0.001	% variance explained by axes			
			0.024	0.021	0.021	0.017
Variables			Biplot scores			
City – Montpellier		0.001	0.5725	-0.818	0.049	-0.0244
City – Warsaw			0.382	0.889	0.227	0.11
Habitat – Urban		0.001	-0.215	-0.119	0.967	-0.073
Sex – Male		0.26	0.14	0.096	-0.033	-0.985
Partial RDA for city	0.012	0.001	% variance explained by axe			
			0.025		0.022	
Variable			Biplot scores			
City – Montpellier		0.001	0.624		-0.781	
City – Warsaw			0.362		0.932	
Partial RDA for habitat	0.004	0.001	% variance explained by axe			
			0.022			
Variable			Biplot score			
Habitat – Urban		0.001	0.999			
Partial RDA for sex	0.0004	0.373	% variance explained by axe			
			0.018			
Variable			Biplot score			
– Urban		0.369	-0.999			

Table S3: Fst estimation between pairs of subpopulations. 95% confidence computed intervals in brackets were computed using StAMPP package with 1000 bootstrap.

	Mont-Rur	Mont-Urb	Var-Rur	Var-Urb	Bar-Rur
Mont-Urb	0.012 (0.011 – 0.013)	-	-	-	-
Var-Rur	0.009 (0.009 – 0.010)	0.018 (0.017 – 0.019)	-	-	-
Var-Urb	0.018 (0.017 – 0.018)	0.026 (0.025 – 0.027)	0.018 (0.017 – 0.019)	-	-
Bar-Rur	0.009 (0.009 – 0.010)	0.019 (0.017 – 0.018)	0.009 (0.009 – 0.010)	0.018 (0.017 – 0.019)	-
Bar-Urb	0.025 (0.024 – 0.027)	0.034 (0.033 – 0.035)	0.025 (0.025 – 0.026)	0.034 (0.033 – 0.034)	0.018 (0.017 – 0.019)

Table S4: Fst estimation averaged on autosomes between pairs of subpopulations. 95% confidence computed intervals in brackets were computed using StAMPP package with 1000 bootstrap.

	Mont-Rur	Mont-Urb	Var-Rur	Var-Urb	Bar-Rur
Mont-Urb	0.017 (0.017 – 0.017)	-	-	-	-
Var-Rur	0.018 (0.018 – 0.018)	0.015 (0.014 – 0.015)	-	-	-
Var-Urb	0.014 (0.014 – 0.015)	0.011 (0.011 – 0.011)	0.015 (0.014 – 0.015)	-	-
Bar-Rur	0.018 (0.018 – 0.018)	0.015 (0.014 – 0.015)	0.018 (0.018 – 0.018)	0.014 (0.014 – 0.014)	-
Bar-Urb	0.011 (0.011 – 0.011)	0.008 (0.007 – 0.008)	0.011 (0.011 – 0.011)	0.008 (0.007 – 0.008)	0.014 (0.014 – 0.014)

Table S5: Fst estimation averaged on Z chromosome between pairs of subpopulations. 95% confidence computed intervals in parenthesis were computed using StAMPP package with 1000

	Mont-Rur	Mont-Urb	Var-Rur	Var-Urb	Bar-Rur
Mont-Urb	0.013 (0.006 – 0.018)	-	-	-	-
Var-Rur	0.016 (0.009 – 0.021)	0.020 (0.013 – 0.025)	-	-	-
Var-Urb	0.025 (0.019 – 0.031)	0.025 (0.020 – 0.032)	0.020 (0.015 – 0.025)	-	-
Bar-Rur	0.015 (0.010 – 0.019)	0.017 (0.011 – 0.022)	0.014 (0.009 – 0.019)	0.018 (0.015 – 0.024)	-
Bar-Urb	0.031 (0.026 – 0.036)	0.041 (0.032 – 0.047)	0.028 (0.022 – 0.034)	0.036 (0.030 – 0.043)	0.014 (0.009 – 0.021)

bootstrap.

Table S6: Redundancy analysis (RDA) performed on the methylation data including the Z chromosome.

	adjusted R-squared	P-value	RDA1	RDA2	RDA3	RDA4
Full RDA	0.007	0.001	% variance explained by axes			
			0.021	0.019	0.018	0.016
Variables			Biplot scores			
City – Montpellier		0.001	0.21	0.259	-0.648	0.685
City – Warsaw			-0.603	-0.77	0.198	-0.067
Habitat – Urban		0.03	-0.256	0.286	0.669	0.637
Sex – Male		0.001	0.736	-0.562	0.263	0.269
Partial RDA for city	0.003	0.001	% variance explained by axe			
			0.021		0.018	
Variable			Biplot scores			
City – Montpellier		0.001	-0.356		0.934	
City – Warsaw			0.986		-0.161	
Partial RDA for habitat	0.001	0.3	% variance explained by axe			
			0.019			
Variable			Biplot score			
Habitat – Urban		0.181	0.999			
Partial RDA for sex	0.003	0.001	% variance explained by axe			
			0.021			
Variable			Biplot score			
Sex – Male		0.001	0.999			

Table S7: Redundancy analysis (RDA) performed on the methylation data without Z chromosome.

769

	adjusted R-squared	P-value	RDA1	RDA2	RDA3	RDA4
Full RDA	0.006	0.001	% variance explained by axes			
			0.021	0.019	0.018	0.016
Variables			Biplot scores			
City – Montpellier		0.001	0.231	-0.204	-0.666	0.679
City – Warsaw			-0.645	0.761	0.2467	-0.058
Habitat – Urban		0.019	-0.261	-0.365	0.632	0.632
Sex – Male		0.001	0.698	0.569	0.326	0.289
Partial RDA for city	0.003	0.001	% variance explained by axe			
			0.02		0.018	
Variable			Biplot scores			
City – Montpellier		0.001	-0.357		0.933	
City – Warsaw			0.987		-0.159	
Partial RDA for habitat	0.001	0.176	% variance explained by axe			
			0.019			
Variable			Biplot score			
Habitat – Urban		0.168	0.999			
Partial RDA for sex	0.003	0.001	% variance explained by axe			
			0.021			
Variable			Biplot score			
Sex – Male		0.001	0.999			

770

Table S8: Results of Type I ANOVA performed on mean methylation level of cytosines in a CpG context per individual, including: habitat (rural vs urban), city (Barcelona, Montpellier and Warsaw) and sex as explanatory variables. The first part of the table shows results of the analysis on autosomes and the second part shows results of the analysis on the Z chromosome only.

Autosomes					
Effect	Df	Sum Sq	Mean Sq	F value	P
Habitat	1	0.416	0.416	0.511	0.478
Sex	1	1.041	1.041	1.278	0.263
City	2	5.408	2.704	3.319	0.044
Habitat* City	2	0.316	0.158	0.194	0.825
Residuals	53	42.174	0.8146		
Z chromosome					
Effect	Df	Sum Sq	Mean Sq	F value	P
Habitat	1	0.012	0.012	0.011	0.915
Sex	1	133.584	133.584	125.077	1.45x10 ⁻¹⁵
City	2	0.979	0.489	0.458	0.635
Habitat* City	2	1.1879	0.594	0.556	0.577
Residuals	53	56.605	1.068		

Table S9: Significantly enriched GO associated with genes overlapping 5kb windows around genomic outliers between forest and urban habitats.

GO.ID	Term	Annotated	Significant	Expected	Weight	Associated Genes
GO:2000618	regulation of histone H4-K16 acetylation	5	3	0.34	0.003	ATG5, AUTS2, SIRT1
GO:1902287	semaphorin-plexin signaling pathway invo...	7	3	0.48	0.009	PLXNA1, PLXNA2, PLXNA4
GO:1901842	negative regulation of high voltage-gate...	3	3	0.21	0.000	DYSF, GEM, RRAD
GO:0090335	regulation of brown fat cell differentia...	10	4	0.69	0.005	FTO, PIM1, PRDM16, SIRT1
GO:0071391	cellular response to estrogen stimulus	6	3	0.41	0.006	AR, CRHBP, ESR1
GO:0071277	cellular response to calcium ion	28	7	1.93	0.002	CRHBP, ECT2, HPCA, RASAL1, SYT1, SYT11, SYT9
GO:0050953	sensory perception of light stimulus	57	6	3.92	0.026	CDH23, CRYBB2, NYX, PCDH15, RPE65, WHRN
GO:0050765	negative regulation of phagocytosis	7	3	0.48	0.009	ATG5, DYSF, SYT11
GO:0048846	axon extension involved in axon guidance	23	4	1.58	0.026	NRP2, PLXNA4, SEMA4G, SLIT3
GO:0048026	positive regulation of mRNA splicing vi...	6	3	0.41	0.006	HMX2, NCBP1, NUP98
GO:0045777	positive regulation of blood pressure	9	3	0.62	0.026	ADRA1D, GRIP2, WNK1
GO:0044351	macropinocytosis	6	3	0.41	0.006	DOCK2, MAPKAPK2, SNX33
GO:0043410	positive regulation of MAPK cascade	197	15	13.56	0.013	ADRA1D, ADRB2, APP, ARAXIN1, EPHA8, FGFR3, MTURN, NRG1, PDGFRB, PLCB1, RET, ROR1, ROR2, SH3RF2
GO:0043087	regulation of GTPase activity	194	17	13.36	0.025	AGAP3, ARHGAP17, ARHGAP45, ARHGEF12, DOCK2, ECT2, IQGAP2, MLST8, PLCB1, PLXNA1, PLXNA2, PLXNA4, RASAL1, RDX, RGS8, SIPA1L1, WNK1
GO:0042535	positive regulation of tumor necrosis fa...	6	3	0.41	0.006	APP, HSPB1, MAPKAPK2
GO:0042311	vasodilation	13	4	0.89	0.010	ADRB2, ATG5, CPS1, NOS1
GO:0035418	protein localization to synapse	26	5	1.79	0.009	BSN, GPC4, GRIP2, HSPB1, NRXN1
GO:0033555	multicellular organismal response to str...	37	6	2.55	0.006	BRINP1, DBH, NOS1, NR4A2, PPP3CA, RET
GO:0021987	cerebral cortex development	56	8	3.86	0.013	CCDC85C, CNTNAP2, DCX, HIF1A, NRP2, PAX5, PLCB1, TRAPPC9
GO:0016339	calcium-dependent cell-cell adhesion via...	15	4	1.03	0.016	CDH13, CDH17, CDH19, CDH20
GO:0009755	hormone-mediated signaling pathway	87	12	5.99	0.014	AR, ASIP, CRHBP, ESR1, ESRRB, KMT2D, NR5A1, PPARG, PRLR, REN, RNF14, SIRT1
GO:0008344	adult locomotory behavior	46	7	3.17	0.006	APP, CHL1, FGF12, NR4A2, NTAN1, PCDH15, SEZ6L
GO:0007274	neuromuscular synaptic transmission	9	4	0.62	0.002	ADARB1, CHRNA1, FCHSD2, SLC5A7
GO:0007171	activation of transmembrane receptor pro...	8	3	0.55	0.014	DGKQ, NRG1, PRLR
GO:0007156	homophilic cell adhesion via plasma memb...	57	11	3.92	0.001	CDH13, CDH17, CDH19, CDH20, CDH23, CNTN4, DSCAML1, IGSF11, MYOT, PCDH15, RET
GO:0007155	cell adhesion	508	53	34.97	0.015	ADAMTS12, ADGRL3, AMIGO3, ASS1, ASTN2, CDH13, CDH17, CDH19, CDH20, CDH23, CHL1, CLDN23, CNTN4, COL8A1, CTNNA1, CTNNA3, DSCAML1, DTX1, DYSF, EGFLAM, EPHA8, GPC4, IGSF11, ITGA11, ITGB2, LAMA3, LSAMP, MACF1, MAD1L1, MEGF9, MYH9, MYOT, NCAM1, NEGR1, NFKBIZ, NRXN1, PARD3B, PARVA, PCDH15, PLXNA1, PLXNA2, PLXNA4, PPP3CA, RDX, RET, STAB2, STK10, TLN1, TMEFF2, TNC, VTN, VWF, ZDHHC2
GO:0003073	regulation of systemic arterial blood pr...	39	6	2.68	0.003	ADRA1D, ADRB2, AR, NCALD, REN, WNK1
GO:0001967	suckling behavior	7	3	0.48	0.009	APP, CNTFR, UBE2Q1
GO:0001778	plasma membrane repair	7	3	0.48	0.009	DYSF, MYH9, SYT11

Table S10: Significantly enriched GO associated with genes overlapping 5kb windows around DMRs between forest and urban habitats.

GO.ID	Term	Annotated	Significant	Expected	Weight	Barcelona	Montpellier	Warsaw
GO:0050804	modulation of chemical synaptic transmis...	190	10	2.92	0.016	CACNG3, CPLX2, OPHN1, PPP1R9B, PRKCB	CDH11	DRD3, EPHB1, NAPB, SNAPIN
GO:0042981	regulation of apoptotic process	619	8	9.52	0.004	DIP2A, HERPUD1, NR4A1, SOD1, ZBTB7A		ANP32E, RSL1D1, UACA
GO:0007626	locomotory behavior	122	6	1.88	0.005	ADCY6, ELAVL4, SOD1		CDH23, DRD3, SEZ6L
GO:0006816	calcium ion transport	173	6	2.66	0.024	CACNB3, CACNG3, PRKCB		CDH23, DRD3, STIM1
GO:0050728	negative regulation of inflammatory resp...	61	5	0.94	0.011	PSMB4, SOD1,	CD276, WFDC1	PBK
GO:0046677	response to antibiotic	91	5	1.4	0.012	ADCY6, SOD1	RSRC1	AOC1, DRD3
GO:0043087	regulation of GTPase activity	237	5	3.65	0.025	GAPVD1, OPHN1, RANBP3L, SOD1	PROM2	PROM2
GO:2000300	regulation of synaptic vesicle exocytosi...	40	4	0.62	0.017	CPLX2	PRKCB	NAPB, SNAPIN
GO:0008217	regulation of blood pressure	74	4	1.14	0.040	SOD1	POMC	ADRA1B, DRD3
GO:0044060	regulation of endocrine process	22	3	0.34	0.012	NR3C1,	POMC	RAB11FIP3
GO:0055072	iron ion homeostasis	42	3	0.65	0.014	NEO1, SOD1	HMOX2	
GO:0071559	response to transforming growth factor b...	113	3	1.74	0.031	NR3C1, ZBTB7A, ZFH3		
GO:0050852	T cell receptor signaling pathway	52	3	0.8	0.046	CACNB3, SKAP1	CD276	
GO:0007200	phospholipase C-activating G protein-	40	3	0.62	0.048	ADRA1B	P2RY1	DRD3

Table S11: Significantly enriched GO associated with genes overlapping 5kb windows around DMRs between sexes.

GO.ID	Term	Annotated	Significant	Expected	Weight	Barcelona	Monpellier	Warsaw
GO:1903706	regulation of hemopoiesis	172	8	2.28	0.050	KAT7, RARA, RUNX1, TJP2	CARD11, KAT7, PTN, RASSF2, SCAP, SOD1, VDR	
GO:1902930	regulation of alcohol biosynthetic process	20	3	0.27	0.039	SCAP		
GO:1902652	secondary alcohol metabolic process	53	4	0.7	0.013	LSS, SCAP	LSS, SCAP, SOD1	FGFR1
GO:0060173	limb development	123	4	1.63	0.028	RARA	PTN, TBX3	FGFR1, TBX3
GO:0060117	auditory receptor cell development	13	3	0.17	0.026	WHRN	SOD1	FGFR1
GO:0046887	positive regulation of hormone secretion	46	3	0.61	0.016			BLK, FGFR1, NMB
GO:0046849	bone remodeling	46	3	0.61	0.023	TPH1	PTN, RASSF2	
GO:0043087	regulation of GTPase activity	237	6	3.15	0.006	MLST8	PROM2, SOD1	LRRK2, PROM2, RGS14, STARD8
GO:0042981	regulation of apoptotic process	619	19	8.22	0.002	DIP2A, RARA, UBE2Z	BAG5, CARD11, CIDEC, DIP2A, GRK5, RASSF2, SOD1, TBX3	ANP32E, BLK, FGFR1, HERPUD1, KDM2B, LRRK2, PDIA3, PHLDA1, TBX3
GO:0035264	multicellular organism growth	93	4	1.23	0.033	RARA, SELENOM	ANKRD11, SOD1	ANKRD11, SELENOM
GO:0032436	positive regulation of proteasomal ubiquitination	46	4	0.61	0.018		CSNK1A1	CSNK1A1, HERPUD1, LRRK2, STUB1
GO:0032091	negative regulation of protein binding	50	3	0.66	0.028		ZFPM1	LRRK2, STUB1
GO:0031647	regulation of protein stability	152	5	2.02	0.048	LSS	BAG5, LSS, RASSF2	LRRK2, STUB1
GO:0009790	embryo development	625	15	8.3	0.049	MDF1, PCDH8, RARA, WHRN	ANKRD11, HOXB9, PTN, SOD1, TBX1, TBX3, ZFPM1	ANKRD11, ASCL2, DNMT3A, FGFR1, KDM2B, PCDH8, TBX3
GO:0009267	cellular response to starvation	85	4	1.13	0.044	PIK3C2B	XPR1	LRRK2, WDR24
GO:0007218	neuropeptide signaling pathway	46	3	0.61	0.023			GALR2, NMB, POMC
GO:0003151	outflow tract morphogenesis	46	3	0.61	0.046	RARA	TBX3, ZFPM1	TBX3
GO:0001666	response to hypoxia	91	4	1.21	0.023	NPEPPS, SCAP	NPEPPS, SCAP	ASCL2, LIMD1
GO:0000122	negative regulation of transcription by ...	454	12	6.03	0.017	RARA	CBFA2T2, SCAF4, TBX1, TBX3, VDR, ZFPM1	ASCL2, DNMT3A, FGFR1, KDM2B, MAF, TBX3

Table S12: Number of DMRs between habitats, found in females and males separately. See Supplementary Analysis 2. “Hypermethylated” and “Hypomethylated” refer to hyper- or hypomethylated DMR in the urban habitat compared to the forest habitat.

	Females		
	Barcelona	Montpellier	Warsaw
Hypermethylated	240	275	205
Hypomethylated	224	133	316
Total	464	408	521
	Males		
	Barcelona	Montpellier	Warsaw
Hypermethylated	107	347	315
Hypomethylated	317	229	218
Total	424	576	533

Table S13: Number of habitat linked DMRs shared between females and males. See Supplementary Analysis 2. “Same direction” refers to hyper- or hypo-methylated in both sexes.

	Number of DMRs found in females and males	
	same or opposite direction	same direction only
Barcelona	37	20
Montpellier	79	15
Warsaw	56	28

SUPPLEMENTARY FIGURES

Figure S1: Results of the Bayescan tests between forest and urban great tits, for each of the three urban-forest pairs. (A) Distribution of F_{ST} values per position in function of q-value. (B) Distribution of q-value in function of $\log(PO)$.

Figure S2: Distribution of genomic RDA loadings of each locus (above), and outliers (below). Vertical red lines represent the thresholds used to select outliers (please see methods for more details).

Figure S3: Semantic space representation for enriched GO terms from genes associated with genetic outlier (the outliers of the RDA analyses). Semantic space coordinates have been calculated based on similarity of GO term word composition using REVIGO. The color gradient indicates the number of genes involved in each GO term.

Figure S4: Sharing among each location of the significant DMRs found between A) forest and urban habitats, and B) females and males great tits. Hypermethylated DMRs (A) in urban, B) in males) are shown in red, hypomethylated in blue and for shared DMRs, grey represent cases where a DMRs was found in multiple locations but for which the direction of methylation was opposite.

Figure S5: Semantic space of enriched GO terms from genes associated with urbanization-linked DMRs. Semantic space coordinates have been calculated based on similarity of GO term word composition using REVIGO. The color gradient indicates the number of genes involved in each GO term and circle size indicates the number of city contributing to each category.

Figure S6: Circos plot of differentially methylated regions (DMRs) identified between females and males great tits in and near Barcelona, Montpellier and Warsaw (from inner to outer circles). Red points show hypermethylated regions in female great tits relatively to males, and blue points show hypomethylated regions. For graphical clarity, only a subset of genes are represented: genes associated with the 10% most extreme DMR. Names of the genes found within 5 kb of the represented DMRs are given.

Figure S7: Semantic space of enriched GO terms from genes associated with sex-linked DMRs. Semantic space coordinates are calculated based on similarity of GO term word composition using REVIGO. The color gradient indicates the number of genes involved in each GO term and circle size indicates the number of city contributing to each category.

Figure S8: Mean methylation difference per base along the CLOCK gene. Vertical dashed lines materialised start and end of the gene.

Figure S9: Mean methylation difference per base along the COL4A5 gene. Vertical dashed lines materialised start and end of the gene.

Figure S10: Mean methylation difference per base along the DRD4 gene. Vertical dashed lines materialised start and end of the gene.

Figure S11: Mean methylation difference per base along the ERG1 gene. Vertical dashed lines materialised start and end of the gene.

Figure S12: Mean methylation difference per base along the FOXP2 gene. Vertical dashed lines materialised start and end of the gene.

SUPPLEMENTARY ANALYSES

Differentially Methylated Regions (DMRs) between Females and Males

A total of 238 DMR were found between sexes: 58 in Barcelona, 81 in Montpellier and 99 in Warsaw (Figure S6). Warsaw presented significantly more hyper than hypomethylated DMR ($\chi^2=5.878$, $P=0.015$), but it was not the case for Barcelona ($\chi^2=0$, $P=1$) nor Montpellier ($\chi^2=1.25$, $P=0.264$). DMR were distributed on 29 chromosomes and 35 unplaced scaffolds.

In total, 181 DMR were located on genes or in 5kb upstream/downstream regions around genes. A gene ontology enrichment analysis performed on the pooled list of genes associated with DMR revealed significantly enriched modules associated with metabolism, development and hormonal function (see details Figure S4B and S6, and Table S11).

Differentially Methylated Regions between associated with urbanization in females and males

To explore the potential sex dependent effects of urbanization on methylation patterns, we searched for DMRs between forest and urban habitats in females and males separately. The numbers of significant DMRs per sex per location are reported in Table S12.

In females, 101 DMRs (7.9%) were shared between 2 or more locations and 45 (3.5%) showed the same direction of variation in all cities.

Similarly, in males 134 DMRs (9.7%) were shared by multiple locations, and 58 (4.18%) showed the same direction of variation.

Within locations, sexes showed shared urban-linked DMRs, see Table S12.