Title: Comprehensive analysis of male-free reproduction in *Monomorium triviale*

(Formicidae: Myrmicinae)

Authors: Naoto Idogawa^{1*}, Tomonori Sasaki^{2,3}, Kazuki Tsuji³ and Shigeto Dobata^{1,4*}

Affiliations:

¹ Laboratory of Insect Ecology, Graduate School of Agriculture, Kyoto University,
Kitashirakawa-Oiwakecho, Sakyo-ku, Kyoto 606-8502, Japan.

² Research and Development Department, Fumakilla Limited, 1-11-13, Umehara, Hatsukaichi,
Hiroshima 739-0494, Japan.

³ Department of Subtropical Agro-Environmental Sciences, Faculty of Agriculture, University
of the Ryukyus, 1, Senbaru, Nishihara, Okinawa 903-0213, Japan.

⁴ Department of General Systems Studies, Graduate School of Arts and Sciences, The
University of Tokyo, 3-8-1, Komaba, Meguro-ku, Tokyo 153-8902, Japan.

***Corresponding authors**

E-mail: idogawa.naoto.na@alumni.tsukuba.ac.jp

E-mail: dobata@g.ecc.u-tokyo.ac.jp

21 **Abstract**

22 We report comprehensive evidence for obligatory thelytokous parthenogenesis in an
 23 ant *Monomorium triviale*. This species is characterized by distinct queen–worker dimorphism
 24 with strict reproductive division of labor: queens produce both workers and new queens
 25 without mating, whereas workers are completely sterile. We collected 333 nests of this species
 26 from 14 localities and three laboratory-reared populations in Japan. All wild queens dissected
 27 had no sperm in their spermathecae. Laboratory observation confirmed that virgin queens
 28 produced workers without mating. Furthermore, microsatellite genotyping showed identical
 29 heterozygous genotypes between mothers and their respective daughters, suggesting an
 30 extremely low probability of sexual reproduction. Microbial analysis detected no bacterial
 31 genera that are known to induce thelytokous parthenogenesis in Hymenoptera. Finally, the
 32 lack of variation in partial sequences of mitochondrial DNA among individuals sampled from
 33 across Japan suggests recent rapid spread or selective sweep. *M. triviale* would be a promising
 34 model system of superorganism-like adaptation through comparative analysis with
 35 well-studied sexual congeners, including the pharaoh ant *M. pharaonis*.

36

37

38 Introduction

39 Hymenopteran insects are characterized by haplo-diploid sex determination systems,
40 in which females are derived from fertilized diploid eggs and males are derived from
41 unfertilized haploid eggs via arrhenotokous parthenogenesis [1]. However, the production of
42 diploid females from unfertilized eggs—known as thelytokous parthenogenesis—has been
43 reported sporadically across the Hymenoptera [2]. Thelytoky has provided an opportunity for
44 empirical testing of evolutionary hypotheses to account for the near-ubiquity of sex among
45 eukaryotes [3]. The last decade has seen an increasing number of known thelytokous
46 Hymenoptera, especially in eusocial species [4]. In studies taking advantage of this simplified
47 mode of reproduction, thelytokous social insects have been acknowledged as a model system
48 in behavioral ecology (e.g., [5–8]) and sociogenomics (e.g., [9]).

49 Previous studies of thelytokous parthenogenesis of ants have identified three distinct
50 categories [10, 11]: (type I) queens produce workers sexually but daughter queens via
51 thelytoky; (type II) the queen caste is usually lost, males are absent, and workers produce
52 workers via thelytoky; and (type III) queens produce both workers and queens via thelytoky,
53 males are usually absent, and workers are sterile.

54 In this paper, we report comprehensive evidence of the third type of thelytokous
55 parthenogenesis, namely obligatory thelytoky by queens of *M. triviale* [12]. The genus
56 *Monomorium* is one of the most species-rich genera among ants [13] and has a worldwide
57 distribution [14]. With the marked exception of tramp species such as the pharaoh ant *M.*
58 *pharaonis* [15], most *Monomorium* species remain to be investigated [16]. *M. triviale* is
59 reported from East Asia, encompassing Japan, South Korea and mainland China [14]. Neither
60 males nor inseminated females have been found to date; this has been referred to in several

61 studies [1, 4, 17–23]. Surprisingly, however, direct evidence of thelytoky is lacking. We
 62 examined the reproductive system of *M. triviale* through multiple approaches, namely
 63 dissection of spermathecae of field-collected queens, direct observation of virgin queen
 64 reproduction, and microsatellite genotyping of mothers and daughters. Furthermore, we
 65 analyzed for the presence of parthenogenesis-inducing microsymbionts and within-species
 66 phylogenetic relationships among populations in Japan.

67

68 **Materials & Methods**

69 **Colony sampling**

70 A total of 333 nests of *M. triviale* were collected from 14 local populations in Japan
 71 (Fig 1, Table 1). No specific permission for sampling was required. In the field, nests were
 72 found in small gaps in dead plant bodies such as rotten roots, dead twigs, hollow bamboo
 73 sticks, and acorns. Because of the small nest size (about 200 workers on average) it was easy
 74 to collect the whole nest. During the field investigation, males of *M. triviale* were never found.
 75 Colonies were transferred into artificial nests in the laboratory as soon as possible and were
 76 kept at 25 °C until the following experiments. In addition, ethanol-preserved samples
 77 originating from three other sites were examined (Fig 1, nos. 15 to 17).

78

79 **Fig 1. Localization of the 14 sampling sites (1 to 14) and the three sites of origin of**
 80 **laboratory-reared populations (15 to 17) in Japan.** Open circles indicate site locations. Site
 81 names, numbers of nests collected, and types of analyses are described in **Table 1**. Map data
 82 by Natural Earth (<http://www.naturalearthdata.com/>).

83

84 **Table 1. Sampling sites of *Monomorium triviale* populations and experiments performed**

	Locality	Latitude	Longitude	Collection dates	No. of nests	Experiments
1	Kyoto City, Kyoto Prefecture	35.060087	135.788488	19/04/2017 to 27/11/2017 (24 times)	176	SP, VR, SEQ
2	Tsuchiura City, Ibaraki Prefecture	36.077927	140.165473	20/06/2017, 09/10/2018	28	SP, VR, SEQ
3	Chofu City, Tokyo Metropolis	35.668182	139.549061	16/06/2017, 27/09/2018	28	SP, VR, SEQ
4	Tsukuba City, Ibaraki Prefecture	36.100415	140.101297	19/06/2017, 08/10/2018	26	SP, VR, SEQ
5	Otsu City, Shiga Prefecture	34.970262	135.956026	10/06/2017	26	SP, VR, SEQ
6	Matsudo City, Chiba Prefecture	35.774787	139.899362	18/06/2017, 28/09/2018	22	SP, VR, SEQ
7	Higashiosaka City, Osaka Prefecture	34.665748	135.671267	09/11/2017	5	SP, SEQ
8	Okazaki City, Aichi Prefecture	34.941529	137.175594	03/09/2017	1	SP, SEQ
9	Sanda City, Hyogo Prefecture	34.914575	135.165764	09/05/2017	1	VR, SEQ
10	Kaizu City, Gifu Prefecture	35.22	136.63	06/06/2018	1	SP
11	Tobishima Island, Sakata City, Yamagata Prefecture	39.19	139.55	20/09/2018	2	SP
12	Takamatsu City, Kagawa Prefecture	34.364349	134.098205	17/06/2017	16	SEQ
13	Matsuyama City, Ehime Prefecture	33.845539	132.765722	19/09/2018	2	SEQ
14	Takashima City, Shiga Prefecture	35.3677	135.9168	30/05/2017	1	SEQ
15	Kanonji City, Kagawa Prefecture	34.12	133.66	Reared in Laboratory	–	SEQ
16	Kagamihara City, Gifu Prefecture	35.40	136.85	Reared in Laboratory	–	SEQ
17	Matsue City, Shimane Prefecture	35.47	133.05	Reared in Laboratory	–	SEQ
Total					333	

85 The nests from Kaizu City (locality no. 10) and Tobishima Island (no. 11) were provided by Dr. K.

86 Ohkawara. Samples from Higashiosaka City (no. 7) and Takashima City (no. 14) were provided by Mr.

K. Sadahiro and Dr. T. Nozaki, respectively. Ethanol-preserved samples from Kanonji City, Kagamihara City, and Matsue City (nos. 15 to 17) were provided by Dr. F. Ito. Experiments on the samples from the 17 sites are abbreviated as follows: SP: dissection of wild queens' spermathecae; VR: observation of virgin queen reproduction; SEQ: phylogenetic analysis based on mtDNA sequencing.

91

92 Dissection of wild queens

93 To confirm the reproductive status of the queens from wild nests, we dissected 63
94 individuals from 38 nests of 10 *M. triviale* populations within 6 months after collection (Table
95 2). First the queen was immobilized by soaking in 70% ethanol for 3 min. The body was then
96 transferred to a 30-mm petri dish filled with distilled water, and the internal reproductive
97 organs were pulled out from the end of the abdomen by using precision forceps under a
98 binocular microscope (SZ40; OLYMPUS Optical, Tokyo, Japan). The mating status of the
99 queen was determined by the presence or absence of sperm in the spermatheca. As an
100 indicator of oviposition, the ovary's yellow body was checked. As a positive control [18], 11
101 queens of a congeneric species, *M. intrudens*, were dissected in the same manner.

102 **Table 2. Dissected queens from each sampling site**

Locality	No. of nests	No. of queens	Inseminati on		Yellow body	
			yes	no	yes	uncle ar
1 Kyoto City, Kyoto Prefecture	3	10	0	10	9	1
2 Tsuchiura City, Ibaraki Prefecture	8	8	0	8	7	1
3 Chofu City, Tokyo Metropolis	8	8	0	8	5	3
4 Tsukuba City, Ibaraki Prefecture	8	8	0	8	3	5
5 Otsu City, Shiga Prefecture	2	9	0	9	9	0
6 Matsudo City, Chiba Prefecture	4	8	0	8	8	0

7	Higashiosaka City, Osaka Prefecture	1	1	0	1	1	0
8	Okazaki City, Aichi Prefecture	1	8	0	8	8	0
10	Kaizu City, Gifu Prefecture	1	1	0	1	1	0
11	Tobishima Island, Sakata City, Yamagata Prefecture	2	2	0	2	1	1
	Total	38	63	0	63	52	11
–	<i>Monomorium intrudens</i> , Kyoto City, Kyoto Prefecture	4	11	10	1	11	0

103 The numbers of individuals possessing a yellow body and presumed to have experienced oviposition are
104 also shown. As a positive control, data for the sexual congener *M. intrudens* are shown in the bottom row.

105

106 Rearing experiment

107 Each wild nest was put into a plastic container (68 × 39 × 15 mm) with gypsum on
108 the bottom. The nests were kept in the laboratory at 25 °C, and water and food were
109 replenished every 3 days. The nests were fed mealworms, *Tenebrio molitor*, cut into
110 approximately 5-mm lengths. During July and August 2017, a total of 44 queen broods (larvae
111 or pupae) from 21 nests of seven populations were produced (Table 3). Each queen was
112 isolated with 10 nestmate workers in a plastic container (36 × 36 × 14 mm). These nests were
113 kept under the same conditions as the source nests. Reproduction by these virgin queens was
114 observed for 6 months. Finally, all remaining queens were dissected and confirmed to have no
115 sperm in their spermathecae (Table 4).

116 **Table 3. Thelytokous worker production in the rearing experiment**

	Locality	No. of nests	No. of queens isolated	No. of queens surviving for 6 months	No. of workers produced
1	Kyoto City, Kyoto Prefecture	6	10	6	17
2	Tsuchiura City, Ibaraki Prefecture	3	7	2	11

3	Chofu City, Tokyo Metropolis	3	5	5	26
4	Tsukuba City, Ibaraki Prefecture	3	5	5	28
5	Otsu City, Shiga Prefecture	3	8	8	52
6	Matsudo City, Chiba Prefecture	2	7	8	28
9	Sanda City, Hyogo Prefecture	1	2	2	15
Total		21	44	36	177

Table 4. Dissection of virgin queens from the rearing experiment

Locality		No. of queens dissected	Insemination		Yellow body	
			yes	no	yes	unclear
1	Kyoto City, Kyoto Prefecture	0 ^a	NA	NA	NA	NA
2	Tsuchiura City, Ibaraki Prefecture	2	0	2	2	0
3	Chofu City, Tokyo Metropolis	5	0	5	3	2
4	Tsukuba City, Ibaraki Prefecture	5	0	5	2	3
5	Otsu City, Shiga Prefecture	8	0	8	3	5
6	Matsudo City, Chiba Prefecture	7	0	7	2	5
9	Sanda City, Hyogo Prefecture	2	0	2	1	1
Total		29	0	29	13	16

^aWe did not examine queens from Kyoto due to their death before dissection.

Microsatellite analysis

To provide genetic evidence of thelytoky, 33 queens (from 17 nests of seven populations; all used in the rearing experiment) were examined (Table 5). They were genotyped at microsatellite locus *Mp-1* (f: GCCAATGGTTTAATCCCTCA; r: TCATACTGCGTGTGCCTTTC), originally developed from *M. pharaonis* [24]. Daughter workers produced from these virgin queens (174 individuals in total) were also genotyped and were compared with their mothers. The thorax of each individual was crushed and placed in a 0.2-mL microtube filled with 100 μ L of a DNA extraction reagent (PrepMan Ultra Reagent, Applied Biosystems, Foster City, CA, USA). The polymerase chain reaction (PCR) cocktail contained 1 μ L of template DNA, 0.3 μ L of 25 mM MgCl₂, 0.3 μ L of 10 mM dNTPs, 1.5 μ L

of 10× PCR Buffer, 0.1 μL of 5 U/μL Taq DNA Polymerase (QIAGEN, Valencia, CA, USA), 0.2 μL of U19 fluorescent dye, and 1.0 μL of each primer pair, to which distilled water was added to make a total volume of 15.2 μL. The PCR program consisted of an initial step of 94 °C for 180 s, followed by 35 cycles of 94 °C for 30 s, 57 °C for 60 s and 72 °C for 60 s, with a final step of 72 °C for 10 min. PCR products were mixed with Hi-Di formamide and GS-600 LIZ size standard and were analyzed by using a 3500 Series Genetic Analyzer and GeneMapper 5.0 software (Applied Biosystems).

Table 5. Microsatellite analysis data

Locality	No. of nests	No. of queens	No. of workers	Genotype	Prob. that all offspring were heterozygous
1 Kyoto City, Kyoto Prefecture	4	4	15	222/234	$(1/2)^{15} = 3.05 \times 10^{-5}$
2 Tsuchiura City, Ibaraki Prefecture	1	2	11	220/232	$(1/2)^{11} = 4.88 \times 10^{-4}$
3 Chofu City, Tokyo Metropolis	3	5	26	222/232	$(1/2)^{26} = 1.49 \times 10^{-8}$
4 Tsukuba City, Ibaraki Prefecture	3	5	28	220/232	$(1/2)^{28} = 3.73 \times 10^{-9}$
5 Otsu City, Shiga Prefecture	3	8	52	220/228	$(1/2)^{52} = 2.22 \times 10^{-16}$
6 Matsudo City, Chiba Prefecture	2	7	27	220/234	$(1/2)^{27} = 7.45 \times 10^{-9}$
9 Sanda City, Hyogo Prefecture	1	2	15	222/238	$(1/2)^{15} = 3.05 \times 10^{-5}$
Total	17	33	174	–	$(1/2)^{174} = 4.18 \times 10^{-53}$

Genotypes at the *Mp-1* locus are shown. Under the assumption of sexual reproduction, the probability that all daughter workers would be heterozygous was calculated.

Microbial analysis

To assess potential infection of *M. triviale* by thelytoky-inducing bacteria, we performed high-throughput amplicon sequencing by using whole bodies of adults. In August 2020, three *M. triviale* nests were collected in Takaragaike Park, Kyoto, Japan (lat 35.060087,

long 135.788488). All colonies were transported to our laboratory and moved to plastic containers (68 × 39 × 30 mm) with gypsum on the bottom. They were maintained at 25 °C, and water replenishment and mealworm-feeding were performed every 3 days. In October 2020, ten queens and 100 workers were picked up from each nest. The queens or workers from each nest were stored as a group in acetone at −30 °C. The pooled individuals represented one biological replicate (i.e., three replicates per caste). Whole bodies in each replicate were air-dried and pooled in a 1.5-mL plastic tube, and DNA was extracted with a QIAamp DNA Micro Kit (QIAGEN). We followed the manufacturer’s instructions and added extra steps of thermal cycling and lysozyme to the protocol to ensure the lysis of Gram⁺ positive bacterial cell walls: After being crushed in 180 µL ATL buffer, samples in plastic tubes first underwent two cycles of −80 °C for 30 min and 50 °C for 5 min; then, under room temperature, we added 2 µL of lysozyme from egg white (Nakalai Tesque, Kyoto, Japan; 20 µg/µL TE buffer) to each sample and incubated the samples at 37 °C for 30 min. 16S rRNA amplicon sequencing and the subsequent data analysis were performed according to in-house workflow by Bioengineering Lab. Co., Ltd. (Sagamihara, Kanagawa, Japan). The V4 region was amplified from 1 ng of template DNA by using ExTaq HS (Takara Bio, Otsu, Shiga, Japan) polymerase and the 515f–806r primer pair. Sequences were determined by using a MiSeq system with a MiSeq reagent kit v3 (Illumina, San Diego, CA, USA), which generated 2 × 300-bp paired-end reads. Demultiplexing (on the basis of a perfect match with the primer sequences used), adapter trimming, and quality filtering (phred score ≥ 20; primer sequences, 50 bp on both 3′-ends, noise and chimeric reads were removed) of the paired-end reads were performed by using a Fastx toolkit (ver. 0.0.14, http://hannonlab.cshl.edu/fastx_toolkit/) and dada2 plugin for

Qiime2 (ver. 2020.8) [25], resulting in representative sequences and operational taxonomic unit (OTU) tables. Assignment of appropriate taxa (confidence level > 0.7) was performed by using the default setting of the feature-classifier plugin for Qiime2 against the EzBioCloud 16S reference database (<https://www.ezbiocloud.net/>).

Phylogenetic analysis

To investigate intraspecific diversity and determine the phylogenetic position of *M. triviale*, phylogenetic analysis was performed. A 639-bp region of the *cytochrome oxidase I* (*COI*) sequence was determined by using PCR with a combination of primers, namely LCO1490 (GGTCAACAAATCATAAAGATATTGG) and HCO2198 (TAAACTTCAGGGTGACCAAAAAATCA) [26]. One worker randomly chosen from each of 15 local populations (Table 1) of *M. triviale* and a single worker of the sexual species, *M. intrudens*, collected in Kihoku-cho, Mie, Japan (lat 34.21, long 136.33), were sequenced. Protocols for DNA extraction and the PCR mix were the same as for the microsatellite analysis. The thermal cycle consisted of an initial denaturation at 94 °C for 1 min, 35 cycles of denaturation at 94 °C for 30 s, annealing at 50 °C for 30 s, extension at 72 °C for 60 s, and a final extension at 72 °C for 10 min. PCR products were ethanol-precipitated and sequenced in both directions by using a BigDye Terminator v3.1 cycle sequencing kit on a 3500 Series Genetic Analyzer (Applied Biosystems). A total of 16 sequences were submitted to the DNA Data Bank of Japan under accession numbers LC592050 to LC592065; see Fig 3). To identify the phylogenetic position of *M. triviale* after recent advances in systematics of the subfamily Myrmicinae [27], a congeneric species, *M. pharaonis* (GenBank accession number GU710434) and two former *Monomorium* species, *Sylophopsis sechellensis* (EF609858) and

Eromyrma latinodis (GU709833) were added to the analysis. The red imported fire ant, *Solenopsis invicta* (HQ928672), was used as an outgroup. The 16 sequences, together with those of the above four species, were aligned by using ClustalW [28] in MEGA 10.0 [29]. Maximum likelihood analyses were conducted as implemented in MEGA 10.0 by using a GTR+I+ Γ model with node support assessed with 1000 bootstrap replicates.

Results

Dissection of wild queens

All 63 dissected queens of *M. triviale* had no sperm in their spermathecae (Fig 2, Table 2). Yellow bodies, suggestive of queen oviposition experience, were identified in 52 individuals across all 10 sampling sites. In contrast, 10 out of 11 *M. intrudens* queens dissected as positive controls had sperm in their spermathecae. Dissection of *M. triviale* workers confirmed their complete sterility (Supporting Information S1 File), indicating that their complete sterility reported in previous studies [30, 31].

Fig 2. Reproductive organs of (a) *Monomorium triviale* and (b) *M. intrudens* queens. (a) *M. triviale*: Translucent (empty) spermatheca (spt), ovarioles with oocytes and obvious yellow bodies (yb) indicating that virgin queens were reproductively mature and laid eggs. **(b) *M. intrudens*:** Opaque (filled with sperm) spermatheca, well-developed ovarioles, and yellow bodies indicating that the queen is sexually reproducing. Yellow arrowheads indicate yellow bodies.

Rearing experiment

Eight queens died before producing daughter workers (Table 3). Thirty-six virgin queens from seven local populations survived, and each produced at least one worker. In total, 177 workers emerged during the experimental period; no males were produced. All dissected queens had empty spermathecae (Table 4). Despite all the queens producing workers, only 13 of 29 queens dissected had obvious yellow bodies.

Microsatellite analysis

Analysis of 33 queens and 174 workers showed that all individuals were heterozygous at *Mp-1* and that the genotypes of all the daughter workers were identical to those of their mothers. In addition, all individuals within the same local population had the same genotype. Comparison of genotypes between mothers and their daughters enables us to infer the occurrence of sexual reproduction. A potential male mate of a heterozygous mother (genotype AB) either shares or does not share the same allele (A or B) of *Mp-1* as the mother. We can rule out the latter possibility on the basis of the daughter genotypes. In the former case, the mother's sexually produced daughters should have two genotypes in the expected ratio of 1:1. One is the same as their mother's (AB) and the other is homozygous at one of the two mother alleles (AA when the father's genotype is A, or BB when the father's genotype is B). Under sexual reproduction, the observed bias toward the daughter's same heterozygous genotypes as their mother would be extremely rare. (The exact binomial probabilities are given in Table 5.) Therefore, we can reject the possibility of sexual reproduction among the individuals that we genotyped.

Microbial analysis

MiSeq sequencing and Qiime2-based analysis yielded 8364 to 46,817 reads per biological replicate and a total of 267 OTUs (summarized in S2 Table). Among them, 263 OTUs (covering 99.9% to 100% of the reads) were classified as bacterial, and 117 OTUs (92.4% to 99.5% of the reads, excepting 54.7% of the reads from one queen replicate from nest B, see below) were classified at least to the phylum level. No reads were assigned to bacterial genera that are known to induce thelytokous parthenogenesis in Hymenoptera, i.e., *Cardinium*, *Rickettsia*, and *Wolbachia* [1].

In two nests (A and C) out of the three replicates for both queens and workers, the most abundant reads were classified as from *Spiroplasma platyhelix* (88.4% and 67.4% of total reads from queen and worker replicates, respectively, of nest A; 91.5% and 91.1%, respectively, for nest C). The replicates obtained from nest B showed a different pattern of *S. platyhelix* abundance: 0% from the queen replicate and 13.2% from the worker replicate. The absence of the *S. platyhelix* sequence from the nest B queen replicate was likely associated with the relatively small number of total reads (8364 vs. >30,000, S2 Table).

Phylogenetic analysis

The 639-bp partial sequences of *COI* were completely identical among all individuals representing 15 populations of *M. triviale* (Fig 3). These individuals were placed in the monophyletic group with *M. intrudens* and *M. pharaonis*. Although the genus *Monomorium* has recently been revealed to be a polyphyletic group [27, 32], our result suggests that *M. triviale* belongs to the genus *Monomorium sensu stricto*.

Fig 3 Maximum likelihood tree of 15 populations of *M. triviale* and related species, based on *COI* sequences. The number at each branch of the phylogenetic tree represents the bootstrap percentage (1000 replicates). GenBank accession codes follow the taxon names. Scale bar: 0.1 substitutions per site.

Discussion

Our comprehensive investigation allows us to add *M. triviale* to the list of parthenogenetic ants. Production of daughters by unmated queens (indicative of type III thelytoky) was corroborated on the basis of multiple lines of evidence: (i) absence of males and inseminated queens in the field-collected nests; (ii) worker production by laboratory-reared virgin queens; (iii) unfilled spermathecae of queens that produced workers; and (iv) identical genotypes of mother queens and daughter workers. These features were common to all the locations we tested, suggesting that thelytoky is a dominant mode of reproduction across Japanese populations of *M. triviale*.

It should be noted here that rare occurrences of males have been reported in other parthenogenetic ants (type II), such as *Ooceraea biroi* [19] and *Pristomyrmex punctatus* [33]. In addition, queens of *M. triviale* retain undegenerated spermathecae [18], suggesting that they have low level of specialization to male-free reproduction. Moreover, geographic variation in sexual and asexual reproduction has been reported in some thelytokous ant species (types II and III), such as *Mycocepurus smithii*, *Myrmecina nipponica*, and *Platythyrea punctata* [17, 21, 34]. Whether and how often sexual reproduction occurs in *M.*

triviale, especially in populations outside Japan, would be an interesting topic for future study.

Our exploratory analysis of bacterial communities in *M. triviale* provides basic information for future studies of the host–symbiont relationship. No evidence was found for infection with thelytoky-inducing bacteria in this species, confirming previous reports of no cases of microorganism-induced thelytoky in ants [10, 24, 36, 37]. *Spiroplasma* has been detected in various ant species [38], including another thelytokous species, *Mycocepurus smithii* (type III thelytoky) [39]. In *Solenopsis*, a genus closely related to *Monomorium*, *Spiroplasma* has been detected as a dominant bacterial taxon in whole bodies of adult workers of *S. geminata* [40], thus drawing parallels with our finding. It is noteworthy that type I thelytoky has recently been found in polygynous colonies of this species [41]. *Spiroplasma* is well known as a sex ratio distorter in *Drosophila* [42], and its role in the host’s reproductive biology deserves further study in ants.

In our phylogenetic analysis, the extremely low levels of diversity observed in the *COI* sequences suggest the possibility of rapid spread or selective sweep in the recent past [43]. Having no mutation in 639 bp of *COI* sequence translates to an estimated divergence time of no more than ca. 45,000 years (assuming a divergence rate of 3.54% My⁻¹ [44]). Our results support the concept that this species is a member of the genus *Monomorium sensu stricto*. This phylogenetic status is advantageous in that the study designs established in *M. pharaonis*, a well-studied model system (e.g., [45–50]), will be applicable to future comparative analyses.

Thelytokous parthenogenesis is often considered as a trait overrepresented among introduced or invasive ant species [51]. Although some studies have categorized *M. triviale* as

invasive [1, 19], no exotic distribution has been reported so far [14] and there is insufficient information to support the possibility of an introduced origin of the Japanese population of *M. triviale*. The known distribution range of this species is far more limited than those of successful invasive congeners such as *M. pharaonis* and *Monomorium floricola* [15, 52]. A preference for disturbed and urban habitats is shared widely among invasive ants [53]. Ito et al. [54] listed *M. triviale* along with *Strumigenys membranifera*, *P. punctatus*, and *O. biroi* as thelytokous ants found in “open, disturbed areas.” Among these species, the life history of *M. triviale* is most poorly known. In our study, *M. triviale* nests were found even in a thicket dominated by deciduous broad-leaved trees, which is not typical of “open, disturbed areas.” Nevertheless, the above information does not rule out the possibility that *M. triviale* will become invasive in the future. Additional studies of this species’ social structure, such as queen numbers and the mode of colony foundation, will help to evaluate the potential invasion risk of this species as an ecological consequence of its life history.

Acknowledgments

We are grateful to Kenji Matsuura who allowed us to use his laboratory. Fuminori Ito and Hroyuki Shimoji provided valuable advice on *M. triviale* biology and microbial analysis, respectively. We thank Kazuya Takeda, Kunio Sadahiro, Riou Mizuno and Yu Hisasue for field assistance. We also thank Hiroyuki Shimoji, Kiyomi Nakagawa, Kyosuke Ohkawara and Tomonari Nozaki for providing us with *M. triviale* nests.

References

- 325 1. Rabeling C, Kronauer DJC. Thelytokous Parthenogenesis in eusocial hymenoptera.
326 Annu Rev Entomol. 2013;58: 273–292. doi:10.1146/annurev-ento-120811-153710
- 327 2. Engelstädter J. Constraints on the evolution of asexual reproduction. BioEssays.
328 2008;30: 1138–1150. doi:10.1002/bies.20833
- 329 3. Neiman M, Schwander T. Using parthenogenetic lineages to identify advantages of
330 sex. Evol Biol. 2011;38: 115–123. doi:10.1007/s11692-011-9113-z
- 331 4. Goudie F, Oldroyd BP. The distribution of thelytoky, arrhenotoky and androgenesis
332 among castes in the eusocial Hymenoptera. Insectes Soc. 2018;65: 5–16.
333 doi:10.1007/s00040-017-0597-0
- 334 5. Hartmann A, Wantia J, Torres JA, Heinze J. Worker policing without genetic
335 conflicts in a clonal ant. Proc Natl Acad Sci. 2003;100: 12836–12840.
336 doi:10.1073/pnas.2132993100
- 337 6. Ravary F, Lecoutey E, Kaminski G, Châline N, Jaisson P. Individual Experience
338 alone can generate lasting division of labor in ants. Curr Biol. 2007;17: 1308–1312.
339 doi:10.1016/j.cub.2007.06.047
- 340 7. Dobata S, Tsuji K. Public goods dilemma in asexual ant societies. Proc Natl Acad
341 Sci. 2013;110: 16056–16060. doi:10.1073/pnas.1309010110
- 342 8. Bernadou A, Busch J, Heinze J. Diversity in identity: behavioral flexibility,
343 dominance, and age polyethism in a clonal ant. Behav Ecol Sociobiol. 2015;69:
344 1365–1375. doi:10.1007/s00265-015-1950-9
- 345 9. Oxley PR, Ji L, Fetter-Pruneda I, McKenzie SK, Li C, Hu H, et al. The genome of
346 the Clonal raider ant *Cerapachys Biroi*. Curr Biol. 2014;24: 451–458.
347 doi:10.1016/j.cub.2014.01.018

- 348 10. Himler AG, Caldera EJ, Baer BC, Fernandez-Marin H, Mueller UG. No sex in
349 fungus-farming ants or their crops. *Proc R Soc B Biol Sci.* 2009;276: 2611–2616.
350 doi:10.1098/rspb.2009.0313
- 351 11. Rabeling C, Lino-Neto J, Cappellari SC, Dos-Santos IA, Mueller UG, Bacci M.
352 Thelytokous parthenogenesis in the fungus-gardening ant *Mycocepurus smithii*
353 (Hymenoptera: Formicidae). *PLoS One.* 2009;4. doi: 10.1371/journal.pone.0006781
- 354 12. Wheeler, William Morton HS. The ants of Japan. *Bulletin of the AMNH.* Bull
355 AMNH. 1906; v. 22, art.
- 356 13. Bolton B. An online catalog of the ants of the world. 2020 (accessed 17 December,
357 2020). Available from: <http://antcat.org>.
- 358 14. Guénard B, Weiser MD, Gómez K, Narula N, Economo EP. The Global Ant
359 Biodiversity Informatics (GABI) database: Synthesizing data on the geographic
360 distribution of ant species (Hymenoptera: Formicidae). *Myrmecological News.*
361 2017;24: 83–89.
- 362 15. Wetterer JK. Worldwide spread of the pharaoh ant, *Monomorium pharaonis*
363 (Hymenoptera: Formicidae). *Myrmecological News.* 2010;13: 115–129.
- 364 16. Pontieri L, Linksvayer TA. *Monomorium*. *Encyclopedia of Social Insects.* Cham:
365 Springer International Publishing; 2019. pp. 1–6.
366 doi:10.1007/978-3-319-90306-4_171-1
- 367 17. Rabeling C, Gonzales O, Schultz TR, Bacci M, Garciad MVB, Verhaaghe M, et al.
368 Cryptic sexual populations account for genetic diversity and ecological success in a
369 widely distributed, asexual fungus-growing ant. *Proc Natl Acad Sci U S A.*
370 2011;108: 12366–12371. doi:10.1073/pnas.1105467108

18. Gotoh A, Billen J, Tsuji K, Sasaki T, Ito F. Histological study of the spermatheca in three thelytokous parthenogenetic ant species, *Pristomyrmex punctatus*, *Pyramica membranifera* and *Monomorium triviale* (Hymenoptera: Formicidae). Acta Zool. 2012;93: 200–207. doi:10.1111/j.1463-6395.2010.00498.x
19. Kronauer DJC, Pierce NE, Keller L. Asexual reproduction in introduced and native populations of the ant *Cerapachys biroi*. Mol Ecol. 2012;21: 5221–5235. doi:10.1111/mec.12041
20. Masuko K. Thelytokous Parthenogenesis in the Ant *Strumigenys hexamera* (Hymenoptera: Formicidae). Ann Entomol Soc Am. 2013;106: 479–484. doi:10.1603/an12144
21. Masuko K. Thelytokous Parthenogenesis in the Ant *Myrmecina nipponica* (Hymenoptera: Formicidae). Zoolog Sci. 2014;31: 582–586. doi:10.2108/zs140050
22. van der Kooi CJ, Schwander T. On the fate of sexual traits under asexuality. Biol Rev. 2014;89: 805–819. doi:10.1111/brv.12078
23. Lee CC, Hsu SF, Yang CCS, Lin CC. Thelytokous parthenogenesis in the exotic dacetine ant *Strumigenys rogeri* (Hymenoptera: Formicidae) in Taiwan. Entomol Sci. 2018;21: 28–33. doi:10.1111/ens.12277
24. The invasion biology and sociogenetics of pharaoh ants. PhD Thesis, University of Copenhagen, København, Denmark.
25. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol. 2019;37: 852–857. doi:10.1038/s41587-019-0209-9
26. Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates.

- 395 Mol Mar Biol Biotechnol. 1994;3: 294–9. Available:
396 <http://www.ncbi.nlm.nih.gov/pubmed/7881515>
- 397 27. Ward PS, Brady SG, Fisher BL, Schultz TR. The evolution of myrmicine ants:
398 Phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae).
399 Syst Entomol. 2015;40: 61–81. doi:10.1111/syen.12090
- 400 28. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: Improving the sensitivity of
401 progressive multiple sequence alignment through sequence weighting,
402 position-specific gap penalties and weight matrix choice. Nucleic Acids Res.
403 1994;22: 4673–4680. doi:10.1093/nar/22.22.4673
- 404 29. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary
405 genetics analysis across computing platforms. Mol Biol Evol. 2018;35: 1547–1549.
406 doi:10.1093/molbev/msy096
- 407 30. Hölldobler B, Wilson EO. The Ants. Cambridge, MA: Harvard University Press;
408 1990.
- 409 31. Ito F, Yamane S. Reproduction by ergatoid queens in the myrmicine ant
410 *Monomorium brocha* (Bolton) (hymenoptera: Formicidae) in West Java, Indonesia,
411 with a description of the male. Asian Myrmecology. 2014;6: 105–113.
- 412 32. Sparks KS, Andersen AN, Austin AD. A multi-gene phylogeny of Australian
413 *Monomorium* Mayr (Hymenoptera: Formicidae) results in reinterpretation of the
414 genus and resurrection of *Chelaner* Emery. Invertebr Syst. 2019;33: 225–236.
415 doi:10.1071/IS16080
- 416 33. Yamada A, Eguchi K. Description of the male genitalia of *Pristomyrmex punctatus*
417 (Smith, 1860) (Hymenoptera, Formicidae, Myrmicinae). Asian Myrmecology.
418 2016;8: 87–94. doi:10.20362/am.008010

- 419 34. Kellner K, Seal JN, Heinze J. Sex at the margins: Parthenogenesis vs. facultative and
420 obligate sex in a Neotropical ant. J Evol Biol. 2013;26: 108–117.
421 doi:10.1111/jeb.12025
- 422 35. Pearcy M, Hardy O, Aron S. Thelytokous parthenogenesis and its consequences on
423 inbreeding in an ant. Heredity (Edinb). 2006;96: 377–382.
424 doi:10.1038/sj.hdy.6800813
- 425 36. Grasso DA, Wenseleers T, Mori A, Le Moli F, Billen J. Thelytokous worker
426 reproduction and lack of *Wolbachia* infection in the harvesting ant *Messor capitatus*.
427 Ethol Ecol Evol. 2000;12: 309–314. doi:10.1080/08927014.2000.9522803
- 428 37. Wenseleers T, Billen J. No evidence for *Wolbachia*-induced parthenogenesis in the
429 social Hymenoptera. J Evol Biol. 2000;13: 277–280.
430 doi:10.1046/j.1420-9101.2000.00168.x
- 431 38. Kautz S, Rubin BER, Moreau CS. Bacterial infections across the ants: Frequency and
432 prevalence of *Wolbachia*, *Spiroplasma*, and *Asaia*. Psyche (London). 2013.
433 doi:10.1155/2013/936341
- 434 39. Sen R, Ishak HD, Estrada D, Dowd SE, Hong E, Mueller UG. Generalized antifungal
435 activity and 454-screening of *Pseudonocardia* and *Amycolatopsis* bacteria in nests of
436 fungus-growing ants. Proc Natl Acad Sci U S A. 2009;106: 17805–17810.
437 doi:10.1073/pnas.0904827106
- 438 40. Ishak HD, Plowes R, Sen R, Kellner K, Meyer E, Estrada DA, et al. Bacterial
439 Diversity in *Solenopsis invicta* and *Solenopsis geminata* Ant colonies characterized
440 by 16S amplicon 454 pyrosequencing. Microb Ecol. 2011;61: 821–831.
441 doi:10.1007/s00248-010-9793-4

- 442 41. Lacy KD, Shoemaker DW, Ross KG. Joint Evolution of Asexuality and Queen
443 Number in an Ant. *Curr Biol.* 2019;29: 1394-1400.e4. doi:10.1016/j.cub.2019.03.018
- 444 42. Paredes JC, Herren JK, Schüpfer F, Marin R, Claverol S, Kuo CH, et al. Genome
445 sequence of the *Drosophila melanogaster* male-killing *Spiroplasma* strain MSRO
446 endosymbiont. *MBio.* 2015;6. doi:10.1128/mBio.02437-14
- 447 43. Avise JC, others. *Phylogeography: the history and formation of species.* Harvard
448 university press; 2000.
- 449 44. Papadopoulou A, Anastasiou I, Vogler AP. Revisiting the insect mitochondrial
450 molecular clock: the mid-Aegean trench calibration. *Mol Biol Evol.* 2010;27:
451 1659–1672. doi:10.1093/molbev/msq051
- 452 45. Beekman M, Sumpter DJT, Ratnieks FLW. Phase transition between disordered and
453 ordered foraging in pharaoh’s ants. *Proc Natl Acad Sci U S A.* 2001;98: 9703–9706.
454 doi:10.1073/pnas.161285298
- 455 46. Jefford CW, Tang Q, Zaslona A, Tang Q, Zaslona A. Short, Enantioselective Syntheses
456 of (-)-Indolizidine 167B and (+)-Monomorphine. *J Am Chem Soc.* 1991;113:
457 3513–3518. doi:10.1021/ja00009a043
- 458 47. Jackson DE, Holcombe M, Ratnieks FLW. Trail geometry gives polarity to ant
459 foraging networks. *Nature.* 2004;432: 907–909. doi:10.1038/nature03105
- 460 48. Robinson EJM, Jackson DE, Holcombe M, Ratnieks FLW. Insect communication:
461 “No entry” signal in ant foraging. *Nature.* 2005;438: 442. doi:10.1038/438442a
- 462 49. Warner MR, Mikheyev AS, Linksvayer TA. Genomic signature of kin selection in an
463 ant with obligately sterile workers. *Mol Biol Evol.* 2017;34: 1780–1787.
464 doi:10.1093/molbev/msx123

50. Oliveira RC, Warson J, Sillam-Dussès D, Herrera-Malaver B, Verstrepen K, Millar JG, et al. Identification of a queen pheromone mediating the rearing of adult sexuals in the pharaoh ant *Monomorium pharaonis*. Biol Lett. 2020;16: 20200348. doi:10.1098/rsbl.2020.0348
51. Tribble W, McKenzie SK, Kronauer DJC. Globally invasive populations of the clonal raider ant are derived from Bangladesh. Biol Lett. 2020;16: 20200105. doi:10.1098/rsbl.2020.0105
52. Wetterer JK. Worldwide spread of the flower ant, *Monomorium floricola* (hymenoptera: Formicidae). Myrmecological News. 2009;13: 19–27.
53. Rabitsch W. The hitchhiker’s guide to alien ant invasions. BioControl. 2011;56: 551–572. doi:10.1007/s10526-011-9370-x
54. Ito F, Touyama Y, Gotoh A, Kitahiro S, Billen J. Thelytokous parthenogenesis by queens in the dacetine ant *Pyramica membranifera* (Hymenoptera: Formicidae). Naturwissenschaften. 2010;97: 725–728. doi:10.1007/s00114-010-0688-5

Supporting information

S1 File. Dissection of workers of *M. triviale*.

S2 Table. Microbial taxa and their relative abundances (% of total reads) per each *M. triviale* sample.

Funding

486 This work was supported by a Japan Society for the Promotion of Science (JSPS) Research
487 Fellowship for Young Scientists to NI (19J22242) and a grant from the Secom Science and
488 Technology Foundation to SD.

a

b