Uvariopsis ebo (Annonaceae) a new, Critically Endangered tree species from the Ebo Forest, Cameroon and a key to the Cameroonian species of *Uvariopsis*

George Gosline¹, Martin Cheek¹, Jean Michel Onana^{2,3}, Eric Ngansop², Xander van der Burgt¹, Leo-Paul Dagallier³

¹ Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AE, U.K.

² IRAD-National Herbarium of Cameroon, P.O. Box. 1601, Yaoundé, Cameroon

³ Faculty of Science, Department of Plant Biology, University of Yaoundé I, Yaoundé, Cameroon

⁴ IRD, DIADE, University of Montpellier, Montpellier, France.

Summary. A new species to science of evergreen forest tree, *Uvariopsis ebo* (Annonaceae) is described, illustrated, mapped, and compared morphologically with the other cauliflorous species of the genus. Restricted so far to a single site in evergreen lowland forest in the Ebo Forest, Yabassi, Littoral Region, Cameroon, this species is Critically Endangered using the IUCN 2012 standard, because the forest habitat of this species remains unprotected, and there exists threats of logging and conversion to plantations. This species adds to the growing list of threatened species resulting from anthropogenic pressure on Cameroon forests.

Observations on the unusual corolla structure of the new species are made. A revised key to the 14 Cameroonian species of *Uvariopsis* is presented. Notes are given on other narrowly endemic and threatened species in the Ebo forest area, a highly threatened centre of diversity in Littoral Region, globally important for conservation.

Key Words: conservation, endemic species, threatened species.

Introduction

The new species reported in this paper was discovered as a result of a long-running survey of plants in Cameroon to support improved conservation management. The survey is led by botanists from the Royal Botanic Gardens, Kew and IRAD (Institute of Agricultural Research for Development)-National Herbarium of Cameroon, Yaoundé. The study has focussed on the Cross-Sanaga interval (Cheek *et al.* 2001, 2006) which contains the area with the highest plant species diversity per degree square in tropical Africa (Barthlott *et al.* 1996). The herbarium specimens collected in these surveys formed the foundations for a series of Conservation Checklists (see below). So far, over 100 new species and several new genera have been discovered and published as a result of these surveys, new protected areas have been recognised and the results of analysis are feeding into the Cameroon Important Plant Area programme (<https://www.kew.org/science/our-science/projects/tropical-important-plant-areas-cameroon>), based on the categories and criteria of Darbyshire *et al.* (2017).

In connection with preparation of a Conservation Checklist of the plants of the Ebo Forest, Littoral Region, Lorna MacKinnon, a volunteer botanical assistant, made a plant collecting expedition to the Ebo Forest in 2008. Among the collected plant specimens and photographs that she made, we identified an *Uvariopsis* Engl. & Diels (MacKinnon 51, Fig. 1 & 2) which resembled no other species known in the genus. In this paper we test the hypothesis that it is a new species to science, and name the new species as *Uvariopsis ebo*.

Fig. 1. *Uvariopsis ebo*. Cauliflorous inflorescences. *MacKinnon* 51(K). Photo Lorna MacKinnon.

Uvariopsis (Annonaceae) is a highly distinctive and easily recognised genus, since most of its species have unisexual flowers, a calyx with two basally connate sepals, and the petals in a single whorl of four (very rarely three, see below). Annonaceae are otherwise characterised by bisexual flowers with trimerous perianths. Most species of the genus are cauliflorous small trees, the flowers being produced from the trunk, although some species are ramiflorous or bear axillary flowers (Kenfack *et al.* 2003, Couvreur *et al.* 2021).

Nineteen species are currently accepted in *Uvariopsis*. Five species have been published in the 21st Century: *Uvariopsis korupensis* Gereau & Kenfack (Gereau & Kenfack 2000), *U. submontana* Kenfack, Gosline & Gereau (Kenfack *et al.* 2003) and *U. etugiana* Dagallier & Couvreur (Couvreur *et al.* 2021) all from Cameroon, *U. citrata* Couvreur & Niang. (Couvreur & Niangadouma 2016) from Gabon and *U. lovettiana* Couvreur & Q. Luke (Couvreur & Luke 2010) from Tanzania. In addition a sixth species, *Uvariopsis tripetala* (Baker f.) G.E.Schatz, was transferred to the genus from the monotypic *Dennettia* Baker f. (Kenfack *et al.* 2003). The genus is centred in Cameroon, where 13 of the 19 species occur, followed in species diversity by Gabon, with six species. The most widespread species of the genus is *Uvariopsis congensis* Robyns & Ghesq. Which occurs from Cameroon to S.Sudan, Zambia and Kenya. Several species are rare, being known from only one or two specimens and have restricted ranges, these include *U. etugiana* (Cameroon endemic) and *U. citrata* (Cameroon & Gabon), both known from two specimens, and *U. sessiliflora* (Mildbr. & Diels) Robyns & Ghesq. endemic to Cameroon and known from a single specimen.

Fig.2. *Uvariopsis ebo*. Trunk apex with cauliflorous flowers and canopy. *MacKinnon* 51(K). Photo Lorna MacKinnon.

The genus is distributed throughout continental tropical African evergreen forests, from Guinea in the West to Tanzania in the east, and as far south as northern Zambia. The species usually occur at low altitude, exceptions being *U. submontana* and *U. lovettiana* which occur in submontane or cloud forest in Cameroon and Tanzania respectively (Kenfack *et al.* 2003; Couvreur & Luke 2010). Species are usually small trees in high quality, undisturbed forest and appear not to be pioneers. They usually occur at low frequency. For example, in the Mefou Proposed National Park of Central Region Cameroon, only a single mature individual with one juvenile of one species of the genus, *U. solheidii* (De Wild.) Robyns & Ghesq., was found in the course of many weeks of botanical surveys by numerous botanists collecting thousands of specimens (Cheek *et al.* 2011: 123). However, in some rare ecological circumstances, some species can become locally dominant e.g., *U. tripetala* in the understorey of maritime lowland evergreen inselberg forest in Guinea (Couch *et al.* 2019: 41), and also *U. congensis* locally subdominant in forests in western Uganda where it flowers synchronously and is dispersed by primates (Dominy & Duncan 2005; Gosline pers. obs. 2016).

Materials and Methods

Herbarium citations follow Index Herbariorum (Thiers *et al.* 2020). Specimens were viewed at EA, K, P, WAG, and YA. The National Herbarium of Cameroon, YA, was searched for additional material of the new species, but without success. Images for specimens at WAG

were studied at <https://bioportal.naturalis.nl/?language=en> and those from P at https://science.mnhn.fr/institution/mnhn/collection/p/item/search/form?lang=en_US. We also searched JStor Global Plants (<https://plants.jstor.org/> accessed March 2021) for additional material, and finally the Global Biodiversity Facility (GBIF, www.gbif.org accessed March 2021). Binomial authorities follow the International Plant Names Index (IPNI, 2020). The conservation assessment was made using the categories and criteria of IUCN (2012). Herbarium material was examined with a Leica Wild M8 dissecting binocular microscope fitted with an eyepiece graticule measuring in units of 0.025 mm at maximum magnification. The drawing was made with the same equipment using Leica 308700 camera lucida attachment. The botanical terms follow Beentje & Cheek (2003), and format of the description follow the conventions of Kenfack *et al.* (2003) and Couvreur & Luke (2010). The map was made using SimpleMappr (<https://www.simplemappr.net>).

Results

Because it has leaves exceeding 15 cm long, cauliflorous flowers with pedicels exceeding 10 mm long, petals exceeding 7 mm long, MacKinnon 51, described below as *Uvariopsis ebo*, keys out in the key to the species of the genus in Couvreur *et al.* (2021) to couplet 12, leading to *U. dioica* (Diels) Robyns & Ghesq. (flower buds globose) and *U. solheidii* (flowers buds conical). Of these two choices, our material is closest to the second of these leads, the buds being narrowly ovoid. The two species are similar, but can be separated using the differential characters in Table 1, and the new species can be separated from all other cauliflorous species of *Uvariopsis* using the key (both presented below).

Character	<i>Uvariopsis solheidii</i>	<i>Uvariopsis ebo</i>
Indumentum of stem, petiole and abaxial midrib	Tomentose	Glabrous
Leaf-blade dimensions	16.6 – 29 x 5 – 9.5 cm	17.7 – 20.3 (– 23) x (6.4 –) 7 – 7.9 cm
Number of secondary nerves on each side of midrib	8 – 13	5 – 8
Number of flowers per inflorescence	2 – 3	4 – 7
Petal dimensions	7 – 10 x 2.5 – 5 mm	(14 –) 16 x (5.5 –) 9 mm
Petal colour	Wine brown	Yellow-green

Table 1. Differential characters separating *Uvariopsis solheidii* and *U. ebo*. Data for *Uvariopsis solheidii* taken mainly from Couvreur *et al.* (2021) and Cheek *et al.* (2011).

Key to the species of *Uvariopsis* in Cameroon

1. - Crushed leaves emitting a strong citrus scent *U. citrata*
- Crushed leaves without citrus scent 2
2. - Leaf blades generally < 15 cm long; pedicel < 7 mm long and petals < 7 mm long 3
- Leaf blades generally > 15 cm long, pedicel > 10 mm long and petals > 7 mm long .. 6
3. - Flowers cauliflorous *U. sessiliflora*
- Flowers ramiflorous 4
4. - Flowers bisexual, petals 3 (but occasionally 4) *U. tripetala*
- Flowers unisexual, petals 4 5

5. - Young branches and petioles densely to sparsely pubescent, petals free .. *U. congensis*
- Young branches glabrous or very sparsely pubescent, petals basally fused ... *U. zenkeri*
6. - Petals basally fused 7
- Petals free 10
7. - Flowering pedicels generally > 10 cm long; petals 3 *U. congolana*
- Flowering pedicels < 10 cm long; petals 4 8
8. - Flower buds globose, monocarps verrucose *U. pedunculosa*
- Flower buds conical to pyramidal, monocarps smooth 9
9. - Sepals 5 – 10 mm long, flowers completely covering base of trunk, generally occurring above 800 m a.s.l. *U. submontana*
- Sepals 2 – 4 mm long, flowers partially covering base of trunk, generally occurring below 800 m a.s.l. *U. korupensis*
10. - Flowering pedicels < 10 mm long 11
- Flowering pedicels > 10 mm long 12
11. - Petals linear, 25–45 mm long, more than 6 times longer than wide *U. bakeriana*
- Petals elliptic to ovate, 10–14 mm long, less than 6 times longer than wide *U. etugiana*
12. - Flower buds globose *U. dioica*
- Flower buds conical 13
13. - Stems and petioles tomentose, lateral nerves 8-13; flowers wine brown *U. solheidii*
- Stems and petioles glabrous; lateral nerves 5-8 on each side of the midrib; flowers green-yellow *U. ebo*

***Uvariopsis ebo* Cheek & Gosline sp. nov.** Type: Cameroon, Littoral Region, Ebo Forest, 4° 20' 44" N, 10° 24' 33" E, 849 m alt. Dicam Trail 2000 m from Bekob camp, male fl. 25 March 2008, *MacKinnon* 51 (holotype K001381842; isotypes YA, MO).

Cauliflorous, probably monoecious understorey tree 3 – 4 m tall. Trunk terete, lacking flutes or prop roots, 1.8 – 2.5 cm diameter at 1.5 m above the ground, bark smooth, dark-brown (Fig. 1), the crown sparsely branched (Fig. 2). Leafy stems with 3 – 4 leaves per season's growth, terete, internodes (1.2 –) 1.5 – 2.8 (– 4.3) cm long, 0.15 – 0.2 cm diam., pale yellow-green, later orangish brown, glabrous. Axillary buds dome-shaped 0.5 – 0.75 x 1 mm, bud-scales numerous, linear, spreading, densely hairy, hairs simple, appressed hairy c. 0.5 mm long, colourless or red brown. *Leaves* distichous, held in the horizontal plane, lacking scent when crushed (collection data, *MacKinnon* 51), blades oblanceolate 17.7 – 20.2 (– 23) x (6.4 –) 7 – 7.9 cm, acumen narrowly triangular (0.5 –) 1 – 1.3 cm, base broadly acute with convex edges, minutely cordate, blade mounted above petiole, margins undulate-sinuuous (live & dried), midrib impressed on adaxial surface, inconspicuous, below a groove; on abaxial surface subcylindric, 1 – 1.2 mm diam., conspicuous; secondary veins 5 – 8 (– 9) on each side of the midrib, brochidodromous, arising at c. 50° from the midrib, initially straight, then curving in the outer third, uniting with the secondary nerve above to form a looping inframarginal nerve, attaining 3 – 4 mm from the margin; intersecondary nerves sometimes present, tertiary nerves raised, conspicuous, forming a reticulum with cells 4 – 5 mm long, quaternary nerves inconspicuous; glabrous (except in bud when densely orange-brown hairy, hairs 0.1 mm long). Petiole stout, shallowly canaliculate, 4 (– 5) mm long, 1.9 – 2.1 mm diam., narrowing at base and apex, the adaxial groove shallow, c. 0.5 mm wide, glabrous.

Female inflorescences unknown. *Male inflorescences* cauliflorous, scattered along the trunk from ground level to the top of the trunk 2.5 – 3 m above the ground, each 3 – 7-flowered (Fig. 1 & 2). Peduncles patent, c. 2 x 2 mm, pale brown, glabrous, bearing sub-umbellate, radiating, 1-flowered partial-peduncles. Partial-peduncles 0.5 – 2 x 0.9 – 1.2 mm, terminating in 1 – 2 bracts subtending a pedicel. Bracts oblong-elliptic, 1.5 x 0.5 – 0.6 mm, apex acute, outer surface about 50% covered in appressed white hairs c. 0.15 – 0.2 mm long, inner surface glabrous. Pedicels 1.8 – 2.5 cm long, 0.1 cm diam., articulated with the partial-peduncle, with (0 –) 1 (– 2) scattered, bracteoles in the proximal few mm. Bracteoles similar to the bracts, ovate-oblong, shortly sheathing, (1 –) 1.25 x 1 mm, outer surface with sparse scattered simple appressed translucent hairs 0.05 – 0.2 mm long. buds narrowly ovoid, 16 x 11 mm. *Sepals* 2, opposite, drying pale brown, reflexed, *Male flowers* semi-orbicular, 1 – 1.5 mm long, 2.1 – 2.5 mm wide, glabrous. *Petals* 4, uniseriate, free, yellow-green when live (Fig. 1), drying black, lanceolate-oblong, (14 –) 16 x (5.5 –) 9 mm, apex rounded, base rounded, outer surface sparsely and inconspicuously hairy, hairs 7 – 9 per mm² (5% of surface covered), simple, translucent, appressed, 0.1 mm long, apices rounded. Inner surface of petals with a shallow elliptic-oblong excavation c. 8 x 5 mm, the margin of the excavation raised, the apex with a ridge extending along the midline to the petal apex, glabrous apart from a few scattered erect, minute white hairs 0.05 mm long at the excavation apex. Staminal dome 3.5 – 4 mm long, 3.5 – 4 mm diam., consisting of stamens and a receptacular torus. Stamens shortly cylindrical-angular, 0.5 x 0.1 (– 0.2) mm, connective surrounded by four longitudinal anther cells, each exceeding the connective, in plan view the stamen is 4-lobed. Apical connective appendage absent. Female flowers, fruit and seed unknown. Fig. 1 & 2.

RECOGNITION. Similar to *Uvariopsis solheidii* (De Wild.) Robyns & Ghesq., differing in the stem, petioles and abaxial midrib glabrous (versus tomentose); number of secondary nerves on each side of the midrib 5 – 8 (versus 8 – 13); petals yellow-green, (14 –) 16 x (5.5 –) 9 mm (versus wine brown, 7 – 10 x 2.5 – 5 mm). Additional differential characters are given in Table 1, above.

DISTRIBUTION. Cameroon (Map1), endemic to the Ebo Forest of the Littoral Region on present evidence.

SPECIMENS EXAMINED. Cameroon, Littoral Region, Yabassi, Ebo Forest, 4° 20' 44" N, 10° 24' 33"E, 849 m alt. Dicam Trail 2000 m from Bekob camp, male fl. 25 March 2008, MacKinnon 51 (holotype K001381842; isotypes YA, MO).

HABITAT. *Uvariopsis ebo* is so far only known from lower submontane forest (849 m elev.). below the elevation for the upper montane forest indicator species *Podocarpus latifolius* (Thunb.) R.Br. ex Mirb. The geology is ancient, highly weathered basement complex, with some ferrallitic areas in foothill areas which are inland, c. 100 km from the coast. Altitude varies from c. 200 m to 1200 m elevation. The wet season (successive months with cumulative rainfall >100 mm) falls between March and November and is colder than the dry season. Average annual rainfall at Bekob measured 2010 – 2016 is 2336 mm (Abwe, Ebo Forest Research Programme, Cameroon pers. comm., Abwe & Morgan 2008, Cheek *et al.* 2018a).

CONSERVATION STATUS. *Uvariopsis ebo* is currently known from a single specimen at a single location inside the mid-eastern part of the Ebo Forest. Less than 50 mature individuals have been observed (Bethan Morgan pers. comm. to Cheek, March 2021), despite the species being highly conspicuous in flower (Fig. 1) and situated on a major footpath close to a research camp used by many biologists over the last 15 years.

Since 2006, botanical surveys have been mounted almost annually, at different seasons, over many parts of the formerly proposed National Park of Ebo. About 2500 botanical herbarium

Map 1. Global distribution of *Uvariopsis ebo*, together with *U. korupensis* and *U. submontana*.

specimens have been collected, but this species has not yet been seen elsewhere in the c. 2000 km² of the Ebo Forest. However, the area outside the two research camps, especially the western edge, has not been fully surveyed for plants. While it is likely that the species will be found at additional sites within the Ebo Forest, there is no doubt that it is genuinely range-restricted as are some other species of *Uvariopsis* in Cameroon (see introduction). Botanical surveys and other plant studies for conservation management in forest areas north, west and east of Ebo, and just over the border in Nigeria, resulting in tens of thousands of specimens being collected and identified have failed to find any additional specimens of this species (Cheek *et al.* 1996; Cable & Cheek 1998; Cheek *et al.* 2000; Maisels *et al.* 2000, Chapman & Chapman 2001, Harvey *et al.* 2004; Cheek *et al.* 2004; Cheek *et al.* 2010; Harvey *et al.* 2010; Cheek *et al.* 2011).

The area of occupation of *Uvariopsis ebo* is estimated as 4 km² using the IUCN preferred cell-size. The extent of occurrence is the same. In February 2020 it was discovered that moves were in place to convert the forest into two logging concessions (e.g. <https://www.globalwildlife.org/blog/ebo-forest-a-stronghold-for-cameroons-wildlife/> and <https://blog.resource shark.com/cameroon-approves-logging-concession-that-will-destroy-ebo-forest-gorilla-habitat/> both accessed 19 Sept. 2020). Such logging would result in timber extraction that would open up the canopy and remove the intact habitat in which *Uvariopsis ebo* is found. Additionally, slash and burn agriculture often follows logging trails and would negatively impact the population of this species. Fortunately the logging concession was suspended in August 2020 due to representations to the President of Cameroon on the global importance of the biodiversity of Ebo (<https://www.businesswire.com/news/home/20200817005135/en/Relief-in-the-Forest-Cameroonian-Government-Backtracks-on-the-Ebo-Forest> accessed 19 Sept. 2020). However, the forest habitat of this species remains unprotected and threats of logging and conversion of

the habitat to plantations remain, and mining is also a threat. *Uvariopsis ebo* is therefore here assessed as Critically Endangered, CR B1+2ab(iii), D.

PHENOLOGY. Flowering has only been observed in late March and early April (Bethan Morgan pers. comm. 2021).

ETYMOLOGY. Named as a noun in apposition for the forest of Ebo.

VERNACULAR NAMES & USES. None are known.

NOTES. An unusual and distinctive feature of *Uvariopsis ebo* is the shape and form of the corolla. The centre of the proximal half of each petal is concave before anthesis, forming a globose chamber for the staminal dome with the other three petals (Fig. 1). On the inner surface of the petal, the concave area is demarcated by an inverted U-shaped, distinct wall-like, broad ridge which seems to be the point of contact with those of the other sepals, sealing the chamber. The distal half of the petals and the margins of the proximal half are flat, wing-like and held against each other (applanate). In section therefore, the distal part of the corolla will appear cross-shaped (see Fig. 1). This seems to be an extreme form of the petal structure seen in the probably closely related Cameroonian species *Uvariopsis korupensis* and *U. submontana* which also occur in the SW quadrant of Cameroon (see Map 1). Additional species of *Uvariopsis* present at Ebo Forest are *U. zenkeri* Engl. and *U. submontana*.

Discussion

The range of *Uvariopsis ebo* and other endemic species in the Ebo Forest area.

Abwe & Morgan, (2008) and Cheek *et al.* (2018a) give overviews of habitats, species and importance for conservation of the Ebo forest to which *Uvariopsis ebo* is restricted on current evidence. Sixty-nine globally threatened plant species are currently listed from Ebo on the IUCN Red List website and the number is set to rise rapidly as more of Cameroon's rare species are assessed for their conservation status as part of the Cameroon TIPAs programme. The discovery of a new species to science at the Ebo forest is not unusual. Numerous new plant species have been published from Ebo in recent years. Examples of other species that, like *Uvariopsis ebo*, appear to be strictly endemic to the Ebo area on current evidence are: *Ardisia ebo* Cheek (Cheek & Xanthos, 2012), *Crateranthus cameroonensis* Cheek & Prance (Prance & Jongkind, 2015), *Gilbertiodendron ebo* Burgt & Mackinder, *Inversodicraea ebo* Cheek (Cheek *et al.* 2017), *Kupeantha ebo* M.Alvarez & Cheek (Cheek *et al.* 2018b), *Kupeantha yabassi* M.Alvarez & Cheek (Alvarez *et al.* 2021), *Palisota ebo* Cheek (Cheek *et al.* 2018a) and *Pseudohydrosme ebo* Cheek (Cheek *et al.* 2021).

Further species described from Ebo have also been found further west, in the Cameroon Highlands, particularly at Mt Kupe and the Bakossi Mts (Cheek *et al.* 2004). Examples are *Myrianthus fosi* Cheek (in Harvey *et al.* 2010), *Salacia nigra* Cheek (Gosline & Cheek, 2014) and *Talbotiella ebo* Mackinder & Wieringa (Mackinder *et al.* 2010).

Additionally, several species initially thought endemic to Mt Kupe and the Bakossi Mts and adjoining areas in the Cameroon Highlands have subsequently been found at Ebo, e.g. *Coffea montekupensis* Stoff. (Stoffelen *et al.* 1997), *Costus kupensis* Maas & H. Maas (Maas-van der Kamer *et al.* 2016), *Deinbollia oreophila* Cheek (Cheek & Etuge 2009), *Microcos magnifica* Cheek (Cheek, 2017), and *Uvariopsis submontana* Kenfack, Gosline & Gereau (Kenfack *et al.* 2003). It is considered likely that additional Kupe species may yet be found at Ebo such as *Brachystephanus kupeensis* I.Darbysh. (Champluvier & Darbyshire, 2009), *Impatiens frithii* Cheek (Cheek & Csiba 2002) since new discoveries are still frequently being made in the Ebo Forest. Therefore, it is possible that *Uvariopsis ebo* might yet also be found in the Cameroon highlands, e.g. at Mt Kupe. However, this is thought to be only a relatively small possibility given the high level of survey effort at Mt Kupe: if it occurred there it is highly likely that it would have been recorded already since it is so spectacular it would be difficult to overlook.

Conclusions

Such discoveries as this new species underline the urgency for making further such discoveries while it is still possible since in all but one of the cases given above, the species have very narrow geographic ranges and/or very few individuals, and face threats to their natural habitat, putting these species at high risk of extinction.

About 2000 new species of vascular plant have been discovered each year for the last decade or more. Until species are known to science, they cannot be assessed for their conservation status and the possibility of protecting them is reduced (Cheek *et al.* 2020). Documented extinctions of plant species are increasing, e.g. *Oxygyne triandra* Schltr. and *Afrothismia pachyantha* Schltr. of South West Region, Cameroon are now known to be globally extinct (Cheek & Williams 1999, Cheek *et al.* 2018c, Cheek *et al.* 2019). In some cases, species appear to be extinct even before they are known to science, such as *Vepris bali* Cheek, also from the Cross-Sanaga interval in Cameroon (Cheek *et al.* 2018d) and elsewhere, *Nepenthes maximoides* Cheek (King & Cheek, 2020). Most of the 815 Cameroonian species in the Red Data Book for the plants of Cameroon are threatened with extinction due to habitat clearance or degradation, especially of forest for small-holder and plantation agriculture following logging (Onana & Cheek, 2011). Efforts are now being made to delimit the highest priority areas in Cameroon for plant conservation as Tropical Important Plant Areas (TIPAs) using the revised IPA criteria set out in Darbyshire *et al.* (2017). This is intended to help avoid the global extinction of additional endemic species such as *Uvariopsis ebo* which will be included in the proposed Ebo Forest IPA.

With only one locality known, *Uvariopsis ebo* represents another narrowly endemic Cameroonian species threatened with extinction due to deforestation for oil palm plantations, small-scale agriculture, mining and logging (Onana & Cheek 2011, Cheek *et al.* 2018a).

Acknowledgements

This paper was completed as part of the Cameroon TIPAs (Tropical Important Plant Areas) project at RBG, Kew, which is supported by Players of People's Postcode Lottery. Ekwoke Abwe and Bethan Morgan and their team at the Ebo Forest programme are thanked hugely for making available the specimen and photos on which this paper is based and for expediting our botanical surveys in the Ebo forest of Cameroon over several years which allowed us to give context about the Ebo forest in this paper. We thank Janis Shillito for typing the manuscript, Lorna MacKinnon for the photos and specimen.

The heads of IRAD (Institute of Research in Agronomic Development)-National Herbarium of Cameroon, Yaoundé, successively Florence Ngo Ngwe, Eric Nana and Jean Betti Lagarde, are thanked for co-ordinating the co-operation with the Royal Botanic Gardens, Kew.

The authors would like to thank two anonymous reviewers for comments on an earlier version of this manuscript.

References

Abwe, E.E, Morgan, B.J. (2008). The Ebo Forest: four years of preliminary research and conservation of the Nigeria-Cameroon chimpanzee (*Pan troglodytes vellerosus*). *Pan Africa News* 15: 26–29. <https://doi.org/10.5134/143494>

- Alvarez, M. G., Cheek, M. Sonké, B. (2021). *Kupeantha yabassi* (Coffeeae-Rubiaceae), a new Critically Endangered shrub species of the Ebo Forest area, Littoral Region, Cameroon. *Kew Bulletin*, submitted.
- Barthlott, W., Lauer, W., Placke, A. (1996). Global distribution of species diversity in vascular plants: towards a world map of phytodiversity. *Erkunde* 50: 317 – 328 (with supplement and figure).
- Beentje, H. & Cheek, M. (2003). Glossary. In Beentje, (ed.), *Flora of Tropical East Africa*. Balkema, Lisse, Netherlands. 115 pp.
- Cable, S., Cheek, M. (1998). *The Plants of Mt Cameroon, a Conservation Checklist*. Kew, Royal Botanic Gardens.
- Champluvier, D., Darbyshire, I. (2009). A revision of the genera *Brachystephanus* and *Oreacanthus* (Acanthaceae) in tropical Africa. *Systematics and Geography of Plants* 79: 115–192. <https://doi.org/10.2307/25746>
- Chapman J, Chapman H. 2001. *The Forests of Taraba and Adamawa States, Nigeria an Ecological Account and Plant Species Checklist*. University of Canterbury: Christchurch, New Zealand. pp. 221.
- Cheek M. (2017). *Microcos magnifica* (Sparrmanniaceae) a new species of cloudforest tree from Cameroon. *PeerJ* 5:e4137 <https://doi.org/10.7717/peerj.4137>
- Cheek, M., Csiba, L. (2002). A new epiphytic species of *Impatiens* (Balsaminaceae) from western Cameroon. *Kew Bulletin* 57(3): 669 – 674. <https://doi.org/10.2307/4110997>
- Cheek, M. & Etuge, M. (2009). A new submontane species of *Deinbollia* (Sapindaceae) from Western Cameroon and adjoining Nigeria. *Kew Bull.* 64(3): 503 – 508. <https://doi.org/10.1007/s12225-009-9132-4>
- Cheek, M. & Williams, S. (1999). A Review of African Saprophytic Flowering Plants, pp. 39 – 49 in Timberlake & Kativu (Eds.) *African Plants. Biodiversity, Taxonomy & Uses*. Proceedings of the 15th AETFAT Congress at Harare, Zimbabwe. Royal Botanic Gardens, Kew.
- Cheek, M. & Xanthos, M. (2012). *Ardisia ebo* sp. nov. (Myrsinaceae), a creeping forest subshrub of Cameroon and Gabon. *Kew Bull.* 67: 281 – 284. <https://doi.org/10.1007/s12225-012-9362-8>
- Cheek, M., Achoundong, G., Onana, J-M., Pollard, B., Gosline, G., Moat, J., Harvey, Y.B. (2006). Conservation of the Plant Diversity of Western Cameroon. In: S.A. Ghazanfar, H.J. Beentje (eds). *Taxonomy and ecology of African plants, their conservation and sustainable use*. Proceedings of the 17th AETFAT Congress, Addis Ababa, Ethiopia: 779 – 791. Kew, Royal Botanic Gardens.
- Cheek M. Alvarez-Aguirre M.G., Grall A., Sonké, B., Howes M.J.R., Larridon I. (2018b). *Kupeantha* (Coffeeae, Rubiaceae), a new genus from Cameroon and Equatorial Guinea. *PLoS ONE* 13(6): e0199324. <https://doi.org/10.1371/journal.pone.0199324>

- Cheek, M., Cable, S., Hepper, F.N., Ndam, N., Watts, J. (1996). Mapping plant biodiversity on Mt. Cameroon. pp. 110 – 120 in van der Maesen, van der Burgt & van Medenbach de Rooy (Eds), *The Biodiversity of African Plants* (Proceedings XIV AETFAT Congress). Kluwer. https://doi.org/10.1007/978-94-009-0285-5_16
- Cheek, M., Etuge, M., Williams, S. (2019). *Afrothismia kupensis* sp. nov. (*Thismiaceae*), Critically Endangered, with observations on its pollination and notes on the endemics of Mt Kupe, Cameroon. *Blumea - Biodiversity, Evolution and Biogeography of Plants*. 64(1): 158 – 164 <https://doi.org/10.3767/blumea.2019.64.02.06>
- Cheek, M., Feika, A., Lebbie, A., Goyder, D., Tchiengue, B., Sene, O., Tchouto, P., van der Burgt, X. (2017). A synoptic revision of *Inversodicraea* (Podostemaceae). *Blumea* 62, 2017: 125–156. <https://doi.org/10.3767/blumea.2017.62.02.07>
- Cheek M, Gosline G, Onana J-M. 2018d. *Vepris bali* (*Rutaceae*), a new critically endangered (possibly extinct) cloud forest tree species from Bali Ngemba, Cameroon. *Willdenowia* 48: 285 – 292. doi: <https://doi.org/10.3372/wi.48.48207>
- Cheek, M., Harvey Y., Onana J-M. (2010). *The Plants of Dom, Bamenda Highlands, Cameroon, A Conservation Checklist*. Kew, Royal Botanic Gardens.
- Cheek, M., Harvey, Y., Onana, J.-M. (2011). *The Plants of Mefou Proposed National Park, Yaoundé, Cameroon, A Conservation Checklist*. Kew, Royal Botanic Gardens.
- Cheek, M., Mackinder, B. Gosline, G., Onana J.-M., Achoundong, G. (2001). The phytogeography and flora of western Cameroon and the Cross River-Sanaga River interval. *Systematics and Geography of Plants* 71: 1097 – 1100. <https://doi.org/10.2307/3668742>
- Cheek, M., Nic Lughadha, E., Kirk, P., Lindon, H., Carretero, J., Looney, B., Douglas, B., Haelewaters, D., Gaya, E., Llewellyn, T., Ainsworth, M., Gafforov, Y., Hyde, K., Crous, P., Hughes, M., Walker, B.E., Forzza, R.C., Wong, K.M., Niskanen, T. (2020). New scientific discoveries: plants and fungi. *Plants, People Planet* 2:371 – 388. <https://doi.org/10.1002/ppp3.10148>
- Cheek, M., Onana, J-M., Pollard, B.J. (2000). *The Plants of Mount Oku and the Ijim Ridge, Cameroon, a Conservation Checklist*. Kew, Royal Botanic Gardens.
- Cheek, M., Pollard, B.J., Darbyshire, I., Onana, J-M., Wild, C. (2004). *The Plants of Kupe, Mwanenguba and the Bakossi Mountains, Cameroon: A Conservation Checklist*. Kew, Royal Botanic Gardens.
- Cheek, M., Prenner, G., Tchiengué, B., Faden, R.B. (2018a). Notes on the endemic plant species of the Ebo Forest, Cameroon, and the new, Critically Endangered, *Palisota ebo* (Commelinaceae). *Plant Ecology & Evolution* 151(3): 434 – 441. <https://doi.org/10.5091/plecevo.2018.1503>
- Cheek, M., Tchiengué, B., van der Burgt, X. (2021). Taxonomic revision of the threatened African genus *Pseudohydrosme* Engl. (Araceae), with *P. ebo*, a new, critically endangered species from Ebo, Cameroon. *PeerJ* 9:e10689 <https://doi.org/10.7717/peerj.10689>

Cheek M, Tsukaya H, Rudall PJ, Suetsugu K. (2018c). Taxonomic monograph of *Oxygyne* (Thismiaceae), rare achlorophyllous mycoheterotrophs with strongly disjunct distribution. *PeerJ* 6:e4828 <https://doi.org/10.7717/peerj.4828>

Couch, C., Cheek, M., Haba, P.M., Molmou, D., Williams, J., Magassouba, S., Doumbouya, S., Diallo, Y.M. (2019). *Threatened habitats and Important Plant Areas (TIPAs) of Guinea, west Africa*. Royal Botanic Gardens, Kew. London.

Couvreur, T. L., & Luke, W. R. Q. (2010). A new species of *Uvariopsis* (Annonaceae), endemic to the Eastern Arc Mountains of Tanzania. *Blumea-Biodiversity, Evolution and Biogeography of Plants*, 55(1), 68 – 72. <https://doi.org/10.3767/000651910X499196>

Couvreur, T. L., & Niangadouma, R. (2016). New species of *Uvariopsis* (Annonaceae) and *Laccosperma* (Arecaceae/Palmae) from Monts de Cristal, Gabon. *PhytoKeys*, (68), 1:1 – 8. <https://doi.org/10.3897/phytokeys.68.9576>

Couvreur, T.L.P., Dagallier, L.-P.M.J., Crozier, F., Ghogue, J.-P., Hoekstra, P.H., Kamdem, N.G., Johnson, D.M., Murray, N. & Sonké, B. (2021). Flora of Cameroon – Annonaceae. *Phytokeys*.

Darbyshire, I., Anderson, S., Asatryan, A., Byfield, A., Cheek, M., Clubbe, C., Ghrabi, Z., Harris, T., Heatubun, C. D., Kalema, J., Magassouba, S., McCarthy, B., Milliken, W., Montmollin, B. de, Nic Lughadha, E., Onana, J. M., Saidou, D., Sarbu, A., Shrestha, K. & Radford, E. A. (2017). Important Plant Areas: revised selection criteria for a global approach to plant conservation. *Biodiversity Conservation* 26: 1767 – 1800. <https://doi.org/10.1007/s10531-017-1336-6>

Dominy, N. J. & Duncan, B. W. (2005). Seed-spitting Primates and the Conservation and Dispersion of Large-seeded Trees. *International Journal of Primatology* 26(3): 631–649. <https://doi.org/10.1007/s10764-005-4370-2>

Gereau, R. E. & D. Kenfack. 2000. Le genre *Uvariopsis* (Annonaceae) en Afrique tropicale, avec la description d'une espece nouvelle du Cameroun. *Adansonia*, ser. 3, 22: 3943

Gosline, G., Cheek, M. & Kami, T. (2014). Two new African species of *Salacia* (*Salacioideae*, *Celastraceae*). *Blumea* 59: 26 – 32. <https://doi.org/10.3767/000651914x682026>

Harvey, Y., Pollard, B.J., Darbyshire, I., Onana, J.-M., Cheek, M. (2004). *The Plants of Bali Ngemba Forest Reserve, Cameroon. A Conservation Checklist*. Kew, Royal Botanic Gardens.

Harvey, Y.H., Tchiengue, B., Cheek, M. (2010). *The plants of the Lebialem Highlands, a conservation checklist*. Kew, Royal Botanic Gardens.

IPNI (continuously updated). *The International Plant Names Index*. Available from: <http://ipni.org/> [accessed Mar. 2018].

IUCN (2012). *IUCN red list categories: Version 3.1*. Gland, Switzerland and Cambridge, U.K., IUCN Species Survival Commission.

Kenfack, D., Gosline, G., Gereau, R.E., Schatz, G. (2003). The genus *Uvariopsis* in Tropical Africa, with a recombination and one new species from Cameroon. *Novon* 13: 443–449. <https://doi.org/10.2307/3393377>

King, C., Cheek, M. (2020). *Nepenthes maximoides* (Nepenthaceae) a new, critically endangered (possibly extinct) species in Sect. *Alatae* from Luzon, Philippines showing striking pitcher convergence with *N. maxima* (Sect. *Regiae*) of Indonesia. *PeerJ* 8:e9899. <https://doi.org/10.7717/peerj.9899>

Maas-van de Kamer, H., Maas, P.J.M., Wieringa, J.J., Specht, C.D. (2016). Monograph of African *Costus*. *Blumea - Biodiversity, Evolution and Biogeography of Plants* 61: 280–318. <https://doi.org/10.3767/000651916X694445>

Mackinder, B.A., Wieringa, J.J., van der Burgt, X.M. (2010). A revision of the genus *Talbotiella* Baker f. (*Caesalpinioideae: Leguminosae*). *Kew Bulletin* 65: 401–420. <https://doi.org/10.1007/s12225-010-9217-0>

Maisels, F.M., Cheek, M., Wild, C. (2000). Rare plants on Mt Oku summit, Cameroon. *Oryx* 34: 136 – 140. <https://doi.org/10.1017/s0030605300031057>

Onana, J.-M. (2011). *The vascular plants of Cameroon, a taxonomic checklist with IUCN Assessments*. Kew, Royal Botanic Gardens.

Onana, J.-M., Cheek, M. (2011). *Red data book of the flowering plants of Cameroon, IUCN global assessments*. Kew, Royal Botanic Gardens.

Prance, G.T., Jongkind, C.C.H. (2015). A revision of African Lecythidaceae. *Kew Bulletin* 70: 6: 13. <https://doi.org/10.1007/s12225-014-9547-4>

Sosef, M.S.M., Wieringa, J.J., Jongkind, C.C.H., Achoundong, G., Azizet Issembé, Y., Bedigian, D., Van Den Berg, R.G., Breteler, F.J., Cheek, M., Degreef, J. (2005). [Checklist of Gabonese Vascular Plants](#). *Scripta Botanica Belgica* 35. National Botanic Garden of Belgium.

Stoffelen, P., Cheek, M., Bridson, D., Robbrecht, E. (1997). A new species of *Coffea* (Rubiaceae) and notes on Mt Kupe (Cameroon). *Kew Bulletin* 52(4): 989–994. <https://doi.org/10.2307/3668527>

Thiers, B. (continuously updated). Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. [continuously updated]. Available from: <http://sweetgum.nybg.org/ih/> [accessed: March 2021].